[redacted] 1780

John Scott

A

GENERAL SYSTEM OF SURGERY,

IN THREE PARTS.

CONTAINING THE

DOCTRINE and MANAGEMENT

I. Of WOUNDS, FRACTURES, LUXATIONS, TUMORS, and ULCERS, of all Kinds.

II. Of the ſeveral OPERATIONS performed on all PARTS of the BODY.

III. Of the ſeveral BANDAGES applied in all OPERATIONS and DISORDERS.

To which is prefixed,

An INTRODUCTION,

Concerning the *Nature*, *Origin*, *Progreſs*, and *Improvements* of SURGERY; With ſuch other PRELIMINARIES as are neceſſary to be known by the YOUNGER SURGEONS.

Being a Work of THIRTY YEARS Experience.

Illuſtrated with Forty COPPER-PLATES, exhibiting all the OPERATIONS, INSTRUMENTS, BANDAGES, and IMPROVEMENTS, according to the Modern and moſt approved Practice.

By Dr. LAURENCE HEISTER,

Profeſſor of *Phyſic* and *Surgery* in the Univerſity of HELMSTADT, Fellow of the Royal Society, LONDON, and of the Royal Academy at PARIS, *&c.*

Tranſlated from the AUTHOR's laſt Edition, greatly improved.

The EIGHTH EDITION, Reviſed and Corrected.

LONDON,

Printed for J. WHISTON, L. DAVIS and C. REYMERS, B. WHITE, H. WOODFALL, R. BALDWIN, W. JOHNSTON, T. LOWNDES, L. HAWES and Co. T. CASLON, Z. STUART, W. LAW, W. KEARSLEY.

MDCCLXVIII.

[redacted] 1768.

BOSTON MEDICAL
DEC 15 1916
LIBRARY.

* 1. Mv. 21.

THE

EDITOR'S PREFACE

to the SIXTH Edition.

THIS moſt excellent and elaborate Work is ſo well known in the learned World, and has every where met with ſo favourable a Reception, that to detain the Reader at preſent with any Encomiums on this Subject, would be vain and ſuperfluous. But, as the judicious Author (who has not only his own Credit, and the Dignity of his Profeſſion, but, what is infinitely more noble, the public Utility ever at Heart) has greatly improved and enlarged his Undertaking in ſucceeding Editions; the Proprietors of this Tranſlation have, therefore, thought it their Duty to give a freſh Impreſſion of it, enriched withall thoſe uſeful and valuable Improvements. Theſe Additions are diſperſed almoſt every where throughout the Work; but are moſt large and copious in the important Branches of Lithotomy and Midwifery.

We have likewiſe inſerted in this Edition two additional Plates engraved from thoſe of the Author's. And, that nothing might be wanting to compleat it, the whole Tranſlation has been reviſed, with great Labour and Attention, and particularly with an Eye to the Accuracy of the Pointing, which greatly aſſiſts the Clearneſs and Perſpicuity of a Work, eſpecially to the young Beginner, not yet verſed in the Subject.—Upon the whole, we flatter ourſelves, that we here offer to the Public, the moſt uſeful Work of its Kind now extant, thus carefully reviſed, and amply improved; which, on its firſt Appearance, not only led the Pupil through each Branch of the Practice, but was confeſſed, by the experienced and judicious Practitioner, to be a moſt complete, diſtinct, and comprehenſive Syſtem of the whole Art of Surgery.

London, November 1, 1756.

THE

TRANSLATOR'S PREFACE.

THE Tranſlation of the Book before us, which now appears in the World, will obviate a Complaint frequently made among the junior Surgeons, and Pupils of this Art in *England, viz.* that they are in Want of a general Syſtem, capable of inſtructing at large one that is a Learner in Surgery, for the Execution of all the Branches of his Profeſſion; and this, till now, might indeed be affirmed with ſome Juſtice. It is true, the ſeveral Branches of Surgery have been tolerably well handled by various Authors, at different Times, and in ſeparate Treatiſes: Some have confined themſelves to Wounds, Fractures, Luxations, Tumours, and Ulcers, which make the Subject of the firſt Part of the preſent Syſtem; others have writen profeſſedly on the Operations, Inſtruments, and Bandages; or miſcellaneous Obſervations appertaining to the Practice of Surgery; and others have given us ſhort Introductions to the whole; but in no one Book, except the preſent, do we meet with all theſe Branches treated in that ample, eaſy, and intelligent Manner, which is neceſſary for the firſt Information of Beginners, or the occaſional Conſultation of the more advanced. We have, in this Work, not only the beſt and moſt modern Methods of Practice uſed by the principal Surgeons of the ſkilfulleſt *European* Nations, but alſo exact Figures of their ſeveral Inſtruments and Bandages, with the Methods of uſing or applying them in all chirurgical Caſes whatever; the whole Doctrine of which is here explained in the minuteſt Circumſtances; and brought down even to the loweſt Capacities. In ſhort, no Character of the Book can ſo well recommend it to the Reader as his own Peruſal, and the Author's Preface following.

London, Oct. 1742.

THE

THE AUTHOR'S PREFACE.

AFTER having ſtudied Phyſic with great Aſſiduity above four Years in our *German* Univerſities, my Affections, being ſtrongeſt for *Anatomy* and *Surgery*, led me to the then celebrated Profeſſors RUYSCH and RAW, at *Amſterdam*, in the Year 1706; whoſe anatomical and chirurgical Demonſtrations I diligently attended for about the Space of a Year. During which Time I was alſo employed in frequent Diſſections, and in trying chirurgical Operations upon dead Subjects; in the mean Time omitting no Opportunities of being preſent at the Performance of any conſiderable Operation by theſe Profeſſors, or by the other eminent Surgeons of the ſame City, as VERDUIN, BORTEL, KOENERDING, *&c.* By which Means, joined with an attentive Reading of the beſt Writers, I acquired a conſiderable Knowledge in Surgery.

But, being deſirous of all Helps to render myſelf ſtill more expert and ſucceſsful in the Practice of this Art, there being at that Time a ſharp War in *Flanders*, betwixt the *French* and *Dutch*, in the Summer following, *viz.* in the Year 1707, I went from *Holland* to the *Dutch* Camp in *Brabant*, that I might inſpect and obſerve the Practice of the *Engliſh*, *Dutch*, and *German* Surgeons, who there attended. Thus, through many Dangers and Hardſhips, I ſpent this whole Summer in the Hoſpitals of the Camp, for the Sake of Improvement. But in Autumn I went from *Brabant* to *Leyden*, and ſpent the whole Winter in attending the Lectures of the then celebrated Profeſſors in that Univerſity, BIDLOO, ALBINUS ſenior, and BOERHAAVE: And thus I continued till the Beginning of the Summer 1708. After which, having taken my Degree of *Doctor*, I returned again to the Camp, where I found large Opportunities of learning and improving myſelf in Surgery from the Multitude wounded, *&c.* in the ſeveral bloody Fights, particularly at the Siege of *Liſle*, and the Battles of *Oudenarde* and *Wynnendale*. Upon the Approach of Winter again, I was determined to ſettle in the Practice of *Phyſic* and *Surgery* in *Holland*, at *Amſterdam*, partly from

from the Delight I had in the Country, and partly through the Solicitations of the famous RUYSCH, who reſpected me as a Son. Here therefore I ſtayed the Winter, and Part of the enſuing Spring, teaching *Anatomy* and *Surgery* to Students and Gentlemen, as RAW had done before me, who was now rejected for his ill Conduct or Miſbehaviour.

The following Summer, in 1709, I had ſtill a ſtrong Deſire to follow the Camp, to become more and more perfect in the Practice of *Surgery*; and, *Tournay* being at that Time inveſted by the confederate Army in *Flanders*, I was, by the Recommendations of my Friend RUYSCH, appointed Phyſician to the Camp Hoſpital for the *Hollanders*. I had now an Opportunity of performing all the chirurgical Operations which offered in the Camps and adjacent Cities, which I generally executed with Succeſs. After the taking of *Tournay*, the confederate Army marched to beſiege *Mons*, near which Place the *French* Army was alſo aſſembled. That, however, did not prevent us from inveſting and taking the City; before which the numerous Army had ſuch a bloody Battle, that the Wounded were brought in upon us in Crowds. Their Number continually increaſing, from the uncommon Heat of the Combat, every Surgeon had now his Hands full of Buſineſs, and infinite Calls for the Practice of his Arts for the Wounded, on the Side of the *Hollanders* only, amounted to above Five Thouſand. I had here therefore an ample Occaſion to extend the Bounds of my Practice, and was obliged to put on that Intrepidity of Mind, which CELSUS requires as an eſſential Qualification in a Surgeon; and for Want of which, ſome, who are, in other reſpects, ſkilful Operators, do frequently miſcarry.

After the Army had entered into their Winter-Quarters, and the wounded Men recovered, I returned again to *Amſterdam*, where I continued my *Anatomical* and *Chirurgical* Demonſtrations this Winter as before. In the mean Time, I never refuſed my Aſſiſtance at the Operations of the other Surgeons there.

But, in the Beginning of the Spring following, 1710, I was, beyond all Expectation, called by the Republic of *Norimberg* to teach *Anatomy* and *Surgery*, as public PROFESSOR in the Univerſity of *Altorf*. Being therefore unwilling to neglect this honourable Invitation, having obtained

Leave

Leave from the Republic, I firſt made a Tour into *Great Britain* where I was, from Spring to Autumn, collecting every thing new in the ſeveral Branches of Phyſic, and then, returning to *Norimberg* and *Altorf*, I aſſumed my new *Profeſſorſhip*.

In this Station, I was under a Neceſſity of teaching publicly, among the other Parts of *Phyſic*, that moſt ancient, neceſſary, and uſeful Branch of it which we call *Surgery*, and which I had before taught privately during the two preceding Winters in *Holland*. But in doing this I was much perplexed for want of a convenient Manual, or compendious Syſtem of the Art, to aſſiſt and inform thoſe Learners who attended my Lectures. To our Want of ſuch a *Compendium* I alſo attributed the general Ignorance and Inſufficiency of the young Surgeons and Students in this Branch of *Phyſic*, which at that Time univerſally prevailed, through *Germany* eſpecially. And, from the ſame Cauſe, the Generality of our Surgeons, being unequal to the more difficult Operations, were content with being able to cure a ſlight Wound, open a Vein or Abſceſs, or at moſt to ſet a Fracture, and reduce a Luxation; leaving thoſe Diſorders and Operations, which require the greateſt Skill, to the Management of daring Quacks and itinerant Operators, with which *Germany* at that Time ſwarmed.

If any one examines the beſt Books, ſuch as the *Microtechnia* of VAN HOORN, the *Operations* of NUCK, *&c.* which were at that Time conſulted not only by our Surgeons, but alſo by our Univerſity-Profeſſors, for teaching and learning the Art, it will readily appear how imperfect and inſufficient they are to give a juſt Notion of any one Branch, much more of the whole Syſtem or Body of Surgery; ſince they deſcribe only a few of the Operations, and thoſe too imperfectly; taking little or no Notice of the Doctrine and Treatment of Wounds, Fractures, Luxations, Tumors, and Ulcers, which make the moſt conſiderable Part of Surgery, and in which a Learner ought to be the moſt fully inſtructed. It is true, the Works of GUIDO CAULIACUS, AQUAPENDENS, PAREY, SCULTETUS, SOLINGEN, and ſome other Writers of the laſt Century, are very full and explicit in all or moſt of the Operations, and the five Kinds of Diſorders beforementioned. But even in theſe we muſt not expect to find the many Improvements, Emendations and Diſcoveries, made by the Moderns: And their Practice being moſtly

moſtly obſolete, they muſt conſequently be allowed to be unfit for the Inſtruction of Learners. And it is an Objection to many of our Books in Surgery, of a more modern Date than the preceding, that they have been either compiled by Phyſicians, little converſant in chirurgical Diſſections and Operations, as thoſe of BARBET, VERDUC, BONTECK, DOLEY, BLANCARD, CHARRIERE, JUNCKEN, VAUGUION, LE CLERC, *&c.* in which many of the old Errors are continued, and not a few Things ſtated otherwiſe than will be found in Practice: Or elſe they have been reſtrained to but one or two Subjects only, as the Bones, Wounds, Tumors, Bandages, Operations, *&c.* beſides their being written either in a rude, or a foreign Language, unknown to moſt of our Surgeons.

Theſe were chiefly the Motives that firſt induced me to attempt the Compoſition of a chirurgical Syſtem, to be ſubſervient to my own Lectures and Auditors: In doing which, I endeavoured to take in all the more uſeful Part both of our ancient and modern Writers in every Branch of Surgery; rejecting what appeared uſeleſs or obſolete, and comparing or correcting the whole conformably to my own Experience, and what I had ſeen in the Practice of the Art under many of the moſt ſkilful Surgeons and Phyſicians. And thus, from Time to Time, I endeavoured not only to correct and compleat my Collections and Remarks, ſo as to take in every, even the minuteſt, Part of Surgery; but alſo I digeſted and diſpoſed the whole in the Method which appeared to me the moſt natural, and the beſt adapted both for the Teacher and Learner.

Theſe my firſt Labours I writ originally in *Latin*, in which Language they were alſo delivered to my Hearers, and permitted to be *tranſcribed* by them: But, conſidering the immenſe Fatigue that this Method of obtaining it gave the Student, with the great Loſs of Time, which he might have otherwiſe employed to more Advantage, I was at length determined to *publiſh* it in *Latin*, in the Manner I had then compoſed it. Yet ſo great was the Ignorance of our *German* Surgeons, at that Time of Day, as well in the *Latin* Tongue as in their own Profeſſion, that my Work being chiefly intended for them, I now judged it would be more uſeful to print the Book in our native *German*; for then both the learned and ignorant of the *Latin* might have the ſame Benefit of it. Accordingly I tranſlated and ſent it to the Preſs in the Year 1717; and in the Year following, 1718, it was publiſhed

as my *Surgery*, in *4to*, at *Norimberg*, being illuſtrated with Copper-plates, exhibiting the beſt Inſtruments, *&c.* And from this time it is that we have had better of more expert Surgeons in *Germany* than before; many or which have ſince often declared to me, that they had drawn moſt of their Knowledge from my Surgery.

I intended, ſoon after, to have publiſhed the Book in *Latin*, for the Sake of *Foreigners*; but, in the Year enſuing, I received a moſt gracious Call to the publick Profeſſorſhip of *Anatomy* and *Surgery* in the *Julian* Univerſity of *Helmſtadt*, from his *Britannic* Majeſty, as Duke of *Brunſwick* and *Lunenburg*, under whom the Univerſity flouriſhes, and is liberally ſupported; ſo that what with the Care and Trouble of packing up, and removing my Goods, and the Fatigue of a long Journey, added to the Multitude of Buſineſs, and many Avocations conſequent on my new Office, I have been obliged to delay the *Latin* Edition of my Surgery much longer than I ever thought or deſigned. However, the *German* Impreſſion was ſold off in a little Time, and the Bookſeller urging for a ſecond Edition, as there were ſeveral Improvements made lately in *Surgery*, particularly in *Lithotomy*, I therefore reviſed, corrected, and enlarged the Book, according to the later Diſcoveries, and my own recent Obſervations ſince made, ſo as to fit it then for a ſecond, and ſome time after for a third Edition. But then this, with other Avocations in the mean time, prevented me from compleating the Work in the learned Language, for the better Sort of Readers, ſo as to make it correſpond to the Performances of foreign Authors, with which our *German* Surgeons were unacquainted.

But being at length ſollicited, as well by many learned Phyſicians and Surgeons of other Nations, as by my Bookſeller at *Amſterdam*, to publiſh my Surgery in *Latin*, for the Advantage of Foreigners; and being unwilling to deny the Requeſt, I have now, notwithſtanding my academical and practical Buſineſs, made ſhift to print it in that Language, in many Places much enlarged and amended beyond any of the preceding Editions. And I hope it may be a Means of inſtructing young Surgeons in all the Branches of their Profeſſion, according to the beſt modern Diſcoveries and Improvements which have been made in the Art. I have here endeavoured to preſent them with the whole Body of Surgery together, that Learners eſpecially may not have their

Knowledge to ſeek in many different Books, by turning over ſome upon Wounds; others upon Fractures, Luxations, Tumors, or Ulcers; and others, again, upon Operations, Inſtruments or Bandages: All which, I think, are here ſufficiently explained, not only for the Inſtruction of Learners, but all the Purpoſes of the more advanced.

Whether I have ſucceeded in this Taſk, muſt be left to the Determination of more prudent and impartial Judges; but this I may be allowed to ſay, that I have uſed my beſt Endeavours to promote the Glory of God, and the public Good, by theſe Labours of their

Devoted Servant,

Helmſtadt,
Jan. 10, 1739.

THE AUTHOR.

CONTENTS

CONTENTS of the FIRST VOLUME.

THE Introduction, *Of the Nature, Conſtitution, Origin, Progreſs, Improvement, and Diviſion of* Surgery; *and of other Things in general, which are principally neceſſary for* Students *in* Surgery *to be acquainted with* Page 1, & ſeq.

BOOK I. *Of WOUNDS.*

Chap.
I. *Of* Wounds *in general* Page 27
II. *Of the Diſorders accompanying* Wounds, *commonly called the* Symptoms *of* Wounds 48
III. *Of* Gun ſhot Wounds 55
IV. *Of* Wounds *of the* Abdomen 63
V. *Of* Gaſtroraphy 65
VI. *Of* Wounds *of the* Inteſtines, *and the Manner of ſtitching them up* 71
VII. *Of* Loſs *of* Subſtance *in the* Inteſtines 74
VIII. *Of the* Falling down *of the* Omentum 76
IX. *Of* Wounds *of other Parts of the* Abdomen 78
X. *Of* Wounds *of the* Thorax 79
XI. *Of* Wounds *of the* Neck 84
XII. *Of* Wounds *of the* Head *in general* 88
XIII. *Of* Wounds *of the* Face 89
XIV. *Of the principal* Wounds *of the* Head 93
XV. *Of* Contuſions 105
XVI. *Of* Venomous Wounds, *and of thoſe that are made by the* Bites *of* Animals 110

BOOK II. *Of FRACTURES.*

I. *Of* Fractures *in general* 116
II. *Of* Diſorders *accompanying* Fractures 126
III. *Of* Fractures *in particular* 131
Chap.
IV. *Of a* Fracture *of the* Jaw Page 132
V. *Of a* Fracture *of the* Clavicles, Sternum, *and* Humerus 134
VI. *Of* Fractures *in the* Ribs, Vertebræ, Os Sacrum, *and* Innominatum 137
VII. *Of* Fractures *in the Bones of the* Humerus, Cubitus, *and* Hands 141
VIII. *Of a fractured* Thigh 143
IX. *Of a* Fracture *of the* Patella, Rotula, *or* Knee-pan 149
X. *Of* Fractures *in the Bones of the* Leg *and* Foot 151
XI. *Of Bones broken by ſharp-pointed Inſtruments, which may be termed* Wounds *of the* Bones 153

BOOK III. *Concerning* LUXATIONS, *or* LUXATED BONES.

I. *Of* Luxations *of the* Bones *in general* 158
II. *Concerning the* Cure *of* Luxated Bones 164
III. *Of* Luxations *in particular*; *and on thoſe of the* Head *and* Noſe 167
IV. *Of a* Diſlocation *of the* Lower Jaw 168
V. *Of* Luxations *of the* Head *and* Spine 169
VI. *Of* Luxations *of the* Os Coccygis, Ribs, *and* Clavicles 173
VII. *Of a* Luxation *of the* Humerus 176
VIII. *Of a* Luxation *of the* Cubitus 180
IX. *Of* Luxations *of the* Hand, Carpus, Metacarpus, *and* Fingers 182
X. *Of a* Luxation *of the* Thigh 183
XI. *Of a* Luxation *of the* Patella *and* Knee, *or* Tibia, *and* Fibula 188
XII. *Of a* Luxation *of the* Foot *and* Ancle 190

BOOK IV. *Of* TUMORS.

Chap.
I. *Of* Tumors *in general* Page 194
II. *Of a* Phlegmon 195
III. *Of* Suppuration *and* Abſceſs 201
IV. *Of* Tumor *and* Inflammation *in the* Breaſts 205
V. *Of* Inflammation *in the* Teſticles 209
VI. *Of an* Eryſipelas 211
VII. *Of a* Furuncle, *or* Boil 213
VIII. *Of the* Bubo *and* Parotis 215
IX. *Of* Peſtilential Bubos, *where alſo of* Carbuncles 216
X. *Of the* Preſervatives *particularly neceſſary to defend and preſerve the Phyſician or Surgeon from* Peſtilential Contagion 220
XI. *Of* Carbuncles *or* Anthraces 223
XII. *Of* Venereal Bubos 226
XIII. *Of* Chilblains 228
XIV. *Of a* Gangrene *and* Sphacelus 231
XV. *Of* Burns *and* Scalds 240
XVI. *Of a* Scirrhus 244
XVII. *Of a* Cancer 248
XVIII. *Of an* Oedema 252
XIX. *Of* Fungous Tumors, *and* Dropſy *of the* Joints 255

BOOK V. *Of* ULCERS.

I. *Of* Ulcers 259
II. *Of the Method of treating* Fiſtulous Caſes 264
III. *Of the Method of treating* Stubborn Ulcers 265
IV. *Of the Method of treating* Venereal Ulcers 269
V. *Of* Callous Ulcers 271
VI. *Of* Ulcers *ſuppoſed to be produced by* Magic *or* Witchcraft 272
VII. *The Method of treating old* Ulcers, *eſpecially thoſe that affect the* Legs 273
VIII. *Of* Caries *of the* Bones 275
IX. *Of the* Spina Ventoſa, Pædarthrocaces, *and* Exoſtoſis, *which may be called* Tumors *of the* Bones Page 281
X. *Of* Ulcers *of the* Head 288

PART II.

Of OPERATIONS.

SECT. I.

Of GENERAL OPERATIONS *practicable in ſeveral different Parts of the Body.*

I. *Of* Phlebotomy *in general* 293
II. *Of* Opening *the* Veins *of the* Arm 295
III. *Of* Phlebotomy *in the* Hand 299
IV. *Of* Bleeding *in the* Foot 300
V. *Of* Bleeding *in the Veins of the* Forehead, Temples, *and* Occiput 301
VI. *Of* Bleeding *in the* Veins *of the* inner Corners *of the* Eyes 302
VII. *Of* Bleeding *in the* Jugular Veins *of the* Neck 303
VIII. *Of* Bleeding *in the* Veins, *called* Ranulæ, *under the Tongue* 304
IX. *Of* Phlebotomy *in the* Penis ibid.

Of the Symptoms *or* Accidents *which attend* Phlebotomy.

X. *Of an* Eccymoſis 305
XI. *Of the* Puncture *of a* Nerve *or* Tendon *in Phlebotomy* 306
XII. *Of* Wounds *of the* Arteries *in* Phlebotomy 307
XIII. *Of* Aneuriſms 310
XIV. *Of injecting Liquors into the* Veins, *and of tranſfuſing the* Blood *of one Animal into another* 323
XV. *Of* Inoculation *for the* Small-Pox 327
XVI. *Of* Scarification *and* Cupping 328
XVII. *Of* Bleeding *by* Leeches 332
XVIII. *Of* Acupuncturation *uſed by the* Chineſe *and* Japoneſe 334

XIX.

Chap.
XIX. *Of* Iſſues Page 334
XX. *Of* Bliſtering *with* Cantharides 336
XXI. *Of* Injections 338
XXII. *Of* actual Cauteries 339
XXIII. *Of* Burning *with* Moxa 340
XXIV. *Of* Cauſtic *and* Corroding Medicines 341
XXV. *Of Opening* Abſceſſes 342
XXVI. *Of* Warts ibid.
XXVII. *The Method of removing* Excreſcences, *fleſhy* Tumors, *and* Marks *from the* Mother 344
XXVIII. *Of* Encyſted Tumors, *and eſpecially* Scirrhi, Atheromata, Steatomata, Meliceres, *and others* 345
XXIX. *The Method of extracting foreign* Bodies *from* Wounds 349
XXX. *Of* Sutures *of* Wounds 350
XXXI. *Of ſeparating* Adheſions *betwixt the* Fingers *and* Toes 351
XXXII. *Of amputating* Diſeaſed *and* Superfluous Fingers 352
XXXIII. *Of amputating ſphacelated* Fingers *and* Toes. 353
XXXIV. *Of amputating the* Hand, Cubitus, *and* Humerus 355
XXXV. *Of amputating the* Foot *and* Leg 363
XXXVI. *Of amputating the* Thigh 366
XXXVII. *Of amputating the* Arm *in its Articulation with the* Scapula 369

PART II. SECT. II.

Of Operations *belonging to the* Head.

XXXVIII. *Of making* Iſſues *upon the* Coronal Suture 373
XXXIX. *Of* Arteriotomy *in the* Temples. 375
XL. *Of the* Hydrocephalus 377
XLI. *Of* Trepanning *the* Cranium 378
XLII. *Of Extracting Bodies fallen into the* Eyes 387
Chap.
XLIII. *Of* Tubercles *and* Excreſcences *on the* Eye-lids Page 388
XLIV. *Of* Warts *on the* Eye-lids 389
XLV. *Of Relaxation and Tumor of the* Eye-lids, *termed* Phalangoſis, *and* Proſis 390
XLVI. *Of the* Trichiaſis, *or Inverſion of the* Eye-lids, *in which the Hairs irritate the Eyes* 391
XLVII. *Of the* Ancyloblepharon, *or* Concretion *of the* Eye-lids 392
XLVIII. *Of the* Everſion *and* Gaping *of the* Eye-lids, *termed* Ectropium *and* Lagophthalmia 394
XLIX. *Of the* Encanthis 396
L. *Of the* Sarcoma *and* Hyperſarcoſis, *or* Excreſcence *formed betwixt the* Eye *and its* Lids 397
LI. *Of* Bleeding *in the* Eyes 399
LII. *Of* Scarifying *the* Eyes 401
LIII. *Of the* Epiphora, *or* Watery Eye 405
LIV. *Of the* Fiſtula Lacrymalis, *and of the Diſorders related to it* 407

Of Operations *on the* Eyes.

LV. *Of* Suffuſions *or* Cataracts 425
LVI. *Of Dilating* Contractions *of the* Pupil 440
LVII. *Of the* Pterygium, *or* Unguis Oculorum 442
LVIII. *Of the* Albugo, Leucoma, Nebula, Nubecula, *and other* Spots *in the* Cornea *of the* Eye 444
LIX. *Of the* Staphyloma 446
LX. *Of the* Hypopyon 448
LXI. *Of Inciding the* Cornea, *to diſcharge Extravaſated* Blood 450
LXII. *Of the* Diſtention *and* Prolapſus Oculi, *alſo the* Fungus *and* Cancer. 451
LXIII. *Of* Artificial Eyes 453
LXIV. *Of the* Strabiſmus, *or* Squinting 454

CONTENTS of the SECOND VOLUME.

Chap.
LXV. *The Apertion of a closed* Meatus Auditorius Page 1
LXVI. *Of extracting foreign Bodies from the Ears* 2
LXVII. *Of Tubercles in the* Meatus Auditorius 3
LXVIII. *Of Cauterising behind the Ears for the Tooth-ach* ibid.
LXIX. *Of Acoustic Instruments to help the Hearing* 4
LXX. *Of boring the Lobes of the Ear* 5
LXXI. *Of a* Polypus *in the Nose* 6
LXXII. *Of an* Ozæna 14
LXXIII. *Of Artificial Noses* 16
LXXIV. *Of Opening the Nostrils preternaturally closed* 17
LXXV. *Of the Hare-Lip* 19
LXXVI. *Of a Cancer in the Mouth and Lips* 24
LXXVII. *Of Opening the Teeth and Jaws which are clinched* 25
LXXVIII. *Of cleansing black and foul Teeth* 27
LXXIX. *Of hollow and decayed Teeth* 28
LXXX. *Of the Chirurgical Methods for easing the Tooth-ach* 29
LXXXI. *Of rectifying Irregularities of the Teeth, which lacerate the Tongue and Cheeks* ibid.
LXXXII. *Of drawing Teeth* ibid.
LXXXIII. *Of Artificial Teeth* 31
LXXXIV. *Of Lancing the Gums in Dentition* 32
LXXXV. *Of* Epulides, *or Excrescences of the Gums* 33
LXXXVI. *O* Parulides, *or Boils and Abscesses of the Gums* 34
LXXXVII. *Of depressing the Tongue* 35
LXXXVIII. *Of dividing the* Frænulum *of the Tongue.* ibid.
LXXXIX. *Of a* Ranula *or* Tumor, *and* Calculi *under the Tongue* 37
XC. *Of a* Scirrhus *and Cancer in the Tongue* 38
Chap.
XCI. *Of Ulcers in the Palate* Page 39
XCII. *Of stopping Perforations of the Palate into the Nose* 40
XCIII. *Of a Tumor and* Prolapsus *of the* Uvula ibid.
XCIV. *Of Scarifying the Tonsils when inflamed in a Quincy* 42
XCV. *Of opening Abscesses in the Tonsils* 43
XCVI. *Of extirpating scirrhous Tonsils* ib.
XCVII. *Of Tubercles and Excrescences in the* Fauces, *or near the Tonsils* 44
XCVIII. *Of extirpating scirrhous, maxillary, and parotid Glands* 45

PART II. SECT. III.

Of DISORDERS *in the* NECK, *curable by the Hands and Instruments.*

XCIX. *The Method of extracting and removing small Bones of Fish and other Animals, Plumb-stones, Pins, and Needles,* &c. *sticking in the* Fauces *or* Gula 48
C. *Of the* Brush *for scowring the Stomach* 49
CI. *Of the* wry Neck ibid.
CII. *Of* Bronchotomy, Laryngotomy, *or* Tracheotomy 51
CIII. *Of strumous or scrophulous* Tumors *and* Bronchocele 54
CIV. *Of* Setons 56

PART II. SECT. IV.

Of DISORDERS *of the* THORAX, *coming under the Province of Surgery.*

CV. *The Manner in which the Nipples of the* Breasts *in Women may be drawn out, extended, and milked* 58
CVI. *Of chapped and sore* Nipples 59
CVII. *Of a Cancer in the* Breasts 60
CVIII. *Of the Paracentesis or Perforation of the* Thorax 65

Chap.
CIX. *Of trepanning the* Sternum Page 68
CX. *Of the crooked or hump-back* 69

PART II. SECT. V.

Of DISORDERS *in the* ABDOMEN, *appertaining to Surgery.*

CXI. *The Method of tying the Navel-ſtring in new-born Infants* 70
CXII. *The Manner of diſcharging the Water contained in the* Abdomen, *in the Dropſy Aſcites by Paracenteſis* ibid.
CXIII. *Of the Cæſarean Section or Birth, being the Method of cutting the Fœtus out of the Womb* 74
CXIV. *Of Herniæ or Ruptures in general, and particularly of the Umbilical, and its Method of Cure* 85
CXV. *Of other* Herniæ, *and particularly thoſe of the Abdomen, or the* Hernia Ventralis 93
CXVI. *Of the* Bubonocele *or* Hernia Inguinalis 95
CXVII. *Of the Hernia Inguinalis incarcerata, or intercepted* Bubonocele 98
CXVIII. *Of the Hernia femoralis, or crural* Rupture 104
CXIX. *Of the Oſcheocele, or Hernia of the Scrotum, and particularly of the* Enterocele, *or Prolapſus of the Inteſtine into the Scrotum* 106
CXX. *Of the* Epiplocele, *or Prolapſus of the Omentum into the Scrotum* 112
CXXI. *Of ſpurious Ruptures, and firſt of the* Sarocele, *and* Caſtration 114
CXXII. *Of the* Hydrocele 117
CXXIII. *Of the* Hæmatocele 124
CXXIV. *Of the Hydrops Scrotalis and Pudendi* ibid.
CXXV. *Of the* Hydro-ſarcocele 125
CXXVI. *Of the* Hydro-enterocele 126
CXXVII. *Of the* Pneumatocele, *or* Hernia Flatulenta ibid.
CXXVIII. *Of the* Ciſſocele, *or* Hernia Varicoſa 127

Chap.
CXXIX. *Of a Cancer and Sphacelus in the Teſticle* Page 129
CXXX. *The Method of treating a Diſorder of the* Penis, *as*, 1. *of a* Phimoſis ibid.
CXXXI. *Of a* Paraphimoſis 131
CXXXII. *Of a* Cancer *and* Sphacelus *in the* Penis 133
CXXXIII. *The Manner of dividing the Frænulum of the* Penis 134
CXXXIV. *Of Warts, and other Excreſcences of the* Penis ibid.
CXXXV. *The Method of opening the imperforated* Glans *or* Urethra 135
CXXXVI. *The Method of curing an Incontinency of* Urine *in Males* 138
CXXXVII. *Of introducing the* Catheter *into the Bladder, in order to ſearch for the Stone, or diſcharge the Urine, when ſuppreſſed* 141
CXXXVIII. *Of* Caruncles *in the* Urethra 148
CXXXIX. *The Method of extracting a* Calculus *in the* Urethra 151
CXL. *Of* Lithotomy, *or cutting for the Stone in Males, particularly by the Old Method, termed* Apparatus Minor; *where we ſhall alſo propoſe ſomething concerning* Nephrotomy 154
CXLI. *Of Lithotomy by the* Apparatus Major 163
CXLII. *Of Lithotomy by the* Apparatus Altus, *or the high Operation of* Peter Francus, *whereby the Stone is extracted by an Inciſion in the hypogaſtric Region, above the* Oſſa Pubis 181
CXLIII. *Concerning the Artifices uſed by Frier* James (Frere Jaques) *in cutting for the Stone; as alſo on the* lateral Operation *of* Raw 205
CXLIV. *Of Puncturing the Perinæum and Bladder* 234
CXLV. *Of* Fiſtulæ *in* Perinæo 240
CXLVI. *The Method of dividing preternatural*

Chap.
natural Cohesions in the genital Parts of Women Page 243
CXLVII. *The Method of Opening the* Vagina, *when obstructed near the Womb* 246
CXLVIII. *Of the* Clitoris *growing too large* 248
CXLIX. *The Method of treating the* Nymphæ *when too much enlarged* ibid.
CL. *The Method of removing Tubercles, Caruncles, and other Excrescences in the* Vagina 249
CLI. *The Method of extracting the Stone in the Bladder of Women* 250
CLII. *The Method of treating difficult Births, the Fœtus being alive* 257
CLIII. *Method of extracting a dead Fœtus* 273
CLIV. *Of profuse* Hæmorrhages, *or Floodings of the* Uterus *in Women with Child* 278
CLV. *The Method of extracting the* Secundines 281
CLVI. *The Method of discharging* Molæ, *or false Conceptions* 287
CLVII. *Of a* Prolapsus Uteri, *or bearing down of the Womb* 288
CLVIII. *Concerning the* Prolapsus, *or bearing down of the Vagina* 294
CLIX. *Of an Incontinency of the Urine in Women* 295
CLX. *Of the* Perinæum *lacerated in Women* 296
CLXI. *Of Disorders and Operations proper to the* Anus, *and of* Clysters 297
CLXII. *Of Suppositories* 299
CLXIII. *The Method of Opening an imperforated* Anus 300
CLXIV. *Of a* Prolapsus Ani 302
CLXV. *Concerning Tumors of the* Anus, *such as the* Condyloma, Crista, Ficus, *and* Fungus 303
CLXVI. *The Method of treating the* bleeding Piles 304
Chap.
CLXVII. *The Method of treating the* Blind Piles Page 305
CLXVIII. *Of* Fistulæ *in the* Anus 309
CLXIX. *Concerning* Abscesses *of the* Anus 320

PART II. SECT. VI.

Concerning Disorders incident to the upper and lower Extremities, particularly to the Hands and Feet.

CLXX. *Of the* Paronychia, *or* Whitloe 324
CLXXI. *Of* Ganglions, *or Knots of the* Tendons 331
CLXXII. *The Suture of Tendons in the Hands* 333
CLXXIII. *Of Disorders belonging to the lower Extremities* 338
CLXXIV. *Of* Varices 343
CLXXV. *The Method of cutting out the Nail of the Great-toe, when it turns into the Flesh* 344
CLXXVI. *Of treating* Corns *in the Feet* 345
CLXXVII. *Concerning the Treatment of Infants that are* Bandy-legged, *with their Feet turning inward or outward* 346

PART III.

I. *Of* Bandages *in general* 350
II. *Of the* Bandages *belonging to the* Head 354
III. *Of* Bandages *for the* Neck 360
IV. *Of* Bandages *pertaining to the* Thorax 362
V. *Of* Bandages *proper to the* Abdomen *and private Parts* 371
VI. *Of* Bandages *for the* Arm *and* Hand 377
VII. *Of* Bandages *for the* Leg *and* Thigh 386
VIII. *Of the Deligation for a compound Fracture of the* Leg 393

BOSTON MEDICAL
DEC 15 1915
LIBRARY.

GENERAL
INTRODUCTION
TO
SURGERY.

Of the Nature, Conſtitution, Origin, Progreſs, Improvement, and Diviſion of SURGERY; *and of other Things in general, which are principally neceſſary for* STUDENTS *in* SURGERY *to be acquainted with.*

I. THE principal End of *Phyſic* is to prevent or relieve the Diſorders of the human Body. This the firſt Phyſicians endeavoured to effect by three Means, either by *Food*, *Medicines*, or the *Appliation of the Hand*[a]; or by all together, if the Caſe required it: which Method, Reaſon and Experience teach us, is abſolutely neceſſary at this Time. And of theſe three Branches of this ſalutary Profeſſion, they call the firſt *Diet*, or *Dietical* (διαιτητικὴ); the ſecond *Pharmaceutical* (φαρμακευτικὴ); and the third *Chirurgical* (χειρουργικὴ). For ſince the End of *Phyſic* could by no means be always obtained by *Diet* and *Medicine* alone (though they are of very great Service in preſerving and reſtoring the Health of Mankind), but manual Operation is alſo found ſometimes to be abſolutely neceſſary; it is plain therefore that this Branch of *Phyſic*, which is called *Surgery*, is very neceſſary to Mankind: more eſpecially, as it appears, that by this means many grievous Diſorders are relieved, as Wounds, Fractures, Luxations, and ſeveral others, where *Diet* and *Medicine* would afford very little, and ſometimes no Help at all. But that the Excellence and Neceſſity of this Art may appear more clearly, it may be neceſſary to obſerve, that other Arts only conduce to the Conveniencies of Life, but the Art of *Surgery* is frequently neceſſary for the Preſervation of Life, and the Continuance of Health, the moſt valuable Treaſure we can be poſſeſſed of. This Neceſſity appears more particularly in dangerous Wounds received in War, Skirmiſhes, or Sieges[b], where many brave Men muſt neceſſarily periſh

[a] See CELSUS, Præf. Lib. I. pag. 3. Edit. Almelov. & Patav.

[b] Of the Uſe and Excellency of Surgery, ſee VESALIUS, in Præfat. Corp. Hum. fabric. GESNER, in Script. opt. Chir. CYPRIAN, Orat. Encom. in Chirurg. SCHELLAMER, Præf. de Tumor. and KESSELRING, in Diſſert. de meth. *Fourbertian*, who treats very learnedly of the flouriſhing State of Surgery, and contends, that it is equal, if not preferable, to Phyſic, from the great Certainty and Rationality of its Practice.

from Loſs of Blood, and other Cauſes, unleſs they were reſtored, and ſnatched as it were from the Jaws of Death, by the Skill of their Surgeons. And no doubt the better Opinion the Soldiers conceive of their Surgeons, the more Spirits have they for the Combat, having good Confidence that the Wounds they receive ſhall be properly treated, and their Lives preſerved. And from hence, becauſe Surgery is chiefly exerciſed in the Treatment of Wounds, it is called by the *Germans* the Cure for Wounds *(Wund-Arzeney)*; not as if Wounds were the ſole objects of *Surgery*, but as it is of more particular and frequent Service in Caſes of that Kind.

Surgery, what.

II. " *Surgery, ſays* CELSUS [a], is that Branch of Phyſic, which informs us how " to cure or prevent Diſorders by the Aſſiſtance of our Hands or Inſtruments, " or by the Application of external Remedies," as Diſorders are frequently prevented by Bleeding, Scarifying, opening of Iſſues, and by Setons, *&c.* Since therefore *Surgery* is properly the Work of the Hand, it is very juſtly called by the *Greeks* Χειρουργία from the two *Greek* Words of that Signification, Χεὶρ and Ἔργον from whence the Perſon alſo ſkilled in this Work was called a *Chirurgeon*. But He, whoſe Office it is to cure Diſorders only by adminiſtring Medicines internally, and by preſcribing Rules for the Regulation of the Diet, is at preſent in *Latin* called *Medicus:* though this is a modern Diſtinction, and unknown to the Ancients, among whom both Offices were performed by the ſame Perſon, called Ἰατρὸς; as appears plainly by the Writings of HOMER, HIPPOCRATES, CELSUS, and many others.

It may be called a *Science* and an *Art*.

III. Some call *Surgery* a *Science*, others an *Art:* but, in my opinion, it will claim either Appellation. For it may be called a *Science*, becauſe the Student in Surgery, before he is ſkilled in the Method of healing, muſt have acquired the Precepts or Foundation of what is to be done towards diſcovering and remedying Diſorders that are to be relieved by the Aſſiſtance of the Hand, from *Anatomy*, *Phyſics*, and *Mechanics*; for without this Knowledge he would not only go very idly to work, but would do more Harm than Good to his Patients, and conſequently to the Public. It alſo well deſerves the Name of an *Art*, when any one is ſo well verſed in the Elements of this Art, that he is able to preſerve the Body ſound, as well as to relieve it when it is otherwiſe. Hence we very properly term thoſe ſkilled in the *Art* of *Surgery*, who are expert in healing Wounds, replacing fractured and diſlocated Bones, and underſtand the right Methods of treating other Diſorders which require the Aſſiſtance of the Hand or Inſtruments. From hence, I imagine, aroſe the Diſtinction which ſome have made between *theoretical* and *practical Surgery*. Thus *Surgery* was conſidered, when ranked under the firſt Denomination, as a *Science:* as when a Man has learnt and underſtands the Rules, and the Reaſons upon which thoſe Rules are grounded, which teach the beſt Methods of treating Diſorders that call for the *Surgeon's* Hand, and in what manner *Operations* (as they are vulgarly called) are to be performed; but never attempts the Performance of any of theſe *Operations*, whether they are dividing, amputating, cauteriſing, or reducing Bones, or of any other Kind. This *Science* we call *Medical Surgery*. And this Branch of *Surgery*, at leaſt, all *regular Phyſicians* ought to be well acquainted with; that they may be of Service to the *Surgeons* and their *Patients*,

[a] Lib. I. Præfat. pag. 3. and Lib. VII. in the Beginning of the Preface.

by

by being able to give prudent Advice in Diſorders of this Kind. *Surgery*, when it falls under the ſecond Denomination, and is termed *practical*, ſignifies the Exerciſe of it, or the *Art* of performing Chirurgical Operations, of replacing, tying, cutting, extirpating, dividing, cauteriſing, *&c.* The *Practical Surgeon* is well inſtructed in the Art of managing his Hands and Inſtruments dextrouſly in the Performance of ſuch Operations as the Neceſſity of the Caſe ſhall require. Much the greater Part of the modern *Phyſicians* have been content with the Knowledge of the former Part of *Surgery*, leaving the Execution of the latter, which is much to be lamented, to unſkilful Quacks and [a] Mountebanks. This happens partly becauſe the Diſorders that are curable by the prudent Adminiſtration of Medicines internally, and a well-regulated Diet, which more immediately come under the Province of the Phyſician, are ſo numerous, and withal ſo intricate, as to be a ſufficient Exerciſe for his whole Study; and partly becauſe Cures which are to be performed by the Hand, eſpecially thoſe which are attended with Danger or Severity in the Execution, require a ſingular Fortitude and Firmneſs of Reſolution; or, as the elegant CELSUS expreſſeth it [b], A Mind intrepid, diveſted of Tenderneſs, and unmoved by the Shrieks of the ſuffering Patient: Which is to be met with in very few, though they may be perfectly well acquainted with every thing that ought to be done. But whoſoever deſires to be a perfect *Surgeon*, muſt be a thorough Maſter of his Profeſſion under both Heads, as a *Science* and as an *Art:* and in ſuch a manner that the *theoretical* Part, or Knowledge of the Elements (in which *Anatomy* claims the firſt place), ſhould preceed the Exerciſe of the Art. For if any one ſhould be bold enough to proceed in the contrary Method, and invert this Rule, by undertaking to perform Operations, eſpecially thoſe of the more difficult Kind, before he had made himſelf well acquainted with *Anatomy*, the Nature of Diſeaſes, and what is proper to be done towards removing them; of Neceſſity he will do great Harm to thoſe entruſted to his Care, and deſtroy more than he will ſave; though this is unhappily every where practiſed by bold daring Fellows, to the great Detriment of Mankind, and to the Diſgrace of this truly noble Art. For "Knowledge [c] ought to direct the Hands, and ſhew them what is proper for "them to perform." Therefore, if any *Surgeon* has been long in Practice, and, as they are fond of terming it, is a *Man of great Experience*, and is not thoroughly verſed in *Anatomy*, and the Inſtitutions of *Surgery*, his Actions are always doubtful and uncertain, and are ever obnoxious to Multiplicity of Dangers. Therefore, it is neceſſary for the good Surgeon to be a thorough Maſter of both; but he whoſe comprehenſive Knowledge takes in all the other Branches of *Phyſic*, as many amongſt the ancient and modern *Phyſicians* have done [d], is by ſo much the abler and more accompliſhed *Surgeon*.

IV. The end of *Surgery*, as appears by what we ſaid above at N°. I. is three-fold: 1. *To preſerve Mankind in a ſound State*, in the manner we explained it at The End of Surgery.

[a] This is very rarely the caſe in *England*, but too common in *Germany*.

[b] Lib. VII. in Præfat. [c] CELSUS ſpeaks more largely of this, Lib. I. in Præfat.

[d] As ÆSCULAPIUS, PODALIRIUS, MACHAON, HIPPOCRATES, GALENUS, CELSUS, ÆTIUS, ÆGINETA, ORIBASIUS, GUIDO CAULIACUS, SALICETUS, VESALIUS, FALLOPIUS, MARIANUS SANCTUS, JO. DE ROMANIS, VAROLIUS, CABROLIUS, FABR. AB AQUAPENDENTE, M. A. SEVERINUS, HILDANUS, SPIGELIUS, GLANDORPIUS, GEIGERUS, SCULTETUS, MARCHETTUS, ROLFINCIUS, WEPFERUS, MURALTUS, SOLINGENIUS, RUYSCHIUS, BIDLOUS, NUCHIUS, GROENVELTIUS, CYPRIANUS, BOHNIUS, BRUNNERUS, RAVIUS, LEUSDENIUS, &c.

N° II. 2. *The Reſtitution of a ſound State*; that is, the Cure of Diſeaſes by the Aſſiſtance of the Hands. Or, 3. *To preſerve the Life of a Man*, though with a maimed and wounded Body, if it is impoſſible to render it entire. This third End is chiefly obtained by the Amputation of ſphacelated, cancerated, or carious Limbs; ſo in Cancers, Schirrus's, old Ulcers, and other ſuch like incurable Diſeaſes, and in ſeveral Diſorders of the Head, eſpecially in Weakneſſes of the Eyes and Ears, to prevent their growing worſe, it is uſual to order Iſſues, Setons, frequent Blood-letting, Bliſtering, *&c.* though a perfect Cure is not perhaps to be looked for. And therefore under this Head may be ranked inveterate *Herniæ*.

Auxiliaries of Surgery what.

V. The *Auxiliaries* or *Means* which *Surgery* makes uſe of to obtain the End we have been diſcourſing of, are chiefly the *Surgeon's Hands* and proper *Inſtruments*. For as often as a fractured or diſlocated Bone is to be reduced, a Vein to be opened, a Stone to be extracted, or a Cataract depreſſed, proper Inſtruments are always neceſſary. But that every thing may go on with more Speed, Eaſe, and Safety, the Adminiſtration of proper internal Remedies, and the Regulation of Diet, will never be neglected, in any of the foregoing Caſes, by a prudent *Surgeon*; which confirms the Saying of CELSUS [a], "That all the Parts "of Phyſic are ſo intimately connected, that it is impoſſible to ſeparate any "one of them entirely from the whole." And in another place [b], "I, *ſays he*, "can eaſily conceive one Man to be capable of performing all the Offices of "Phyſic, and, where they have been divided, think him praiſe-worthy that "unites them in himſelf."

Origin of Surgery.

VI. The ſtrong Connection that there is between *Phyſic* and *Surgery*, is, in my Opinion, a perſuaſive Argument that the Origin, Progreſs, and Fate of both, were always the ſame. Though, to ſay Truth, I cannot help believing with CELSUS [c], and others, that *Surgery* is more ancient than any other Branch of *Phyſic*, and near coeval with Mankind, and therefore the true Parent of *Medicine*. The nearer Mankind was to its firſt Original, at ſo much the greater Diſtance were they, as CELSUS obſerves, from Luxury and Debauchery, and of conſequence ſo much the farther removed from internal Diſeaſes. The native Strength of Man, as yet unhurt by Intemperance, ſtood in no Need of internal Aids. But on the other hand, even in the earlieſt Times, Men were as liable, as we are to this day, to external Injuries, which require the Aſſiſtance of the *Surgeon's* Hand. For who in thoſe days was ſecure from falling, or from Fractures of the Bones, which are the Conſequences of ſuch Accidents; from the Bites of wild Beaſts; or from the Wounds of an open or an inſidious Enemy? Since in the very firſt Ages Men waged War with each other, can it be reaſonably ſuppoſed that they were always free from Bloodſhed, fractured and diſlocated Bones, *&c.*? As therefore it cannot be doubted, but that, by the Direction of Nature, who taught them to extract Thorns, and to tie up Wounds, to prevent a large Effuſion of Blood, they by degrees were uſed to receive Aſſiſtance from the Hand of ſome Kind of Inſtruments; and if by chance, after many repeated Experiments of this Kind, any thing ſhould be found to anſwer the deſired End, diligent Men would certainly retain it in their Memories, and mark

[a] In Præfat. Lib. V. item SCRIBONIUS LARGUS, Cap. LXVIII.
[b] In Præfat Lib. VII.
[c] In Præfat. Lib. I.

it

it down, which being repeated with Succeſs in ſimilar Caſes, was handed down to Poſterity. So this ſalutary Profeſſion took its Riſe from ſmall, and thoſe rude, Beginnings and vulgar Experiments, till by degrees it received Improvements, and was brought to its preſent Perfection by the Induſtry and Sagacity of ingenious Men.

VII. By as much as we can collect from ancient Hiſtory, the *Chaldeans* and *Egyptians*, who were the firſt Cultivaters of Science, as we learn from the Scriptures, found *Surgery* naked and in her Infancy, enriched her with new Experiments, and laid her down Rules and Inſtitutions to walk by. And afterwards [a] *Surgery* was ſtill much farther enriched by the *Greeks*, thoſe ancient and noble Patrons of Knowledge. APOLLO and his Son ÆSCULAPIUS were chiefly celebrated as Surgeons in thoſe Ages, who, for their Sagacity in cultivating this Science, gained to themſelves ſo great Applauſe, that they were reckoned among the Number of the Gods. After theſe came PODALIRIUS and MACHAON, two ſons of ÆSCULAPIUS, who accompanied AGAMEMNON to the *Trojan* War, and were of great Service to the Army. But HOMER never takes notice of them as being ſerviceable in the Plague or other Kinds of Diſtempers, but only as Perſons ſkilful in healing Wounds by the Application of Inſtruments and Medicines. From whence it appears, that they were only expert in *Surgery*, and that it is the moſt ancient Branch of *Phyſic*. We read of CHIRON the *Centaur*, and other *Surgeons* after them, who equalled them in Reputation, but the Monuments of thoſe days are long ago entirely defaced by time. HIPPOCRATES the *Coan* ſeems to have far exceeded all the reſt in Sagacity and Succeſs; CELSUS declares of him, "that he was not only celebrated "for Wiſdom and Art, but for Eloquence alſo." He inherited *Surgery* by Deſcent, being ſprung from the Race of ÆSCULAPIUS. With no leſs Judgment than Aſſiduity he formed a compleat Syſtem of the Experiments and Rules of his Anceſtors, with their various and elaborate Methods of Cure; which he greatly improved, through the Aſſiſtance and Directions of DEMOCRITAS, by his conſtant and indefatigable Attention to the Study of *Human Anatomy* [b]. For which reaſon they are by no means deceived who have pronounced HIPPOCRATES the Father of all Branches of *Phyſic*, but more particularly of *Surgery*. The Writings of this great Man, notwithſtanding they are the moſt ancient, ſo far exceed all the reſt, that at all times they have been laid down as examples to all *Profeſſors* of *Phyſic*.

Improvements of *Surgery* in *Greece*.

VIII. The *Greeks*, by their ſtrenuous Application to the ſtudy of *Surgery*, excited a Deſire in the *Romans*, and at the ſame time in the *Egyptians*, to give encouragement to the ſame Art. "[c] About this time, a little before the "Birth of CHRIST, PHILOXENUS was eminent in *Surgery*, who, according "to CELSUS, wrote ſeveral Volumes upon this Branch of *Phyſic*. GORGONUS "alſo and SOSTRATUS, and HERONES, and the two APOLLONIUS's, and AM"MONIUS ALEXANDRINUS, and many other famous Men, all enriched this "*Science* with ſomething new. At *Rome* alſo, *ſaith the ſame Author*, there "were Profeſſors of great note, eſpecially TRYPHON the Father, and EUEL"PISTUS the Son of PHLEGES, and as we may gather from his Writings, the

Proficiency of the *Romans*, *Greeks* and *Arabians* in *Surgery*.

[a] Vide CEL. Lib. I. Præf. [b] As CELSUS teſtifies, Lib. I. Præf. [c] See CELSUS in Præf.

"principal of all, MEGES; all of them, by various Emendations, ſtill added "Improvements to this *Science*." But the Writings of theſe Men are not tranſmitted to us. In the Ages next after CHRIST, CELSUS acquired the greateſt name among the *Latin Writers* (on whom we have beſtowed repeated Encomiums), but among the *Greek Writers*, GALEN, PAULUS ÆGINETA, ÆTIUS, and ORIBASIUS; whoſe Works are ſtill extant. But after this, in the ſubſequent Ages, the barbarous Nations began to over-run the whole Earth, and *Surgery* was ſo far from encreaſing, that it received the ſame Fate with all other Branches of *Science*, and ſuffered under the common Calamity. Therefore it is no wonder that thoſe Times produced no one to whom *Surgery* was indebted, if you except only RHAZES, HALY ABBAS, ALBUCASIS, and AVICENNA, who flouriſhed in *Arabia* about the XIth or XIIth Century. It is to be obſerved though by the way, from GUIDO DE CAULIACO[a], the *Phyſicians* at this time firſt refuſed to undertake the Performance of any *manual Operation*, though indeed in CELSUS's time there were ſome who ſeparated *Surgery* from *Phyſic*.

Induſtry of Surgeons of a later Date.

IX. In the XIIIth and XIVth Centuries, when the Clouds that had overſhadowed all Science began to diſperſe, *Surgery* alſo again emerged with the reſt, and was cultivated both by *Phyſicians* and *Surgeons*. At firſt aroſe BRUNUS, THEODORICUS, SALICETUS, LANFRANCUS, ARNOLDUS DE VILLA NOVA, and many others equally eminent: but afterwards, in a ſtill more conſpicuous Light, ſhone that true Reſtorer of *Surgery* GUIDO DE CAULIACO, DE LARGELATA, JO. DE VIGO, VESALIUS, FALLOPIUS, ANDREAS A CRUCE, ARCAEUS, MARIANUS SANCTUS, ANGELUS BOLOGNINUS, BERENGARIUS CARPUS, ALPHONSUS FERRIUS, JOANNES TAGOLTIUS, BARTHOLOMAEUS, MAGGIUS, PARAEUS, SCHILLHANS, GERSTOFF, BRUNSVIC, RYFF, and others, who greatly contributed, as appears by their Writings, to the Improvement of *Surgery*.

Of the Moderns.

X. At length in the laſt and preſent Age, by the Induſtry firſt of the *Italians*, *French*, *Germans*, and more latterly alſo of the *Engliſh*, *Surgery* has been ſo wonderfully enrich'd with extraordinary Inventions and Obſervations in *Anatomy*, *Mechanics*, and *Phyſics*, and with elegant Inſtruments and new Methods of Curing, that it ſeems to want little or no Addition to raiſe it to its higheſt State of Excellency and Perfection. But although I purpoſed now to give a regular Account of thoſe by whoſe Labours *Surgery* has gained the Fruits it at preſent enjoys, yet ſince the Number of thoſe is ſo large, let it ſuffice for the preſent to reckon up the principal of them; leaving the Enumeration of the reſt to another opportunity. In this Rank we may reckon FABRICIUS AB AQUAPENDENTE, FABRICIUS HILDANUS, M. A. SEVERINUS, SPIGELIUS, MARCHETTUS, GLANDORPIUS, JO. SCULTETUS, FELIX WURTZIUS, GUILLEMAU, CÆSAR MAGATUS, CASP. TALIACOTIUS, GOUSMELINUS, RONHUYSIUS, VAN NEEKEREN, CORN. SOLINGEN, NUCHIUS, BURMANNUS, MAURICEAU, TOLET, VERDUCCIUS, BIDLOUS, RUYSCHIUS, BOHNIUS, CYPRIANUS, RAUIUS, MASSIERUS, DIONIS, PETIT, WISEMAN, DOUGLAS, CHESELDEN, GARENGEOT, MARINUS, TURNER, MORAND, LE DRAN, and many others whom you will find among the *Chirurgical* Writers.

[a] See his Chirurgical Works.

XI. Be-

XI. Before we proceed farther, I think it will be of Service to the Students in Surgery, to inform them of the beſt Writers that have treated of particular Parts of *Surgery*, and have either handled theſe ſeparately, or at leaſt with ſuperior Succeſs: in deſcribing of theſe I ſhall obſerve, as near as I can, the ſame Order, in which this Book is diſpoſed. And firſt, the following Authors have treated of *the five principal Parts of Surgery*, to wit, *Wounds*, *Fractures*, *Luxations*, *Tumours*, *and Ulcers*; VESALIUS, TAGULTIUS, ANDR. A. CRUCE, FABRIC. AB AQUAPENDENTE, then CORTESIUS, PECCETIUS, WISEMAN, MUNNICK. Writers on particular Parts of Surgery.

XII. The following Writers upon *Wounds* in general well merit reading; PARÆUS, ARCÆUS, FABRICIUS AB AQUAPENDENTE, GLANDORPIUS, MAGATUS, BELLOSTIUS. Upon *Wounds of the Head* in particular; HIPPOCRATES, CELSUS, CARPUS, ARANTIUS, PAVIUS, MILLERUS, SCHULTZIUS, WALTHERUS, and ROHAULT, a modern *Frenchman*. On *Diſeaſes of the Eyes*; FALLOPIUS, Jo. LANGIUS in *Ephemerid. N. C. Cent.* & VI. ST. YVES. On *Wounds of the Breaſt*; FUMANELLUS, PECHLINUS. *On Gunſhot Wounds*; PLAZZONUS, MAGGIUS, FERRIUS, ROTA, PARÆUS, FALLOPIUS, GUILLEMEAU, HILDANUS, BOTALLUS, BURMANNUS, TASSIN, VERDUC, VAUGUION, CHARRIERE, and, tho' laſt, not the leaſt eminent, LE DRAN. Of *Tents*; BAIETUS. Of the *Abuſe of Tents in Wounds*; MAGATUS, BELLOSTIUS, and a late *French* Piece of CHABERT's, and of LUPUS in *Italian*, who have maintained that Wounds ſhould ſeldom be kept open by Tents. Uſeful *Obſervations on Wounds* have been publiſhed by SCULTETUS, BELLOSTIUS, SCHWARTIUS, DE LA MOTTE, CHABERT, LE DRAN: The beſt Diſcourſes on *Mortal Wounds*, and the Method of diſcovering them to be ſo, have been written by BOHNIUS, TEYCHMEYERUS, ZACCIAS, AMMANNUS, VALENTINUS, ZITTMANNUS, FRID. HOFEMANNUS: To the ſame purpoſe is a Book whoſe Title is, *The Art of forming Prognoſtics in Surgery*, in *French*, and BLEGNIUS upon the ſame Subject. Authors on Wounds.

XIII. On *Fractures* and *Luxations*; PARÆUS, AQUAPENDENS, HILDANUS, VERDUC, in a particular Volume on this Subject; LE CLERC in his *Oſteology*; PETIT's *Art of curing the Diſeaſes of the Bones*, in *French*; PALFYNUS, in *Dutch*. On *Fractures of the Cranium*; HIPPOCRATES, CELSUS, CARPUS, BERENGERIUS, CORTESIUS, PAAW, and the Authors above recited, who have diſcourſed on Wounds of the Head. On Fractures and Luxations.

XIV. On *Tumours*; INGRASSIUS, FALLOPIUS, ARANTIUS, SAPORTA, M. A. SEVERINUS, SCHELHAMMER, CALVERS, MAUBEC, in *French*. On *Suppuration*; LAZERME. On *Abſceſſes*; SEVERINUS. On the *Carbuncle and Peſtilential Bubo*; FALLOPIUS, GEMMA. On *Œdema and Schirrus*; HARRIS. On *Fungous Tumours of the Limbs*; SLENOGTIUS. On *Gangrene and Sphacelus*; HILDANUS, KOENERDING, HARRIS. On *Burns*; HILDANUS. On a *Cancer*; ALLIO, GENGRON, HELVETIUS, HARRIS, and much earlier TEXTOR. On *Ulcers*; TAGAULTIUS, BONONINIUS, FALLOPIUS, AQUAPENDENS, VERDUC, LE CLERC. On *Caries of the Bones*; PETIT. On a *Spina Ventoſa*; SEVERINUS, PANDOLPHINUS, MARCHETTUS, and WALKER, in *High Dutch*. Of Tumours and Ulcers.

XV. The beſt Authors on *Chirurgical Operations* in general, are CELSUS, ÆGINETA, PARÆUS, FABR. AB AQUAPENDENTE, SOLINGEN, NUCHIUS, VERDUC, VAUGUION, CHARRIERE, DIONIS, PALFYNUS, MASSIERUS, GARENGEOT, MARINUS, LE DRAN, SHARPE. On Chirurgical Operations.

XVI.

Of Blood-letting, and Operations of the like Nature.

XVI. On *Bleeding* in particular, beſides many others, you will find BOTALLUS, P. P. MAGNUS, SCHMID, JONDOT, VERNA, MELLIUS, CRONE, HARRIS, SILVA, CHEVALIER, HECQUET, QUESNAY, MARTIN. On *Bleeding oftener in the Jugular Vein*; TRALLES. On the *Aneuriſm*; BARTHOLIN, HORN, HARRIS. On *Infuſing Humours into the Blood*; MAJOR, ETMULLER, ELSHOLZIUS. On *Transfuſion*; LOWER, STURMIUS, SANTINELLUS, MAFRIDUS, MARKLINUS, BURMANNUS.

Of Operations which are performed on various Parts.

XVII. Of *Inoculation of the Small Pox*; MAITLAND, PILARINUS, LE DUC, VATERUS, WREDEN, HARRIS. Of *Cupping and Scarifying*; CELSUS, GALEN, MAGNUS, BOTALLUS, MANNUS, MELLUS. Of the *Abuſe of Cupping in Putrid Fevers*; AQUAPENDENS. Of the *Egyptian Method of Scarifying*; ALPINUS, STAHLIUS. Of *Leeches*; GALEN, MAGNUS, HEURNIUS, STAHLIUS. Of *Puncture with a Needle after the Manner of the Japoneſe*; RHYNE, and KOEMPFLERUS. Of *Iſſues*; GALVANUS, in *Italian*; GLANDORPIUS, RESTAURANT, and SCHORETUS, in *High Dutch*. Diſſertations on this Subject have been written, by ALBINUS, SCHELLHAMMERUS, SCHACHERUS, FR. HOFFMANUS, HILSCHERUS, and others. On *Cantharides*; GEYERUS, ALBINUS, WEDELIUS. On the *Uſe of Bliſters*; CAIUS, NENTERUS, FR. HOFFMANNUS, LAETIUS A FONTE, and HERCULES SAXONIA. On *Cauteries*; ALBUCASIS, CAPIVACCIUS, GAVASSETIUS, SEVERINUS, COSTÆUS, MAGNUS, FALLOPIUS, FIENUS, BARTHOLINUS, BAUHINUS, SLEVOGTIUS. On the Method of uſing the *Indian Moſs (Moxa)*; TEN RHYNE, CLEYERUS, VALENTINI, LE TEMPLE. Of *Atheromata* and *Steatomata*; GORTESIUS, JO. LANGIUS, ELSHOLSTIUS. Of the *Meliceris*; HILDANUS, SEBIZIUS. Of *Encyſtated Tumours*; SLEVOGTIUS. Of *Extracting foreign Bodies from Wounds*; BIDLOO. Of *Amputation of the Limbs*; FIENUS, HILDANUS, HOFFMANUS, HILSCERUS. Of *a new Method of taking off Limbs*; JONGE, VERDUIN, RUYSCHIUS, KOENERDINGIUS, SALZMANNUS.

Operations on the Head.

XVIII. Of *an Iſſue upon the Coronal Suture*; SLEUOGTIUS. Of *Arteriotomy*; FIENUS, SEVERINUS, ALPINUS, SCHEURLIUS. Of the *Hydrocephalus*; CORTESIUS. Of *Trepanning*, and particularly of the Difficulties that attend that Operation; FIENUS, BOHNIUS, COSCHWITZIUS.

On the Eyes.

XIX. Of *Diſorders of the Eyes*; BARTISCHIUS, who has very accurately delineated many of the Diſeaſes of the Eyes; GUILLEMEAU, READ, COWARD, MAITRE JEAN, KENNEDY, ST. YVES. Of the *Trichiaſis*; HEISTER. Of *Scarification of the Eyes*; MANCHATUS, PLATNERUS. Of the *Fiſtula Lacrymalis*; ANELLUS, HEISTER, MELLIUS, in *Italian*, PLATNERUS. Of a *Cataract*; MAITRE JEAN, BRISSEAU, WOLHUSIUS, HEISTER, WIDEMANNUS, MARINUS. Of the *Hypopion*; BIDLOUS, and MAUCHARTUS, who likewiſe publiſhed Treatiſes on the *Ectropion*, the *Fiſtula in the Cornea*, the *Empyeſis*, and other Diſeaſes of the Eyes.

On the Noſe and Mouth.

XX. Of a *Polypus of the Noſe*; GLANDORPIUS. Of the *Hair Lip*; MARINUS. Of *Diſorders of the Teeth*, and the Methods of remedying them; GUILLEMEAU, STROBELBERGERUS, CRONE, and FRAUCHARD, a *Frenchman*; who lately wrote a Treatiſe called *Le Chirurgien Dentiſte*. Of the *Epulis and Parulis*; SCHELLHAMMERUS,

On the Neck and Breaſt.

XXI. Of *Laryngotomy*; CASSERIUS, MOREAU, FIENUS, DEKKERUS, MONAVIUS, FONTANUS, MASSIERUS. Of *Strumæ and Scrophulæ*; LAURENTIUS, BROWNE, GIBBS. Of *Setons*; GULVANUS, JO. FRANCUS, WEDELIUS, METZGERUS.

GERUS. Of the *Cancer of the Breaſts*; ſee above, under the Head *Cancer*. Of *Gibboſity*; WEDELIUS.

XXII. Of a *Paracenteſis*; there are ſeveral academical Theſes extant upon this Subject, by MEIBOMIUS, ALBINUS, SLEVOGTIUS, HENNINGERUS. Of the *Cæſarean Birth*; ROSSETUS, BAUHINUS, DOERING, HILDANUS, BULEAU, RAYNAUDUS, FIENUS, LANKISCHIUS, CYPRIANUS, SLEVOGTIUS. Of *Herniæ*; PETRUS FRANCUS, GEIGERUS, LE QUIN, LAUNAY, BERENGER, VAN HAMMEN, WIDEMANNUS; HARRIS, HOUSTON, in *Engliſh*; JO. SERMES, in his Book of *Lithotomy*, in *Dutch*, and divers academical Theſes; in particular upon the *Hernia incarcerata*, by MAUCHART; on the *Crural Hernia*, by KOCHIUS; on the *Enterocele*, by ROLFINCIUS and PETERMANNUS; on the *Sarcocele*, by MARINUS; on the *Hydrocele*, by the ſame; and on the *Abuſe of Kelotomy*, by HEISTER. On the Abdomen.

XXIII. Of a *Phimoſis* and *Paraphimoſis*; WEDELIUS. Of the *Cloſure of the natural Paſſages*; WIERUS. Of *Imperforations*; WEDELIUS. Of *Hypoſpadiacs*; LAVATERUS. Of *Paſſing the Catheter*; MEIBOMIUS, MARINUS. Of a *Stone in the Urethra*; MARINUS. Of a *Caruncle in the Meatus Urinarius*; FERRIUS, LACUNA, BENEVOLUS. Of *Fiſtulæ in the Urethra*; HILDANUS, MARCHETTUS, BECKERUS. On the Parts of Generation.

XXIV. Of *Lithotomy, and particularly of what they call the great Apparatus*; MARIANUS SANCTUS, HILDANUS, TOLETUS, GROENFELD, ALGHISIUS, MARINUS, CALLOTUS. Of *the leſſer Apparatus*; CELSUS, ALBUCASIS, CAULIACO; but it was afterwards laid aſide; yet not long ſince MARINUS, an *Italian*, defended it in ſome particular caſes, though by others it is altogether rejected. Of *the high Apparatus*; PETRUS FRANCUS, ROSSETUS, Jo. DOUGLAS, CHESELDEN, MIDDLETON, MORAND, J. SERMES, PROEBISCHIUS, and HEISTER. Of FRERE JACQUES's *Method*; MERYUS, LISTERUS, DIONIS. Of RAW's *Method*; ALBINUS, HERTIUS, and JAC. DENYSIUS. Of the *lateral Operation*; JAMES DOUGLAS. Of *the different Methods of cutting for the Stone*; PYE, an *Engliſhman*, and LE DRAN, a *Frenchman*; and SCHEFFERUS, and HERTIUS, in their academical Theſes, and others. Of FOUBERT's *Method*; KESSELRING. Of *the Methods of curing the Stone*, invented by FOUBERT, GARENGEOT, PARCHET, LE DRAN, and LE CAT; GUNZIUS. Of *Abuſe of Tents after Lithotomy*; HILDANUS. Of *the Puncture of the Bladder in a Suppreſſion of Urine*; MARINUS, MEYERUS. Of Lithotomy.

XXV. Of *the Art of Midwifry*; among the Ancients, RUPEUS, RUEF, RHODIO, PARÆUS; among the Moderns; SCIPIO MERCURIUS, MAURICEAU, PEU, PORTAL, VIARDEL, VOELTERUS, SIGISMUNDA, a Midwife of *Brandenburg*, DEVENTER, DIONIS, MELLIUS, ST. AMAND, DE LA MOTTE, HOORN, SUECUS, WIDEMANNUS. Of *the Method of extracting a dead Child*; HIPPOCRATES, SOLINGEN, FONTANUS, and the Authors we have juſt recited. Of *the Bearing down of the Womb*; BECKIUS. Of the Art of Midwifry.

XXVI. Of *Clyſters*; LANZONUS, SWARTZIUS. Of *the Fiſtula of the Anus*; MARCHETTUS, LE MONNIER, GLADBACCIUS, BASSIUS. Operations on the *Anus*.

XXVII. Of *the Paronychia*; GLANDORPIUS, WEDELIUS, ALBINUS. Of *the Suture of the Tendons*; KISNERUS. Of *Clefts in the Feet*; WEDELIUS. Of *Ingrafting*; TALIACOTIUS, SALTZMANNUS. On the extreme Parts.

On Bandages.

XXVIII. Of *Bandages*; GALEN, tranſlated by VIDO VIDIUS, with Figures; PARÆUS, in his Surgery, Part III, VERDUC on *Bandages* in *French*, and SOLINGEN; but the beſt Writers of all are LE CLERC, in his *Appareil Commode*; and BASSIUS in *High Dutch*, and ULHORNIUS. Of *Chirurgical Inſtruments* you may conſult ORIBASIUS, PARÆUS, and SCULTETUS.

Writers of Obſervations.

XXIX. Of *Obſervations in Surgery*; the beſt are related by PARÆUS, HILDANUS, SCULTETUS, MARCHETTUS, TULPIUS, MEEKREN, ROONHUSIUS, LAMBSWERDIUS, RUYSCHIUS, BELLOSTIUS, PURMANNUS, SAVIARDUS, DE LA MOTTE, CHABERT, LE DRAN.

Miſcellaneous Writers.

XXX. Of *the principal Controverſies in Surgery*, conſult FIENUS. On *the Duties of a Surgeon in the Army*, read FRANC. DE ROMA, MURALTUS, SCHMID, TASSIN, PURMANNUS, BELLOSTIUS, ABEILLE. Of *Surgery in the Time of a Plague*; PURMANNUS. Of *Chirurgical Anatomy*; GERGA, CHESELDEN, PALFINUS. Of *Medicines uſed in Surgery*; HOLLERIUS, PIGRÆUS, WURTZIUS, HILDANUS, in his Tract *de Ciſtâ Militari*, ETMULLER *de Chirurgiâ Medicâ*, LE CLERC, VERDUC DE FASCIIS, and BELLOST *in Pharmaciâ Chirurgicâ*. *Chirurgical Inſtruments* are beſt deſcribed by ALBUCASIS, ANDR. A CRUCE, HILDANUS, GUILLEMEAU, FABR. AB AQUAPENDENTE, PARÆUS, SCULTETUS, SOLINGEN, MASSIERUS, DIONIS, HEISTER, and GARENGEOT.

Knowledge of Languages neceſſary to a *Surgeon*.

XXXI. Since many of the moſt valuable Treatiſes in Surgery have been publiſhed in the learned as well as in the modern Languages, it will eaſily appear of what great Service it will be to the Surgeon, to be well verſed in thoſe Languages, eſpecially the *Latin* and *French*, ſince without this Aſſiſtance, they will reap very little Advantage from the Inventions of others; but whoever is moderately verſed in the *Latin* Tongue, I would adviſe him to procure the academical Theſes upon Chirurgical Subjects which are yearly publiſhed, for the Expence is trifling, and the Advantage that accrues from reading them, is by no means ſo; for they frequently contain many new and uſeful Obſervations, Deſcriptions of Inſtruments and Machines, and new Methods of Cure, that are not to be met with in larger Volumes.

Diviſion of *Surgery* into its Parts.

XXXII. Hitherto we have treated of the Nature and End of *Surgery*, deſcribed the Aids that are neceſſary to it, and related the Fortunes it has met with in different Ages; Order therefore now requires us to proceed to its Diviſion, which is very different according to different Authors. There are many Profeſſors of *Surgery* who divide this Art into ſix Parts, and diſtinguiſh each of them with a *Greek* Name. Theſe are, 1. *Syntheſis*. 2. *Diæreſis*. 3. *Exæreſis*. 4. *Aphæreſis* 5. *Proſtheſis*, and 6. *Diorthoſis*. On the other hand, ſome divide it into five, ſome into four, ſome into three Parts, whilſt others aſſert that it may be comprehended under two of theſe Diviſions. But ſince Perſons ignorant of the *Greek* Language are eaſily puzzled with *Greek* Terms, and beſides that the Diſtinctions are not juſt, as not comprehending all Parts of *Surgery*, it ſeems to be high Time to aboliſh them, as we live in an Age more inquiſitive after Things than Words: more eſpecially as theſe Terms would perplex the Memory of young Students in Surgery, who for the moſt part are unacquainted with the *Greek* Language. Some, laſtly, have been fond of dividing Surgery into five Parts, the firſt treating of Wounds, the ſecond of Ulcers, the third of Fractures, the fourth of Luxations, the fifth of Tumors. Though even this

Method

Method of dividing by no means ſatisfies me, ſince the whole Art cannot be clearly explained, by ſpeaking to each of theſe Heads.

The Author's Diviſion of Surgery.

XXXIII. Wherefore in my Judgment, it is beſt to divide *Surgery* into the three following Parts, by which means the whole Art may be laid down and taught with Clearneſs. The firſt, which is called *Pentateuch* by FABRICIUS AB AQUAPENDENTE, from the Number of Chapters it is compriſed in, treats of the Diſorders that are moſt common to the Human Body, and takes up five Books. 1. *Of Wounds.* 2. *Fractures.* 3. *Luxations.* 4. *Tumors*, and, 5. *Ulcers.* The ſecond Part treats of *Chirurgical Operations* (as they are commonly called), deſcribing at the ſame Time all ſuch Diſorders of the Human Body as are to be relieved by the Aſſiſtance of the Hand, and could not properly be deſcribed in the firſt Part. Laſtly, *Chirurgical Bandages* will be the Subject of the third Part, which we ſhall deſcribe in ſo clear a Manner, that it will be very eaſy to learn not only how each of them is to be made, according to the Nature of the Diſeaſe or of the Limb, but alſo how they are to be applied, to the Benefit of the Patient; for though we find that *Surgeons* have paid very little Regard to the Deſcriptions of Bandages in their Writings, it is nevertheleſs not only extremely uſeful, but abſolutely neceſſary. Sometimes Accidents happen of ſuch a Nature, as Luxations, Fractures, Hæmorrhages, Herniæ, as only to admit of Help by Bandages, and where without ſuch Aſſiſtance the Cure would be extremely doubtful or deſperate; beſides this, by a neat and dextrous Application of a proper Bandage, the Surgeon not only gains the Admiration of the Standers by, but his Patient alſo puts more Confidence in him, which very often forwards the Cure wonderfully.

The Author deſcribes the Method that he intends to follow in Writing

XXXIV. Leſt any one ſhould be ignorant of the Method which I intend to obſerve in expounding the Chirurgical Doctrines which I am going to lay down, I ſhall give a brief Deſcription of it in this Place: that thoſe who are deſirous of acquiring a thorough Knowledge of *Surgery* may not be diſappointed. I ſhall not, according to the Cuſtom of many others, content myſelf with ſolely deſcribing the Inſtruments and Machines that are made uſe of by Surgeons to relieve ſuffering Nature, neglecting at the ſame Time the Hiſtory of Diſeaſes, and the Regulations that are to be obſerved with regard to Diet and Medicine, as if they were Things unneceſſary for the Surgeon to be acquainted with; but, on the contrary, I ſhall uſe the utmoſt Diligence to explain, as clearly as it is poſſible, 1. the proper Nature and Diſpoſition of the Diſorder. 2. what Parts of the Body are liable to be affected by this or that Diſorder. 3. what the peculiar Symptoms of each Diſorder are, and how to form a proper Prognoſtic by them. 4. I ſhall deſcribe the principal Chirurgical Inſtruments which are beſt adapted to each Caſe, of which you will find Copper-plates, for the moſt part of the ſame Size with the Inſtruments which they repreſent. 5. I ſhall not only ſhew the beſt Method of performing all Operations in Surgery; but, 6. in what Manner the Patient is to be treated after the Operation, ſo as to recover his Health in the moſt ſpeedy, ſafe, and pleaſant Manner; and this not only with regard to the Dreſſing and Bandages which are to be applied to the Part, but alſo with reſpect to the Medicines which are proper to be adminiſtred, and the Rules which are to be obſerved as to his Diet.

The Knowledge of Inſtruments is recommended, and a Supply of them premiſed.

XXXV. We declared above, that a Surgeon's Hands would be of little Service to him, if he was not ſupplied with Variety of Inſtruments, which he ought to be very well inſtructed in, that ever hopes to arrive at a proper Uſe of them in the Cure of Diſeaſes. Therefore that we may the more readily form our Surgeon, it will be well worth our while to treat briefly of the neceſſary Apparatus of Inſtruments which he is to be furniſhed with, before we are ſolicitous about teaching him the Manner in which they are to be uſed. I cannot deny but that there are a great Number of Chirurgical Inſtruments to be found in Chirurgical Authors; but, at the ſame time, I can with Truth affirm, that many of them are obſolete and uſeleſs, and many of excellent Uſe have been omitted (eſpecially at the Time when I firſt publiſhed my Book of Surgery in the *German* Language in the Year 1718). therefore it ſeems neceſſary to publiſh a Deſcription, not only of the moſt modern Chirurgical Inſtruments, but of thoſe beſt adapted to Uſe, keeping up to their proper Size as much as poſſible in the Plates. Whether our Plates have ſatisfied this End or not, let others judge: this I am certain of, that I have made it my Study to ſave Students in Surgery the Labour of having Recourſe to many Volumes to ſearch after proper Inſtruments, and to exhibit to their View all the beſt and moſt general Inſtruments, in one Book; and in ſome Places they will find Copies of Inſtruments which are not to be found in other Authors. GARENGEOT publiſhed a Book in *French* on Chirurgical Inſtruments, in which he exhibited many new and correct ones, but delineated in too ſmall a Size, which eaſily led Surgeons and Workmen, who endeavoured to imitate them, into Errors; the chief of theſe I have copied into this Book, and, wherever my Page would admit of it, I have given you the true Dimenſions of the Inſtruments, in order to render them more uſeful. But as it is of much more Service to examine the Inſtruments themſelves than the Plates of them, therefore a Surgeon ought to neglect no Opportunities of examining and contemplating upon the beſt he can lay his Hands on, and eſpecially the neweſt invented. For my own part, when I read Chirurgical Lectures, I always ſhew my Pupils all Kinds of Inſtruments that are uſed in Surgery, and point out the Defects of the Ancients, and the Improvements of the Moderns.

Pocket Inſtruments deſcribed.

XXXVI. But in the firſt place, as they are more immediately neceſſary, and are in conſtant Uſe, I ſhall deſcribe the Inſtruments in their true Dimenſions, which a Surgeon ought always to carry about him in a proper Caſe, and are therefore called *Pocket Inſtruments*. To this place belong thoſe Inſtruments in particular, which are deſcribed in Plate I, under the Letters A and B; *two Lancets* of different Sizes. Theſe are uſed, eſpecially the ſmaller Sort, in opening Veins, for which Reaſon the *Greeks* called them *Phlebotoma*; but the larger Sort are uſed to open Abceſſes with, and are therefore called by the *French Lancettes à l' Abſcé*. The Letter C ſhews a Pair of *ſtraight Sciſſars*, fit for many Uſes; the Surgeon ſhould have ſeveral Pair of theſe at home, of different ſizes, as they are neceſſary in different Diſorders. D, a Pair of *crooked Sciſſars*, proper to be uſed in dividing *Fiſtulæ*, and in many other Caſes. E, a Pair of *Forceps* furniſhed with Teeth at one End; theſe are uſed to remove Dreſſings, and ſometimes to extract Splinters or Thorns; they are alſo ſerviceable to the Surgeon in his Anatomical Exerciſes. *Forceps* of this Kind are commonly made of Steel, but thoſe of Silver are much neater. F, a *Razor*; G, a *ſtraight Inciſion*





A
B
P
X

Plate I.




D
F
G
H
I
K
L
M
N
O
P
S
T
V
X

Knife, by the *French* called *Biſtouri*; H, a *crooked Inciſion Knife*; I, a *ſtraight double-edged Inciſion Knife*; K, a *Probe*, which the *French* call *Une Sonde*; one End of which is broad and thin, for diſcovering a Fiſſure in the *Cranium*, and other Uſes; the other End is rounded with a Knob, to examine the Depth and Situation of Wounds and Ulcers; for which Uſes alſo the Probe at Letter L may ſerve. The neateſt Probes are made of Silver, though they are frequently alſo made of Steel, Ivory, or Whalebone. M, repreſents a *grooved Probe* or *Director*, to direct the Edge of the Knife or Sciſſars in opening *Sinus's* or *Fiſtulæ*, that by this means the ſubjacent Veſſels, Nerves and Tendons may remain unhurt; the Ornament at the upper Part of it is for a Handle, though ſometimes that End is made in the Form of a Spoon, as you may ſee in the Figure at N, to contain a Powder to ſprinkle upon Wounds or Ulcers; ſometimes alſo it is forked at the End to divide the *Frænum* of the Tongue, as at the Letter O. Nor muſt we here omit the *Spatula*, as deſcribed at the Letter P. The Uſe of this Inſtrument is to depreſs the Tongue, in order to examine the State of the Tonſils, Uvula, and Fauces, when they are affected with any Diſorders; it is alſo uſed to ſuſpend the Tongue, when the *Frænum* is to be divided; for which Purpoſe it has a Fiſſure at its Extremity, and ſhould therefore be rather made of Silver than of any other Metal. The following *Spatulæ* alſo, at Q and R, ſomewhat reſemble this. Theſe are chiefly uſed in ſpreading Plaſters, Ointments and Cataplaſms, ſometimes with their ſulcated Extremity they are of Service in raiſing up fractured Bones of the *Cranium*. Here likewiſe, in the laſt Place, we muſt remember different Sorts of *Needles*, *ſtraight* and *crooked*, for ſtitching up of Wounds, taking up of Arteries, and many other Uſes; I have given you crooked ones of different Sizes at the Letters S, T, V, X.

What Medicines a Surgeon ought to be furniſhed with.

XXXVII. What I have ſaid concerning the Inſtruments that are immediately neceſſary for a Surgeon to be provided with, is ſufficient; I ſhall proceed now to deſcribe other Things, with which he is equally obliged to be furniſhed as certain Medicines; ſuch as *Unguentum digeſtivum commune*, *Unguentum ægyptiacum*, *aut fuſcum Wurtzii*; for cleanſing or digeſting foul Ulcers: and ſome vulnerary Balſam, as the *Linimentum Arcæi*, *Balſamum Samaritanum*, *Peruvianum*, *Capyvæ*, *de Mechâ*, *Unguentum Baſilicum*, *Oleum Terebinth.* or *Balſamum Sulph. Terebinthinatum*, *&c.* To theſe muſt be added a Plaſter or two, as *Emplaſtrum Diapalmæ*, or *ſtypticum Crollii*, ſince they will almoſt always be required. Neither ſhould a Surgeon ever be unfurniſhed with a Piece of *Vitriolum Romanum*, to take down luxuriant Fleſh, and ſtop Hæmorrhages; but if you are without Vitriol, its corroſive Intention will be anſwered by *Alumen uſtum*, *Mercurius præcipitatus ruber*, or *Lapis infernalis*, or any other corroſive Medicine, which will alſo ſerve to make Iſſues or open Abceſſes, or to perform any Work of that Kind. But the Surgeon ſhould always have in Readineſs a certain Quantity of ſcrap'd Lint, that he may be able to give immediate Aſſiſtance to wounded Perſons; ſince, if he is unprepared, they may be eaſily taken off with an Hæmorrhage, which Circumſtance ought alſo to prevail ſtrongly with a Surgeon, never to be entirely unprovided with *Bandages*.

XXXVIII.

Neceſſary Qualifications for a Surgeon.

XXXVIII. We obſerved above (N. 2.) that it was the Buſineſs of an able Surgeon readily to apply a Remedy to the Diſorders of the human Body by manual Operations with the Aſſiſtance of Inſtruments. We have now deſcribed the principal Inſtruments as well as Medicines with which a Surgeon muſt of Neceſſity be provided. It remains therefore to examine into the Qualifications he ought to be Maſter of to render him accompliſhed in his Profeſſion. The Agility of Body and Reſolution of Mind that are neceſſary to a Surgeon, are elegantly deſcribed by CELSUS: "[a] A Surgeon (ſays he) ought to be in his full "Vigour, to have a ſtrong, ſteady Hand, never given to tremble, and, to be "as ready with his left-hand as his right; to have a quick, clear Sight, an "intrepid Mind, void of all Tenderneſs, ſo as not to be at all moved by the "Outcries of his Patient; to uſe no more Haſte than the Caſe requires; nor "to cut leſs than is neceſſary; but he ſhould act in all Reſpects as if he was en-"tirely unaffected by his Patient's Complaints." But at the ſame Time I would have him behave with ſuch Caution as not to proceed raſhly or cruelly, and particularly avoid giving unneceſſary Pain.

Agility of Body. Reſolution of Mind.

Skill in Phyſic and Anatomy.

XXXIX. The two Qualifications that I have juſt recited, are by no means ſufficient of themſelves to compleat the Surgeon; but there are others alſo which CELSUS has omitted, which are highly uſeful, and conſequently neceſſary. No one will excel in Surgery, unleſs he is firſt furniſhed with a good natural Genius, to which he muſt join a well-grounded Knowledge in Anatomy and Medicine; if he is furniſhed with theſe Gifts, he will not only with great Sagacity judge of the Cauſes and Circumſtances of the Diſorders upon which he is conſulted, but will with great Readineſs make uſe of the beſt Methods, both with regard to the Adminiſtration of Medicines, and the Choice of proper Inſtruments for their Relief; or, if Occaſion require, invent new ones, and apply them with Succeſs: whilſt, on the contrary, they who are not Maſters of theſe Qualifications, will daily be guilty of capital Errors.

Frequent Exerciſe in Hoſpitals.

XL. When the ſolid Foundations for Surgery are laid and the Qualifications attained, which we have here recommended, our Student muſt by no means omit a proper Attendance upon the Lectures of Profeſſors, and a due Diligence in reading Chirurgical Authors. Thoſe therefore who deſire a thorough Knowledge in Surgery, are not ſatisfied with viſiting Caſes that may accidentally occur to them in their private Practice, but diligently frequent all the Hoſpitals they can get Admittance to, and by this means they ſee more in one Year, than they could otherwiſe do perhaps in the whole Courſe of their Lives. But in order to make the greater Proficiency in theſe Schools of Surgery, it will be worth while to diſtinguiſh the different Kinds of Diſorders that fall under your Inſpection, after what Method, and with what Succeſs they are treated by Maſters of the greateſt Experience. Being prepared by repeated Obſervations of this kind, aſſiſted by the Advice of Maſters, you may at length try your Hand, at firſt upon dead Bodies, and afterwards, when you have Opportunity, upon diſeaſed Perſons; for this trite Saying will always have its Force: *The Artiſt is not made by reading, meditating, or diſputing, but by Practice.*

[a] Vid. Lib. VII. Præfat.

XLI. Laſtly

XLI. Laſtly, that the Surgeon may not appear diſagreeable or terrible to his Patients, eſpecially if they are Perſons of Diſtinction or Quality, he ſhould diligently avoid the Appearance of Roughneſs in his Behaviour, or Naſtineſs in his Dreſs: for good Breeding and Cleanlineſs have their proper Effect in all Parts of Life; but the Surgeon gains a particular Confidence with his Patient by his Addreſs, which has no ſmall Share in the Succeſs of his Endeavours. Good Manners and Cleanlineſs.

XLII. The Surgeon being endued with theſe Principles and Qualifications, may ſafely apply himſelf to the Practice of his Profeſſion; but that he may ſucceed the better in the Execution of it, it is proper he ſhould be acquainted with what is his Duty in every Step of it. As ſoon as ever he is introduced to his Patient, he ought in the firſt place (as HIPPOCRATES well adviſes) to enquire of him, or his Friends or Domeſticks, what ails him? where is the Seat of his Complaint? from what Cauſe it proceeds? and how long it has been upon him? If there is no particular Objection, he ſhould examine the Part himſelf, and diligently weigh all that he has heard or ſeen that may give him any Light into the Caſe, that he may come at a thorough Knowledge of the Nature of the Diſorder. The Surgeon's Duties to his Patient. Firſt, he is to examine the Caſe.

XLIII. Having finiſhed his Examination the next Thing to be done is to conſider under what Claſs of Diſorders it is to be ranked, and whether it be curable or not? If it is deemed curable, whether it will be a Caſe of Time and Difficulty or not? whether it is curable by Medicines alone? or whether the Aſſiſtance of the Knife be neceſſary? for the ſafeſt and moſt gentle Methods, as CICERO with great Propriety obſerves, muſt always be preferred to harſh and dangerous ones, and are always to be tried firſt, that the Patient may not ſuffer unneceſſary Tortures, nor his Life be hazarded through the Raſhneſs of the Surgeon; but to Diſorders of a violent Nature, dangerous and even doubtful Remedies are to be applied: agreeable to the ſaying of HIPPOCRATES (*Aph.* 6. Sect. 8.), "Where Medicines fail, Inſtruments ſucceed." They are to be highly condemned, therefore, who, after the Methods of [a] Mountebanks, condemn their Patient who labour under *Herniæ*, without regard to Age or Habit of Body, to the Operation of the Knife, when far the greater Part of them might be cured by a ſafer and eaſier Method. But if you ſhall find it impoſſible to ſave your Patient by gentle Methods, you ſhould declare the Danger to the Patient, or rather to thoſe about him, leſt, if the Diſorder ſhould get the better of your Art, you ſhould be ſuſpected of Ignorance, or perhaps, of Knavery. Whether curable or not, and by what means.

XLIV. If the Surgeon ſhall find the Diſorder to be curable, but to be of ſuch a Nature as to require the Knife, he ſhould declare this in due Time to the Patient, and ſhould have his Approbation or Conſent before he undertakes it; for a Surgeon is not only to take care to ſtop the Fury of the Diſeaſe, and allay preſent Pain, but alſo to provide againſt Accidents that ariſe from Delay; ſuch as may aggravate and enhance the Diſtemper, and at laſt render the Caſe incurable. In very difficult Caſes, where Danger conſiſts not ſo much He ſhould undertake the Cure with great Circumſpection.

[a] I ſaw an Inſtance of this in a Mountebank, who undertook the Cure of a Boy of about ſix Years of Age, for a *Hernia*, and not only performed the Operation, but caſtrated him: when I aſked him in private, why he uſed this hazardous Method without trying a Truſs, ſince his tender Age would eaſily have admitted of it, he ingenuouſly confeſſed he did it for Profit, for he would have been paid but a Crown for the Truſs, whereas the Operation brought him Ten, if not Twenty.

in Delay as in the Manner of Treatment, the Surgeon not only provides for his Patient's Good, but his own, if he calls in other ſkilful Practitioners as well in Phyſic as Surgery, with whom he may conſult maturely and deliberately before he proceeds to any Operation; for by this means he will ſave himſelf from all Blame of having proceeded raſhly or ignorantly, eſpecially when he is concerned for Perſons of Diſtinction; if Things ſhould go otherwiſe than he could wiſh, he will avoid the Charge of having loſt a Patient through his Indiſcretion, whom perhaps no Art could have ſaved: which very Reaſon ſhould always induce a prudent Surgeon, in Caſes attended with Difficulty and Danger, to deſire the Aſſiſtance, if it can be procured, of the ableſt and moſt experienced of his Fraternity.

He ſhould be furniſhed with proper Inſtruments before he begins.

XLV. Having proceeded ſo far, with the Cautions that I have adviſed, every thing ſhould now be carefully provided which is neceſſary for Inciſion, Dreſſing, or any other Action, before the Operation be entered upon; but this Apparatus of Inſtruments and Dreſſings ſhould never be got ready in your Patient's Chamber, or in his Sight, leſt they ſhould ſtrike him with a ſudden Fear, and bring on fainting Fits and other Accidents, which would very much diſturb the Operation. For the ſame Reaſon a Crowd of uſeleſs Spectators ſhould never be admitted into the Room, becauſe, beſides the Diſturbance that they create to the Patient, it is to be feared they will very much annoy the Operator, by intercepting the Light, and filling up the Room: beſides, ſhould any one rudely preſs upon him whilſt he is performing any nice Operation, it might be of the utmoſt ill Conſequence.

He ſhould encourage his Patient.

XLVI. When the Surgeon is entering upon the Operation, he ought to uſe his utmoſt Endeavours to encourage the Patient, by promiſing him in the ſofteſt Terms to treat him tenderly, and to finiſh with the utmoſt Expedition; and indeed he ſhould uſe Expedition but not Hurry, and ſhould be very careful to give no unneceſſary Pain, but at the ſame Time to leave no Miſchief unremedied if he obſerves theſe Rules, he will be ſure to gain Credit with the Standers by.

After the Operation the Wound is to be dreſſed.

XLVII. The Operation being now over, the Surgeon is to conſider what remains to be done towards ſupporting his Patient, and confirming the Cure; the Hæmorrhage occaſioned by it is to be ſtopped, the Wound to be dreſſed, the wounded Part is to be placed in the moſt convenient and eaſy Situation; and it is now Time not only to think of preventing any new Diſorder falling upon the Part, but to uſe all Endeavours for reſtoring Health itſelf.

Proper Diet is to be adviſed.

XLVIII. It is his Duty now to preſcribe a proper Regimen for his Patient's Diet to provide him a commodious Apartment in a healthy Air, to encourage him to reſt, and to avoid all Paſſions, and Reflections upon any Thing that may diſturb his Mind; and if any freſh Inciſion, or other Operation, be neceſſary, he ſhould be adviſed readily to ſubmit to it. Every Thing ſhould be carefully avoided that may ruffle the Patient, for Diſturbances of the Mind are great Enemies to the Health of the Body.

Impertinent Viſits ſhould be prevented.

XLIX. Frequent and impertinent Viſits to the Sick from his Friends or others, ſhould be carefully prevented, for they will undoubtedly fatigue and diſturb him; but we don't mean by this to cut him off from all Converſe with Mankind; a little chearful Company now and then would rather give him Eaſe, and make him forget his Pains; but I had much rather he ſhould divert himſelf by attending to others, than by ſpeaking himſelf.

L. CELSUS declared Physic to be a conjectural Art; these Conjectures therefore must be made with the utmost caution, and the Surgeon also should use the same caution in delivering his Prognostic, when he is called upon, and not, like bold Quacks, promise all will go well, whether the Case is curable or not. For should the Case turn out contrary to your Prognostic, you will either be accused of Knavery or Folly: So if we listen to Reason and [a] CELSUS, it is the Part of a Mountebank to aggrandize a small Performance: an honest Surgeon will always be very careful to avoid both Extremes; it is the Part of a prudent Man to declare from his Conscience what he takes to be the true State of his Patient's Case; whether he believes it to be curable or incurable; what Hopes he entertains of his Recovery; and it is particularly incumbent on him to take the utmost care, that he encrease not a Disorder, which appears slight in the beginning, by treating it negligently. In doubtful Cases, where there is reason for great Fear, but not for certain Despair, he should declare his Reasons both for Hope and Fear; but where the Case is extremely dangerous, he should do it to the Relations. Sometimes it is better not to be concerned with a Patient, when it is impossible to be of any service to him, lest you should be said to have killed him, who died by his Disease [b]: But where you are concerned, let the Case be ever so desperate, it is always the Duty of a prudent Surgeon, to cherish the Patient with sweet Words, and give him Hopes of his Recovery; for some Disorders are very much aggravated by Fear, whereas the Expectation of Health and Ease is always so comfortable, that, though it will not cure a Disease, it will at least make it easier to be born.

Great Caution is to be used in prognosticating.

LI. We have already declared what are the principal Duties of a Surgeon; but since the *First*, which is strictly to examine the Case, and the *Sixth*, which concerns the dressing of the Wound, are more immediately necessary, we shall more largely explain what Methods are to be observed both in examining and dressing Wounds. In examining and discovering dangerous and difficult Disorders, the Surgeon requires many Assistances; at first his *Eyes* are necessary to him, by the use of which he will distinguish Wounds, Ulcers, Tumours, Fractures, Cataracts, and most Disorders of the Eye, and a thousand others; but if the Case is of such a Nature that it escapes the Sight, or is not wholly discoverable by it, the Hands are to be called in aid. This happens frequently in Fractures, Luxations, Abscesses, Herniæ, *&c.* *Instruments* also are sometimes required in this place, especially *Probes*, in discovering the Situation of Wounds, Ulcers, Fistulæ, Fractures of the Skull, Stone in the Bladder, and the like. The *Ears* also are required to give their report of some Disorders; Fractures of the Bones are frequently discovered by the noise which their Extremities make when they are rubbed together; the Sense of Hearing is of so eminent Service in discovering of Stones in the Bladder, that unless the Extremity of the Catheter is heard to strike against the Stone, we are never sufficiently justified in determining a Stone to be there. Some Disorders are discovered by the *Smell*. By the benefit of this Sense we discover the State of Malignity of an Ulcer; and in difficult Births, the *Fœtus* is discovered to be dead by the great Stench that proceeds from the Womb, and this is the only Method we have of being certain in this case. We are assisted also by this Sense in acquiring an

The Senses and proper Instruments of Service in examining Disorders.

[a] Lib. V. Cap. 26. [b] Ibidem.

eaſier Knowledge of a Caries of the Bones, an ulcerated Cancer, and Diſorders of this ſort, which carry with them a peculiar Smell. If, therefore, the Surgeon has theſe Faculties in Perfection, Seeing, Feeling, Hearing, and Smelling; with the proper Exerciſe of theſe, and the Aſſiſtance of Inſtruments, he will ſeldom be at a loſs in diſcovering the Diſorder.

And Reaſon itſelf.

LII. But Caſes in Surgery ſometimes happen, where the external Senſes, aſſiſted by Inſtruments, will by no means yield ſufficient Light to their Diſcovery; but *Reaſon* and *Judgment* are alſo required: the true Nature of a Diſeaſe is traced by reaſoning upon its various Symptoms. HIPPOCRATES, the common Parent of Phyſic, ſeems to have regarded this, when he ſaid [a], whatever eſcapes the Reach of our external Sight, ſhould be ſearched for and overtaken by the Eyes of the Mind. So when any one has had a violent *Concuſſion of the Brain*, from a Fall or a Blow, without receiving any external Hurt, he will lay ſenſeleſs, as if he were in a profound Sleep; Reaſon in this caſe will eaſily inform us, that there is an Extravaſation of Blood in the Cavity of the Cranium, and that proper Methods muſt inſtantly be uſed to make a Paſſage for it externally. Our Reaſon is of equal Service to us in an *Empyema*; for tho' in this caſe Matter is formed in the Cavity of the Thorax, from a previous Inflammation of ſome of its Contents, yet we ſhall meet with great Difficulty in diſcovering this to be the caſe, by our external Senſes; but by comparing the preſent Symptoms with the Diſorder that was previous to them, we find it neceſſary to treat the Caſe as an *Empyema*; and of this kind there are many inſtances.

Of the neceſſary Apparatus for Dreſſing.

LIII. We are next to treat of what principally belongs to the Method of dreſſing the diſordered Parts. In this place we are firſt to ſpeak of [b] *Lint*, which is the Scrapings of fine Linen, by the *French* called *Charpie*. This may be made into various Forms, which acquire a different Name, according to the difference of their Figure; thoſe that approach neareſt to an oval or orbicular Form are called *Pledgits*, (by the *French Plumaceau*) ſee *Table* II. *Letters* A *and* B. Lint made into a Cylindrical Form, or reſembling the Shape of Dates or Olive Stones, is called a *Doſſil* (in *French*, *Bourdenets*); their ſize is very different as appears *from the Figures at* C D E. Sometimes they are ſecured by a Thread tied round their Middle, as it is expreſſed *by the Figures at the Letters* F G. It requires a good deal of Time and Experience, to require a proper Expertneſs in making up theſe Forms.

Uſes of ſcraped Lint.

LIV. Theſe different Forms of ſcraped Lint, eſpecially the cylindrical, are required for many Purpoſes; for they are applied, 1ſt, *To ſtop Blood in freſh Wounds*, by filling them up with dry Lint before you apply the Bandage; but if you have not ſcraped Lint at hand, you may tear a fine piece of Linen into ſmall Rags, and apply it in the ſame manner, and perhaps with a better Effect; but in very large Hæmorrhages they ſhould firſt be dipt in ſome Styptic Liquor, Alcohol, or Oil of Turpentine; or ſprinkled with a Styptic Powder; but of this we ſhall preſently treat more largely. 2dly, *To agglutinate and heal Wounds*, to which end ſcraped Lint is very ſerviceable; if it is ſpread with ſome digeſtive Ointment or Balſam, or dipt in ſome vulnerary Liquor, they alſo yield us great Aſſiſtance. They are often ſerviceable, 3dly, *In drying up Wounds and Ulcers*, and forwarding the Formation of the Cicatrix. They are uſed

[a] In Lib. de Arte. [b] CELSUS, Lib. V. Cap. 26. Num. 21.

alſo with Succeſs, 4thly, *In keeping the Lips of Wounds at a proper Diſtance*, that they may not haſtily unite, before the Bottom is well digeſted and healed. 5thly, and laſtly, They are highly neceſſary *to preſerve Wounds from the Injuries of the Air.* The ſmall portions of Lint that are tied round with a Thread (*ſee Tab.* II. *Letter* F *and* G) are chiefly uſed in dreſſing Wounds and Ulcers that are of the deeper kind, and are always applied to the Bottom of ſuch Wounds, the remaining Cavity being filled up with other Portions of Lint, not ſupplied with a Thread, and by this means we do not only provide for the immediate Removal of theſe Dreſſings, when we ſhall think it neceſſary, but at the ſame time prevent a Poſſibility of leaving any Part of them concealed in the Bottom of the Wound. In very large Wounds, and eſpecially in Amputations of the larger Limbs, which Operations are frequently required in the Army and Navy, at times when Lint is very ſcarce, it will be ſufficient to dreſs the bare Bone and Face of the Wound with ſcraped Lint, filling up the Cavity with *Tow*, covering all with a large *Compreſs*; Figures of which you will ſee at the *Letters* H *and* I, *Plate* II. The Surgeons in former Ages formed Compreſſes of Sponge, Feathers, Wool, or Cotton, Linen being a ſcarce Commodity with them; but Lint is far preferable to all theſe, and is at preſent univerſally uſed; excepting, that in Wounds of the Thorax or Abdomen, the uſe of a Sponge may ſometimes be neceſſary to ſuck up the Blood ſpilt in thoſe Cavities.

LV. Beſides the different Forms of Lint that we have deſcribed, there remains another, which is ſometimes uſed in dreſſing of Wounds, called *Tents*, made of Lint worked into the ſhape of a Nail, with a broad flat Head; they differ in Thickneſs and Length according to the Size of the Wound for which they are intended, as appears by the Figures *in Plate* II, *at the Letters* KLMN. Theſe Tents are chiefly uſed in deep Wounds and Ulcers. They are of Service, 1. Not only in conveying Medicines to the inmoſt Receſſes and Sinuſes of the Wound; but, 2. To prevent the Lips of the Wound from uniting before it is healed from the Bottom; to which we may add, 3. That by their Aſſiſtance grumous Blood, Sordes, *&c.* are readily evacuated. They are to be made extremely ſoft, that the Cure of the Wound may not be retarded by the Pain they would otherwiſe bring on: but that the Wound may not be kept open too long, I would adviſe the Surgeon, as ſoon as he has cleanſed the Part ſufficiently, and finds the Sinuſes heal up, to leſſen the ſize of his Tents by degrees, and, as ſoon as he can conveniently, entirely lay them aſide. I am not at all ſurprized, that many Surgeons of good Name (amongſt which are CÆSAR MAGATUS, BELLOSTE, and others) have entirely forbid the uſe of Tents; ſince to beſure it proceeded from a total neglect of this caution in their uſe, amongſt too many of their Brethren.

Of Tents compoſed of Lint.

LVI. But there is another kind of Tents, differing from that which we juſt now deſcribed, made of Linen Rags, not ſcraped, worked up into a Conical Form, to the Baſis of which is faſtened a ſtrong Thread; the Apex of it muſt be a little unravelled to make it ſofter, that it may not become painful. The Thread is faſtened to the Baſis that it may be recovered with the greater Eaſe, if by any Accident it ſhould be forced into the Cavity of the Thorax or Abdomen (*See Plate* II, *Fig.* O.); for it is to be obſerved here, that the Tents we now deſcribe are chiefly uſed to keep open Wounds that penetrate into the

Of Tents made of Linen Rags.

Cavity of the Thorax or Abdomen, in order to make way for the proper Diſcharge of Blood, Matter, *&c.*

Of Spongy Tents.

LVII. A third ſort of Tents remains to be deſcribed, whoſe principal Office is, not only to keep open, but to enlarge by degrees the Mouth of any Wound or Ulcer, which ſhall be thought too ſtrait, that by this Means a freer Paſſage may be procured for the Blood and Matter that was confined, and that proper Medicines may find a more ready Admittance. Theſe Tents are made either *of Sponge*, prepared in a certain Manner, or of dried *Roots of Gentian*, *Calamus Aromaticus*, *&c.* for theſe kind of things imbibe the Matter that flows to them, and being preſently enlarged, dilate the Lips of the Wound. Not much unlike Tents, are the ſmall *Silver or Leaden Tubes*, which are frequently uſed to draw off Blood, Matter or Water from Wounds or Ulcers of the ſmalleſt Orifices; ſometimes in extracting Water from dropſical Patients, or evacuating the Urine in a Puncture of the Bladder: They are made of all Sizes and Shapes, as you may ſee in *Plate* II, *at the Letters* P Q R S T V X. What farther concerns the Uſe of theſe Tubes, you will ſee more largely treated of, when we ſhall deſcribe the Diſorders that more immediately call for their Aſſiſtance.

Of Plaſters.

LVIII. Your Apparatus for Dreſſings will be very deficient if you are not furniſhed with *Plaſters*. The meaning of the Term is ſo well known, that I ſhould appear ridiculous if I went about to explain it. But there are different Kinds of Plaſters without number; the principal of theſe, and the manner of making them, may be learnt from various Books, as particularly, the *Auguſtan Diſpenſatory*, the *London*, the *Pruſſian*, and that of *Lemery*. Theſe Plaſters are ſpread upon Linen or Leather, according to the different Circumſtances of the Wound, Place, or Patient. If the Part upon which the Plaſter is to be laid is naturally hairy, it muſt be ſhaved; that it may ſtick the cloſer, and be removed without pain to the Patient: But for the better Application to different Parts of the Body, the natural Shape of the Part muſt be conſulted, and the Plaſter formed accordingly: Therefore ſome Plaſters aſſume a Round, Square, Triangular, Elliptical, or Lunar Form; others the Shape of the Letter T, *&c.* as will clearly appear at *Plate* II, *Numb.* 1, 2, 3, 4, 5, 6, 7, 8. Others there are which are divided at one or both Ends, *See Numb.* 9, *and* 10. To theſe we may add thoſe kind of Plaſters which are perforated near the Middle, ſome with a ſingle, ſome a double Perforation, which are of frequent Uſe in Fractures attended with a Wound; for by this Contrivance the Wound may be cleanſed and dreſſed without removing the Plaſter, *See Number* 11. Such Plaſters are uſed too, eſpecially thoſe with the ſingle Perforation, in making Iſſues or removing Warts and other Excreſcencies, by corroſive Medicines.

The Size and Uſe of Plaſters.

LIX. The *Size*, as well as Form of Plaſters, is very various, ſince it muſt always correſpond with the Part which is bruiſed or wounded. Their *Uſe* alſo is of great Advantage in defending Wounds and Ulcers from the external Air, or from any Filth which they might otherwiſe contract; for they are not only ſerviceable in ſecuring the Dreſſings, but they alſo forward the Maturation of the *Pus*, agglutinate and heal Wounds, unite broken Bones, heal Burns, aſſuage Pain; and, laſtly, ſtrengthen the weaker Parts.

LX. It

LX. It is frequently the Cuſtom, after the Plaſter and other Dreſſings are applied, to cover all with a *Compreſs*, which is made of the ſofteſt old Linen, four, ſix, or eight times doubled, without Seam or Hem; theſe are of ſervice, not only by preſerving the Parts more ſafe from the Injuries of the external Air, but alſo for the better ſecuring and fixing the Plaſters and other Dreſſings. Compreſſes are alſo frequently applied, where no Plaſter is made uſe of, and that, ſometimes dry, ſometimes wetted with certain Liquors, which are ſuppoſed to be ſtrengthening, reſolving, lenient, emollient, or cooling; they are frequently dipped in Decoctions of certain Herbs, into Wine, Spirit of Wine, Water, Vinegar, or Oxycrate, and ſometimes into Lime Water; and theſe are either adminiſtred cold or hot, as the Circumſtances of the Caſe ſhall require. The antient Phyſicians called them *Splenia*, from their Shape, frequently reſembling the Spleen. Of Compreſſes.

LXI. When you come to enquire after the *Figure* and *Size* of *Compreſſes*, you will find as great variety as you did amongſt Plaſters; many of them are *Square*, (*See Plate* II. *N.* 12.) others are *Oblong*, and not unlike the Spleen, *N.* 13) again, others *Triangular*, (*N.* 14.) others reſemble the Form of a *Croſs*, (*N.* 15. according to their Situation, ſome are called *Streight*, others *Oblique*, others *Tranſverſe*, others *Annular*, as when they ſurround the Arm, or Foot. There are others again in the Form of an *Aſteriſm*, (*N.* 16.) ſome are divided either on one or on both Sides, as far as the Middle, (*N.* 17, 18.) ſometimes they form a *Hexagon*, (*N.* 19.) or are *Round*, or *Globular*, reſembling a Ball; theſe are uſed in Luxations of the *Os Humeri*, and are placed under the *Axillæ*, (*N.* 20.) ſometimes Compreſſes of a much ſmaller Size are required, which are either *Square*, (*N.* 21.) and are uſed in Wounds of the Blood-veſſels, to reſtrain Hæmorrhages; or *Taper*, (*N.* 22) when they are called for in Sutures of Wounds, or in Ligatures of the Arteries. But all of them, of what Shape ſoever, ſhould be ſomething larger than the Plaſters they are deſigned to cover. The Shape and Size of Compreſſes.

LXII. Compreſſes of all Kinds are intended for theſe Purpoſes; 1. To preſerve and cheriſh the natural Heat of the Body. 2. To ſecure the Dreſſings that lie under them. 3. To convey liquid Remedies to Parts wounded, or otherwiſe diſordered, and to prolong the Uſe of them. 4. To fill up any Cavities or Depreſſions of the Parts, that the Dreſſings (eſpecially in Fractures) may be applied with greater Security. 5. To prevent Bandages from bringing on a troubleſome Itching, or other Pain or Uneaſineſs upon the Skin. And laſtly, 6. to ſtop Hæmorrhages. Uſe of Compreſſes.

LXIII. But it is now high Time to ſpeak of *Bandages*, ſince they are ſo neceſſary a Part of the *Apparatus* in dreſſing and binding up of Wounds. They are not only of greater Service than Compreſſes and Plaſters in ſecuring the other Dreſſings, but are alſo of excellent Uſe in reſtraining dangerous Hæmorrhages, and in joining fractured or diſlocated Bones. Though I have ſet aſide the third and laſt Part of this Work purely for the Deſcription of Bandages, where you will find them more fully and accurately treated, I thought it nevertheleſs neceſſary to touch ſlightly upon theſe things that are principally neceſſary to a Surgeon, by way of Introduction. Of Bandages.

LXIV. Almoſt all Bandages, that are uſed in Dreſſings of Wounds, Ulcers, fractured or diſlocated Bones, ſhould be made of clean Linen Cloth, ſoftened by Wearing, but ſtrong. They ſhould be of a proper Length and Breadth; and, that they may be the ſtronger, examine the Courſe of the Threads, and tear the Cloth. Of what Materials they are to be formed.

Cloth lengthways; Darns, Seams, and large Hems in the Linen ſhould be avoided as much as poſſible, that no Inconvenience may be brought on by the Roughneſs and Irregularity of the Roller. The proper Size of Bandages we ſhall deſcribe more fully below.

Some Sorts of Bandages deſcribed.

LXV. There are different Sorts of *Bandages* for different Uſes. Some are *common*, others *proper*; theſe are only applied to particular Parts, thoſe may be applied to any Part. So we may diſtinguiſhed them into *ſimple* and *compound*; the *ſimple* are thoſe that are formed of one intire Piece of Linen, the *compound* of ſeveral Pieces of Linen ſewed together in different Manners. The moſt ſimple of all is not rolled up, and is the Bandage uſed in Phlebotomy, *See Letter a, Plate* II. That at *Lett. b*, ſeems next to this, which is rolled up at one End, and is from thence called the *ſingle-headed Bandage*, as thoſe are called *double-headed* which are rolled up at both Ends, *See Plate* II, *Letter c*. Next to theſe come other Bandages which are made out of one Piece of Linen, but divided at both Ends almoſt as far as the Middle. *See Plate* II, *Lett. d*. Theſe are called by the Surgeons *four-headed Bandages*. The Bandage at *Letter e* is ſomewhat ſhorter and narrower, and is divided at one End, and perforated at the other; this is generally uſed in Dreſſings that are applied to the Penis, or one of the Fingers. The *Letter f* deſcribes a double headed Bandage, divided about the Middle, which is called the *uniting Bandage*, from its Uſe, for it ſerves to unite Wounds that are made lengthways, without Suture. The *ſcapular Bandage*, which (as appears at *Letter g*) is provided in the Middle with an opening, through which the Head may eaſily be paſſed, the extreme Parts of the Bandage hanging one over the Breaſt, the other over the Back. The chief uſe of this Bandage conſiſts in this, that in dreſſing Wounds of the *Thorax* or *Abdomen*, it is capable of ſupporting another Bandage that is ſomewhat wider, made of a Cloth four or ſix Times doubled, and bound round the Breaſt or Belly; as will appear more clearly from what you will read below.

Of the T Bandage of *Heliodorus*.

LXVI. There remains ſtill to be deſcribed a compound Bandage, made of two Pieces of Cloth, almoſt in the form of the Letter T, as you ſee it deſcribed at *Letter h*; its upper Part is brought round the Belly and faſtened by a Knot, but the lower Part paſſes under the Body between the Thighs, and being brought up again, is faſtened to the upper Part upon the Back. Theſe Bandages plainly appear to be deſigned for the Security of ſuch Dreſſings as ſhall be thought proper to be applied to the *Anus*, or *Parts of Generation*. Some, from the Inventor, call it *Heliodorus's Bandage*; from its Shape it is called the T *Bandage*; and ſometimes, from the Diviſion that is frequently made in the lower part of it, it is called the *double* T.

The Explanation of the Second Plate, *which exhibits thoſe Things which are principally required in Dreſſings, taken chiefly from* Dionis.

1. *Of* Pledgits, Tents *and* Compreſſes.

A and B, *Scraped Lint*, commonly called *Pledgits*, of an orbicular or oval Figure.

C D E, *Doſſils*, which are compoſed of Lint, worked into the Likeneſs of Olives, or Dactyle Stones.

F and G, the ſame, with the Addition of a Thread tied round them.

H and I,





Plate. 2.
Vol. P. 22.
A
F
H
K
L
O
T
c
d
h
16
22
22

Plate 2.




1
2
3
11 *
4
5
6
7
11 * *
8
9
10
11
a
b
c
d
e
f
g
h
12
13
14
15
16
17
18
19
20
21
22
22

H and I, *larger Pledgits* made of Tow.
K L M, repreſent *Tents* of different Sizes made of Lint.
N, ſhews you a very large *Tent*, with a Thread annexed to it.
O, a *Conical Tent*, ſtill larger than the former, made alſo of Lint.
P Q R S T V X, *Tubes* of different Kinds made of Silver or Lead.
Number 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11. different Forms of *Plaſters*.
Num. 12, 13, 14, 15, 16, 17, 18, 19. different Sorts of *Compreſſes*.
Num. 19. three Sorts of *Compreſſes*, reſembling the Form of an Aſteriſm.
Num. 20. Balls of Lint, which are ſometimes uſed as *Compreſſes*.
Num. 21. A ſmall ſquare *Compreſs*.
Num. 22. Several ſmall ſlender *Compreſſes*.

Of Bandages.

c, A ſimple *Bandage*, not rolled up.
b, A *Bandage of one Head*; that is, rolled up at one end.
c, A *double-headed Bandage*, rolled up at both ends.
d, A *four headed Bandage*.
e, A ſmall *Bandage*, particularly intended for the Security of Dreſſings that are applied to one of the Fingers, or the Penis.
f, The *uniting Bandage*, which is perforated in the Middle.
g, The *Scapular Bandage*.
h, *Heliodorus's*, or the T *Bandage*.

The moſt common Bandage for the Head.

LXVII. Though Surgeons have formerly invented different Kinds of Bandages, for every Wound that could be inflicted upon the Head; yet there is but one Form that ſeems neceſſary, and that will anſwer every End that can be propoſed from this kind of Application. This is made in the following Manner: Take a Handkerchief, Napkin, or any ſquare Piece of Linen, double it up in a triangular Form, and apply it, as we frequently do in hot Weather, when we lay aſide the uſual Coverings of the Head, to moderate the exceſſive Heat of the Sun. The Bandage which is ſo much in Uſe amongſt the modern Surgeons, called by the *French*, *Le grand covrechef*, differs very little from this, and is commonly made of a Napkin, or ſome ſoft Piece of Linen, in a ſquare Form. It is doubled in ſuch a Manner, that the lower Part is about four Fingers Breadth wider than the upper; the middle Part of this Cloth is placed ſo upon the Head, that the fore Part may reach almoſt as far as the Eyes, the four Extremities or Corners of it hanging over the Cheeks. The two Corners of the upper or narrower Part are to be tied under the Chin, at the ſame Time the Corners of the lower or wider Part are to be brought towards the back Part of the Head, and tied together, or faſtened with a Needle and Thread. The fore Part that was extended towards the Eyes, is turned back as far as the Crown of the Head; the two Parts that hang over the Neck almoſt to the Shoulders are alſo to be turned back, and faſtened behind the Ears with a Needle and Thread. This kind of Bandage, when it is neatly made, ſticks cloſe to the Head, and is an excellent Contrivance to preſerve it from the Injuries it might receive from cold Air; for which reaſon it is at preſent in great Uſe and Eſteem. You may in ſome Meaſure form an Idea of the Appearance it makes upon the Head by conſulting *Plate* III, *Fig.* 1, *Letter* A.

But

But the Method of applying it must be learnt from some skilful Artist; for it will easily appear, from this one Instance, how difficult it is to describe the Art of applying Bandages, by Words, and how impossible it is to learn it from such Descriptions.

Application of the Scapular Bandage.

LXVIII. *Letter* B, *Plate* III, *Fig.* 1, describes a Bandage which is generally used to secure Compresses and other Dressings that are applied to the Breast or Belly. The manner of preparing this Bandage is described above at *Sect.* 65; therefore in this Place it remains only to shew the most convenient Method of applying it. After the Wound is dressed, take a double Cloth, and wrap it round the Abdomen or Thorax, sewing not only the Ends of the Cloth strongly together, but fastening it also in the same Manner to the Extremities of the Scapular Bandage, to prevent it from slipping down; the Manner in which it is done appears very plainly in *Plate* III, *Fig.* 1, *Lett.* B *and* C.

The Bandage for Phlebotomy.

LXIX. The *Letter* D shews the Bandage or Ligature that is used to Veins of the Arm; E, to those of the Foot; but we shall treat more largely of the Manner of preparing and applying them in the third Part of our Chirurgical Institutions.

Names of the most simple Windings of the Bandage.

LXX. We have this farther to add, concerning simple Bandages; they assume different Names, according to the different Windings that they form in the Manner of applying them: For instance, if a simple Bandage with one Head surrounds an injured Part with one direct Course, it is called *annular*, *orbicular* or *circular*. On the contrary, if the Windings of the Bandage ascend or descend equally in a spiral Manner, they are called *obtuse* or *spiral*; this frequently happens in Fractures, and other Kinds of Disorders, and is of very eminent Service. But when the Limbs which are to be bound in this Manner are of different Thicknesses in different Parts of them, which is the Case of the *Tibiæ*, it requires a good deal of Art to prevent the Windings of the Bandage from hanging loose. The Bandage is to be applied to the *Tarsus*, and to be brought upwards so as to cross the *Malleoli*, rolling it round the *Tibiæ* in a spiral Manner; but when you are come up to the Calves of the Legs, each round of the Roller must be turned in a particular Manner, and tightened according as the Case requires. It is much easier to demonstrate this Manner of turning in the Roller at each Round, than to describe it in Words. Consult in this Place *Plate* III, *Fig.* 1, *Lett.* F. But from what has been said, you will easily conceive the Reason why the Windings of the Bandages that we have been describing, are generally said to be inverted, and by the *French* are called *Renversées*. These Bandages are so managed that the Windings of the Roller are contiguous to each other; but there is another Method of rolling in Use, where the Windings of the Bandage are not so frequent, and keep a greater distance from each other, and are therefore called *creeping Bandages*, in the *French* Schools *Rampans*; an Example of which you may see in the left Arm of the last mentioned Figure, at *Letter* G. These creeping or serpentile Bandages are used to secure Compresses, Fomentations, or Cataplasms upon a diseased Part. Lastly, there are Bandages called by the Surgeons, *Containing*, *Uniting*, *Dividing*, or *Expelling*, from their several Uses and Intentions.

Where the Bandage ought to begin and end.

LXXI. But lest any one should be ignorant of the neatest and most proper Way of applying these Bandages, you are diligently to observe what follows; to wit, when the Arm is to be dressed, the beginning is formed by two or three circular

circular Windings on the Wriſt, aſcending by looſe Spires to the Cubit or Shoulders as the Caſe ſhall require; but when the beginning is to be on the Foot, it is to be formed by three or four circular Windings of the Bandage round the *Tarſus* and *Metatarſus*, then proceeding in a ſerpentine Courſe up to the Knee, or, if the Caſe requires it, up to the Head of the Thigh, and then, as it ſometimes happen, deſcending again. But we ſhould not neglect to mention in this Place, that the beginning of the Bandage is ſometimes applied even to the diſeaſed Part, as in ſeveral Kinds of Fractures; ſometimes near it, above it, or below it, and ſometimes at a great diſtance from it, according to the Diſpoſition of the Wound. On the contrary, the Extremity of the Bandage is ſcarce ever faſtened upon the diſeaſed Part, but rather upon a ſound one, to avoid giving Pain[a]. Upon the whole, we muſt inculcate this general Admonition; that, whatever the Caſe be, whether Wound, Fracture, Luxation, or Amputation, your Bandage be neither too looſe, nor too tight; for looſe Bandages do not confine the Parts enough: and, when they are too tight, they may cauſe not only violent Pains, but Inflammations, Tumors, Gangrene, and Mortification.

Of Cords, Bands, Ligatures and Strings.

LXXII. The neceſſary Apparatus for Dreſſings ſeems by no means complete without *Ligatures*, *Cords*, *Bands* and *Strings*, and theſe of different ſorts, ſome fine, others coarſe, ſtrong, made either of Flax, or Hemp, or Cloth, or Silk, or Horſe-hair, according to the nature of the Diſorder; for theſe things are almoſt conſtantly required. We uſe them to replace, or extend Bones that are broken or diſlocated, to tye the Patients down, in Lithotomy, Amputations, and Operations of that kind; to tye up the Veins in Phlebotomy, to tye up Arteries after Amputations, or in large Wounds; to ſecure the Splints that are applied to Fractures; to tye up the Proceſſes of the Peritoneum with the Spermatic Veſſels in Caſtration; and laſtly, in taking off Warts and other Excreſcences by Ligature, and in all other Operations of this kind, as we ſhall more fully explain below.

The Study of Surgery is very difficult,

LXXIII. What we have already ſaid concerning the Qualifications which every Surgeon ought to be indued with, and of the Inſtruments with which it is neceſſary for him to be furniſhed, is ſufficient for this place, by way of Introduction to the following Work. We may evidently draw this concluſion from the foregoing Diſcourſe, that Surgery is no eaſy Art, but affords a large Field for Enquiry, and is not to be attained without great Aſſiduity and Labour. The Surgeon has not only a vaſt number of Diſorders to encounter, but the means by which every kind of Diſorder is to be ſubdued are almoſt infinite, the particular nature of which muſt be known to the greateſt Exactneſs. But I by no means diſcourage any one from theſe Studies by the Difficulties that I here ſpeak of, for there is nothing, according to the old Proverb, but what is to be overcome by Induſtry. I would rather adviſe Students in Surgery to have the moſt famous of the Ancients in this Art always in their eye; and to conſider that we not only enjoy all the Advantages they had, but far greater; for we have been ſo largely aſſiſted by the Inventions of ingenious Men in theſe latter days, that if we equal our Anceſtors in Induſtry, we ſhall eaſily exceed them in Skill.

But nevertheleſs extremely neceſſary.

LXXIV. But altho' the Attainment of Surgery had been ſtill more difficult than it is, yet as we do not enquire into the Difficulties, but the Honours and Uſes that attend the Acquiſition of an Art or Science, before we make choice of it; this is ſo far from being a Diſcouragement to generous Minds, that it is rather an Incitement to their Induſtry. That Surgery is extremely neceſſary for

[a] See CELSUS, Book V. Chap. 26. Number 24.

the Preſervation of Life, does not only appear from what we have already laid down, but from the Neceſſity the Phyſicians frequently lye under of calling for the aſſiſtance of this Art, not only in external Diſorders (to which ſome would impertinently confine Surgery) but in internal Complaints alſo, where Medicines, and a proper Diet, are in no wiſe equal to the Cure, as in the Cataract, Stone in the Bladder, Empyema, Dropſy, Suppreſſion of Urine, difficult Births, and an infinite Number of other Caſes; in which they are often obliged to fly to Surgery, as their ſole Anchor; as appears from the Teſtimony of the great HIPPOCRATES [a]. Amongſt the great numbers that have been Scoffers and Deriders of Phyſic, there have been very few ſo hardy as to reject Surgery as an uſeleſs Art; for indeed he muſt be entirely given up to Impudence and Folly that would pretend this to be an uſeleſs Art, by whoſe Aſſiſtance the moſt grievous Diſorders that the Body is ſubject to are relieved; to wit, Wounds, and the Loſs of Blood that is conſequent upon them, Fractures or Luxations of the Bones, Stones in the Bladder, Suppreſſion of Urine, and an infinite Number of others.

Surgery the moſt certain part of Phyſic.

LXXV. I would have no one be ſurpriſed at the Aſſertion, that Surgery ſurpaſſes all other Branches of Phyſic in point of Certainty; what CELSUS ſaid formerly upon this Occaſion is very true [b], "The effects of Surgery are more "evident than thoſe of any other Branch of Phyſic, ſince in many Diſeaſes (the "internal eſpecially), Nature or Accident may do much, and the ſame Me-"dicines have ſometimes a good Effect, and ſometimes no Effect at all, ſo "that it becomes Matter of Doubt, whether Health be the Effect of the Me-"dicines that have been adminiſtred, or of a good natural Conſtitution of the "Body; but in Diſorders that are relieved by the Aſſiſtance of the Hand, it "is very evident from whence the good Effects proceed." Whatever good Effects we produce by ſtopping violent Hæmorrhages, by taking off Tumors and Excreſcences, by curing Herniæ, by cutting for the Stone, by couching Cataracts, by drawing forth ſuppreſſed Urine, by changing the croſs Poſition of the Infant in the Womb, and bringing it into the World, by ſetting broken Bones, and reducing luxated ones, and by relieving other Diſorders of this Sort; for all this we are evidently obliged to the Hand of the ſkilful Surgeon.

Students in Surgery incited to Induſtry.

LXXVI. Having premiſed this by way of Introduction, we cannot avoid again and again exciting all Students of this moſt noble Art to Diligence and Induſtry, and not to reſt ſatisfied with being able to ſhave, ſpread a Plaſter, or open a Vein; for I would have them know, that not only a good natural Sagacity, but great Labour and Study alſo, are abſolutely required to qualify a Man for ſo great a Truſt as that of taking care of the Health of Mankind. Students in Surgery ſhould not only be furniſhed with Strength of Body, but Conſtancy of Mind alſo, that they may remain unmoleſted and unmoved by the Stench, Blood, Pus, and Naſtineſs that will naturally occur to them in their Practice. They ſhould conſider, that by frequent Exerciſe theſe Things will become cuſtomary to them, and they will acquire, as it were, a ſecond Nature. And a Surgeon ſhould ſubmit to the greateſt Inconveniences, rather than neglect any thing that might be for the Benefit of his Patient; for then he will have performed his Duty properly, and have ſatisfied his own Conſcience, when he has done every thing that comes within the Compaſs of his Art for the Service of his Patient.

[a] Sect. 8. Aph. 6. [b] See Lib. VIII. Præfat. and HIPPOCRATES de Arte, V.

INSTITU-

INSTITUTIONS OF SURGERY.

PART the FIRST.

Of the Five Kinds of Diſorders of the Body.

BOOK the FIRST.

Of WOUNDS.

CHAP. I.

Of WOUNDS *in general.*

A Wound, what.

I. WE were perſuaded by two Reaſons to begin theſe Inſtitutions of Surgery, with an Enquiry into the Nature of Wounds; for Wounds are not only more common than any other external Injuries, but the Nature of them alſo is more eaſily explained in our Schools of Surgery. And indeed when we are thoroughly acquainted with the Nature of a Wound, we ſhall with much greater Eaſe and Clearneſs comprehend all the other Doctrines of Surgery. What a *Wound* is, the moſt unſkilful are acquainted with; but it is frequently defined to be a *violent Solution of the Continuity of the ſoft external Parts of the Body, made by ſome Inſtruments, whether ſharp or obtuſe.* Others take a greater Latitude in defining it, *and call every external Hurt of the Body, by what Cauſe ſoever produced, a Wound.* For Inſtance, they reckon violent Strokes upon the Head, Thorax, or Abdomen, under the Title of Wounds, though no external Parts are divided, as will eaſily appear from what we ſhall ſay below, when we come to treat of mortal Wounds.

Differences of Wounds, in relation to the inflicting Inſtrument.

II. On the other hand, ſome are of Opinion, that unleſs the injured Parts of the Body are divided by ſome ſharp Inſtrument, as by a Sword or Knife, it is by no means to be called a Wound; though it plainly appears, from what has been already ſaid, that thoſe Injuries which are produced by blunt Inſtruments, may properly enough be called Wounds; under this Head are Gun-ſhot Wounds, Wounds inflicted by Stones, Clubs, or that come by violent Falls: Therefore we may conſtitute two Differences of Wounds; the one made by *acute*, the other by *blunt Inſtruments.*

On what Parts of the Body Wounds are inflicted.

III. Wounds are generally inflicted upon the *ſofter Parts of the Human Body*, ſuch as the Skin, Fat, Muſcular Fleſh, Ligaments, Blood-veſſels, and Nerves, and Parts that are compoſed of theſe, as the Viſcera and Inteſtines; but whilſt we are aſſerting this, we muſt by no means entirely exclude *the more ſolid Parts of the Body, as the Bones*; ſince the Bones themſelves afford frequent Examples of Injuries received from ſharp Inſtruments. The Parts, therefore, that are ſubject to theſe Injuries will afford us two Diſtinctions of Wounds; one, *Wounds of the ſoft Parts*; the other, *Wounds of the Bones.*

Cauſes of Wounds.

IV. As *Cauſes of Wounds*, all Inſtruments of what kind ſoever, whether blunt or ſharp, may properly be reckoned, provided they are of ſuch a Nature, that, upon the violent external Application of them, they are capable of producing a Solution of Continuity in the Parts of the Body upon which they are inflicted: For a Solution of the external Parts from an internal Cauſe is not called a Wound, but rather an *Abſceſs*, or *Ulcer*. So when the harder Parts of the Body, to wit, the Bones, are broken by a Fall, or by a violent Blow received from a blunt Inſtrument, we do not call that a Wound, but a *Fracture*.

Effects of Wounds.

V. *The Effects which are produced by Wounds*, beſides the Diviſion of the ſofter Parts, are generally *Profuſions of Blood*, though they are ſometimes attended with much greater Miſchiefs than theſe: For it can ſcarcely happen, but that the divided Parts muſt in ſome meaſure, if not totally, loſe their natural Functions, according to the different Uſes for which the Part is intended, and according to the different Degree of Injury that it receives. The greater Number of Uſes a Part is intended for by Nature, the worſe will be the Conſequence of a Wound upon that Part. This Principle is ſo extenſive, that we are always guided by it in forming our Prognoſtic, whether the Wound will prove mortal or not. He therefore that is beſt ſkilled in Anatomy, that is beſt inſtructed in the Situation of Parts, and their Uſes, will be enabled to form the moſt accurate Judgment of the Conſequences that will neceſſarily attend a Wound upon any particular Part.

Different ſorts of Wounds.

VI. What we have taught of the different Situations and Cauſes of Wounds, ſufficiently demonſtrates, that there are many different kinds of Wounds: ſome are brought on by *Puncture*, ſome by a *Stab*, and ſome again by a *Blow*: ſome are *curable*, others *incurable*: ſome are made with *ſharp Inſtruments*, others with *blunt* ones: To which Claſs may be referred all Gun-ſhot Wounds, all that are occaſioned by a Blow, or a Fall, and which the Surgeons diſtinguiſh by the Name of Contuſions. With regard to their *Figure*, ſome form a *right Line*, others are *curve*, *tranſverſe*, or *oblique*: with reſpect to their *Situation*, ſome are ſeated in the *Head*, others in the *Neck*, *Thorax*, or *Abdomen*; and of theſe, ſome are *external*, others *internal*. Variety of different kinds of Wounds ariſe from the great diverſity of Condition that Wounds are left in: For in ſome Wounds the inflicting Inſtrument, or Part of it, remains; for inſtance, a Leaden Bullet, a Piece of Glaſs, or of a Grenade, the Points of Swords or Arrows: But in ſome Wounds nothing of this kind is left. Sometimes *Fractures of the Bones* accompany Wounds, which we almoſt always find to be the Caſe in Wounds of the Head, and in Gun-ſhot Wounds. Some Wounds alſo are attended with Poiſon, as thoſe which are made with poiſoned Arrows, or other Inſtruments. Under this Head we may very properly rank the Bites of Animals, but more particularly of mad or venomous Animals. Some are of Opinion,

that Wounds which are made with Copper or Silver Inſtruments ſhould be reckoned in this Claſs, the Poiſon of which, if there is any, is owing to the Vitriol that is mixed with theſe Metals.

What appears after a ſlight Wound.

VII. In *ſlight Wounds*, that is to ſay, where no conſiderable Vein, Artery, Nerve or Tendon is concerned, you will uſually remark the following *Appearances:* At firſt ſight, the Wound appears to us as a red Line drawn upon the Part, but upon being dilated the Blood inſtantly guſhes out, in greater or ſmaller Quantities, in Proportion to the Size and Number of the Blood-veſſels that are injured. The Hæmorrhage, after a ſhort Continuance, ſtops of its own accord, or by ſome external Application, and the Blood concreting in the Wound, forms a Cruſt; The Lips of the Wound now begin to look red, and ſwell, and are attended with ſome Degree of Pain and Inflammation. If it is a large Wound, a Fever, that is to ſay, an univerſal Heat and quickneſs of Pulſe almoſt always enſues. Upon the third or fourth Day, ſooner or later, a whitiſh glutinous Humour, not unlike white Oil, appears; and this is known to the Surgeons by the Name of *Pus*, or *Matter*. Upon the Appearance of Matter, the Redneſs, Tumor, Pain, Inflammation, and Fever diſappear entirely, or at leaſt are ſenſibly abated; and theſe are the Signs of a Wound inclining to heal: For under the Matter we have deſcribed, new Fleſh ſprings up from the wounded Veſſels, which having by Degrees filled the Wounds, dries upon its upper Part, and forms a Cicatrix.

What after dangerous Wounds.

VIII. In *dangerous Wounds*, that is, where any conſiderable Blood-veſſel is wounded or divided, there generally enſues ſo violent an *Hæmorrhage*, that the wounded Perſon is in an inſtant ſenſible of great loſs of Spirits, and Weakneſs, and faints away; and when the larger Arteries are wounded, whether they are internal or external, he dies upon the Spot. Although ſomewhat leſs Danger is apprehended from Wounds that are inflicted upon the Veſſels, which are ſituated upon the external Parts of the Body (ſome few excepted) becauſe they will admit of the Ligature, and other Means for reſtraining the Violence of the Hæmorrhage: Nevertheleſs it is almoſt impoſſible to prevent the Limbs which lie below the Diviſion of the Artery, and are uſed to receive their Nouriſhment by that Channel, from becoming paralytic; nay, ſometimes, from mortifying. This is almoſt conſtantly the Caſe when the Trunk of the Brachial or Crural Artery is divided.

What follows the partial Diviſion of a Blood-veſſel.

IX. The Conſequences we have juſt related, follow upon the total Diviſion of a conſiderable Vein or Artery: It remains now that we conſider what will follow upon a partial Diviſion of them. Whenever a large Artery is wounded, and not entirely divided, the wounded Fibres inſtantly contract themſelves; by this Means they dilate the Orifice of the Wound, and render it difficult to ſtop the Flux of Blood; and though the Hæmorrhage be ſtopped for a little Time, yet it will burſt out again on a ſudden violently, or at leaſt produce a dangerous Tumor, called an *Aneuriſm*. This will frequently be the Caſe, when only the external Coat of the Artery is wounded: For by this Means the internal Coat of the Artery is left to ſuſtain the whole *Impetus* of the Pulſe, which being unequal to, it is forced by Degrees into a Tumor like a Bag, whence frequently enſue the moſt calamitous Conſequences. But of this Caſe we ſhall treat more fully in another Part of this Work.

X. Upon

Conſequences following from a wounded Nerve.

X. Upon *the Diviſion of a Nerve*, the Limb to which that Nerve was extended becomes inſtantly rigid, void of Senſation, and withers; ſo it is no wonder that a Man inſtantly expires upon the Diviſion of thoſe Nerves that are ſent to the Heart, or *Diaphragm.* A Wound alſo is attended with great Danger, where the Nerve is only partially wounded, and not entirely divided; for the wounded Fibres contract themſelves, and thoſe that remain undivided ſuffer too great Extenſion, which will bring on moſt violent Pain, Spaſms, Convulſions, Inflammations and Gangrenes, and ſometimes Death itſelf.

Conſequences of a wounded Tendon.

XI. When a *Tendon is wounded or divided*, the Part to which it belongs loſes its Motion; but if it is divided only in Part, it will produce much the ſame Symptoms with a Nerve in the ſame Circumſtances. The conſequence of Wounds upon the internal Parts, you will find more fully explained when we come to treat of the *Diagnoſis* and *Prognoſis* of Wounds.

Of the *Diagnoſis* of Wounds in general.

XII. The *Diagnoſis* of Wounds is for the moſt Part extremely eaſy, for the Size, Situation, and Nature of the Wound, generally lies open to the Sight. Nevertheleſs there are ſome Caſes that are not very uncommon, where it is ſomewhat difficult to diſcover the true Nature of the Wound. But in order to make the more eaſy Diſcovery, whether the Wound is deep or ſuperficial, whether any of the internal Parts are wounded or not, the Surgeon ſhould always be careful upon the firſt Viſit to clean the wounded Part with a Sponge ſqueezed out of warm Wine or Water, that he may have a clear View of the Bottom of the Wound. But whenever the Flux of Blood from the Wound is very violent, it muſt be inſtantly dreſſed up, and the cleaning of it in this Manner deferred, till it is in a quieter Diſpoſition.

What internal Parts are wounded may be diſcovered, 1. By Anatomy.

2. By the Poſture of the wounded Perſon.

XIII. In deep Wounds we are to examine, whether the fat and fleſhy Parts are the ſole Objects of the Wound, or whether ſome conſiderable Blood-veſſels, or other internal Parts, are not Partakers of the Injury. We are aſſiſted in this Examination by ſeveral Means. Our firſt Aſſiſtance we receive from the Knowledge of *Anatomy*, ſince by that Science we are taught the Situation of each particular Artery, Nerve, Tendon, Viſcus, and Inteſtine. The *Poſture of the wounded Perſon* at the Time he received the Injury is alſo to be diligently conſidered, whether he was ſtanding upright, or lying down, and to which Side he inclined; by this Means we may with ſome Certainty judge what Parts were Sufferers by the Wound, and how far the Weapon penetrated. We are alſo to conſider of *the Poſture, Manner, and Force uſed by the wounding Perſon*; for the greater degree of Force there was in dealing the Blow, ſo much the larger and deeper will the Wound be. Nor ſhould we neglect here to enquire after *the Shape of the Weapon* by which the Wound was inflicted, ſince by conſidering its Size, and obſerving the Quantity of Blood that adheres to it, we may in ſome Meaſure judge of the Depth of the Wound.

3. By the Poſture and degree of Force uſed by the wounding Perſon.

4. By viewing the Weapon.

5. By the Diſturbance given to Actions of particular Parts.

XIV. In a word, there is nothing will give you truer Light into the Nature and Conſequence of a deep Wound, than a due Conſideration of what natural Actions of the Body are impeded or diſturbed thereby. For inſtance, in Wounds of the Breaſt, when the Patient draws his Breath with Shortneſs and Difficulty, and is at the ſame time attended with an Hæmoptyſis and Hiccoughs, we may fairly conjecture that the *Lungs* or *Diaphragm* are wounded; though the Hiccoughs often ariſe too from Wounds in the Stomach, in the Bladder, and other internal Parts of the Body: So in Wounds of the Abdomen, when Chyle is voided,

voided, it is a plain Indication that the *Stomach*, the *ſmall Inteſtines*, or the *Lacteal Veſſels* are wounded. When Excrements paſs by the Wound, the *large Inteſtines* are wounded. In the ſame manner, an Effuſion of Bile ſhews the *Liver* or *Gall-Bladder* to be divided. If Urine paſſes by the Wound, the *Bladder*, or *Ureters* are injured; but bloody Urine denotes a Blow on the *Kidnies*, or a Wound of the *Bladder*; but when the Diſcharges of Blood this Way are violent, it is a ſtrong Indication that ſome of the larger Blood-veſſels muſt be wounded. Vomiting of Blood, for the moſt part, declares the *Stomach* to be the injured Organ; *violent Pains*, attended with *convulſive Twitches*, ſhew that a Nerve is wounded, or that ſome foreign Subſtance is left in the Wound. Whenever the Senſes are diſordered after a wound received upon the Head, a Concuſſion of the Brain is much to be feared.

XV. What we have laid down concerning the general Method of forming the *Diagnoſis* on Wounds, will alſo ſerve us in forming their *Prognoſis*, or Judgment of the Conſequences that will attend them: For after a due Conſideration of the Nature of a Wound, and the Symptoms attending it, it will be no very difficult matter, to determine whether it be attended with great danger or not, whether the Cure will be difficult or eaſy, whether it will be a perfect or imperfect Cure. We may remark in general, that ſlight Wounds admit of an eaſier Cure than deep ones: Young Patients, or thoſe of a ſound Conſtitution, are eaſier cured than the old or diſeaſed, particularly than Hydropical, Conſumptive, Scorbutical, or Pocky Perſons. The Cure is eaſier performed in *a temperate Air*, than in a cold or hot Climate. There are alſo greater hopes of Succeſs where there are no violent Symptoms attending, as profuſe Hæmorrhages, large Tumors, vehement Pains, Convulſions, Inflammation, Fever. But HIPPOCRATES has very rightly remarked [a], "Where a large Wound is made, it is a very bad ſign "if no Tumor ſucceeds." This CELSUS has explained in a much more elegant manner [b]: "It is of bad conſequence for a Wound to be attended with a large "Tumor, but it is of the laſt conſequence if it is attended with no Degree at "all of Tumor; the firſt is an Indication of great Inflammation, the laſt of "Mortification." Some Degree of Tumor therefore is beſt. *Of the Prognoſis of Wounds.*

XVI. We come now to enquire what *Wounds admit of Cure*, and what are *incurable* or *mortal*. The Knowledge of this Point is no leſs uſeful and neceſſary to the Phyſician and Surgeon, than it is difficult to attain: And more eſpecially as the Law inflicts a very heavy Puniſhment upon Murderers, it is of very great Conſequence to be able to diſtinguiſh what Wounds are of themſelves mortal, and what only become ſo by Accident or Neglect; that the Guilty may receive their juſt Sentence, and the Innocent be freed: On which Account the moſt eminent Phyſicians and Surgeons are often called in, and, with great Reaſon, by the Adminiſtrators of Juſtice, in all Caſes of Difficulty or Uncertainty. In order to enable the Surgeon to anſwer Queſtions upon this Head with greater Readineſs and Certainty, we ſhall be very particular in this Article. Therefore in this View we ſhall divide Wounds into three Sorts. Some Wounds, 1. Are abſolutely of themſelves mortal: other, 2. Are in their own Nature mortal, if not relieved by timely Aſſiſtance: others, laſtly, 3. Become mortal by Accident or imprudent Treatment, though they were otherwiſe curable. *Whether Wounds are curable or incurable.*

[a] HIPP. Aphoriſm 66. Sect. V. [b] Book V. Chap. 29.

XVII. We

1. Mortal Wounds.
2. Where there is an Hæmorrhage not to be ſtopped by Art.

XVII. We properly ſtyle thoſe *Wounds mortal, which are not to be remedied by all the Art and Induſtry of Man.* So thoſe Wounds are juſtly deemed mortal that are attended with ſo violent an Hæmorrhage as to produce inſtant Death. In this Claſs are reckoned Wounds that penetrate the Cavities of the Heart, and all thoſe Wounds of the *Viſcera* where the larger Blood-veſſels are opened. Such are large Wounds of the Lungs, Liver, Spleen, Kidnies, Stomach, Inteſtines, Meſentery, Pancreas, Uterus, Aorta; of the Iliac, Cœliac, Renal, Meſenteric, Carotid, and Crural Arteries, (eſpecially if they are wounded near their Origin) of the Subclavian alſo or Vertebral, of the Vena Cava, the Iliac Vein, internal Jugular, Vertebral, Renal, Meſenteric, of the Vena Porta, and of other large Veins that lie deep in the Body, becauſe their Situation will not admit of proper Applications to reſtrain the Flux of Blood. I think therefore I may very juſtly reckon theſe amongſt the Wounds that are abſolutely incurable, ſince they are not remediable either by Aſtringents, Ligature, or Fire. We may refer alſo to the ſame Claſs the Wounds of the Brachial Artery, if near its Origin; for the large Effuſion of Blood generally deſtroys the Patient before Aſſiſtance can be procured; more eſpecially if the Artery and Vein are both wounded together,

2. Where the Communication is cut off between the Head and the Body.

XVIII. Thoſe Wounds are no leſs mortal than the former, *which obſtruct or entirely cut off the Paſſage of the Animal Spirits to the Heart.* Such are Wounds of the Cerebellum, Medulla Oblongata, and all violent Strokes of the Brain itſelf. There is Reaſon to apprehend very great Danger, when the ſmall Veins or Arteries which are contained in the Cranium are injured; for the Blood flowing from them into the internal Sinuſes of the Brain, either produces too great a Preſſure upon thoſe very tender Parts of the Brain, and ſo obſtructs the Courſe of the Blood and Spirits; or being corrupted, putrefies the Brain, if it cannot be evacuated by the Aſſiſtance of the Trepan, which is the Caſe when this Accident happens at the lower Part of the Cranium, or in the Sinuſes of the Brain. Nor is there leſs Danger, where *the Nerves which tend to the Heart*, or the *Cerebellum*, are wounded, or entirely divided; for after this, it is impoſſible for the Heart to continue its Motion.

3. Where the Power of Breathing is taken away.

XIX. To this Claſs alſo are to be referred *all Wounds that entirely deprive the Animal of the Faculty of Breathing.* Therefore there is great Danger where the *Aſpera Arteria* is completely divided; for where is is only divided in part, it may be healed again by the Aſſiſtance of an expert Surgeon. I have many [a] Hiſtories of Cures of this Kind, both by myſelf and others. To this Place alſo belong violent Shocks of the *Bronchia*, *Mediaſtinum*, and Diaphragm, eſpecially the tendinous Part of it; or any Injuries of the principal Nerves, as the *Par Vagum*, or *Intercoſtals*.

4. Where the Courſe of the Chyle is interrupted.

XX. Thoſe Wounds, alſo, *which interrupt the Courſe of the Chyle to the Heart*, are no leſs incurable than the former: Such are the Wounds of the Stomach, Inteſtines, Receptacle of the Chyle, Thoracic Duct, and larger Lacteals: to which we may add Wounds of the Œſophagus, if they are large; though Death is not ſo ſudden an Attendant upon theſe Wounds, but for want of Nouriſhment they are greatly weakened by Degrees, and die conſumptive.

[a] See BOHNUS *de Vuln. renunc.* Pag. 21. though he reckons theſe among incurable Wounds.

XXI. In

XXI. In this Place we muſt by no means omit to ſpeak of *Wounds which are inflicted upon the interior membranous Parts that contain ſome ſecreted Fluid*, as on the Bladders, either for the Bile or Urine, the Ureters, Stomach, Inteſtines, Receptacle of the Chyle, and Lacteal Veſſels. The Fluids contained in theſe Parts, when once they are let looſe into the Cavity of the Abdomen, cannot be properly diſcharged, and therefore eaſily corrode the internal Parts of the Body; and the Membranes that contained them are generally ſo fine, that they will not admit of Agglutination, eſpecially ſince no Medicine from without can be applied: a few, indeed, have recovered after ſlight Wounds in theſe Parts, but ſince that Number is but few, and the Cure was accidental, and not performed by the Surgeon's Art, I think I am ſufficiently juſtified in adding theſe to the Number of incurable Wounds.

Where the abdominal Fluids which are contained in Membranes are extravaſated.

XXII. We have hitherto been treating of Wounds that were curable by no Art or Induſtry; we proceed now in order to deſcribe thoſe *which prove fatal, if neglected and left to Nature.* By theſe we mean thoſe Wounds that produce inſtant Death, unleſs relieved by preſent Aſſiſtance, but are curable by a good Surgeon called in time; ſuch are Wounds of the larger external Blood-veſſels, which might be remedied by Ligature, by the Application of aſtringent Medicines, or of the actual Cautery. Of this kind are Wounds of the brachial or crural Artery, unleſs they are too near the Trunk of the Body. Wounds in the large Arteries of the Cubit or Tibia, of the Branches of the external carotid and temporal Arteries, are of this kind; to theſe may be added Wounds of the Jugular, and other Veins ſituated upon the external Parts of the Body; but in theſe Caſes we always ſuppoſe that Help is called for before there has been a vaſt Profuſion of Blood.

Wounds proving mortal, if left to themſelves.

XXIII. *Wounds are properly ſaid to become mortal by Accident, where the Patient's Death is occaſioned either by the ill Conduct of the Patient himſelf, or by the Ignorance or Neglect of the Surgeon, the Wound itſelf being deemed curable.* Under this Head are to be reckoned, 1. *Thoſe Wounds which the Surgeon has neglected to cleanſe ſufficiently, though he had it in his Power to do it*; as when ſome foreign Body, which might eaſily have been extracted, is left in the Wound by the Careleſſneſs of the Surgeon, and produces Inflammations, Hæmorrhages, Convulſions, and at laſt Death itſelf. So in Wounds of the Thorax and Abdomen, if the Surgeon does not uſe his utmoſt Diligence to evacuate the grumous Blood, it will corrupt there, and by drawing the neighbouring Parts into Conſent, will expoſe the Patient to inſtant Death [a]: therefore great Care muſt be taken that

What Wounds become mortal by Accident.

[a] There are ſome Caſes where the Surgeon finds all his Attempts to evacuate the Blood fruitleſs, and there he is in no wiſe to be blamed, but the Wound is to be looked upon as mortal. Take the following Caſe by way of Example: In the Year 1725, a Man received a Wound by a Sword: the Sword entered about half an Inch below the right Pap, between the fifth and ſixth Ribs, and paſſed downwards through the Diaphragm into the Cavity of the Abdomen. Now although a conſiderable Quantity of Blood was diſcharged by the Wound for the three firſt Days, yet it was impoſſible that the Blood which was extravaſated in the Cavity of the Abdomen, ſhould be diſcharged by the Wound at the Breaſt, the Patient therefore died on the eighth Day. His Body being opened, we found a large Quantity of grumous Blood under the Liver, which adhered ſo ſtrictly to its concave Part, that we found it difficult to ſeperate them with our Fingers. Upon clearing away the Blood, we perceived a Wound through the Body of the Liver, about half an Inch wide, and a Wound anſwering to that in the muſcular Part of the Diaphragm. There were two or three Ounces of Blood found in the lower Part of the Abdomen, but none in the Cavity of the Thorax. From

the Lips of the Wound do not close, till the Blood which is collected in the Cavity of the Body be all evacuated, if possible, which you will easily perceive by the Difficulty of Breathing, and other bad Symptoms being removed [b]. But if any of the larger internal Vessels are wounded, then all Attempts to discharge the Blood are vain; for the Violence of the Hæmorrhage takes off the Patient. 2. *Wounds also are reckoned mortal by Accident, which are treated or searched in too rough a manner by the Surgeon*; for if you handle Wounds roughly, that are full of nervous Parts or large Blood-vessels, there is great Danger of bringing on Hæmorrhages, Convulsions, Inflammation, Gangrene, and Death itself. The Case is also the same, 3. *in external Wounds which are slight of themselves, but the Patient is lost by the Violence of the Inflammation, which is brought on, and increased by the Surgeon's injudicious Treatment.* or, 4. *when any one is taken off by the Violence of the Hæmorrhage from a Wound of the Hand or Foot*; for in this Case a Surgeon might easily have stopped the Blood by the Application of proper Remedies, or by the actual Cautery, or Ligature. or, 5. *where the Patient is guilty of any Intemperance in Eating or Drinking, of Excess of any Passion, of exposing himself to the cold Air, or of using any violent Exercise:* for by this means, Wounds, more especially those of the Head, by being liable to fresh Hæmorrhages, and other dangerous Accidents, frequently become mortal, notwithstanding the Surgeon uses his utmost Care and Skill: under this Head also are to be reckoned, 6. *those Wounds of the Head where the Patient is lost by the vast Quantity of Blood which is extravasated in the Cavity of the Cranium, and confined there; but where he might have been relieved if the Trepan had been used in Time*; for though Wounds of this kind generally prove incurable, yet as there is a Possibility of saving a Person in these Circumstances by the Use of the Trepan, this may properly be reckoned amongst the doubtful Cases, and not deemed absolutely mortal: lastly 7. *a bad Habit of Body frequently prevents the Cure of Wounds, which would admit of an easy Cure in an healthy Subject.* So you frequently see the slightest Puncture in the Hand or Foot of an hydropical, consumptive, or scorbutical Person, shall produce a Gangrene, and prove mortal, though the Surgeon neglects no proper Application to prevent it. I know very well that some Physicians reckon all Wounds of this kind as absolutely mortal; but I think they are much better justified who pronounce a milder Sentence, and deem them of the doubtful Kind.

the Impossibility that appeared of discharging the extravasated Blood, and the Largeness of the Wounds of the Vessels, I pronounced this Wound mortal: but, to my great Surprize, some Physicians declared it so only *per accidens*, for which reason the Murderer was acquitted. Whose Opinion was most justifiable, I leave to others to determine. See *Fr. Hoffman. Consult.* Tom. I. p. 376. *& seq.*

[b] The Surgeon is not to be blamed if he is sometimes deceived in this Point; of which I will here give you a notable Instance. In the Year 1726, a Man at *Helmstadt* was wounded in such a Manner under the right Pap, that the Blood did not only flow in great Quantities from the Wound, but discharged itself also by the Mouth: but in two Days Time the Discharge of Blood both at the Wound and by the Mouth, through the Application of proper Medicines, entirely ceased, and the Patient found himself in so good Order, that he expected in a very short Time to get abroad: he breathed so freely, that he easily prevailed upon me to remove the Tent that I had put in to keep the Wound open. But behold the Consequence! after remaining in this Manner entirely easy for two Days; on the third he died suddenly. Upon opening the Thorax, we found at least a Pound of concreted Blood, which could by no means have been discharged, since there

It is difficult to form a Judgment concerning the Fate of Wounds.

XXIV. We have laid down theſe Principles to guide Phyſicians in giving their Opinions in Courts of Juſtice, concerning the neceſſary Conſequences and Fate of Wounds. Although all Wounds ſhould be examined upon theſe Occaſions with great Circumſpection, yet none require more careful looking into than Wounds of that Claſs which are deſcribed under *N.* XXII, becauſe there are great Diſſenſions amongſt the learned upon this Head. Some are of Opinion, that the Wounds mentioned at *N.* XXII, are to be referred to the third Claſs, and ſo are to be reckoned mortal only by Accident, and by this Means they frequently acquit a Murderer. How they ſupport this Opinion I cannot tell: for my own Part, whenever I have found a Man loſe his Life by receiving a Wound in an Artery, at a Time of Night when a Surgeon could not be called, I have always determined that Wound to be mortal, and that the offending Party was guilty of the Murder: on the other Hand, where a Wound of the ſame Kind has been received in the Day-time, and the Patient has loſt his Life by the Neglect of the By-ſtanders, in refuſing to call proper Aſſiſtance, or by the Ignorance of the Surgeon; in theſe Circumſtances I have always declared the Wound to have been mortal only *per accidens*, and have given my Opinion, that the accuſed Perſon ought to be aquitted, and the Surgeon indicted. But in order to form a proper Judgment in theſe Caſes, it is neceſſary that we ſhould be well informed of all the Circumſtances [c].

What is to be done in doubtful Caſes.

XXV. In very doubtful Caſes, to be ſure, the mildeſt Sentence ought to take Place, according to the old Saying, *It is better to let ten guilty Perſons eſcape, than to puniſh one innocent Man:* for to be too rigid in theſe Caſes, will not only burden the Conſcience of the Judge, but be alſo injurious to the Public.

The Form of delivering-in a Surgeon's Opinion.

XXVI. For the uſe of the younger Surgeons, I ſhall here ſubjoin the Form which I always uſe in giving my Opinion into Court, concerning the Nature of a Wound.

appeared no Symptoms which could give room to ſuſpect that there remained any extravaſated Blood concealed. Beſides, BELLOST, DE LA MOTTE, and ſeveral other celebrated Surgeons amongſt the Moderns, abſolutely forbid keeping Wounds of the Breaſt open by the uſe of Tents, though I doubt much whether this Advice is always to be followed. But I leave this to the Determination of others.

[c] As an Example of this, take the following Relation. In the Year 1733, a Woman living in the Suburbs near *Brunſwic*, walking out in the Evening juſt before the Gates of the City were ſhut, received a Blow on the Head from a Man with a large Club, which laid her flat upon the Ground, and left her quite ſenſeleſs; when the Fellow ſaw this, he took to his Heels, and nobody was left near her, but her Huſband and three ſmall Children; the Man, frighted out of his Wits, ran about to ſee if he could get People to aſſiſt him to his carry Wife home (for ſhe was a very large Woman) but the Night coming on he could prevail with no one, and the City Gates being ſhut, it was impoſſible to bring a Surgeon to her: the Woman therefore was left upon the Ground all Night, without any Aſſiſtance, and died the next Morning. When the Phyſicians and Surgeons came to examine her, they found a Fiſſure in the Cranium, and, upon raiſing the Scalp, they found a large Quantity of extravaſated Blood under the *Dura Mater*, lying upon the right Lobe of the Brain, and therefore very judiciouſly determined it to be a mortal Wound. The Advocate for the Criminal oppoſed this Verdict, becauſe there was no Surgeon called to treat her in a proper Manner, by which ſhe might poſſibly have been ſaved; upon this Difference of Opinions, I was called upon to determine this Matter. I declared as my Opinion, that if the Woman had been within the City, where ſhe might have had the Aſſiſtance of Phyſicians and Surgeons, and had loſt her Life through their Neglect or Ignorance, then the Wound ought to have been deemed mortal *per accidens*; but in the preſent Caſe it was impoſſible ſhe ſhould have had any ſuch Aſſiſtance; therefore her Death was occaſioned by the Blow ſhe received, and the Wound ought to be judged mortal *per ſe*.

"I the under written, having this Day diligently examined the dead Body of "A. B. in the Preſence of C. D. E. *&c.* found it to have received the follow- "ing Wounds: that is to ſay, in the back Part of the Body, under the right "Shoulder, I diſcovered a Wound of the Width of one Inch, through which I "could paſs my Finger with great Eaſe, between the Ribs, into the Cavity of "the Body. Upon opening the Breaſt, almoſt the whole right Side was found "full of coagulated Blood, upon removing which, I found a Wound alſo pene- "trating into the right Lobe of the Lungs, which not only pierced through "this Lobe, but alſo divided ſome of the larger Branches of the pulmonary "Veſſels, with the *Bronchiæ* themſelves. The Heart and all its Veſſels were "entirely empty: no Miſchief appeared either in the Head or Abdomen. The "Effuſion of Blood, which was occaſioned by dividing the Veſſels in the Lungs, "could not but bring on inſtant Death: therefore I hereby declare this Wound "to have been the Occaſion of his Death. In Teſtimony of the Truth whereof "I have hereto ſet my Hand."

N. N.

Done at the Day of
in the Year of our Lord

Some general things with Reſpect to theſe Forms.

XXVII. Forms without Number may be made from this, by varying the Circumſtances: but above all I would adviſe the young Surgeon to be very careful in examining the State not only of the wounded Parts, but alſo of the Contents of the Cranium, Thorax, and Abdomen, that he may obſerve whether any thing preternatural has happened in either of thoſe Cavities. If any one is deſirous of being more thoroughly inſtructed in the Method of examining the Bodies of murdered Perſons, and in the proper Forms of making a Report, let him conſult a *French* Treatiſe upon this Subject, entituled, *L'Art de faire Rapport en Chirurgie.*

The CURE *of* WOUNDS.

Cure of Wounds.

XXVIII. Since a Wound is a Solution of the Continuity of the Parts of the Body, the Re-union of thoſe Parts ſeems to be the principal Intention. But ſince Wounds are of very different Kinds, ſome ſlight, and others of great Conſequence, in Proportion to this Difference ſo will the Manner of proſecuting this Intention differ.

Cure of ſlight Wounds.

XXIX. The Cure of ſlight Wounds is generally performed with great Eaſe, by applying a ſmall Portion of Lint to the Part, well ſaturated *cum Spiritu Vini, Oleo Ovorum, Terebinthinæ, Hyperici, Linimento Arcæi, Balſamo Copaibæ, de Mechâ, Peruviano, &c.* ſecuring the Dreſſings with a [a] Plaſter to keep the Wound clean. The Dreſſings ſhould be renewed once in a Day or two, and the Lips of the Wound will preſently agglutinate: therefore, in Caſes of this Kind, a Surgeon is very rarely applied to.

Dangerous Wounds how to be treated.

XXX. Wounds which are attended with ſome Danger, where the Subſtance of the Part wounded is not impaired, are to be treated as follows. If there be too copious a Diſcharge of Blood, the Hæmorrhage muſt be ſtopped at the firſt Dreſſing: if not, the Wound in the firſt Place is to be cleanſed from all extravaſated Blood, Sordes, *&c.* in the next Place, if a Bullet,

[a] The Plaſters I chiefly uſe are *Empl. Diachyl, S. Diapalm.* or *Stypticum Crollii.*

the Point of a Sword, any Part of the Cloathing, a Piece of Glaſs, Splinter, or any other foreign Body, ſhall remain in the Wound, it is to be removed with the Fingers, or with proper Inſtruments, as ſhall be explained more fully below; then the divided Parts are to be brought as near each other as poſſible, and their Situation is to be ſo maintained, by proper Bandages, that the Cicatrix which is left may appear even.

Method uſed in cleanſing Wounds.

XXXI. Foreign Bodies are removed from Wounds either by the Surgeon's Fingers, or by ſuch Inſtruments as we have deſcribed at *Plate* III, *Fig.* 3, 4, 5, 6, 7, 8, having firſt enlarged the Orifice of the Wound, if there be Occaſion. but where there are no extraneous Bodies to be removed, and the Hæmorrhage is not large, the grumous Blood is to be wiped away with a ſoft Sponge, or ſome fine Lint, wrung out of hot Wine or Brandy; having done this, you muſt proceed to dreſs, and laſtly to agglutinate the Lips of the Wound.

Method of extracting foreign Bodies.

XXXII. Before a Surgeon attempts the Removal of extraneous Bodies from a Wound, it behoves him well to examine whether this is to be done inſtantly, or whether it is not beſt to wait for a more convenient Time: for if the Patient is become extremely faint, from the Loſs of Blood which he has already ſuſtained, it will be neceſſary here to ſtop the Hæmorrhage, and to endeavour in ſome Meaſure to revive him with moderate Draughts of warm Broths, white Wine Whey, or of ſome cordial Medicine; for if ſome ſuch Precautions are not taken, the Patient may not unlikely die in the Operation: ſo where you have Reaſon to apprehend, that, in extracting the broken Point of a Sword or Spear, you are in Danger of wounding a large Blood veſſel or Nerve, it is better to wait a little till the Patient comes to himſelf, or till the Wound is ſomewhat enlarged by the Suppuration of the Parts. All theſe Circumſtances will be well weighed by the prudent Surgeon.

What Inſtruments are uſed in extracting Bodies from Wounds.

XXXIII. Foreign Bodies, as the Points of Swords, Spears, *&c.* ſhould always be extracted from Wounds by the Hand if poſſible; and this ſhould be done with all the Expedition, Tenderneſs, and Care that may be, taking great Care not to wound the neighbouring Parts; but if there are any Bodies that cannot be removed by the Hands, then you muſt have Recourſe to ſuch *Forceps* as we have deſcribed in *Plate* III, *Fig.* 3, 4, and 5. the ſame Aſſiſtances alſo we make Uſe of in extracting Bullets, Broken Pieces of Steel, Glaſs, *&c.* We ſhall ſpeak more clearly of the Method of extracting Bullets, whenwe come to treat of Gun-ſhot Wounds. Where the Wound is too narrow to admit of the Extraction of a foreign Body without lacerating the Parts, it muſt be dilated with the Knife, according to the Direction of the muſcular Fibres. The Extraction will admit of no Delay, but for Reaſons of great Moment, (*N.* XXXII.) beſides, whilſt the Wound is recent, and the Lips of it not ſwelled, it will ſuffer leſs Pain in handling; and the Patient, from a ſtrong Deſire of living, will at this Time endure more than afterwards, when he comes to reflect.

Of uniting dangerous Wounds.

XXXIV. The Wound being cleanſed from Blood, and all extraneous Bodies, and the Hæmorrhage ſtopped, it now becomes the Buſineſs of the Surgeon to cloſe the Lips of the Wound, and to conſider what is proper to be done to keep them in that Situation, that the Parts may ſpeedily unite. Different Methods are uſed in proſecuting this Intention, according as Wounds differ in their Conſequences, and in the Number and Degree of Symptoms attending them: for ſimple and ſlight Wounds require not the ſame Treatment as thoſe which

are

are attended with dangerous Symptoms, or where the wounded Parts are torn and mangled: again, Wounds, which penetrate into the Cavities of the Body, eſpecially if any of the Viſcera are injured, demand a different Method of Cure from thoſe which are inflicted on the external Parts: and, laſtly, another Diſtinction muſt be conſidered, whether the Wound was made by a Stab, or a Puncture.

Method of treating a Puncture.

XXXV. Amongſt the Number of the moſt ſimple Wounds, we reckon thoſe which are made by Puncture, or Stabbing, upon the external Parts, and not penetrating deep. In theſe Wounds, after the Blood has been ſtopped at the firſt Dreſſing, by the Application of dry Lint, the common Digeſtive, or *Balſamum Arcæi*, or any of the vulnerary Medicines recommended in the Introduction, N°. XXXVII. is to be ſpread upon a Pledget, and applied once every Day; or if the Diſcharge is but ſmall, every other Day, covering the Dreſſings with a Plaſter and Compreſs, and ſecuring the whole with a proper Bandage. At every Dreſſing you ſhould be careful to remove every thing that will give way readily; the Pus, or Sanies is to be gently wiped off with fine Rags. It may be remarked in general, that too frequent Dreſſings do more Harm than Good, unleſs a more than ordinary Diſcharge of Matter, particularly in the Summer Time, or any other bad Symptom, require it: the Truth of this is atteſted by CÆSAR MAGATUS, in his Book, *De rarâ Vulnerum Deligatione*; by BELLOSTE, in his *Hoſpital Surgeon*, and others amongſt the Moderns; not to mention my own Experience upon this Head. The firſt Dreſſings that are applied, eſpecially where there has been a Flux of Blood, ſhould by no means be removed forcibly, but be left till they fall off of themſelves, which they will do when the Suppuration is formed; by this Means much Pain, and perhaps a freſh Hæmorrhage, may be avoided. But when a punctured or ſtabbed Wound penetrates verp deep, the Cure is attended with many Difficulties, eſpecially if a Nerve or Aponeuroſis is lacerated, if the Wound is made perpendicularly down, and has no depending Orifice; for in this Caſe the Blood and Matter are eaſily collected at the Bottom, protract the Cure, and frequently form Fiſtulæ. To prevent theſe Conſequences, it will be proper to preſs the Wound from the Bottom upwards; to apply a Compreſs towards the Fundus of the Wound externally, and to apply what is called the *expelling Bandage* over all, which preſſes much tighter upon the lower than the upper Parts.

A new Opening is frequently required.

XXXVI. But if all this Precaution ſhould prove of no Effect, which is frequently the Caſe, it will be beſt to make a large Opening at the Bottom of the Wound before any Fiſtulæ are formed. In order to make this Opening to the greater Advantage, it will be proper to get a particular Sort of Probe or Needle, very blunt at Top, as at the Letter A; but at the other End provided with a large Eye or Hole through which a Linen Rag may be paſſed, (*See Plate* V, *Fig.* 1.) This Probe is to be paſſed to the Bottom of the Wound, and the blunt Part of it preſſed outwards towards the Skin, till you can feel it with your Finger: when you have felt it, cut down upon it, if you can ſafely, and make a large Opening; ſpread the Rag that you have run through the Eye of this Probe with ſome vulnerary Balſam, and draw it through the Wound after the Manner of a Seton, eſpecially in Gun-ſhot Wounds, and leave it there, dreſſing up both the Orifices with the ſame Balſam, covering the Dreſſings with Plaſters and proper Bandages. In every ſucceeding Dreſſing, the Part of the Rag

Rag that is left out of the Wound is to be ſpread with freſh Ointment, and the lower Part drawn down till this takes it's Place; and this Method is to be continued till the Wound is well cleanſed, the Diſcharge greatly diminiſhed, and all in a Readineſs to heal; the Seton is then to be removed, and the Wound healed as uſual.

Another Method of doing it.

XXXVII. GARENGEOT deſcribes a triangular Inſtrument, invented by PETIT, for this Purpoſe, which the *French* call [a] *Trois quarts:* with this he makes an Opening at the Bottom of the Wound or Fiſtula, and introduces a Rag, which is paſſed through the Eye of this Inſtrument, and then through the Wound or Fiſtula (*See Plate* IV, *Fig.* 1.) but as this Inſtrument is ſtraight, and I have frequently met with Caſes where that Form would not anſwer the Purpoſe, therefore I invented another, long before GARENGEOT's Book came out, for the Uſe of a Nobleman, who had a large Abceſs in the fore Part of the *Abdomen*, which opened near the Navel on the right Side, but penetrated as far as the Groin on the ſame Side. The Situation of the crural Veſſels, in this Caſe, would by no means admit of a new Opening, being made by a ſtraight Inſtrument: I invented therefore a crooked one, ſomewhat like the Inſtrument that is uſed to draw Water off in hydropical Caſes, but longer, becauſe the Fiſtula was of a great Length; (*See Table* IV, *Fig.* 2.) by the Aſſiſtance of which, whilſt I directed the Apex towards the Skin, I eaſily made a new Aperture, without endangering the crural Veſſels: and that I might at the ſame Time introduce the Seton, I contrived a Sulcus near the End, to which I faſtened a ſtrong Thread, and by drawing back the Inſtrument, I eaſily introduced the Seton through the Fiſtula. When the Seton was near all uſed, I ſewed new Cloth to the old, and ſo introduced it through the Wound, cutting off the foul Part, going on in this Manner till the Wound was ſufficiently cleanſed, and ſo preventing the Neceſſity of frequently introducing the Inſtrument.

Cautions concerning Healing.

XXXVIII. It is to be remarked here, that although, in ſome Wounds, it is no Matter how ſoon you ſuffer the Opening to heal; in this Caſe, on the other Hand, you muſt take great Care that the Orifices are not healed before the Bottom of the Wound. This may be done by the Aſſiſtance of a Cloth ſomewhat twiſted, by the *French* called *Bourdonet*, or a ſhort ſoft Tent: but when it is healed from the Bottom, you may remove the Tent, and heal the Orifices. How Wounds of this kind, which penetrate into the Cavity of the Thorax or Abdomen, are to be treated, will be taught below in the Vth and X^th^ Chapters.

Method of treating a Cu

XXXIX. Wounds which are made by a cutting Inſtrument, where no Part of the Fleſh is taken off, and the Accident happens to the external Parts of the Body, and does not penetrate deep, after they are cleanſed, ſhould be dreſſed with ſome [b] vulnerary Balſam, and the Lips of the Wound ſhould be cloſed and kept in that Situation; this is done after different Methods, according to the Difference of the Wound. 1. This is to be obtained by *placing the wounded Part*

[a] *Traité des Inſtrumens*, Tom. I. pag. 391.

[b] Beſides the Medicines, which we have recommended above, at N°. XXIX, we may add here *Eſſentia Succini, Terebinthina, Maſtichis, Myrrha & Aloes, Gemmarum Populi*, &c. We muſt obſerve, too, that where a Contuſion is added to the Wound, which is the Caſe in Wounds made by Glaſs, Saws, *&c.* the mildeſt vulnerary Oils and Balſams are to be applied, as *Ung. Digeſtivum*, or *Balſam. Arcæi*; but in thoſe made by Knives, Swords, *&c.* the Eſſences and Balſams which we have juſt deſcribed are to be preferred, as being more aſtringent and drying.

in

in a proper Poſture; as ſoon as the Wound is dreſſed, the Part ſhould be placed in ſuch a Situation, that the divided Parts may be moſt likely to be in conſtant Contact, repeating the Dreſſings once a Day, as we obſerved before, N°. XXXV, or at leaſt every other Day. 2. by *proper Bandage*; tying up the Parts ſo that the Lips may meet, and ſo eaſily unite. This is attended with the greateſt Succeſs in Wounds that are made lengthways, as in the Fore-head, Abdomen, Arms or Legs; for in theſe Caſes the *uniting Bandage at Plate* II, *Lett.* F. anſwers the End completely. 3. by *a proper Suture*, which differs according to the Difference of the Wound, but may be generally divided into the dry and bloody Suture. The *dry*, or, as ſome call it, the *baſtard Suture*, is the Application of ſticking Plaſters to keep the Lips of the Wound united: the *bloody*, or *true Suture*, is performing the ſame Thing with a Needle and Thread.

What Wounds require a Suture.

XL. All Wounds are not to be united by the Needle; but thoſe only that are oblique, tranſverſe, or angular, and at the ſame Time very large and deep; or in Caſes where a Part is near cut off, as in the Noſe, Ear, Chin, Cheeks, *&c.* if a Wound is ſo circumſtanced, that it cannot be kept in a proper Situation by Plaſters and Bandages. 1. Wounds that are to be ſtitched ſhould be in their recent State, and properly cleanſed from extravaſated Blood, and all extraneous Bodies. 2. There ſhould be no Loſs of Subſtance, except in thoſe fleſhy Parts that are eaſily elongated, as the Lips. 3. There ſhould be no Inflammation or Contuſion. In theſe Caſes the Lips of the Wound are cloſed more elegantly and more ſucceſsfully by Suture: on the contrary, Wounds of long ſtanding, rancid or foul, attended with Venom, or that have their Seat in the Breaſt; or, laſtly, where the larger Arteries, Veins, or Nerves are injured, cannot be ſewed up without imminent Danger.

When and in what Manner, the dry Suture is to be performed.

XLI. The dry Suture is to be uſed in ſlight Wounds, and eſpecially when they happen in the Face, and indeed wherever you think it is of Force enough * to keep the Lips together: as it gives no freſh Pain, and occaſions no Scar; it is much fitter for Wounds of the Face than the Needle, eſpecially as the Needle, beſide the Pain and Scars it occaſions, often produces no ſmall Inflammation. The Plaſters which are to form the dry Suture ſhould be of a ſufficient Length, and ſhaped like the Part to which they are to be applied, ſo as to ſurround the greateſt Part of it; but not the whole, leſt they ſhould retard the Circulation of the Blood, and bring on Tumors and Miſchiefs of that Kind. They muſt alſo ſtick very faſt: which Purpoſe is excellently well anſwered by the *Emplaſtrum* ANDREÆ A CRUCE, *vel Stypticum* CROLLII, *vel Diachylum, vel Diapalmæ, Terebinthinâ probe ſubactum.* The Hæmorrhage being ſtopped, and the Wound well cleanſed, ſome tenacious vulnerary Balſam, ſuch as *Eſſentia Maſtichis, Succini, Balſami Peruviani*; or the *Balſamum Præfecti Equitum Melitenſium*, which you will find deſcribed in LEMERII *Pharmacopœia Univerſalis*, under the Title of *Balſamum Equitis Sancti Victoris.* Theſe, and indeed all Balſams of the gummy Kind, beſt anſwer the Intention in this Place, for they preſently form a ſticky balſamic Cruſt, which denies all Entrance to the Air, and preſently brings on the deſired Union; but over this a ſticking Plaſter is to be laid, adapted to the Size of the Part; you may apply two or more according as

* Where the Finger has been cut almoſt off, ſo as to hang by a Piece of Skin, and the Surgeons have adviſed it to be taken off, I have cured it by this Suture frequently, and the Bones have united.

you

you ſee Occaſion, leaving a Space between. The manner of applying them you will ſee at *Plate* IV, *Fig.* 3, 4, 5. they are to be ſecured in their Situation by the Application of proper Boulſters and Bandages.

Other Methods of making the dry Suture.

XLII. According to PETIT's Method, the ſticking Plaſters ſhould have one, two, or more Openings in the Middle, *See Plate* II, *Fig.* 11. or in the Manner of thoſe at *Plate* IV, *Fig.* 7. that you may diſcover through theſe, as by the Spaces left between, in the former Method, whether the Lips of the Wound were properly united or not: And that you may alſo be able to apply proper Remedies to the Part, without removing the Plaſters. Theſe Plaſters are applied in the ſame Manner as the former, and left on till the Work is completed. But the dry Suture may be formed alſo after another Manner; to wit, make two Plaſters after the Preſcription of ANDREAS A CRUCE, ſpread upon ſtrong Cloth, anſwering in Size to the Wound; to the Sides or Margin of theſe faſten three or four Tape-ſtrings, according to the Length of the Wound, and then, after warming the Plaſters, apply them on each Side of the Wound, about the Diſtance of a Finger's Breadth from it, after the Manner deſcribed at *Plate* IV, *Fig.* 8. After this bring the Lips of the Wound together, dreſs it up in the Manner we have deſcribed above, and whilſt an Aſſiſtant keeps the Lips of the Wound in their proper Situation, let the Surgeon tie the Ends of the Tapes, firſt in a ſingle Knot, and then in a ſlip Knot, to keep the Parts in Contact. Over each ſhould be laid an oblong Compreſs, and over all of them a large ſquare one, the whole to be bound up with a proper Bandage. On the next Day the Wound is to be examined, and if the Tapes are looſened they muſt be drawn tighter again; but if they are not looſened, let them remain untouched, only moiſten the Parts with a few drops of Balſam, covering them up again with the Compreſſes and Bandage as before. If they are too tight, and a violent Inflammation ſucceeds, they may be relaxed at Pleaſure; but on the Decreaſe of the Inflammation they muſt be tightened again. Some, in the Room of Tape, uſe Claſps made of Steel or Braſs, as we have deſcribed them at *Plate* IV, *Fig.* 9. and 10. But this Method is leſs convenient than the former, and therefore in very little Uſe.

The bloody Suture what.

XLIII. In large Wounds, eſpecially tranſverſe ones, as their Lips cannot be maintained in their Situation by the dry Suture, which is frequently the Caſe in Wounds of the Thigh, as you may ſee at *Plate* III, *Fig.* 1, *Letter* H; or in the Abdomen, Nates, or Arms; or where Pieces hang from the wounded Part, as in the Forehead, Cheeks, Noſe, or Ears; or when large Wounds are made in an angular or cruciform Manner, as at *Plate* IV, *Fig.* 12, 13, 17. here you muſt uſe the Needle, which Operation is called the *bloody*, or *true Suture*. The true Suture is diſtinguiſhed again into the *ſimple* and *compound*. The *ſimple Suture* is that which is performed only by the Aſſiſtance of the Needle and Thread; to this Claſs belong the *interrupted Suture*, the *Glover's Suture*, and the *twiſted Suture*. The laſt is ſeldom uſed but in the Hare Lip; the ſecond only in Wounds of the Inteſtines, under which Head we ſhall treat of it more largely; but the firſt is in common uſe for all Wounds that require the true Suture, therefore we ſhall begin with the Deſcription of that before the reſt. The *compound Suture* is that which requires other aſſiſtances beſides the Needle and Thread. Of that below.

How to perform the interrupted Suture.

XLIV. The beſt Method of making the *interrupted* or *knotted Suture*, I take to be the following one: Take a double Thread well waxed, paſs it through a ſtrong

ſtrong crooked Needle, as you may ſee *Plate* I, *Lett.* S, T, V, or *Plate* VI, *Fig.* 5 or 6. When the Lips of the Wound are brought together, and held firm in that Situation by an Aſſiſtant, with one Stroke pierce through them both, paſſing your Needle through the lower Lip from without inwards almoſt to the Bottom, and ſo on from within outwards, obſerving to make the Punctures at a Finger's Breadth from the Wound, (which in this Caſe we will ſuppoſe to be in Length two Fingers) varying this according to the Size of the Wound. After taking off the Needle, tie the Ends of the Thread, firſt in a ſingle Knot, and then in a ſlip Knot, covering all with the Dreſſings which we preſcribe in the dry Suture. But if the Wound is of ſuch a Length, that one Stitch will not be ſufficient, then you may make two, three or more after the ſame Manner that we have now deſcribed, always obſerving a Finger's Breadth Diſtance between each Stitch. *See Plate* IV, *Fig.* 11 and 16. But to prevent the Knots from bringing on any Miſchief, lay a ſmall Linen Compreſs (*See Plate* II, *Fig.* 22) over the ſingle Knot, and make the ſlip Knot over that; which, if any Pain or Inflammation ſhould ſucceed, may be eaſily looſened.

Some Cautions.

XLV. We proceed in this Manner in oblique or tranſverſe Wounds. But where there are Angles, as in a triangular Wound, *Plate* IV, *Fig.* 13. you are to proceed in the ſame Manner as before; only the Suture muſt begin at the Angle A; then the Sides of the Wound muſt be ſtitched about the Middle at B and C. If the Wound is quadrangular, or has two Angles like the *Greek* Letter Π, which ſometimes happens in the Face (*See Plate* IV, *Fig.* 14.) then the Sutures muſt be made in both the Angles A A. But when the Wound is ſo large, that theſe are not ſufficient, then as many more as are neceſſary muſt be made in the middle Way between the Angles B B. When you meet with a cruciform Wound, as at *Fig.* 6 and 12. and the Lips of it cannot be kept in Contact by the Uſe of Plaſters, the Needle, as at *Fig.* 12. muſt be paſſed in at A, and come out again at B; it muſt enter again at C, and come out again at D; the Extremities of the Threads muſt then be tied in the Manner we have before directed, between A and D. How the Wounds are to be treated afterwards we ſhall explain below.

The compound Suture.

XLVI. Some of the Surgeons amongſt the Ancients uſed a *compound* Suture for large Wounds in the Room of the interrupted Suture, which was made of two Pieces of Wood, by the *French* called *Chevelle*; and from thence the Suture was ſtyled *Enchevillée*. And they preferred this, becauſe it prevented the Lips of the Wound from being lacerated, which ſometimes happened when the other Method was uſed, which not only prevented the Wound from uniting, but frequently brought on other grievous Diſorders. And though this Method has of late Years been rejected as inconvenient, and particularly by DIONIS in his Surgery; yet it is not at this Day without its [a] Advocates, who highly commend it, and prefer it to the interrupted Suture in many Caſes: But they uſe it with this Difference, that inſtead of two Pieces of Wood, they uſe Pieces of Plaſter rolled up in a cylindrical Form, of the Length of the Wound, and about the Size of a Gooſe Quill, from whence it is by ſome called the quilled Suture. *See Plate* IV, *Fig.* 17 and 18. This Method prevents Tumours, Pain, and Inflammations, that might be brought on by the Hardneſs and Preſſure of the

[a] As PALFYNUS, in Chirurgiâ, Cap. VI. de Suturis; and ſince, GARENGEOT in Chirurg. Cap. de Suturis.

Wood.

Wood. PALFYNUS performs this Operation, in deep Wounds of the muſcular Parts (as the Thighs, Buttocks, Legs, Arms, *&c.*) with a large ſtrong, crooked Needle, furniſhed with a ſtrong double Thread well waxed (*See Plate* IV, *Fig.* 15.) which makes a Bow at one End. The Needle being paſſed through both Lips of the Wound, in the Manner we have before deſcribed, and a ſecond and third paſſed in the ſame Manner, as is ſhewn at *Fig.* 17. a Roll of Plaſter is to be introduced into the bow Ends of the Thread, which are left hanging out at B B. Then when the Needle is taken off at the other Side, another Roll is to be placed between the Ends of the Thread; and the Lips of the Wound being brought together, theſe Ends are to be gently tied over the Roll, firſt in a ſingle, and then in a ſlip Knot, as at CCC. If there are three Threads, you are to tie the middle firſt, and then the reſt, treating the Wound afterwards, as we ſhall ſhew below.

Another Method by GARENGEOT.

XLVII. GARENGEOT performed this Operation much after the ſame Method we have juſt deſcribed (*See his Book of Operations in Surgery*, Chap. iii. *on Sutures*), but with this Difference, inſtead of a double Thread, he made ſeveral Ligatures of ſix or eight Threads (according to the Size of the Wound) joined together and waxed, always obſerving not to make them ſo big, that when they ſhould be doubled they ſhould exceed the Size of the Needle, leſt they ſhould create Pain, by not paſſing readily after the Needle. When a ſufficient Number of theſe Ligatures are paſſed through the Lips of the Wound, he makes a Knot upon each of the Ends that hang out of the upper Lip; *See Plate* IV, *Fig.* 18. DDD, and then unravels the Threads that compoſe the Ligature, between the Knot and the Lip of the Wound; and by this Means forms a Paſſage through which he can introduce the cylindrical Roll of Plaſter. After this he claps two Fingers upon the lower Lip of the Wound, near the Punctures which were made by the Needle, and with the other Hand draws back the Ligature gently, beginning in the Middle, if there are more than two, till the Wound is exactly cloſed: Then he divides the Threads of each Ligature into [a] two Parts, with which he ties the other Roll as before, nicely joining again the Lips of the Wound. In tying theſe Ends, great Care ſhould be taken not to make the Knots too tight at firſt, leſt they ſhould bring on Pain and Inflammation. The Wound is now to be covered with vulnerary Balſams ſpread on Lint, but eſpecially with the *Balſamum Præfecti Equitum Militenſium*, which I have commended before, as it ſoon forms a healing agglutinating Cruſt, denies all Acceſs to the external Air, and brings on the deſired Union, to which you muſt add a Compreſs, a proper Bandage, and a convenient Poſture of the Part affected.

What is further to be obſerved.

XLVIII. On the firſt Days, after whatever Method the Suture is performed, the Bandage and Compreſs are to be gently removed, and the State of the Wound examined. If every Thing looks well, and there is little or no Pain or Inflammation, the Sutures are to be let alone for ſix or ſeven Days, or longer, and the Wound be dreſſed up again as before, till it appears that there is a ſtrict Union procured. But if the Stitches ſhould appear to be too looſe, the Knots ſhould be tightened; if they are too tight, they muſt be looſened a little. When the Lips of the Wound appear to be enlarged or bruiſed, they ſhould be dreſſed with a digeſtive Ointment, or with the *Balſamum Arcæi*, the con-

[a] GARENGEOT here orders them to be ſeparated into three Parts, but what Uſe he puts the third Part to I can't comprehend; I am apt therefore to imagine, that there is ſome Omiſſion in this Place.

tinuance of which will preſently remove all theſe Symptoms. But when the Wound is attended with great Inflammation and Fever, the Stitches ſhould be ſomewhat looſened, the Patient ſhould be let blood, and live upon a thin Diet, and the Body ſhould be kept open. Theſe Symptoms being removed, the Stitches ſhould be again tightened by degrees and the Wound dreſſed as above. But if theſe Applications ſhould prove fruitleſs, and the Complaints ſhould increaſe, ſo as to threaten Danger, the Stitches muſt be cut, and the Wound treated as if there was a loſs of Subſtance, which Method we ſhall explain below.

What is to be done after the Wound is healed.

XLIX. On the other hand, if the Wound heals by the Aſſiſtance of the Suture, which you will be ſure of, not only from obſerving the Lips of the Wound to lie cloſe together and unite, but by the Relaxation of the Threads or Ligature upon the diſordered Part; the Threads or Ligatures are to be cut near the Knots with Sciſſars, the lower Lip of the Wound is to be ſuſpended with one hand, whilſt the Threads are gently drawn out with the other. The Punctures that are left will eaſily heal by the Application of a vulnerary Water, called by the *French l'Eau d' Arquebuſade*, or by injecting *Aqua Calcis*, or *Spiritus Vini*, and laying on Compreſſes dipped in the ſame Liquors. But larger Wounds are to be dreſſed with one of the beforementioned Balſams, and the Lips kept firm together with ſome ſticking Plaſter, till a firm Cicatrix is formed.

Of healing Wounds where there is loſs of Subſtance.

L. Where there is loſs of Subſtance, the Wound will not unite either by the help of Plaſters or Suture, till it is filled up with new Fleſh. For this Purpoſe you will find Lint dipt in Oil, or ſpread with with ſome vulnerary Ointment or Balſam, and applied to the Bottom of the Wound, very ſerviceable, covering it with a Plaſter, Compreſs, and proper Bandages. This Dreſſing is to be repeated daily: though it is a very vulgar Error, to ſuppoſe that theſe Applications generate Fleſh, which is produced by the circulating Fluids, that in a wonderful Manner are continually bringing ſomething new to the wounded Parts. Yet it muſt be owned that Medicines of this ſort conduce very much to the Generation of the Fleſh, and to remove every thing that might hinder that End; therefore it is no wonder they are called *ſarcotic Medicines*. There ought to be a balſamic and emollient Quality in theſe Medicines, that they may not only reſiſt Putrefaction, but may alſo ſoften the young Fleſh, ſo that it may eaſily receive Additions from the Blood, and ſuffer itſelf to be elongated. Of this kind are the Oils, Balſams, and Ointments, which we took notice of at *N.* XXXV, *and* XXXIX.

How the Air is to be kept from Wounds.

LI. As hot or cold Air is very hurtful to Wounds, ſo it muſt by all Means be kept from them, for nothing will ſooner corrupt the Juices, or ſhorten and dry up the Veſſels, and hinder the growth of new Fleſh, than the Air. In order to prevent Inconveniencies from this Cauſe, the Surgeon ſhould be careful not to remove the old Dreſſings till the freſh ones are got ready, and to be as expeditious as poſſible in applying them. Then the Wound muſt be filled up with Lint, dipped in Oil, or ſome emollient Balſam; to this muſt ſucceed ſome vulnerary Plaſter, a Compreſs upon the Plaſter, and laſt of all a Bandage, to bind and ſecure the whole.

How the Wound ſhould be entirely healed.

LII. After this, when a white, ſmooth thick Matter appears in the Wound, the Wound ſhould be dreſſed, as you ſee occaſion, every Day, or every other Day, the ſuperfluous Matter ſhould be wiped away with a very light Hand,

and

and it is better to leave ſome behind than to treat the Wound roughly. For wiping the Wound roughly hinders the Growth of new Fleſh: but a little Matter being left, performs the Office of Oil or Balſam, keeping the Parts moiſt. Theſe Rules being obſerved, new Fleſh will preſently ſpring up, and the Wound unite.

How a Cicatrix is to be formed.

LIII. But that nothing may be omitted which may ſeem neceſſary towards the perfect Cure of the Wound, the Surgeon ought to be induſtrious to procure an even Cicatrix. To this End it will be proper to dry by degrees, and to harden the Surface of the new Fleſh, by the Application of dry Lint, covered with a tight Compreſs and Bandage. But when this is not ſufficient, through a great Redundancy of Humours, it may be proper to uſe ſome of the drying Eſſences, or native Balſams at *N.* XXXIX, or drying Powders, ſuch as *Tutia*, *Lapis Calaminaris*, *Maſtichis*, or *Colophonium*. Rectified Spirit of Wine is frequently uſed for this Purpoſe with great Advantage, which carries a great aſtringent and drying Virtue with it.

How foul Wounds are to be treated.

LIV. When you perceive any Uncleanneſs or Foulneſs in a Wound, that is, if the Fleſh is putrid, fungous, black, pale, or livid, it muſt be well cleanſed before you attempt to heal. Different Methods have been propoſed to execute this Intention; the Antients uſed Honey in this Caſe, *See* CELSUS, *Lib.* V. *Cap.* 26. *N.* 22. But the Moderns apply a digeſtive Ointment, made *ex Terebinthinâ Vitell. Ov. q. ſ. ſubactâ cum Mell. Roſar. q. v. admiſt.* But where this is not ſtrong enough for their Purpoſe, they ſubſtitute *Unguentum Ægyptiacum, vel Vini Spiritu dilutum, vel digeſtivo admiſtum.* Some in the room of this uſe *Unguentum Euſcum Wurtzii.* To theſe digeſtive Ointments you may very properly add a ſmall Quantity of *Aloës* or *Myrrh*, or, if you require ſtill more Strength, *Mercurius præcipitatus ruber.* But the uſe of *Aqua Calcis* is well known to be very beneficial as a Detergent, eſpecially if you add to a Pint of this *Mercurii ſublimati gr.* xx. *vel.* xxx. which, from its known efficacy for this Intention, is called by the Surgeons *Aqua Phagædenica.* Applications of this kind are to be continued till the Wound is entirely clean; and then you are to have recourſe to the vulnerary Balſams, and the Method preſcribed at *N.* L.

How fungous Fleſh is to be taken down.

LV. If the new Fleſh ſhould be luxuriant, and riſe up ſo as to prevent the Formation of an even Cicatrix, it muſt be taken down by the *Vitriolum Cæruleum*; or in the room of this you may uſe a Powder compoſed *ex Alumine uſto, Mercurioque rubro præcipitato*; at the ſame time making a proper Preſſure with the Plaſters, Compreſſes, and Bandages, till the Parts are even.

Rules to be obſerved by the Patient.

LVI. The Patient ſhould particularly obſerve a ſtrict Regimen, with regard to his Diet and manner of Living; that by avoiding every Thing, that produces Crudities or Acrimony, the Blood may be rendered pure and uncorrupted. For nothing forwards the Cure ſo much as a good Habit of Body; which may be procured by obſerving a ſtrict regularity with regard to Diet, conſulting which is the moſt proper Air to live in, keeping the Paſſions under, and neither indulging in too much Sleep, nor ſuffering too great Watchfulneſs. The greater Tendency there is in a Patient to a diſeaſed State of Body, ſo much the ſtricter Courſe of Life ought he to obſerve.

What Air is beſt.

LVII. As to the Air, it ought to be temperate, and the Chamber ſhould be equally guarded from Exceſſes either of Heat or Cold; for this Regulation is of Conſequence in all Wounds, but moſt wonderfully ſo in thoſe of the Head.

If

If the Patient is in any Danger of ſuffering from the Dampneſs of his Situation, it will be very proper to burn Amber, Frankincenſe, and Maſtich round him, to dry the Chamber. If he is in Danger from the Heat of his Situation, the Floor ſhould be frequently ſprinkled with Water.

What Diet is beſt.

LVIII. All Intemperance in Eating and Drinking is moſt diligently to be avoided. That Sort of Food is beſt which is moſt readily digeſted, for it makes a thin light Chyle and good Blood, which wonderfully aſſiſts the Wound in healing. For this Intention various Sorts of Broths may be recommended to the Patient, particularly thoſe that are made *ex Hordeo, Avenâ, Mannâ, Oryzâ, Scorzonerâ, Lactucâ, Endiviâ, Chærophyllo, Petroſelino, Cichorio, Aſparago.* He may eat Veal or Lamb, Pullets or Capons, Ale thickened with the Yolks of Eggs, ripe Fruits, particularly Apples, Cherries, or Plumbs; Vegetables alſo of ſeveral ſorts well boiled, to wit, *Spinachia, Lupulus, Aſparagi, Cinaræ, Lactucæ,* and moſt Pot-herbs. But Perſons of ſtrong athletic Conſtitutions, that cannot be ſatisfied with Diet of this Kind, may be indulged in a more nouriſhing one, if they are attended with no violent Symptoms: But wherever there is any Degree of Inflammation, the Patient muſt entirely abſtain from Fleſh, and all ſolid Food. Wounded Perſons ſhould conſtantly avoid admitting any thing *ſharp, ſalt,* or *ſpicy* into their Diet: For they give a Sharpneſs to the Blood, and increaſe its Heat and Motion, and conſequently occaſion Hæmorrhages, Fevers, and Inflammations. They ſhould therefore abſtain, eſpecially if they are of a hot Conſtitution, from every Thing that is ſeaſoned, from Muſtard, Horſe-radiſh, and Onions. *All Meats that are difficult to digeſt, and breed a thick Blood,* ſhould alſo be denied; ſuch are all fat Meats, Lard, Bacon, Geeſe, Beef, either ſalted or cured in the Smoke, Peas, Beans, and Lentils, eſpecially after they are dried, and all Things of this Kind.

What Drink is beſt.

LIX. The Patient's *common Drink* ſhould never be ſtrong: Therefore he ſhould be forbid the Uſe of Wine, Spirituous Liquors, Mead, Strong Beer, *&c.* The ſmaller his Drink is, by ſo much is it the wholſomer. But in this Caſe we muſt always have a Regard to the Conſtitution and Cuſtom of the Patient, and the Nature of the Wound. If he has been uſed to drink Water, he may go on in the conſtant Uſe of it, or drink in its Stead a Decoction of Bread or Barley, mixed with Liquorice, Anniſeed, Fennel, or Citron Peel. Thoſe who diſlike Water may be indulged in good Small Beer, that is neither too new nor too ſtale: But if the Patient is in great Danger, and of a weak Habit of Body, you may preſcribe him a particular *vulnerary Drink*, to correct the vitiated Fluids. But of theſe we ſhall treat more largely below, at *N.* LXIII, *and* LXIV.

Of Reſt, Motion, Watchfulneſs, and Sleep.

LX. The beſt Remedy for a wounded Perſon is *Reſt.* Therefore he ſhould be indulged in it, eſpecially with regard to the lower Limbs: For to walk, or even to move, is very pernicious. There are many Inſtances of wounded Perſons, who have not only ſuffered grievous Injuries, but even Death itſelf, by violent Motions of the Body. Nor is too great Watchfulneſs of leſs Conſequence to the Patient: Therefore, if Nature denies neceſſary Reſt, it muſt be procured by the Aſſiſtance of Medicines. To anſwer this Intention you may very properly preſcribe *Syrupi Papaveris albi* ℥ ſs *ad* ℥ i *ex Aq. Primulæ veris, vel Ceraſorum nigror. vel ex Emulſione Papaveris Semine, & Amygdalis dulcibus confecta.* When this appears to be too weak for the deſired End, you may give *Theriaca Venet.*

Venet. vel Confectio Mithridatis ad ʒi. *vel* ʒii. *vel Opii puri gr:* i. in one of the Vehicles we mentioned above.

LXI. *The Bowels ſhould by all Means be kept open*, eſpecially in thoſe who have received a Wound in the Head, for they are ſubject to great Heat of Body, and are very apt to be bound. But obſerve in this Place, that ſtrong cathartic Medicines are to be avoided, for in ſo weak a State of Health they are of very ill Conſequence. But it is not only ſafe, but adviſeable to eat and drink thoſe Things, that may at the ſame Time nouriſh and keep open the Body. To this End the Patient may drink plentifully of *Tea* or *Coffee*, or may eat ſtewed Prunes; roaſted Apples alſo and Raiſins may be eaten for the ſame Purpoſe: but hard Meats of all Kinds are to be forbid. Where the Patient is ſo bound up, that a Diet of this Kind has no Effect upon him, it will be neceſſary to have recourſe to Medicines, but to thoſe of the mildeſt Kind: you may here give a gentle Clyſter, or uſe a Suppoſitory, or preſcribe a Solution of an Ounce or two of *Manna, or ſome purging Salts*, in warm Broth, or a Draught compoſed of Tamarinds, Sena and Manna. But you muſt carefully avoid all reſinous and heating Medicines. The Bowels ſhould be kept open.

LXII. *Violent Paſſions of the Mind*; ſuch as Anger, Fear, Sorrow, Penſiveneſs, and particularly Luſt, ſhould diligently be avoided; and a quiet, ſerene, eaſy, chearful State of Mind preſerved; the contrary of which will never fail to bring on dangerous Symptoms. The Mind ſhould be free from Care.

LXIII. Whenever the Violence of the Wound, or the ill Habit of the Patient, require the uſe of *internal Remedies*, vulnerary Drinks will be found to be of the greateſt Conſequence in this Place; in compoſing of which, the Conſtitution of the Patient, and the Nature of the Complaint, ſhould be diligently conſulted: For they are in a great Error, who, according to the Cuſtom of common Surgeons, give one Kind of vulnerary Potion for all ſorts of Wounds, and in all Habits of Body. For if your Patient is of a phlegmatic Habit of Body, cold, pale, naturally ſubject to Tumours, then the vulnerary Decoction ſhould be compoſed of Herbs that will attenuate and divide the Blood; ſuch as the *Radices quinque aperientes, Rad. Caryophyllat. Fœnicul. Gramin. &c. Herb. Sanicul. Alchymyll. Agrimon. Betonic. Veronic. Philoſellæ, Pervinc. Virgæ Aureæ, Sophiæ Chirurgorum, Semen Aniſi, Fœnicul. Dauci, &c.* The Drink is preſcribed in the following Manner: Take two or three Handfuls of any of the beforementioned Ingredients, boil them gently for a few Moments in ſix Pints of Water, ſtrain it, and ſweeten it with ſome proper Syrup, ſuch as the *Syrupus Tunic. Betonic. Capill. Ven. Rad. quinque Aperient. De Cinnam. &c.* Give a Draught of this three or four times in a Day. You may alſo give Infuſions of the ſame Herbs, and made after the manner of Tea, ſweetened with Sugar. What internal Medicines are to be given.

LXIV. Some Perſons have a thin, ſharp Blood; in this Caſe, it will be proper to adviſe Decoctions of viſcous glutinous Plants; ſuch as the *Rad. Symphyt. Liquorit. Polypod. Scorzoner. Sarſaparill.*; *Herb. Malv. Althææ, Verbaſc. Parietar. Mercurial.*; *Flor. Malv. Althææ, Verbaſc. Dactyli, Ficus, Jujubæ*; which may be prepared in the Manner we have juſt deſcribed, with the Addition of ſome of the *Syrup. Alth. vel de Symphyt. Liquoritiæ, vel Papaveris*, to give it an agreeable Taſte, if the Patient have no Averſion to Sweets. But if he is afflicted with great Pain or Wakefulneſs, then, beſides the Methods which we lay down at *Chap.* II. to alleviate Pain, you may give an Ounce or two of the Vulnerary Drinks for a thin, ſharp Blood.

Syrupus

Syrupus Papaveris albi vel de Meconio, mixed with the beforementioned vulnerary Drink, or with Emulſions *ex Amygdalis & papavere albo.*

Remedies againſt Acrimony and Heat.

LXV. If the Patient ſhould be troubled with any Acidity, you may give him Powders every Day *ex Lapid. Cancrorum, vel ex Matre Perlarum, vel ex Conchis præparatis*, or any other Abſorbents. But when you perceive a Quickneſs of Pulſe, and an extraordinary Heat, they are ſure Signs of a Symptomatical Fever: To relieve or take off which, the following Remedies will be found of Service. Give Barley Water with the Addition of ſome Tamarinds, and *Syrupus Mali Citrei vel Ribeſiorum*; or ſome of the Powders mentioned above, ſaturated with Citron Juice, with the Addition of a ſmall Quantity of Nitre. But in this Place it will be very proper for the Patient to loſe ſome Blood, more particularly if he is young and full of Blood, or if the Pulſe is ſtrong and hard. In theſe Circumſtances a *Phyſician* is more proper to be conſulted than a *Surgeon.* But if the Patient is robuſt, and of a ſound Habit, the beſt common Drink that can be preſcribed, is Barley Water, or good Small Beer. What has here been ſaid with regard to the Regimen to be obſerved by the Patient, as well with reſpect to Diet as Medicine, I think is ſufficient; and I heartily recommend the Obſervance of theſe Rules to all wounded Perſons, but more particularly to thoſe who are to undergo ſevere Operations in Surgery; ſuch as Trepanning, Lithotomy, Extirpation of the Breaſt, Amputation of a Limb, or large Tumors. Whenever we ſhall have Occaſion below to ſpeak of the Regularity that Patients ought to obſerve in their Diet, I hope the Reader will endeavour to recollect what has been ſaid upon that Head, that we may not be obliged to make tedious Repetitions.

CHAP. II.

Of the Diſorders accompanying WOUNDS, *commonly called the* SYMPTOMS *of* WOUNDS.

I. *Of an* HÆMORRHAGE.

An Hæmorrhage how to be ſtopped.

I. PROFUSIONS of Blood attending Wounds, all ariſe from Injuries of the Veins or Arteries. The Violence of the Hæmorrhage will be in Proportion to the Size of the wounded Veſſel. Whoever conſiders this, will no longer wonder at the dreadful Conſequences attending this Symptom, unleſs there be immediate Aſſiſtance; ſuch as great Weakneſs, fainting Fits, and ſometimes inſtant Death. No Surgeon therefore ought to be without a preſent Remedy to ſtop Blood. Though there are ſome Caſes where it is by no means proper to reſtrain the Hæmorrhage inſtantly: For in a young, plethoric Habit, or where the Wound has been received in a drunken Fit, or in a Fit of Paſſion, it is beſt to let the Blood run, as long as it continues to do ſo without bringing on any Inconvenience upon the Patient: For by a moderate Loſs of Blood, the Inflammation, Tumor, Pain, and Fever are prevented, or much leſſened.

1. By dry Lint.

II. There are various Methods propoſed to ſtop an Hæmorrhage. If none of the larger Veſſels are wounded, you have your Remedy at hand, to wit,

ary

dry Lint, which you are to fill the Wound with pretty closely, covering it over with large Compresses, and making a proper Degree of Pressure over all with Bandages, and with your Hands: For more Service is frequently done in this Case by making a proper Pressure upon the Part with the Dressings, and with your Hands, than could be effected with more violent Remedies. But you must avoid too strict a Pressure; which often produces violent Pains, Inflammation, and at last even a Gangrene.

III. But if the Hæmorrhage is too large to be stopped by the Application of dry Lint, then *astringent Medicines* are to be called into Use. With this Intention the Ancients applied Rags to the Wound, which were dipped in cold Water or Vinegar, and covered them with Compresses wet with the same Liquors. Amongst the Surgeons of later Date, a certain Fungus called *Lycoperdon*, or vulgarly *Lupi Crepitus*, has been highly extolled for this Purpose: The Wound is filled with this in the room of dry Lint, and afterwards dressed up in the same manner as we directed above. The most common Remedy at present is *Spirit of Wine highly rectified*; this is applied cold to the Wound, filling it up with Dossils dipped in the same Spirit, and covering it with large Compresses wrung out of the same Liquor, making a proper Pressure over all with the Bandage. The same Virtues used to be ascribed to *Oil and Spirit of Turpentine*, applied in the same Manner as the Spirit of Wine. To this End also strong Solutions of *Alum*, *Vitriol*, or *Saccharum Saturni in Aquâ Plantaginis* were recommended by many. Some dissolved Alum and the Vitriol together in the same Water, or, where they would have it of more Force, in Phlegm of Vitriol. Others make a styptic Liquor *ex Vitriol Alb.* ℥ i, & *Aceti fortissimi* ℥ iii, applying it in the foregoing Manner. In this Place we are by no Means to omit the Mention of *astringent Powders:* such as are made *ex Bolo Armenâ*, *Lapide Hæmatite*, *Sanguine Draconis*, *Croco Martis astringente*, *Terrâ Japonicâ*, *Aloë*, *Olibano*, *Mastiche*, *Granat*, *Corticibus*, *Alumine*, *Saccharo Saturni*, *Terra Vitrioli dulci*, *Gipso*, *Hepate Vitulino tosto*, and several other Medicines of this Kind, either alone or mixed in different Proportions, and sprinkled plentifully upon the Wounds, dressing them up with Lint, Compresses, and Bandages as above. 2. By astringent Medicines.

IV. When Vessels of a larger Size are divided, it is usual to apply *caustic Medicines*, which act by their great Astringency. The Medicine chiefly used with this Intention, and indeed the safest is *Vitriolum Romanum*, which being coarsely powdered and sprinkled upon Cotton, is easily applied to the Wound, dressing up with Dossils, Compresses and Bandage. The *Liquor Stypticus Weberi* is also used here, and others of the like Kind, which have Oil of Vitriol in their Composition: But those Dressings only which are applied to the Bottom of the Wound are to convey these Medicines, otherwise the neighbouring Parts would suffer too great Corrosion. Those Medicines which are endued with a stronger caustic Quality than these, such as *Mercurius Sublimatus*, *Lapis Causticus*, *Oleum Vitrioli*, *&c.* can never be used with Safety, because they are constantly attended with violent Symptoms, from their too corrosive Faculty. 3. By caustic Medicines.

V. But if these Applications prove fruitless, it will be proper to *divide entirely the Arteries which are only divided in part*, and occasion the Hæmorrhage: For by this means they will contract and hide themselves under the muscular Flesh, and the Orifices will be choaked up; at least they will more readily 4. By dividing the Arteries.

yield to the Force of the Medicines recited above. This Method of Treatment is principally neceſſary in Wounds of the temporal Arteries, and of thoſe of the Cubits and Tibiæ.

5. By the actual Cautery.

VI. If this Method ſhould alſo fail, you muſt have Recourſe to the *actual Cautery*. The Orifices of the Veſſels being burned, a Cruſt is formed over them, and this Method is ſo effectual, that it is ſcarce poſſible for an Hæmorrhage to happen in Wounds of the external Parts, but what may be ſtopped by it. You ſhould in this Caſe always have two Cauteries ready, that if one ſhould be extinguiſhed before the Operation is finiſhed, you may be prepared with another. Cauteries are made of very different Shapes and Sizes, according to the Parts to which they are to be applied: I have given you eight different Sorts, for different Uſes, in *Plate* III, *Fig.* 9 to 16. There are two Inconveniencies which generally attend the Uſe of the Cautery, and ſometimes force us to neglect it. For firſt, not only the Patient is wonderfully terrified at the Apprehenſion of it, but Mankind in general look upon it as a Piece of Barbarity to adviſe the Uſe of it: When, to ſay the Truth, it does not occaſion ſuch violent Pains as are uſually apprehended from it; and what Pain there is in the Operation, is inſtantly over. But it is alſo attended with another Inconvenience of greater Conſequence; that is, the Eſchar, which is brought on by the Cautery, frequently falls off in two or three Days, eſpecially in the larger Arteries, from whence a freſh Hæmorrhage ſucceeds, and moſt likely a deadly one. To prevent this, two things are to be obſerved: Firſt to handle the Wounds tenderly at the Time of dreſſing; and ſecondly to be provided always with a freſh Cautery, to repeat the Operation if neceſſary. This Caution is to be obſerved in the larger Arteries for fourteen Days: After this there is no great Danger of a Return of the Complaint. But where the crural or axillary Arteries are wounded, the Cautery will be of no Service.

6. By Ligature.

VII. In very dangerous Wounds of the large Arteries, ſuch as the crural and axillary, and in Amputations of the Limbs, the ſafeſt Method is that of making a *Ligature* round the Veſſels. If this is performed by paſſing a ſtrong waxed Thread under the Artery by the Help of a crooked Needle, the Blood is preſently ſtopped, and the Orifices of the Artery coaleſce. Or it is ſometimes taken up with a Forceps, the Thread wound round it, and the Artery is compreſſed.

7. By Inſtruments.

VIII. Laſtly ſeveral Inſtruments have been contrived to ſtop Hæmorrhages in different Parts of the Body. Formerly a large Iron Ring, furniſhed with a Screw, was in great Uſe among the Surgeons; which they applied in ſuch a manner to the wounded Limbs, that by tightening the Screw which preſſed upon the Compreſſes, and other Dreſſings, it cloſed the Mouths of the Veſſels, and ſtopped the Flux of Blood. You may ſee Deſcriptions of this Inſtrument in SCULTETUS. But as this was a very inconvenient Inſtrument, and could only be applied to the Limbs, the Surgeons found themſelves under a Neceſſity of inventing a more convenient Inſtrument, that might be applicable alſo to the Arteries of the Neck or Head. An Inſtrument of this Kind you may ſee in *Plate* V, *at Fig.* 2. the Conſtruction of which is as follows; A Braſs Plate of three Fingers in Length, and two in Breadth, AA, is perforated in the Middle to admit a ſtrong Screw, BB, which is provided at the lower End with a ſmall round Plate, C: a Piece of Leather is ſtrongly faſtened to one End of the Braſs

Plate,

BOSTON MEDICAL LIBRARY DEC 15 1916

Plate, of equal Breadth with it; EE, FF. In violent Hæmorrhages this Inſtrument is fitted to the wounded Part, and the End F is, by Means of Holes that are made in it, faſtened to the Hooks GG; ſo that the ſmall Plate C may preſs exactly upon the Compreſſes and Dreſſings that cover the Wound. All things being thus prepared, the Handle of the Screw, D, is to be turned round gently till a ſufficient Preſſure is made to ſtop the Blood; and then it is to be left in that Condition for a Day or two. But it muſt be entirely left to the Diſcretion of the Surgeon, when he ſhall think it prudent to alter the Poſition, or entirely to take off the Inſtrument. An Inſtrument of this Kind, with a longer Belt, will ſerve in Wounds of the Head and Temples.

IX. When we are ſpeaking of Inſtruments that are uſed to ſuppreſs Hæmorrhages, we muſt not omit the *Tournequet*, which we uſe with great Succeſs after Amputations. There are ſeveral Things required to form this properly. The firſt thing to be enquired after is a ſmall Roller of a Thumb's Breadth, and about a *Paris* Ell in Length; in the next Place a little cylindrical Stick; then a conglomerated Bandage, two Fingers thick and four long; ſome Compreſſes of a good Length, and about three or four Fingers in Breadth, to ſurround the Legs or Arms. Laſtly, a ſquare Piece of ſtrong Paper or Leather, about four Fingers wide. 8. By the *Tournequet*.

X. We are now acquainted with the Nature of the *Tournequet*. It remains that we enquire which is the moſt convenient Manner of applying it. The rolled Bandage is to be applied to the Trunk of the wounded Artery lengthways, covering it in a contrary Direction with Compreſſes, ſurrounding the Leg, Foot, or Arm, as it were with a Ring. The Roller muſt be paſſed twice round theſe Applications, and faſtened in a Knot, but ſo looſely, that you may eaſily introduce your Hand between it and the injured Part: The Leather or thick Paper muſt be nicely placed under it upon the external Part of the Leg, or Arm, and the Roller tightened by Degrees by turning the Stick round, (which is to be introduced into the Knot) till the Hæmorrhage is entirely ſtopped. The Stick muſt be kept in this Situation till the Wound is properly treated, and the Return of the Hæmorrhage prevented by proper Remedies, or by taking off the Limb. When this End is acquired, the *Tournequet* is to be looſened, or entirely taken off, as the Surgeon ſhall think convenient. But where it is applied to the Arm, the rolled Bandage is to be placed near the *Axilla*, in the internal Part of the *Humerus*, and the Stick in this Caſe is to be faſtened on the oppoſite Side; the Situation of the Artery requires this Poſition, *See Plate* III, *Fig.* 1. *Letter* K. When the Hæmorrhage happens in the Thigh, the Bandage is to be applied to the upper Part of the Thigh, or juſt over the Knee, as the Circumſtances ſhall require, in the ſame Manner as before, *See Lett.* L, M, N. But that you may have a clearer Idea of the Figure and Poſition of the *Tournequet*, we have given you a Draught of it, at *Plate* III, *Fig.* 2. How the *Tournequet* is to be appointed.

XI. PETIT, a Surgeon of the firſt Rank in *Paris*, invented another *Tournequet* in the room of this, which is well enough known by the Name of the Inventor. It is ſaid to have this Advantage over the other, that it will preſerve its Situation without requiring the Attendance of an Aſſiſtant: And beſides, that it may be left upon the Limb any given Time, without impeding the Circulation of the Blood: Whereas the common one entirely interrupts the Circulation of the Blood, and therefore cannot be kept on long. The Deſcription that I have ſeen of it is ſo ſhort and imperfect, eſpecially as the Parts of which it is compoſed PETIT's *Tournequet*.

are not deſcribed ſeparately, that in many Places I could not underſtand it. GARENGEOT, *Tom.* II. *de Inſtrument. Chirurg.* differs a little in his Deſcription of it, but he is by no means clear.

Our Improvement upon it.

XII. Therefore I have taken ſome Pains to correct it in the Manner you may ſee at *Plate* V, *Fig.* 6. A A deſcribes the upper Part, B B the lower, and C the Screw; all in their natural Size, made of ſome ſtrong Wood. At the Extremity D D there are two ſmall Iron Screws, to which a ſtrong Silk Roller is to be fixed, of the ſame width with the Inſtrument, but about twenty Fingers in length, that it may be long enough to encompaſs the largeſt Part of the Limbs, and be faſtened at the ſmall Hooks deſcribed at E. Both Extremities at F F are to be hollowed, that the Roller may lye quiet and firm. G deſcribes an Iron Plate which is placed there to ſtrengthen the Wood. The Wound therefore being properly dreſſed up, and the lower Part of the *Tournequet* guarded with a Bolſter, is to be placed on the Side oppoſite the Wound; the Silk Roller is to be brought round the Limb, and being drawn very tight, is to be fixed to the Hooks E; and then by turning the Screw C, till a ſufficient Preſſure is made upon the Parts to ſtop the Flux of Blood, it muſt be left upon the Limb in this Situation, as long as the Surgeon ſhall deem it neceſſary. By means of this *Tournequet*, I have ſtopped an Hæmorrhage in a Wound of the crural Artery, and recovered the Patient without any Ligature or Amputation, in the preſence of many Witneſſes.

MORAND's *Tournequet.*

XIII. GARENGEOT, in the ſecond Edition of his Book of Chirurgical Inſtruments, deſcribes another *Tournequet*, invented by MORAND of *Paris*, of which he has given us a *Plate at page* 360. This reſembles the former in many Circumſtances, but differs from it chiefly in this, that in the room of a ſimple Screw, MORAND has ſubſtituted a compound Screw, that takes Place ſooner; this he makes always of Steel; and it acts more in one Turn than the other can in two or three. This you may ſee more largely deſcribed, if you conſult the Author himſelf. But GARENGEOT makes ſome Exceptions to this Inſtrument, and prefers PETIT's.

Another.

XIV. Some Years ſince, when I attended the Army, I was called to an Officer of Rank, who was dangerouſly wounded. I ſaw there a kind of *Tournequet* made of Iron, and very heavy, that much reſembled MORAND's, but differed from it in ſome things, I do not know by whoſe Direction; But as I have never ſeen it deſcribed before, I have given you a Plate of it, *See Plate* V, *Fig.* 7. A is the lower Part pierced all round the Edges with ſeveral *Foramina*, by which means it will admit of a Bolſter or Cuſhion to be ſewed to it. B is an Iron Barrel to receive the Screw. C is the upper Part. D is another Barrel fixed upon that, for the Reception alſo of the Screw. E E are the Extremities of the upper Plate, one of which is ſupplied with ſmall Hooks, the other with large Hooks, and with an Opening alſo to paſs the Roller through and faſten it, almoſt in the manner we have deſcribed it in ours of *Fig.* 2. *and* 6. F is a kind of Ring, ſurrounding the Screw, above the upper Plate. G is a ſquare Body made like a female Screw, for the Reception of the ſmall Screw H, and the great Screw I K, which would otherwiſe fall down, but by this means is eaſily kept up in the Box D. L is an Iron Cylinder, which is firmly fixed in the lower Plate, but is looſe in the upper: This keeps the two Plates in the ſame Situation, with

with each other, and at the ſame Time admits the upper Plate to ſlide up and down freely, as occaſion ſhall require.

XV. I endeavoured to improve this Inſtrument, and ordered one to be made of Braſs after the Manner deſcribed at *Plate* VI, *Fig.* 1. In this the upper Plate A A, is much ſhorter than the lower C C; the Belt D D, is fixed at one End, and after it has been brought round the Limb, is faſtened to the other End by Holes made for that Purpoſe F F. The Inſtrument is by this Contrivance always kept even, and does not change its Poſture upon the Action of the Screw B. The Reader may chuſe which of theſe Inſtruments he thinks fitteſt for his Purpoſe; they will all anſwer the Intention they were made for; one does it ſooner, the other takes a little more Time. But this Proverb will always have its Force, *Sat citò, ſi ſat benè.* How the *Tournequet* is to be applied in Amputations of the larger Limbs, we ſhall ſhew in the proper Place. An Amendment of my own.

XVI. Before we take Leave of this Article, it may be proper to inform you, that in Wounds of the large Arteries, *the internal Uſe of aſtringent Medicines will be of no Service*; beſides, they frequently occaſion Pain, Inflammation, Fever, and other Diſorders, by making Obſtructions in the Lacteals, Meſenteric Glands, and other Veſſels; therefore it is beſt to lay them entirely aſide. What is to be done by internal Aſtringents.

II. *Of* PAIN *in* WOUNDS.

XVII. Pain may be reckoned amongſt the moſt grievous Symptoms that uſually attend Wounds: For great Watchfulneſs, Weakneſs, Convulſions, Inflammations, Gangrene, and even Death itſelf, ariſe frequently from this Cauſe. The *Cauſes* of Pain are many. 1. Sometimes an extraneous Body is left in the Wound, which occaſions great Irritations, eſpecially in nervous Parts of the Body. 2. Corroſive Medicines, which are ſometimes applied to ſtop the Hæmorrhage. 3. Or a large Obſtruction of the Blood may happen near the Wound, and bring on Tumor and Inflammation: This frequently happens in Plethoric Habits of Body, or in Gun-ſhot Wounds, becauſe in theſe Wounds there is uſually but a ſmall Diſcharge of Blood. 4. Laſtly, Wounds, or Tenſion of Nerves or Tendons may well be reckoned amongſt the principal Cauſes of Pain. Of Pain in Wounds.

XVIII. It will be well worth our while to conſult the Cauſe of Pain, that we may remedy it with the greater Eaſe: For all Pain will not admit of the ſame Remedy. Therefore, 1. *If any extraneous Body is left in the Wound*, the firſt Intention is to remove it, in the Manner we taught at *N.* XXXI, XXXII, XXXIII. 2. *If the Pain ariſes from the Application of any corroſive or aſtringent Medicine*, it muſt be removed, or at leaſt moderated. This Intention will be anſwered by warm Milk, the Decoctions *ex Malvâ, Althæâ, Floribus Chamæmel. Sambuc. Melilot. Verbaſc. Sem. Lin. Papav. &c.* The Wound ſhould be cleanſed with a Sponge expreſſed from Decoctions of this Kind, till nothing corroſive remains in it, and till the Pain is removed. Cataplaſms may be applied warm to the Wound, made of the foregoing Herbs. There are other Medicines alſo which Phyſicians preſcribe to be given internally to aſſuage Pain, as anodyne Emulſions. 3. *When the Pain ariſes from the Violence of the Inflammation*, which is frequently the Caſe, it will be proper to bleed as largely as the Strength of the Patient will allow: But if you cannot draw a ſufficient Quantity, you muſt ſcarify the Part, as near the Wound as is convenient, eſpecially in Gun-ſhot Wounds. Of Remedies for Pain.

By

By this Method the ſtagnaing Blood is ſet at Liberty, and the Inflammation and Pain are inſtantly relieved. In the mean Time, you may foment the Wound *cum Oxycrato vel Spiritu Vini Camphorati*; or, which is much better, *cum Aquâ Calcis Vivæ modica portione Spiritus Vini Camphorati commiſt.* Emollient Cataplaſms, and ſuch Applications as we ſhall more largely treat of when we come to ſpeak expreſsly of Inflammations, take place here. Abſorbents ſhould be taken inwardly, ſuch as *Lapis Cancrorum, Conchæ præparatæ, Antimonium Diaphoreticum*, mixed with a moderate Proportion of *Nitre.* All Things ſhould be forbid that increaſe the Circulation. Laſtly, 4. *Where the Pain ariſes from an Injury of the Tendon or Nerve*, the Cure is very difficult: For this Caſe is always attended with violent Inflammations and Convulſions. To prevent ill Conſequences that may happen in Wounds of this kind, it will be proper to dreſs with *Balſam. Peruv. Balſam. Copaib. Ol. Terebinth. vel cum miſturâ ex Ol. Terebinth. & Aq. Regin. Hungar. confect.* Theſe Medicines ſhould be moderately warmed before they are applied to the Wound, laying a Cataplaſm over the Dreſſings, compoſed *ex Herb. Scord. Abſinth. Abrotani, Flor. Sambuc. Chamæmel. &c. Vin. q. ſ. decoctis.* Internal *antiſpaſmodic* Medicines ſhould by no means be neglected in this Caſe. If the Pain is not leſſened by theſe Remedies, there is great Reaſon to deſpair, unleſs the wounded Part of the Nerve be inſtantly divided; for although this Method deprives all the Part of the Limb that lyes below the Diviſion of the Nerve of Senſe and Motion, yet in ſuch a deſperate Caſe it is better to loſe the uſe of a Limb than Life itſelf.

III. *Of* SPASMS *and* CONVULSIONS.

Convulſions from whence.

XIX. Spaſms and Convulſions are brought on many ways: For they not only ariſe from all the Cauſes that occaſion Pain, but frequently from too great loſs of Blood. This appears from the many Examples of Men, and other Animals, that have died by the Violence of the Hæmorrhage. All theſe before they expire fall into ſtrong Convulſions and Diſtenſions of the Nerves. HIPPOCRATES mentions this as the worſt of Prognoſtics, *Aph.* 2. *Sect.* 5. "Convulſions "ſucceeding a Wound are mortal."

How Convulſions are to be cured.

XX. In order to remedy theſe Diſorders, it is neceſſary firſt to diſcover their Cauſe. Whenever Convulſions are occaſioned by extraneous Bodies, by corroſive Medicines, or by wounded Nerves, the ſame Methods are to be followed, which we adviſed for the Relief of Pain from the ſame Cauſes at *N.* XVIII. If they are occaſioned *by Inflammation or Fullneſs of Blood*, Blood-letting will generally bring Relief, eſpecially if we uſe at the ſame time the emollient Remedies adviſed at *N.* XVIII. If they are occaſioned *by an immoderate loſs of Blood*, Blood-letting is to be avoided, notwithſtanding ſome amongſt the *French* adviſe it in convulſive Diſorders ariſing from what Cauſe ſoever, *See* GARENGEOT, *in his Chirurgie*, Chap. 2. In this Caſe it will be better, by the Methods before adviſed, to ſtop the Blood, and to give the Patient warm Broth, warm Milk, and Draughts of warm Ale thickened with Yolks of Eggs, and ſweetened with Sugar; by this Method the Veſſels are filled again by degrees, and the Cauſe ceaſing, the Convulſions go off. In the mean time ſtrengthening Medicines ſhould by no means be neglected, particularly [a] Wine, Emulſions, and ſtrengthening Drinks.

[a] See CELSUS, B. V. Ch. 26. N. 25.

IV. *Of*

IV. *Of the* SYMPTOMATICAL FEVER.

XXI. If the Patient has a quick Pulſe, and an increaſed Heat, we ſay he has a *ſymptomatic Fever*. This Fever is of very dangerous Conſequence, and will quickly prove mortal, if not timely relieved by the Aſſiſtance of the Phyſician. Symptomatic Fever, what.

XXII. In order to cure Fevers of this Sort, the Phyſician ſhould forbid the Uſe of every Thing, both in Medicine and Diet, that may encreaſe the Heat; and order ſmall Liquors to be drank plentifully, ſuch as Barley Water, thin Gruels, Ptiſans, *&c.* cooling Powders mixed with Nitre and Camphire ſhould be preſcribed. The Bowels ſhould be kept open with Clyſters, if they do not anſwer naturally. Where the Patient has loſt but a ſmall Quantity of Blood and is of a plethoric Habit, it will be right to open a Vein on the oppoſite Side to the Part wounded. A very thin Diet is to be adviſed, as *Decoct. Hordei*, &c. and in ſmall Quantities at a Time. Fleſh, and all ſolid Diet and Spices, ſhould be abſolutely forbid. Cure of the Symptomatic Fever.

CHAP. III.

Of GUN-SHOT WOUNDS.

I. GUN-SHOT Wounds are attended with much worſe Conſequences than Wounds that are made by ſharp Inſtruments: For, beſides the Effuſion of Blood, the Parts are more ſhattered and torn, eſpecially when the Shot falls upon the Joints, Bones, or any conſiderable Part. Gun-ſhot Wounds.

II. The Antients were entirely unacquainted with Wounds of this Sort, as they fought chiefly with ſharp Weapons, as Swords and Spears; or with Clubs, *&c.* For the Uſe of Gun-powder was not known till about three Centuries ago. Although many of the Ancients make mention of Bullets, and of their being uſed in Engagements, yet they had neither Guns nor Gun-powder: On which Account they could not drive them with ſuch Force as the Moderns: For they either hurled them with a Sling, or ſhot them from a Croſs-bow. Whether known to the Antients.

III. Wounds of this Kind have a Cruſt or Eſchar formed upon them, and therefore are attended with little or no Hæmorrhage at firſt, unleſs ſome conſiderable Veſſels are wounded. But as ſoon as the Eſchar falls off, the Hæmorrhage is ſometimes ſo violent as to endanger the Life of the Patient, unleſs a Surgeon is at hand. For the five or ſix firſt Days there is little or no Diſcharge of Matter: Therefore it is not to be admired at, if Gun-ſhot Wounds, while the Veſſels are compreſſed by the Eſchar, exceed all others in Violence of Symptoms, ſuch as Inflammation, Pain, Gangrene, *&c.* Diſcharge ſmall quantities of Blood and Matter.

IV. The Eſchar which is formed upon theſe Wounds is not occaſioned, as many have imagined, ſo much by the Heat of the Bullets, as by the Rapidity with which they deſtroy the Parts, and the Violence of the Symptoms is owing chiefly to this Rapidity, whence enſues a violent Contuſion; and to the extravaſated Blood being long confined under the Cruſt. Formerly they were of Opinion, that there was ſomething poiſonous in Wounds of this Sort, but in this alſo they were miſtaken, for nothing poiſonous enters the Compoſition, either of the Powder or Ball. So far from it, that the Powder is uſed medicinally in acute Fevers.

V. Gun-

How many Differences.

V. Gun ſhot Wounds are ſome of them deep; ſome ſhallow, which only affect as it were the Surface of the Body, and perhaps take off a Piece of Skin or the Fat at fartheſt: Theſe are attended with leſs Danger, and generally with leſs Pain. In ſome the muſcular Parts alone are wounded, in others the larger Veins and Arteries, which often diſcharge ſuch a Quantity of Blood, as to endanger the Life of the Patient. Sometimes the Ball paſſes clear through; ſometimes it remains fixed in the Wound, and frequently carries Part of the Cloaths or Wadding with it. From the Difference of theſe Circumſtances different Symptoms ariſe. In others, again, the Bones are injured either by Colliſion or Fracture; and that ſometimes in the Body of the Bone, ſometimes in the Extremes or Joints. Whence commonly ariſe the worſt of Symptoms; and the Patient fares well, if he loſes only the Limb, and not his Life into the Bargain, eſpecially if the Joints of the Tarſus, Cubitus, Knee, Shoulder, or Thigh be violently ſhattered; unleſs the wounded Part, which is indeed almoſt the only Remedy, be cut off in Time. Gun-ſhot Wounds, which happen to the Viſcera, or the Contents of the Head, the Thorax or the Abdomen, as they fall on the nobler Parts of the animal Œconomy, generally end in Death, if not inſtantly fatal.

Of Wounds in the Cranium.

VI. Gun-ſhot Wounds, which affect the Cranium or Scull and the temporal Muſcles, are for the moſt part attended with great Danger: For even thoſe, that appear very ſlight externally, frequently bring on terrible Symptoms, by the Concuſſion of the internal Parts which they occaſion. For dangerous Fiſſures are often produced, of worſe Conſequence than even Fractures themſelves: Becauſe by this means the internal Laminæ of the Cranium are ſhattered; or the Veins and Arteries of the Brain burſt, in which Caſe the extravaſated Blood has no Vent, nor can the Splinters of the Cranium be extracted. Death therefore muſt be the Iſſue, unleſs prevented by the Trepan. If the Ball is lodged in the Brain, the Patients almoſt univerſally die: But, if it has pierced only one Side of the Brain, they may recover; which has been ſeen in Practice by myſelf and others. Yet all Injuries whatever of the Cranium and temporal Muſcles, are attended with great Danger, on Account of the Nerves and Arteries in thoſe Parts, and are therefore to be treated with great Care and Circumſpection.

Of Wounds in the Thorax and Abdomen.

VII. In the Thorax, if the Heart, or its Auricles, *&c.* if the pulmonary Artery or the Aorta, the Trunk of the Vena Cava or the pulmonary Vein, or any of their larger Veſſels are burſt, they are generally mortal. But if no large Veſſel is wounded, you need not deſpair of a Cure; though the Patient is often troubled with a Difficulty of Breathing. If any Viſcus of the Abdomen is pierced by a Ball, as the Ventricle, Inteſtines, Liver, Spleen, *&c.* or any large Blood-veſſel, in its Contents, Death muſt be expected. But if the Liver, Spleen or Kidneys are only ſuperficially wounded, the Patient frequently recovers. So, if the Bladder ſhould be injured in the Part ſituated without the Peritonæum, theſe Wounds are generally healed. But, if the Ball reſt in the Bladder, it occaſions ſuch an Accretion of calculous Matter, that the Patient cannot be cured without cutting.

Cure of Gun-ſhot Wounds.

VIII. In the Cure of theſe Wounds, the Surgeon muſt in the firſt Place examine, whether the Wound be ſlight, or of a dangerous Kind. We term it ſlight, if the external Parts only are hurt, ſuch as the common Integuments, or part only of a Muſcle, wherever ſituated; if the Bones are not affected, and if it

it be pervious. In these Cases if there be no Hæmorrhage, the Crust is to be removed by Suppuration. To effect this, the best Method is, first to dress the Wound with soft Lint and a Compress; the second or third Day to fill it up gently with a digestive Ointment, or with Honey, as a most excellent Suppurative; then cover it with a Plaster and Compress, or a Compress alone: and lastly, to secure it with a Bandage. If such a Wound should be received in a Part, where Compresses and Bandages cannot conveniently be applied (as for Instance, on the Face) it will be sufficient to secure the Lint with a Plaster only. Let this Dressing be continued either every, or every other Day, till the Crust is separated, the Wound cleansed and incarned; and the Cicatrix will be happily formed by the Application of dry Lint.

IX. When the Crust is removed, they generally apply balsamic Medicines; as all native Balsams, or Oil of Turpentine, or the *Eau d'Arquebusade*, which Method we by no Means disapprove; though indeed Wounds of this Kind are easier healed by being left to Nature, especially in sound and robust Constitutions. If any of the soft Parts are taken off by a large Ball in the Surface of the Thigh, Calf, Side, or Arm, the same Method of Cure is to be followed. If the Suppuration be too abundant, or the Flesh luxuriant, these Mischiefs must be removed by burnt Allum and red Precipitate; and the Wound must be dressed with Oil of Turpentine, *Balsam. Copaiv.* or some Balsamic Essence, as Amber, Myrrh, *&c.* or sometimes with dry Lint only. There are Cases, where the proud Flesh may be removed with the Finger: but in larger Wounds, when the Crust is cleared, you should avoid digestive Medicines, and apply nothing but spirituous Balsamics. What further to be done.

X. If there happen in these Cases a violent Contusion or Inflammation, there is no better Method (especially if the Patient has lost little or no Blood by the Accident, and is of a plethoric Habit) than to make a wide and deep Incision in the Wound, the Neck only excepted, on Account of the large Vessels. After a sufficient Discharge of Blood, the Incision should be dressed with fine dry Lint, Compresses well saturated with warm Spirit of Wine, and a proper Bandage. If there has not been a sufficient Discharge, Venesection must be applied in Plethoric Constitutions. Inflammation.

XI. But if an Hæmorrhage arise from a Wound in the fleshy Parts, it is a certain Sign that some large Vein or Artery is injured; for the small Vessels seldom bleed. In this Case styptic and balsamic Medicines, nay, the *Alcohol Vini*, is always prejudicial; for they constrain the bruised Parts, check the Circulation, and of consequence either cause an Inflammation, or increase it; and too commonly pave the Way to a Gangrene and Mortification: or, if Styptics should perhaps be particularly necessary, I would advise the *Nodolus ex Vitriolo*, or a small Compress dipped in the Styptic of Weber or Rabelius to be applied only to the Lips of the wounded Vessel, and pressed down with the Finger, till it produces a Scar, and the Blood is staunched: for these Wounds will bear the Application of much Lint, or the Stricture of Bandages: but the best and safest Way, is to close up the Mouth of the wounded Vessel with a Needle and Thread. Hæmorrhage.

XII. In Wounds of the large Arteries (the Situation of which the skilful Anatomist is not unacquainted with) that the Patient may not be lost by a violent Hæmorrhage, it is expedient, first to compress the Artery with your Thumb, Of Wounds in the large Arteries.

then apply the *Tourniquet* to the wounded Limb, and by conſtringing the Trunk of the Veſſel, ſtop the Diſcharge: after this, you muſt take it up by means of the crooked Needle (*See Ch.* II. N°. VII.) but if, from the Narrowneſs of the Wound you cannot get at the injured Veſſel, it muſt be enlarged with the Scalpel, in order to reach the Artery and bind it up, or ſtop the Hæmorrhage by the Application of Styptics. The Wound ſhould afterwards be dreſſed with dry Lint, Compreſſes, and a Bandage: nor ſhould theſe Dreſſings be renewed till the third or fourth Day, nor the Lint be removed, for Fear of a freſh Hæmorrhage; but let it remain, till it quits of its own Accord.

Gun-ſhot Wounds of worſe Conſequence.

XIII. But there are Gun-ſhot Wounds of worſe Conſequence than theſe: to which Claſs may be referred, 1. Thoſe where the Balls have not paſſed through; or where they, and other Bodies forced in with them, remain in the Part. 2. Where the Bones are at the ſame Time broken or ſhattered: and laſtly, where the Viſcera, and larger Veſſels, within or without their Cavities, are wounded; from whence ariſe very dangerous Hæmorrhages.

The general Method of treating theſe Wounds.

XIV. In the Treatment of theſe Wounds we muſt have Regard both to the Wounds themſelves, and the Accidents attending them. As to the Wounds themſelves, ſix Particulars are to be obſerved: as (1) If we perceive that the Ball has not paſſed through, which is plainly demonſtrated by a ſingle Aperture; in this Caſe we muſt endeavour with all Speed to extract the Ball, or other foreign Bodies forced in with it, ſuch as Cloaths, Wadding, *&c.* The ſame alſo is to be regarded with reſpect to Splinters of Bones: for, before theſe are removed, you in vain attempt to heal. (2) If there be a violent Hæmorrhage, it muſt be ſtopped by the Method abovementioned; but if it be ſlight, and from the ſmaller Veſſels, it is ſcarce worth attending to: for in the firſt Place it is often ſerviceable in plethoric Habits, by leſſening the Quantity of Blood, and thereby preventing Tumors and violent Inflammations; and it generally ſtops of its own accord; or is eaſily checked by applying dry Lint or a gentle Aſtringent. (3) Bruiſed and corrupted Fleſh ſticking in and about the Wound (which is called a Cruſt or Eſchar, if but a little) is eaſily taken off by a digeſtive Ointment, and Spirit of Wine, impregnated with *Sal Ammoniac:* if much, it cannot be removed without deep Scarifications and ſuppurating Medicines. (4) The Void of the Wound muſt be filled up with new Fleſh: (5) an even Cicatrix be formed: and (6) if any Bones are broken, they muſt be united.

Extraction of Balls and extraneous Bodies.

XV. With regard to the Extracting of Balls or other foreign Bodies, the Surgeon ſhould immediately enquire, except there be a violent Hæmorrhage, if any thing of that Kind remain in the Wound, and where. But here we muſt obſerve, that Balls, Stones, and other hard Bodies are eaſier diſcovered by the Finger, or Probe, than Fragments, or Cloaths or Wadding: for theſe laſt, from their Softneſs, and the Redneſs occaſioned by the Blood, are very difficult to be diſtinguiſhed from the membranous and muſcular Parts, either by the Sight or Touch. But, to be ſurer in this Caſe, it is always uſeful, nay, generally neceſſary, to enlarge the Wound, and ſcarify; carefully avoiding the larger Veins and Arteries, but not regarding the ſmaller, the Nerves, or the Muſcles. This you muſt do, till you come at the foreign Bodies: the Extraction of which ſhould be performed, if poſſible, with the Hand; or, if that cannot be done, with a Forceps, or Hook, *See Plate* III, *Fig.* 3, 4, 5, 6, 7, 8. They are eaſieſt removed at firſt: for after ſome Delay the Tumor and Inflammation of the

Parts

Parts render it difficult and painful. Beſides, Bullets will by Degrees work themſelves deeper, and be buried under the Muſcles: which will occaſion Fiſtulæ, Rigidity of the Limbs, and other Inconveniences. In extracting Balls that lie deep, you muſt take great Care not to lay hold of Blood-veſſels, Nerves, or Tendons; which Accident will be avoided by introducing the Forceps ſhut, and not opening them till you feel the Ball.

Of Balls in the Bones or Joints.

XVI. When a Bullet is lodged in a Bone, we muſt endeavour to extract it with the Forceps or Hooks, (*See Plate* III, *Fig.* 8.) If it lie too deep for theſe, we muſt have Recourſe to the Trepan, as deſcribed *Plate* III, *Fig.* 7. or *Plate* VII, *Fig.* 7. *Lett.* B: but if this will not avail, LE DRAN adviſes immediate Amputation: yet in my Opinion, unleſs in Caſes of great Extremity, we had better leave the Ball, till the Parts ſuppurate, and ſet it at Liberty. Sometimes the Ball is to be removed by trepanning: ſometimes it remains a long while in the Bone without any Danger or Inconvenience: a remarkable Inſtance of which, I remember in an Officer, who for many Years carried a Bullet in the Middle of the Tibia, that gave him no Trouble; whence it appears, that we ſhould not be too haſty in proceeding to Amputation. But if a Ball, or any other extraneous Body, be lodged in the Joint between two Bones, CELSUS (*Book* VII, *Ch.* 5.) has very judiciouſly adviſed, to extend the Joint with your Hands, or by the Means of Bandages and Slings, as in the Caſe of Luxations: by which the Ligaments being looſened, Room is made between the Bones, for the eaſier Extraction of the foreign Body. I have often wondered that this excellent Precept of that great Phyſician ſhould be ſo generally neglected by the Moderns, as a Thing of no Conſequence.

Of Extraction on the oppoſite Side.

XVII Whenever the Ball has penetrated ſo deep into the wounded Part, as particularly in the Arm, Thigh, Thorax, or Abdomen, that you can eaſily feel it with your Finger on the Side oppoſite to the Wound, the Surgeon ſhould examine nicely whether it is ſafeſt to bring it back by the Way it went in, or to make an Opening upon it, and draw it out at the oppoſite Side. When the Wound cannot be enlarged without Danger of injuring the neighbouring Parts, you ſhould ſearch for it with the Probe; and endeavour, but with great Caution, to extract with the Forceps, or ſome other convenient Inſtrument.

Of the Cure of fractured Bones.

XVIII. If the Wound is attended with fractured Bones, after the neceſſary Inciſions have been made, what Fragments are looſe, whether in the Cranium or elſewhere, ſhould be gently removed. If theſe Fragments ſhould adhere to the principal Bone, by their fine Membranes, they muſt be cut off; and the larger Parts of the Bone, that are not much injured, be reduced to their natural Poſition, and retained there by proper Dreſſings, as in other Fractures. Where any Splinters ſtick out, which hinder the Reduction, and hurt the neighbouring Parts you ſhould break them off with the Forceps. If the Bones of the Tibia or Femur are broken, after they have been cleared of the extraneous Bodies, and reunited in the beſt Manner, they are to be dreſſed with the eighteen-headed Bandage, and repoſed in the Straw-couch, (Fr. *Fanones*) or in PETIT's Machine, as in other complicated Fractures. Where a Wound is inflicted by a ſpent Bullet, or the Fragment of a Grenade, and the outward Integuments are not penetrated, though the Bone be broken; in this Caſe worſe Conſequences are to be expected from the Violence of the Contuſion: therefore, to free a Paſſage for the extravaſated Blood, we muſt make deep Inciſions in the Integuments and

Muſcles, but not quite to the Bone. When the Blood is thus diſcharged, the broken Bone ſhould be replaced, and, if poſſible, dreſſed with the eighteen-headed Bandage. But Wounds of this Kind, eſpecially from large Pieces, generally take off the whole Limb, and too often the Patient himſelf.

Further Obſervations on Wounds in the Bones.

XIX. If the Joint be contuſed, and there be no Wound nor Fracture of the Bone, leſt dangerous Symptoms ſhould ariſe from any Injury of the Ligaments, Nerves, and Tendons, your Inciſion muſt be only in the Integuments and fleſhy Parts: but if a noxious Humour be found in the Joint, the Ligaments alſo muſt be cut to let it out. Veneſection and a proper Regimen are here very neceſſary, not without the Uſe of internal reſolving Medicines, and powerful Externals. But if the Bones in the Joint are broken or ſhattered, the Limb can ſcarce be ſaved, nor the Patient himſelf, unleſs you take it off a little above the Wound. In Contuſions of the Bones without a Fracture, if, after proper Inciſions, a violent Pain remains in the Bone; if the neighbouring Fleſh is pallid, the Bone of a darkiſh Colour, with a large Suppuration; in this Caſe either the Limb muſt be taken off, or, as LE DRAN adviſes, the Bone muſt be trepanned upon the Wound, to open a Paſſage for the confined Matter. If, in this Caſe, the Pains are very acute, unleſs you amputate, the Patient generally dies convulſive. The broken Bone, when rejoined and properly dreſſed, ſhould be conſtantly kept in an elevated Poſture.

In Caſe of Delay.

XX. If a Surgeon be not timely called in, and the extraneous Bodies remain long in the Wound, from whence ariſe Tumors, Inflammations, violent Pains, and other bad Symptoms; the Lips of the Wound muſt be greatly enlarged and deep Inciſions made in the Tumors. By this Means you raiſe a freſh Hæmorrhage, the Swelling and Inflammation abate, and the foreign Bodies are eaſily extracted. But in all theſe Inciſions, eſpecially in the Limbs, the *Tourniquet* ſhould be applied to the Part.

Of more extraneous Bodies.

XXI. But as two Balls are often concealed in the ſame Wound, after the Removal of one, the Surgeon ſhould diligently ſearch for another, or for any other extraneous Body that may be forced in with it: for unleſs every thing foreign be firſt removed, it is in vain to expect the Cure of the Wound.

Other neceſſary Obſervations.

XXII. When you attempt the Extraction of a Ball or other extraneous Body, you ſhould endeavour to place your Patient in the ſame Situation he was in at the Time of receiving the Wound: for by frequent Changes of the Situation, the Ball will eaſily bury itſelf, and get out of your Reach. If the Wound cannot eaſily be enlarged, nor the Balls extracted without great Pain and Danger, they muſt be left in the Wound, either till the Pain is abated, or the Paſſage rendered ſo eaſy by Suppuration, that they work themſelves out. On the other hand, extraneous Bodies are inſtantly to be removed, where there is Danger of bringing on Convulſions, Pain, or Inflammation, by being left behind.

Ball lodged in the Thorax or Abdomen.

XXIII. If a Ball has paſſed into any Cavities of the Body, where the Extraction of it cannot be attempted with Safety, it is beſt to leave it where it has lodged, and heal the Wound: for there have been Variety of Inſtances, where Perſons have carried Balls in them for many Years, nay, for the beſt Part of their Lives, without ſuffering any Inconvenience. It ſometimes happens that they work themſelves out into ſome other Part of the Body, from whence they may be extracted with Safety.

Balls lodged in the Viſcera.

XXIV. But if a Ball be lodged in a Viſcus, as the Brain, Lungs, Liver, Spleen, &c. that you can neither ſee, nor reach it, the Caſe is generally mortal.

If

If you can ſee it, it is only on the Surface. Therefore by enlarging the Wound of the Viſcus, as far as is convenient, you will make Room for the Pliers or Forceps to extract the Ball, which muſt be introduced with the utmoſt Caution.

The Treatment of the Wound, when cleanſed.

XXV. When the Wound has been cleared of the extraneous Bodies, and the Hæmorrhage ſtopped, you muſt endeavour to remove the Eſchar by Suppuration. In the next place you attend to the bad Symptoms ariſing from it, ſuch as violen Swellings and Inflammation, Fever, Gangrene, and Mortification, exceſſive Weakneſs, Nauſea, *&c.* ſome of which are to be prevented, and others remedied. To prevent therefore violent Swellings and Inflammations, which are here always dangerous, and often bring on a Gangrene and Mortification; beſides proper Inciſions, you ſhould apply externally *Spirit Vini ſimp.* if the Wound be ſlight: if not very ſlight, add to each lb of the *Spirit.* ℥ſs of *Sal. Ammoniac.* or *Aq Calc.* with about a fourth Part of *Spirit. Vini Camphor.* with a little of the *Sal Ammonic.* Thick Compreſſes dipped in this Mixture ſhould be applied warm to the affected Part ſeveral Times in the Day. If in Parts that are contuſed and greatly inflamed, the Blood ſhould be concreted under the common Membranes of the Muſcles; theſe Membranes muſt be ſcarified without Reſerve, not only in their longitudinal, but in every Direction: for otherwiſe the ſtagnated Blood cannot be removed, and of Conſequence a Gangrene and Mortification would ariſe, or at leaſt very dangerous Abſceſſes. And ſometimes you muſt cut even the Tendons eſpecially in Wounds of the *Tarſus* and *Metatarſus*, where you have acute Pains, that may be attended with Danger.

A great Corruption of the Parts.

XXVI. Where there is a very great Corruption of the Parts, after repeated Scarifications apply the digeſtive Ointment; to which you may add a little Myrrh or Aloes *Unguentum fuſcum*, or red Precipitate. Let the Wound be dreſſed with theſe, till the injured Parts are ſeparated, and the reſt well cleanſed: then let it be treated as a ſimple Wound.

Of Setons in nervous Wounds.

XXVII. In deep Wounds where the Ball has gone quite through (eſpecially if they happen in the Buttocks or Thigh, and the Paſſage of the Ball is oblique) a particular Method is to be uſed. After ſcarifying the Lips of the Wound, a Skein of Thread is to be drawn through the Eye of a long blunt Needle, (*Plate* V. *Fig.* 1.) and, being well ſaturated with the Ointment we have preſcribed, paſſed through the Wound in the Manner of a Seton. It ſhould be kept there, till you diſcover by the Redneſs of the Wound that the corrupted Parts are caſt off, and the whole is in a Readineſs to heal. LE DRAN abſolutely rejects the Setons; but I have often ſeen the good Effects of them. You ſhould take particular Care that the Thread be very ſoft.

Symptoms of theſe Wounds.

XXVIII. Gun-ſhot Wounds, though in the ſtrongeſt Conſtitutions, generally produce the worſt of Symptoms; as exceſſive Weakneſs, Faintings, Tremors, Palpitations, Convulſions, Hiccoughs, *&c.* after which ſucceed inſtantly dangerous Fevers, nauſeous Vomitings, and the like: all which require the particular Attention of the Surgeon.

1. Weakneſs.

XXIX. Amongſt the many terrible Symptoms attending theſe Wounds, the firſt is an exceſſive Languor, which is ſoon followed by fainting Fits, partly occaſioned by the Hurry of their Spirits, and partly by the Effuſion of Blood. If from the firſt Cauſe, Draughts or Mixtures compoſed of Cordial Waters, Cardiac Powders, and a ſtrengthening Diet, are to be preſcribed. If from the latter, to recruit their Strength and repleniſh their Veins, let them indulge in all

all things nouriſhing; as the *Decoct. Hord. vel Aven. Corn. Cerv. Citrat.* for their common Drink, with a ſmall Quantity of Wine or generous Beer, Milk, Emulſion, good Broths, and comfortable Juleps.

2. Nauſea. XXX. Some are ſeized with a violent Nauſea, or Abhorrence of Food. This ariſes partly from the Terror they are under, and partly from too great a Repletion before the Accident. It is too cuſtomary with military Men to eat and drink freely, as if by this means they acquired freſh Strength: hence, on a ſudden Fright, the Motion of the Stomach being inverted, and the Digeſtion diſturbed, a Nauſea is the Conſequence. In ſuch a Diſorder, as Experience teaches that thoſe who vomit ſpontaneouſly receive great Benefit, we ſhould here apply a gentle Emetic, and then ſettle the Patient's Stomach with a Cordial Draught. By this Method his Appetite and his Strength generally return; and the Cure of the Wound is happily promoted. If the Patient is averſe to an Emetic, you ſhould give ſome proper Purgative in its ſtead.

Vulnerary Fevers. XXXI. Intermitting Fevers in vulnerary Caſes are to be treated like others of the ſame kind: but be ſure to give an Emetic at the Beginning. But if an acute Fever come on, attended with frequent Horrors, as they indicate ſtrongly internal Inflammation, we muſt bleed the Patient in Proportion to his Strength and Fullneſs of Blood: then order him a gentle Doſe of *Ipecac.*; ſome thin Liquor for his ordinary Drink; cooling Powders and Draughts, with Camphire, and a moderate Diet. He ſhould take largely of the *Peruvian Bark*: but he muſt particularly abſtain from Pork, from all ſalt Meats and of hard Digeſtion.

Pains and Convulſions. XXXII. Pains and Spaſms are generally produced, either (1.) By extraneous Bodies remaining in the Wound, which prick and vellicate the nervous Parts: or (2.) By large Tumors and Inflammations, which are always viſible: or (3.) by the violent Colliſion and Laceration of the nervous and tendinous Parts. If theſe Symptoms ariſe from foreign Bodies, we muſt endeavour to remove them as ſoon as poſſible: if from exceſſive Tumor and Inflammation, theſe muſt be diſcuſſed by the means above-mentioned: but if there be Danger of a Gangrene and Mortification, beſides the deep Inciſions, frequently repeated for the Diſcharge of the putreſcent Blood, and beſides the Medicines we before recommended, large Quantities of *Peruvian Bark* muſt be given inwardly; and Lint, well ſaturated with Spirit of Turpentine, be applied to the Wound. They are both excellent Remedies in this Caſe.

Amputation where neceſſary. XXXIII. But if all theſe Remedies fruſtrate our Hopes, there remains one only, and, as CELSUS obſerves, a deplorable Refuge, the Amputation of the corrupted Part. But, whenever violent Pains, Spaſms, and Convulſions are occaſioned by the Colliſion or Laceration of the nervous or tendinous Parts; it is adviſeable at the firſt, or very early at leaſt, to cut the Tendons, Aponeuroſes, and common Membranes of the Muſcles that are injured, juſt above the Wound. For if theſe Symptoms are not ſpeedily removed, Death muſt be the Conſequence. But if this avail nothing, it is better at once to amputate the Limb, than by unſeaſonable Delays deſtroy the Patient: for theſe partial Convulſions ſoon become univerſal.

Freſh Hæmorrhages. XXXIV. About the ſeventh or eighth Day, ſooner or later, from the inflicting of the Wound, freſh Hæmorrhages uſually ſucceed [a]: to ſuppreſs which, we

[a] When the *Eſchar* ſeparates.

muſt

muſt proceed, as above, and earneſtly adviſe the Patient to indulge Reſt, and ſtir as little as poſſible: otherwiſe, theſe Hæmorrhages are too apt to return about the fourteenth Day.

In Caſe of a Looſeneſs.

XXXV. About the ſame Time you have generally a Looſeneſs, which ſome think not only unſalutary, but very prejudicial: but in general it ſhould be eſteemed ſalutary and critical; eſpecially in corpulent Patients, and of a bad Habit of Body: for thereby many bad Symptoms are abated; and a Wound, which before was unpromiſing, takes a new Turn, and a different Aſpect. Therefore we ſhould by no means check it with Aſtringents; but gently aſſiſt Nature with warm, diluting, lubricating, and mucilaginous Draughts; with *Ol. Amygdal. dulc.* with lenient Clyſters, till it ſtops ſpontaneouſly. But if it continue too long, we muſt then have Recourſe to the aſtringent Powders; particularly *Pulv. Rhabarb. cum Cort. Caſcarill.* Theſe ſhould be repeated often in the Day; and at Night give a Doſe of *Diaſcordium*, or *Theriaca ex aquâ Menthæ vel Cinnam. Cydoniat.* nor muſt a proper Regimen and Diet be omitted.

Venereal Symptoms, Fiſtulas, Caries and Atrophy.

XXXVI. If the Wound prove obſtinate from any venereal Taint, which will eaſily appear from other Symptoms of that Diſorder, antivenereal Remedies muſt be applied. If there remain Fiſtulas, which are commonly occaſioned by Fragments of the Bones, or extraneous Bodies being left behind; or if there is a Caries; theſe Symptoms are for the moſt Part eaſily removed by removing the Cauſes. But if the Wound is inflicted on the Breaſt, and Part of its Subſtance deſtroyed by it, the Caſe is too generally fatal. A Caries often requires Length of Time and Patience: yet Nature, if aſſiſted with proper Medicines, frequently conquers it. It is no uncommon thing to ſee an Atrophy in the Limbs after violent Wounds, where the Parts are ſo lacerated, and the Circulation of the Blood ſo impeded, that they want their due Nutriment. In this Caſe, all ſtrengthening and emollient Medicines, all Ointments and Fomentations, are extremely efficacious, and particularly the Uſe of natural Baths.

To extract Gunpowder and ſmall Shot.

XXXVII. In Gun-ſhot Wounds ſeveral Grains of Gunpowder frequently penetrate the Skin of the Face, and occaſion a Deformity, if not removed: this may be done with a Pen, or an Inſtrument like an Ear-picker, *See Plate* VI, *Fig.* 14. But if they have penetrated too deep to be picked out in this manner, the Skin muſt be laid open with a fine ſmall Lancet, that you may remove them with the Inſtruments we have deſcribed. Great Care muſt be taken not to break the Grains in picking them out; for they will occaſion very foul Spots. The ſame Method is to be obſerved with regard to ſmall Shot. If you would be further informed of Gun-ſhot Wounds, and the Writers on that Subject, turn to the XIIth Number of the Introduction to this Work; and conſult particularly the excellent Le Dran.

CHAP. IV.

Of Wounds *of the* Abdomen.

The Intent of this Chapter.

I. WE have hitherto conſidered what was in general to be attended to with regard to any ſort of Wound, whether made by Cutting, Stabbing, or by the Exploſion of a Gun. We come in the next Place to explain fully the

Nature of each particular Wound, and ſhall ſpeak diſtinctly of Wounds: 1. of the *Abdomen*. 2. of the *Thorax*. 3. of the *Neck*. 4. of the *Head*.

Differences of Wounds in the Abdomen.

II. *Wounds of the Abdomen* only affect the common Integuments and Muſcles, or penetrate into the Cavity of the Abdomen. Thoſe that penetrate into the Cavity of the Abdomen, are inflicted lengthways, obliquely, or tranſverſely, and in theſe the Bowels either burſt out through the Wound, or preſerve their natural Situation. Theſe Differences of Wounds in the Abdomen ought to be diligently attended to by the Surgeon, ſince they require a different Kind of Treatment.

How thoſe Wounds are to be diſcovered.

III. Theſe Wounds may be conveniently enough examined; 1. by the Eye; 2. by paſſing the Finger or the Probe; or laſtly, 3. by injecting warm Water into the Wound. If the Water meets with no Obſtruction, you are ſure that the Wound penetrates; but if it returns back upon you, and the Probe meets with Reſiſtance, the Abdomen is not entirely opened.

Wounds that do not penetrate.

IV. Wounds which do not penetrate the Cavity of the Abdomen, are attended with much the leaſt Degree of Danger. They are generally divided into two ſorts. 1. either the Wound is only upon the common Integuments; or, 2. the Muſcles alſo of the Abdomen are divided, as far as the Peritonæum. The firſt of theſe is too ſlight to require a diſtinct Method of Cure from other Wounds: but Wounds of the laſt Claſs are extremely dangerous; becauſe the Inteſtines, in this Caſe, eaſily fall through the Wound. If the Wound is large, great Skill is required in the Surgeon, eſpecially if it is made in a tranſverſe or oblique Direction: but if it is lengthways, by cleanſing the Wound, applying the vulnerary Balſam, and a healing Plaſter, with the large uniting Bandage, as at *Plate* V, *Fig.* 8. all will go well; if carefully dreſſed according to theſe Directions, and the Patient indulge Reſt, and obſerve a proper Regimen, the Cure generally is effected without a Suture. But ſhould the Wound be tranſverſe or oblique, in this Caſe, to prevent a Rupture, the Suture is neceſſary to keep the gaping Lips of the Wound together, as we ſhewed partly above at *Chap.* I, *N.* XLIV. The manner of performing this we ſhall deſcribe below in a Chapter upon *Gaſtroraphy*. Having taken theſe Precautions for preſerving the Peritonæum and Inteſtines in their natural Situation, the Surgeon ought to dreſs up the Wound with vulnerary Balſams, and an adheſive Plaſter: to give the Patient Reſt, to order him a ſoft Clyſter if his Bowels are not naturally open, and to enjoin Abſtinence.

How Wounds are to be ſearched when they penetrate.

V. When the Surgeon diſcovers that the Wound penetrates into the Abdomen, he ought, before all things, to examine well, whether it be direct or oblique, and whether any of the Contents of the Abdomen partake of the Injury. He will eaſily determine in the Negative, if it ſhall appear; 1. that there is no great Degree of Weakneſs, Hæmorrhage, Pain, Fever, *&c.* 2. if, upon laying the Patient upon the wounded Side, there is no Diſcharge of Chyle, Gall, Excrement, or Urine. 3. if Milk, being injected warm, returns without any Alteration of its Colour. 4. if the inflicting Inſtrument is not very ſharp; and laſtly, 5. if there is no Vomiting nor Diſcharge of Blood by the Mouth, Stool, or Urine, nor Swelling and Hardneſs of the Belly. But as the Operation of *Gaſtroraphy* is ſometimes extremely neceſſary, and always attended with Danger, if it is not performed with the greateſt Accuracy, I have thought it my Duty to deſcribe it carefully in the following Chapter.

CHAP.

CHAP V.
Of GASTRORAPHY.

I. GASTRORAPHY is the Suture of Wounds of the Abdomen. This Operation is unneceſſary; 1. when the Wound is only in the muſcular Part; or, 2. is not very large, eſpecially if it is made lengthways: for if the Wound ſhould penetrate into the Cavity of the Abdomen, and even let out Part of the *Omentum* or *Inteſtines*; yet, where it is very ſmall, as Wounds generally are which are made by Puncture, or happen lengthways, upon returning the Parts which are puſhed out, ſtopping the Wound up with a ſoft Tent, and ſecuring all with a proper Bandage, it may be healed without the Help of the Needle. Beſides, in fat Perſons, this Operation is very difficult, and it would be an Act of great Cruelty in a Surgeon to perform the Operation upon a Man, when he might be cured after an eaſier Method. When Gaſtroraphy is unneceſſary.

II. But there are two Caſes where Gaſtroraphy is abſolutely neceſſary. The firſt is, where the Wound is ſo large, that there is no Poſſibility of retaining the Inteſtines by any other Method; for as the Inteſtines are continually puſhing forwards in the Act of Inſpiration, by the Action of the Diaphragm and the Abdomen, the Falling down of the Inteſtines in this Caſe is unavoidable, eſpecially in large Wounds from a Cut, *See Plate* III, *Fig.* 1, *Lett.* O, and therefore the Operation neceſſary: but there is another Caſe alſo where this Operation is required; to wit, in large tranſverſe Wounds of the Abdomen where the Muſcles are divided, but the Peritonæum is not concerned. *See above Chap.* IV, *N.* IV. When it is neceſſary.

III. In Wounds of the Abdomen the chief Enquiry is, *Whether the Omentum or Inteſtines are let out?* If none of theſe have burſt through the Wound, the Lips of the Wound ſhould be kept as cloſe together as poſſible with the Hands, and the Patient kept with his Head laying downwards till the Wound is ſufficiently ſecured from letting out the Contents of the Abdomen. But when the Inteſtines are already fallen out, they muſt be returned with the greateſt Expedition, leſt they ſhould receive any Injuries from the external Air. But we ſhould firſt examine whether they have received any Wound or not, and whether they preſerve their natural Warmth and Colour. For where they are cold, livid, and dry, or wounded, they are not to be returned ſuddenly, but treated in the Manner we ſhall deſcribe below. Of the falling down of the Inteſtines.

IV. You will eaſily perceive that there is ſome Hurt in the Inteſtines, though the Wound does not immediately appear, if there is a more than ordinary Flaccidity in them: when this Symptom appears, it will be proper to pull the reſt of the Inteſtines gently forward, till you come at the Injury, and when you have found it, you may treat it as we ſhall ſhew you in *Chap.* VI. How to diſcover a Wound in the Inteſtines.

V. When you find the Inteſtines uninjured, they muſt be inſtantly returned, to prevent them from receiving any Injuries from the external Air: in order to do this with the greater Eaſe, put the Patient in the ſupine Poſture which we deſcribed at *N.* III, only placing him upon the Side oppoſite to the Wound: the Patient being thus ſituated, an Aſſiſtant ſhould endeavour to return the Inteſtine with his two Fore-fingers, taking Care not to take off one Finger till the other is upon the Gut: the Patient ſhould be encouraged all the while to How to return the Inteſtines.

hold his Breath, and the Aſſiſtant ſhould bring the Wound together with his Fingers, or with Hooks, *Plate* VIII, *Fig.* 2, 3.

How the Inteſtines are to be treated when they are dry and cold.

IV. Hitherto we have deſcribed the Method of returning the Inteſtine whilſt it was warm and unwounded: it remains that we teach the Method of treating the Inteſtines when they are cold and dry. In this Caſe it is beſt to foment them with warm Water or Milk before you return them; or, where you can have that Opportunity, get the Cawl of a Calf, a Lamb, a Hog, or of any other Animal juſt killed, wrap this round the Inteſtines whilſt it is reeking, and keep them in it till they recover their natural Heat and Colour: but, if they do not return, all medicinal Efforts are vain: the Patient muſt die. If this Dryneſs or Coldneſs of the Parts is very ſmall, and the Inteſtines are not at all corrupted, it is beſt to return them inſtantly into the Body, where the Heat and Moiſture of the neighbouring Parts, being natural to them, will give them a more ſpeedy and natural Refreſhment, than can be obtained for them by any artificial Means.

How the Inteſtines are to be returned through a ſmall Orifice.

VII. When the Inteſtines are forced through a ſmall Wound, and are afterwards ſo diſtended with *Flatus*, that they cannot conveniently be returned, it will be proper to pull the Inteſtine gently forward, that more of it may come out, that ſo the *Flatus* being divided may take up leſs Room in any one Part. An Aſſiſtant ſhould now gently dilate the Wound with his Hands, or two Hooks, *Plate* VIII., *Fig.* 2, *or* 3 fixed in the internal Membrane, that the Surgeon may return the Inteſtines, which when he has done in ſuch a Manner that each Part may recover its natural Situation, (*See N.* V,) the Wound ſhould be ſecured firſt with his Hand, that the Bowels may not burſt out again: then it ſhould be filled up with ſome Doſſils, or, where there is a conſiderable Quantity of Blood ſpilt in the Abdomen, with a ſoft [a] Tent, *Plate* II, *Lett.* L, M, N, O; dreſſing it up with proper Plaſters, Compreſſes, and Bandage: the Patient is to be kept as ſtill as poſſible, lying as much as he can upon the Wound: after this the Wound is to be dreſſed daily, or, where there is a large Diſcharge of Matter, twice every Day, with ſome vulnerary Balſam; and, if we proceed in this Manner, where the Wound is not very large, the Patient may be excuſed from the Pain, and the Surgeon from the Trouble of making the Suture.

How ſmall Wounds are to be enlarged.

VIII. But if the Wound is ſo narrow, that we can neither bring the Gut forward nor reduce it, it muſt be enlarged with a Knife, or the grooved Probe, called *Conductor*, beginning the Diviſion at that End of the Wound which is moſt convenient, taking great Care not to wound the *Linea alba*, the Veſſels which lie under the *Muſculi Recti*, or, laſtly, the Inteſtines themſelves. Some Surgeons, in the Room of the Inciſion Knife and Conductor, uſe in this Place the *Syringotomus*, whoſe Point is guarded with a Button, and which is uſed in *Fiſtula Ani* (*See Plate* XXXV, *Fig.* 4, 5.) Some are fond of other Inſtruments; but I

[a] See CELSUS, Book VII, Chap. 16.

Some of the modern Writers in Surgery, particularly GARENGEOT, forbid the Uſe of Tents in all *Wounds of the Abdomen*. In the Year 1734, a young Surgeon in my Neighbourhood obſerved this Rule, when he was called to a Man that had received a Wound between the Navel and the Penis, the Wound penetrated the Abdomen; for the firſt two Days the Symptoms were favourable, but upon the fourth Day he died: upon opening his Body, we found a large Collection of Matter in the Abdomen, with the Omentum putrified. If a Tent had been uſed, the Matter would have been diſcharged, and the Patient's Life ſaved.

think

think the beſt Inſtrument by far in this Caſe, is the Knife which I invented for this Purpoſe, and have given you a Deſcription of at *Plate* V, *Fig.* 3. or one of thoſe at 4 and 5. The Knife is never to be uſed, till the Aſſiſtant has applied a warm Omentum to the Inteſtines that are already extra-abdominal, to prevent them from Injuries. But where the Inteſtines are ſo inflated, that it is impoſſible to get the Probe End of the Knife, or a Conductor, into the Abdomen, then hold back the Inteſtines with the Left-hand, and with the Right make an Inciſion through the common Integuments and Muſcles as far as the Peritonæum; ſponging up the Blood as you go on: the Wound will moſt likely be ſufficiently relaxed by this, to make Way for the Re-admiſſion of the Inteſtines, at leaſt it will admit the End of the Knife to divide the Peritonæum, ſo that you may enlarge your Wound at Pleaſure, and return the Gut as directed at *N.* V.

How to return them without enlarging the Wound.

IX. If any hardened Excrement lies in the Inteſtine, and impedes its Reduction, emollient Fomentations and Cataplaſms ſhould be applied, and more of the Inteſtine ſhould be pulled out; for by this Means the Fæces may be divided by the Hands, and the Inteſtine returned conveniently. PARÆUS and other Surgeons have recommended a particular Method of returning the inflated Inteſtine without enlarging the Wound, by making ſmall Punctures in the Inteſtine with a Needle, through which Punctures the Wind will certainly eſcape, and the Sides of the Gut ſubſide: and this, they affirm, is attended with no Danger: nevertheleſs, for my own Part, I prefer the Enlargement of the Wound to making theſe Punctures, and to the pulling out of a greater Share of the Inteſtine to divide the Contents; eſpecially ſince many Surgeons affirm, that theſe Punctures are neither ſafe, nor uſeful for the End to which they are directed. BLANCARD has given us an Inſtance where they failed, in his *Collect. Medico-Phyſic. Part. ult. Obſ.* I.

When and how to heal the Wound without the Operation.

X. When the Inteſtines are returned, if the Wound is not large, and is made lengthways, there will be no Occaſion to perform the Operation; which is always of dangerous Conſequence, and therefore ſhould never be attempted but in Caſes of the greateſt Emergency. If the Suture is not abſolutely neceſſary, paſs a ſoft Tent into the lower Part of the Wound, and apply ſticking Plaſters to the Sides of it, covering them with long thick Bolſters, ſecuring theſe Dreſſings with a uniting Bandage, ſuch a one as you will find deſcribed in *Plate* V, *at Fig.* 8. When the Patient is thus dreſſed, draw ſome Blood from the Arm, to prevent an Increaſe of the Inflammation; adviſe him to keep very ſtill, and obſerve a ſtrict Regimen with regard to his Diet. The Dreſſings are not to be removed, unleſs ſome violent Symptom requires it, before the third Day; and afterwards only once a Day, or rather every other Day, leſt the Union of the Wound ſhould be retarded by frequent Handling: on the other Hand, if the Wound is large, and made in an oblique or tranſverſe Manner, as deſcribed *Plate* III, *Lett.* I, O, ſo that the Inteſtines cannot be kept within the Abdomen by this Method, the Operation muſt be performed without Delay.

How to perform the Operation with two Needles.

XI. The Operation may be performed in the following Manner: paſs a ſtrong double or quadruple Thread, well waxed, through two crooked Needles; (*See Plate* VI, *Fig.* 5 *and* 6. or another, which was communicated to me by a Friend, at *Fig.* 7.) with theſe ſtitch up both Ends of the Wound, beginning at one End with the upper Lip of the Wound, paſſing the Needle through the

Peritonæum, Muſcles of the Abdomen, and the common Integuments, from within outwards, leaving about a Thumb's Breadth between the Stitches and the Mouth of the Wound, that they may take the ſtronger Hold, obſerving the ſame Method in paſſing the other Needle through the lower Lip. Whilſt you are paſſing the Needle with one Hand, it will be proper to ſupport the Lips of the Wound with the other, to prevent the Inteſtines from being wounded: it will frequently be very difficult to hold the Needle ſteady with the naked Hand; to remedy this Inconvenience, the modern Surgeons have invented an Inſtrument to receive the Needle, and form a Handle for it, which the *French* call *Portaiguille*. *See Plate* VI, *Fig.* 2, 3, *and* 4.

With a ſingle Needle.

XII. If you are not provided with two Needles, the Operation may be performed with one; for, after you have ſtitched up one End of the Wound in the Manner we have deſcribed, you may take off the Needle, and perform the ſame Operation on the other End, and proceed afterwards as uſual. Likewiſe, inſtead of Thread, a ſmall kind of Lace [a], compoſed of ſix Threads, may be judiciouſly ſubſtituted; as the Threads themſelves are liable to break, or they may ſometimes cut through the Lips of the Wound, and deſtroy the Suture.

Large Wounds require many Stitches.

XIII. In a Wound of a middling Size, that is to ſay, of about two Fingers Breadth, one Stitch in the Middle will be ſufficient: but in larger Wounds, the Stitches muſt be repeated in Proportion to their Size, leaving a Thumb's Breadth between each of the Sutures, the Extremities of the Thread hanging down on each Side, as we have ſhewn you in *Plate* III, *Fig.* 17. and in *Plate* IV, *Fig.* 15. Having made the proper Number of Sutures, an Aſſiſtant ſhould keep the Lips of the Wound together, whilſt the Surgeon faſtens the Ends of the Threads in Knots.

How the Knots are to be made.

XIV. Both Ends of the Thread are to be taken up, and to be tied firſt in a ſingle, then in a ſlip Knot, as we taught above in *Chap.* I. *N.* XLIV. *and* XLV. paſſing a ſmall Bolſter between the two Knots, (*Plate* II, *Fig.* 22.) to prevent the Skin from being hurt: where there are more Sutures than one, you muſt begin at the upper Part of the Wound, tying them down in Order, that, before the laſt is tied, a ſoft Tent, of the Size of a Finger, with a Thread faſtened to the End of it, may be introduced into the lower Part of the Wound: this Tent will keep a Paſſage open for the Evacuation of grumous Blood or Matter, which may be collected in the Cavity of the Abdomen: ſome of the modern Surgeons, particularly GARENGEOT, abſolutely forbid the Uſe of Tents in theſe Wounds; and aſſert, that the Spaces left between the Sutures will afford a ſufficient Paſſage for the Diſcharge of Matter from the Abdomen; but I believe this frequently proves to be very falſe: ſee the Obſervation which we have added by way of Note to *N.* VII. *of this Chapter:* this one Fact has more Weight with me than all the ingenious Reaſons that can be brought to ſupport the contrary Opinion.

How the Wound is to be dreſſed after being ſtitched.

XV. The Stitches being all tied, and the Tent paſſed into the lower Part of the Wound, the Wound ſhould be well anointed with ſome vulnerary Balſam, and covered with Pledgits of Lint, a ſticking Plaſter, and Bolſters, ſecuring all with the ſcapulary Bandage. *See Plate* III, *Fig.* 1, *Lett.* B, C. At every Dreſſing the Surgeon ſhould be very cautious in removing the Bandage, Bolſters,

[a] See PALFINUS's Surgery, *Cap. de Gaſtroraphia.*

&c.

&c. the Tent ſhould be taken out and the Patient turned upon the wounded Side, that if any Matter is collected in the Cavity, it may be eaſily diſcharged. Where there is a large Collection of Sordes, it will be proper to prepare a *vulnerary Injection, ex Decocto Herbæ Agrimoniæ, Saniculæ vel Hyperici, admixto Roſarum Melle:* this Medicine ſhould be thrown by a Syringe moderately warm into the Cavity of the Abdomen twice or thrice at every Dreſſing, turning the Body afterwards upon the Wound, that the Blood and matter, which are mixed with the Injection, may be evacuated with it: having proceeded in this Manner, paſs a new Tent into the Wound, moiſtened with ſome digeſtive Ointment, and dreſs up as before: this Method of cleanſing the Parts, and dreſſing the Wounds, is to be repeated daily, till there remain no Signs of any Foulneſs within: after this the Tent may be removed, and the Wound healed after the uſual Methods. To forward this Intention, Reſt and proper Regulations in Diet are very ſerviceable, adviſing the Patient to lye as much as poſſible upon the Wound, placing a ſoft Pillow immediately under it; for, by this Poſture, the Matter meets with a more ready Diſcharge; and the Lips of the Wound are induced to heal, from conſtant Preſſure.

Another Caſe which requires this Operation.

XVI. Thoſe Wounds alſo require the Suture, which are extended as far as the *Peritonæum*, though they do not break through it into the Cavity of the Abdomen: for in this Caſe the *Peritonæum* is in conſtant Danger of being too much diſtended, from the vehement Motions of all the Parts of the Abdomen, in Breathing, Riſing, Walking, at the Expulſion of the Excrements, *&c.* Upon the Relaxation of this Membrane the Inteſtines would ſoon make their Way between the Muſcles, and bring on very bad Symptoms and dangerous *Herniæ*. Theſe Miſchiefs cannot better be prevented than by performing the Operation deſcribed above at *N.* XI, XII, XIII. but we muſt obſerve in this Caſe that as the *Peritonæum* is not wounded here, the Needle muſt paſs only through the Muſcles, and common Integuments.

The quilled Suture uſed here.

XVII. The Surgeons for many Years performed this Operation of ſtitching up Wounds of the Abdomen with the interrupted Suture, and preferred that to all other Methods: but ſeveral amongſt the Moderns, as we hinted above, prefer the quilled Suture (by the *French* called *Enchevillee*) in all large deep Wounds, but more particularly in Wounds of the Abdomen: for as the Muſcles of the Abdomen, above all other Parts, are ſubject to violent Motions in Breathing, Sneezing, Coughing, and from many other Cauſes, by which Motions the Threads have ſometimes burſt through the Lips of the Wound, and great miſchiefs have enſued; ſome modern Surgeons therefore, and particularly DIONIS, have introduced the quilled Suture again in this Caſe, which had been before rejected: but to prevent, the Lips of the Wound from ſuffering by the Preſſure of the Pieces of Wood, which were formerly uſed in this Suture, he ſubſtitutes Rolls of Silk ſpread with ſome Plaſter in their Room, as we ſhewed above, *B.* I, *Chap* I, *N.* XLVI. and at *Plate* IV, *Fig.* 16. by which Method the Cure will be effected more ſucceſsfully, and with more Eaſe to the Patient: in large tranſverſe Wounds of the Abdomen, which do not paſs through the *Peritonæum*, PALFINUS adviſes the Uſe of this Suture, which is to be performed according to the Method I have deſcribed above in the firſt Chapter, *N.* XLVI.

XVIII. GA-

GARENGEOT's Method.

XVIII. GARENGEOT prefers this Suture to all others, even in Wounds that penetrate into the Cavity of the Abdomen, and recommends the following Manner of performing it: inſtead of a double Thread, he twiſts ſix or eight ſtrong Threads together, like a Lace, and waxes them well, paſſing them through the Eye of a large crooked Needle, ſuch an one as is deſcribed in *Plate* VI, *at Fig. 5 or 6.*. The Surgeon takes hold of the Needle at the blunt End with his Right-hand, and paſſes the Thumb of his Left-hand into the Wound, raiſing the upper Lip with it, whilſt he fixes the Fingers of the ſame Hand upon the external Part of the Lip: he then introduces the Point of the Needle into the Abdomen, and raiſing it up about two Fingers Breadth from the Opening of the Wound, pierces through the *Peritonæum*, Muſcles, and common Integuments: then taking off the Needle, he fixes it to the other End of the Thread, and lifts up the lower Lip of the Wound, by introducing his fore and middle Fingers under it, and clapping his Thumb upon the external Part of it; and pierces it with the Needle in the ſame Manner he did the upper Lip: if the Wound is four Fingers long, it will be neceſſary to make two Stitches, at equal Diſtances from each other, and from the Extremities of the Wound; if it is leſs[a], one will be ſufficient; if it is larger, more than two are required: the Threads are to be unravelled and divided into three equal Parts at each End; through two of theſe Parts the Rolls of Plaſter are to be paſſed on each Side of the Wound, and to be faſtened on with Bow Knots: then the *Balſamum Arcæi* muſt be laid on with ſome Lint, and that faſtened again, though not too tight, with the remaining Part of the Threads, in the ſame Kind of Knots: the Abdomen is to be well bathed *cum Oleo Roſarum calido, pauco Spiritu Vini admiſto:* this Embrocation is to be uſed chiefly upon the Parts near the Wound, and about the Region of the Navel; a large Bolſter wet with the ſame Medicine is to be applied to theſe Parts, and over this another dipt *in Oxycrato calido.* Theſe Applications are to be covered with Flannel well ſoaked in an emollient Decoction: the whole is to be ſecured with the Scapulary Bandage and Napkin; the Napkin is ſuſpended by the Scapulary, which in this Caſe ought to come ſomewhat lower than uſual. *See Plate* III, *Fig.* 1, C.

How to heal the Wound.

XIX. When the Lips of the Wound about the Sutures appear to be united, which you will perceive by the looſening of the Threads, you may cut the Knots, one after another, either at the ſame Time, or on different Days, as you ſhall ſee Occaſion: and when you have gently drawn them away, as we taught you at *N.* XLIX, the reſt of the Cure will eaſily be performed by the Aſſiſtance of ſome vulnerary Balſam and ſticking Plaſters. You muſt take great Care not to draw the Stitches too ſoon, for by that Means the Lips of the Wound would burſt open again, and bring on grievous Miſchiefs: beſides, a Bandage muſt be continued on the Abdomen for a conſiderable Time.

EXPLANATION *of the* THIRD PLATE.

Fig. 1, *Letter* A, deſcribes how the *Grand Capital Bandage* is to be applied after the Operation of the Trepan, or after Wounds of the Head.

[a] GARENGEOT, *Operat.* Tom. I. p. 220 Edit. 2d. But I wiſh he had been more accurate in deſcribing how the quilled Suture could be performed by one Puncture; for two at leaſt are required to keep the Quills firm.






Plate 3.


Fig: 1.
A
K
D
C
I
B
O
G
L
M
H
N
Fig: 2.
F
E
Fig: 3.
Fig: 4.
Fig: 5.
A
Fig: 6.
Fig: 8
C
Fig: 7.
b
c
a


Fig. 6.
A
Fig: 7.
C
D
B
Fig: 9.
10.
11.
12.
13.
14.
15.
16.




Fig: 2.

B, The *Belt* or Napkin, which is to ſurround the Body in Wounds of the Thorax or Abdomen, to bind the Applications to the Part affected.

C, The *Scapulary*, to ſupport the Belt.

D, The Method of making the Ligature after Bleeding in the Arm.

E, The Manner of tying up the Foot after Bleeding, which, from the Similitude it has with a *Stirrup*, is called by that Name.

F, Shews the ſpiral Manner in which the Bandage ought to aſcend, when it is applied to the Leg or Arm; which is ſometimes called *the expelling Bandage.*

G, A ſerpentile Bandage, where the Convolutions are not ſo frequent.

H, A large Wound in the Thigh, which requires the true or bloody Suture.

K, The Part where the *Tourniquet* is to be applied to the Arm, and the Manner of applying it.

L, The Manner of applying it to the upper Part of the Thigh; the Bandage which is rolled up, and applied as a Bolſter lyes upon the crural Artery at M.

N, Shews how the *Tourniquet* is to be applied to the lower Part of the Thigh, in which Caſe the rolled Bandage is to be applied to the back Part of the Thigh.

O, A large Wound of the Abdomen, with the Inteſtines falling out.

Fig. 2. The common *Tourniquet*, before it is applied.

Fig. 3. Crooked Forceps furniſhed with Teeth at the End, called the *Crane's Beak.*

Fig. 4. A Pair of ſtrait Forceps.

Fig. 5. The *Duck's Bill* Forceps, furniſhed with a moveable Ring at the lower End.

Fig. 6. The *Gooſe Bill* Forcepts.

Fig. 7. The Inſtrument invented by BARTHOLOMÆUS MAGGIUS to extract Bullets that are fixed in a bony Part.

Fig. 8. A Hook to extract Bullets.

Fig. 9, 10, 11, 12, 13, 14, 15, 16. Different Sorts of Cauteries.

Fig. 17. Shews in ſome Meaſure the Manner of performing the Operation called *Gaſtroraphy*, or the Suture of the Abdomen. The Letters *a a* deſcribe the Wound; *b b*, two crooked Needles with the Threads hanging to them; *c c c c*, two Threads drawn through the Lips of the Wound, and cleared from their Needles.

CHAP. VI.

Of WOUNDS *of the* INTESTINES, *and the Manner of ſtitching them up.*

I. WHEN a large Wound is made in the Cavity of the Abdomen, that not only lets out the Inteſtines, but alſo divides ſome Part of them, (*See Chap.* V, *N.* IV.) the Surgeon ought always to ſtitch up the wounded Parts of the Inteſtines, before he returns them. By this Means we may not only expect the Wound to heal more readily, but the Diſcharge of Chyle and Fæces into the Cavity of the Abdomen, which would bring on great Miſchief, is prevented: and although Wounds of the Inteſtines, eſpecially of the ſmall Guts, admit of little or no Hopes of a Cure; yet as the great Guts, as CELSUS obſerves,

When the Suture of the Inteſtines is to be performed.

obſerves, *Lib.* VII, *Cap.* 16. ſometimes admit of the Suture to Advantage, it is better to uſe a doubtful Remedy than none: therefore the Surgeon ſhould never neglect examining whether the Inteſtines are injured, that he may uſe all probable Means of healing them. *See above Chap.* V, *N.* IV, V.

When the Suture is not to be attempted.

II. Small Wounds of the Inteſtines, that do not exceed in Size the Diameter of a Gooſe Quill, ſhould by no Means be ſtitched, but are beſt left to Nature. If they are left to themſelves, they will frequently unite much ſooner than if they are irritated by the Suture: for Stitching uſually brings on great Pain, Inflammation, and other bad Symptoms: therefore it will be much better to return them inſtantly, (*See Chap.* V, *N.* V.) and to bleed the Patient to prevent Inflammation, adviſing him to Reſt and Abſtinence; for it is better by Induſtry and Care to cheriſh even ſmall and glimmering Hopes, than through Fear and Negligence to give the Patient over.

How the Opening is to be performed.

III. But large Wounds of the Inteſtines, though they ſeldom admit of Cure, are to be ſtitched up with *the Glover's Suture*, before the Inteſtine is returned: to perform this, you ſhould be provided with a fine Needle threaded with Silk. An Aſſiſtant ſhould take Hold of one Part of the Gut, with a fine Piece of Linen well aired before the Fire; whilſt the Surgeon ſhould hold the other Part in his left Hand, and ſew up the whole Wound after the Glover's Manner, leaving very ſmall Spaces between each Stitch; to wit, little more than a Mathematical Line. The laſt Stitch ſhould be faſtened with a Knot, but the other End ſhould hang about a Foot out of the Abdomen, by which the Silk may be drawn out when the Inteſtine is healed, *See Plate* IV, *Fig.* 20. Some in this Caſe prefer the *interrupted Suture*, becauſe it is performed with fewer Punctures, and therefore it is not liable to bring on ſo great Inflammation; though the Threads, which are very ſmall, ſhould be left behind. GARENGEOT propoſes another Method of performing *the Glover's Suture*, *in Operat. Chirurg. Artic. de Gaſtroraphia.* But, to ſay Truth, Experience ſhews us, that very few are ſaved, whatever Suture is made Uſe of.

What is to be done afterwards.


IV. After this Operation is performed, the Wound of the Abdomen is next to be taken Care of, and ſtitched up, as we have ſhewn in the former Chapter upon that Subject, always obſerving the Caution I there laid down, which I cannot inculcate too frequently, of keeping the depending Part of the Wound open with a Tent, till all the præternatural Fluids are diſcharged from the Cavity of the Abdomen, and until the Union of the Wound in the Inteſtine ſhall render it proper to draw out the Silk with which it was ſtitched up.

How the Wound is to be dreſſed.

V. There is no Neceſſity for explaining to you the Method of cleaning, dreſſing, and healing the Wound. We have already ſpoken ſufficiently on that Subject in *Chap.* V, *N.* XIV. and the following Numbers. Only I would furniſh the Surgeon with this farther Caution, that where two Threads hang down from the Belly, one belonging to the End of the Tent, the other to the Suture of the Inteſtine, it will be proper to diſtinguiſh them by different Colours, to prevent miſchievous Miſtakes.

An eaſier Method of healing Wounds of the Inteſtines.

VI. As the modern Surgeons have found by Experience, that ſcarce any are ſaved who have received Wounds in the Inteſtines, and that in thoſe few who do recover, the wounded Parts, from the Fineneſs of the Coats of the Gut, do not properly unite, but rather adhere to the inner Part of the *Peritonæum*, or to the *Omentum*, or to ſome of the other Inteſtines; it is no Wonder, therefore, that they entirely



Tabl. 4.

Fig. 1.




Fig. 3.



Fig. 4.



Fig. 5.




Fig. 6.




Fig. 7




Fig.






Fig. 11.




Fig. 12.




Fig. 13.




Fig. 14.



Fig. 15.




Fig. 16.




Fig. 18.




Fig. 19.




Fig. 2.



Fig. 8.




Fig. 9.





Fig. 10.






Fig. 16.




Fig. 17.

Fig. 20.



Fig. 21.





Fig. 22.






Fig. 3.



Fig. 4.




Fig. 10.




Fig. 11.




Fig. 14.



Fig. 17. E



Fig. 18.




Fig. 22.

intirely laid aſide the Practice of ſtitching up the wounded Parts of the Inteſtine, eſpecially with uninterrupted Stitches, like the *Glover's Suture:* which by the Frequency of the Puncture brings on a violent Inflammation, the moſt acute Pains, Convulſions, nay, ſometimes Cancer or Mortification, and Death itſelf: but they rather chuſe now to deal more tenderly with the Patient, and to ſubſtitute a gentler Method of Cure; in Conſequence of which, the preſent Practice is to paſs a waxed Thread through a fine Needle, and with this to faſten the wounded Part of the Inteſtine to the internal Orifice of the Wound of the Abdomen. The Thread that hangs out of the Abdomen is to be ſo firmly fixed by the Application of ſticking Plaſters to the Wound, that the Inteſtine cannot recede from the Part to which it was faſtened, nor can it evacuate any of its Contents into the Cavity of the Abdomen. When this Operation is well performed, the Inteſtine eaſily adheres to the internal Part of the Abdomen, and the Patient ſuffers infinitely leſs Pain and Hazard than from the former Method of making the Sutures. The ſame Regulations in Diet, and the ſame Methods of Dreſſing, and the bleeding the Patient, which we adviſed above at *Chap.* V, *N.* XIV. *and the following Numbers*, are to be obſerved. The ſame Method of Cure will alſo ſerve for Wounds of the Stomach, where they are within the Reach of the Hand, and it is ſometimes crowned with Succeſs. *See* BOHNII *lib. De Renunciatione Vulnerum, Sect.* II. *Chap.* V.

EXPLANATION *of the* FOURTH PLATE.

Fig. 1. PETIT's *triangular Needle*, for making a new Aperture in the Part oppoſite to the Wound, which the *French* call *Contre-ouverture.*

Fig. 2. My Improvement upon PETIT's *Needle*, which will take Place where a ſtrait Needle cannot ſafely be uſed, *See Book* I, *Chap.* I, *N.* XXXVII.

Fig. 3. AA, repreſent a Wound, the Lips of which are to be united by the ſticking Plaſter indented on both Sides at B B.

Fig. 4. Shews a Wound to which two ſticking Plaſters are applied.

Fig. 5. A Wound of the like Nature, to which are applied two ſticking Plaſters without Indentations.

Fig. 6. A Wound made croſs-ways, AAAA, united by two Plaſters laid croſs-ways, BBBB.

Fig. 7. A Wound AA, to which a ſticking Plaſter is applied, with two Openings in the Middle, B B.

Fig. 8. A Wound united by the Application of two Plaſters, with Tapes fixed to each of them, which are drawn together and faſtened with ſlip Knots, *a a a.*

Fig. 9. The ſame Wound with Plaſters of the ſame Kind, furniſhed with Hooks, *a a a*, inſtead of Tapes, by which, with the Aſſiſtance of Threads tied to them, the Lips of the Wound are drawn together.

Fig. 10. Another Method of doing the ſame Thing, uſed by the Antients.

Fig. 11. A tranſverſe Wound, AA, united by the *interrupted Suture*, B B.

Fig. 12. Shews in what Manner a croſs Wound is to be ſtitched up, and the Lips of it brought together by drawing the Threads tight, ABCD.

Fig. 13. Where the Stitches are to be made in a triangular Wound, ABC.

Fig. 14. How a Wound with two Angles is to be ſtitched with the *interrupted Suture*, firſt at the Angles, AA, and then, if it is neceſſary, on each Side at the Letters, B B.

Fig 15. *A large crooked Needle, for ſtitching large Wounds, with a double Thread,* to make the *quilled Suture.* A is the Needle; B the double Thread; C the Bow-end of the Thread.

Fig. 16. A large tranſverſe Wound, AA, united by a triple *interrupted Suture*, BBB.

Fig. 17. The ſame Kind of Wound, DD, which beſides the Threads at *Fig.* 16. is furniſhed alſo with ſmall cylindrical Rolls of Silk ſpread with ſome Wax or Plaſter, AA and BB: the Threads on the upper Lip of the Wound are tied in ſlip Knots, CCC, whilſt the Roll that lies on the under Lip is confined between the Bow-ends of the Threads, EEE. In a Word, this ſhews Palfinus's Method ofmaking the *quilled Suture.*

Fig. 18. Shews you another Method of making the *quilled Suture* in large tranſverſe Wounds, particularly in thoſe of the Belly, which is called *Gaſtroraphy*, *See Book* I, *Chap.* V, *N.* XLVII; *and Chap.* V, *N.* XVIII. AA, the Wound. BB, the upper Roll. CC, the lower Roll. DDD, the ſingle Knots which confine the Lace, compoſed of ſix or eight Threads, and the upper Roll. EEE, the ſlip Knots which ſecure the lower Roll.

Fig. 19. Celsus's *Suture*, which he deſcribes at *Lib.* VII, *Chap.* XVI, for performing the Operation of *Gaſtroraphy* with two Needles. But this is a bad Method, and out of Practice. AA, the Stitches. BB, the End where they are faſtened in a Knot.

Fig. 20. The *Glover's Suture*, uſed for uniting Wounds of the Inteſtines. AA, the Inteſtine. BB, the Wound. C, the Beginning of the Suture, with Part of the Thread hanging out. D, the End of the Suture, where it is faſtened in a Knot.

Fig. 21, 22. *The Suture for the Hare Lip*, which is made with two or three Needles. AA, the deſcending Wound. BB, Needles paſſed through the Lips of the Wound. CCC, the Thread twiſted round the Needles.

CHAP. VII.

Of Loss of Substance *in the* Intestines.

What is to be done when there is a Loſs of Subſtance.

I. WHERE any Part of the Inteſtine is carried away, the Caſe ſeems to be plainly deſperate. It was therefore wonderful that Perſons thus wounded did not all die upon the Spot, or in the Operation of making the Sutures: till [a] Hildanus, [b] Blegny, [c] Dionis, [d] Palfynus, [e] Jo. M. Hoffman, [f] Schacher, [g] Vater, [h] Cheselden, Heister, and others, obſerved, that the Lips of Inteſtines ſo wounded, would ſometimes quite unexpectedly adhere to the Wound in the Abdomen; and therefore there ſeemed to be no Reaſon why we ſhould not take this [i] Hint from Nature. Whenever therefore a Surgeon is called

[a] Obſerv 74. Cent. I. Obſ. 72. Cent. VI. [b] Zodiac. Med. Gall. An. 2. pag. 123. [c] In Chirurg. cap. de Gaſtroraphia. [d] In Chirurg. cap. de Gaſtroraph. [e] Diſq. Corp. Hum. Anat. Path. [f] In Diſſert. de Morb. ex ſitu Inteſtin. [g] In Diſſertat. de Vuln. in Inteſtin. lethal. [h] Lib. de alto apparatu.

[i] A Surgeon tried this firſt with Succeſs upon a Dog. *See* Bleng *Zodiac. Gall. An.* 2. p. 143. afterwards it was performed upon a Man. *See Miſcell. Natur. Curioſ. Dec.* 2. *An.* 8. *Obſ.* 229.

to

to a Case of this Kind, after he has diligently examined the State of the upper Part of the Intestine, which has suffered a Loss of Substance, he should stitch it to the external Wound, either by the continued or interrupted Suture: for by this Means the Patient may not only be saved from instant Death, but there have been Instances where the wounded Intestine has been so far healed, that the Fæces, which used to be voided *per Anum*, have been voided by the Wound in the Abdomen; which, from the Necessity of wearing a Tin or Silver Pipe, or keeping Cloths constantly upon the Part to receive the Excrement, may seem to be very troublesome: but it is surely far better to part with one of the Conveniencies of Life, than to part with Life itself: besides, the Excrements that are voided by this Passage are not altogether so offensive, as those that are voided *per Anum*.

How a mortify'd Gut is to be treated.

II. The same Method of Cure may conveniently enough be put in Practice, where any Part of the Intestine is mortified by being forced out of the Abdomen. For in this Case, if you tie up the mesenteric Arteries, the corrupted or mortified Part of the Intestine may be cut off, and the remaining sound Part made to adhere to the Wound of the Abdomen: for it is better to try this Method, though but few should be saved by it, than to suffer all to perish, as Celsus observes; it is wiser to attempt a doubtful Remedy, than absolutely to despair: I once published a Cure of this Kind in a Dissertation containing various Observations, printed at *Helmstadt*.

How concealed Wounds of the Intestines are to be treated.

III. When the Intestines are wounded, but not let out of the Abdomen, and therefore their Wounds are out of Reach, the Surgeon can do nothing but keep a Tent in the external Wound, according to the Method of Dressing laid down at *Chap.* V, *N.* XIV. and after this, bleed the Patient, if his Strength will admit of it, advising him to rest, to live abstemiously, and to lie upon his Belly: the rest is to be left to divine Providence, and the Strength of his Constitution: but the Question may be asked here, whether a Surgeon may not very prudently, in this Case, enlarge the Wound of the Abdomen, that he may be able to discover the injured Intestine, and treat it in a proper Manner? truly I can see no Objection to this Practice, especially if we consider, that, upon the Neglect of it, certain Death will follow; and that we are encouraged to make Trial of it by the Success of others. Schacherus, *in Programmate publico, Lipsiæ edit.* 1720, mentions a Surgeon who performed this Operation successfully: so Cheselden of *London* gives us an History where in the *Hernia incarcerata* he laid open the Abdomen, returned the Intestines, and perfectly cured his Patient. *See his Treatise on the high Operation*, pag. 180. *and his Anatomy*, 3d edit. pag. 283.

Of Clysters in this Case.

IV. But what Assistance are we likely to receive from Clysters in Wounds of the Intestines? Some Physicians are very high in their Commendation, whilst others, of equal Credit, absolutely prohibit the Use of them: for my own Part, I see no Reason for carrying either Prejudice to so great a Length: the Use of Clysters is very prudently forbid in Wounds of the great Guts, but no less Judgment is shewn in prescribing them in Wounds of the small ones: in the first Case, the Clyster will make its Way through the Wound, into the Cavity of the Abdomen, to the great Detriment of the Patient; whereas in the latter, they will always prove beneficial: for the Inconvenience which attends the other is prevented in this Case by the Valve of the *Colon*; and the Benefits that accrue from

this Application are very obvious: the uſeleſs Fæces are carried off, an equable Courſe of the Blood is reſtored, the Fever and Inflammation are much abated by it, if not entirely taken off, and the Pains greatly aſſwaged.

CHAP. IX.

Of the FALLING-DOWN *of the* OMENTUM.

Falling down of the *Omentum.*

I. IN large Wounds of the Abdomen, the *Omentum* will frequently protrude itſelf through the Wound, either alone or with ſome Portion of the Inteſtines: whenever this is the Caſe, the Surgeon's firſt Inquiry is, whether the protruded Part preſerves its Heat, Moiſture, and natural Colour? if it is not found faulty in any of theſe Circumſtances, it muſt be gently returned: but where the Straitneſs of the Wound forbids this, the protruded Part muſt be taken off cloſe to the Wound, and the Wound healed according to the common Form: the *Omentum* in this Caſe will adhere to the internal Part of the Wound, without bringing any Diſorder upon the Patient: but where the Inteſtines fall out at the ſame Time, the *Omentum* is to be fomented, by an Aſſiſtant, with warm Milk and Water, till the Inteſtines are returned, and then the *Omentum* muſt be replaced.

What is to be done when the *Omentum* is corrupted.

II. If any Part of the protruded *Omentum* is cold, dry, livid, mortified, or putrid, the diſcoloured corrupted Part muſt be entirely cut off before the reſt is returned; leſt the neighbouring Parts ſhould be brought into Conſent, which ſhould inevitably prove fatal to the Patient: GARENGEOT would have the vitiated *Omentum* returned wholly into the Abdomen, without Ligature or Inciſion; but that this Procedure is wrong, I ſhall ſhew more fully in the following Numbers.

How the corrupted Part is to be taken off.

III. The corrupted Part of the *Omentum* may be taken off in this Manner: take a ſtrong waxed Thread, paſs it two or three Times round the ſound Part of the *Omentum*, near the Place where it is injured, and faſten it with a Knot, to prevent any Hæmorrhage enſuing after the Reduction of it: when you have made a ſecure Ligature, take off the corrupted Part with the Knife or Sciſſars; and return gently what remains ſound, leaving at leaſt the Length of a Foot of the Ligature hanging out of the Wound in the Abdomen, till it ſlips off from the ſound Part of the *Omentum*.

How the Wound is to be dreſſed.

IV. What remains with regard to the Cure of the Wound is ſufficiently treated of above at *Chap.* V, *N.* XIV. *and the following Numbers:* the depending Part of the Wound ſhould be kept open with a large Tent, ſuch a one as is deſcribed *Plate* II, *Lett.* O, that a Paſſage may be left for an Evacuation of the Sordes from the Cavity of the Abdomen: it will be proper to give two different Colours to the Thread that hangs from the Tent, and that which belongs to the Ligature of the *Omentum*, to prevent Confuſion, as we adviſed above, *Chap.* VI, *N.* V.

How the Thread is to be drawn out.

V. At every Dreſſing, after the firſt ſix or ſeven Days, you may draw the Thread which hangs out of the Abdomen gently forwards, till it ſhall by Degrees ſlip quite off the *Omentum*: but this ſhould be done without any Violence: when the Thread is drawn out, and you can perceive no Diſcharge from the Cavity of the Abdomen, you may remove the Tent, and uſe proper Means to heal

heal the external Wound: after which you ſhould take away ſome Blood, unleſs there has been a ſufficient Diſcharge before; and recommend Reſt and Abſtinence to the Patient.

VI. What ſhall we ſay to the unwarranted Opinion of DIONIS? who adviſes Surgeons never to take off any Part of the *Omentum*, but rather to follow the Example of MARESCHAL, *firſt Surgeon to the French King*, who, according to our Author's Account, has very frequently returned the *Omentum* without making either Ligature or Inciſion, and never ſaw any bad Conſequence from this Practice: but I will venture to pronounce this Relation of DIONIS's to be very faulty, and not delivered with that Accuracy which is required in a Matter of Fact of this Conſequence: we cannot learn, by this Account of his, whether the *Omenta*, which were returned in this Manner by MARESCHALL, were large or ſmall, whether they were entirely ſound, or corrupted in Part: if they had received no Injury, DIONIS ſpends his Time idly, when he ſo earneſtly entreats all Surgeons to follow the Steps of MARESCHALL in this Point: nobody ever adviſed the contrary: but if they were in Part corrupted or mortified, which DIONIS does not aſſert, it is much to be admired that the Patients felt no Inconvenience from this Practice; and what became of the corrupted Parts after they were returned, is to me Matter of great Wonder. Therefore DIONIS is by no means to be attended to upon this Point, till he ſpeaks to it in a clearer Manner: and more particularly ſo, becauſe PALFINUS gives us the Hiſtory of a Caſe, in his Surgery, where MARESCHALL made a Ligature and Inciſion upon the *Omentum*, and ſeparated the corrupted Part from the ſound, before he returned it: and this he declares to be the Practice of Surgeons of the firſt Name in PARIS. DIONIS's Advice in this Caſe.

VII. GARENGEOT declares himſelf of the ſame Sentiments with DIONIS, though he makes no Mention of his Name: this Author is far from being clear in deſcribing how large a Portion of the *Omentum* was affected, which MARESCHALL, or any other, returned, without Injury to the Patient: I do not deny but that a very ſmall Portion of the *Omentum* may be digeſted in the Abdomen, without bringing on any conſiderable Miſchief: but I can by no Means be perſuaded that this can ever be the Caſe, when a large Portion of the *Omentum* is affected, except I ſhould be confronted with many Inſtances of it: if by Chance one Inſtance ſhould be produced, this will not put the Matter out of Doubt, much leſs ſerve as an Example worthy of Imitation: for miraculous Events happen now and then in very dangerous Wounds: and ſince grievous Symptoms are brought on by letting Sordes remain even in external Wounds, what may we not fear from the ſame Incident in internal Wounds, from whence they cannot poſſibly be diſcharged? A large Degree of Suppuration is to be expected when a large Portion of corrupted *Omentum* is returned into the Body: but when a Ligature is made upon the *Omentum*, and the corrupted Part ſeparated from the ſound, no ſuch Accident can happen: the Suppuration in this Caſe will be very inconſiderable; and the ſmall Quantity of Matter that is made after Reduction, will be eaſily diſcharged through the external Wound that is kept open for that Purpoſe by a Tent: whereas GARENGEOT forbids the Uſe of Tents promiſcuouſly (which this very MARESCHALL uſed with great Succeſs) and adviſes you to heal the Wound as ſoon as poſſible: I am of Opinion, therefore, that you ſhould very carefully diſtinguiſh between a great and ſmall Degree of Suppuration, becauſe this is of greater Conſequence than GARENGEOT ſeems GARENGEOT's Advice.

ſeems to imagine. Since this Matter is left doubtful, and GARENGEOT no where pretends to have had Experience of the good Effects of the Practice which he eſpouſes; but on the other hand, PALFINUS, who was an Eye-witneſs, contradicts him; I think we may very ſafely imitate the Examples of many excellent Surgeons, in making a Ligature upon the *Omentum*, and ſeparating the corrupted Parts of it from the ſound, before we attempt to return it into the Abdomen.

CHAP. IX.

Of WOUNDS *of other Parts of the* ABDOMEN.

IF you can diſcover by your Eye, or by the Touch, that any other Part or *Viſcus* ſituated in the Abdomen, ſuppoſe the Liver, Spleen, or Kidney, has received a Wound from a ſharp Inſtrument, it will be adviſeable, at the firſt Dreſſing, to fill the Wound as tenderly as poſſible, with a good Quantity of Lint, well ſaturated with high rectified Spirit of Wine, or Spirit of Turpentine, ſecuring the Dreſſings with Compreſſes and Bandage: by this Means the Hæmorrhage will be ſtopped, if no large Blood-veſſel is divided; but you muſt obſerve that the Lint be not removed at the firſt Dreſſings; it ſhould remain till it drops off itſelf: when you have gained this Point, the Wound may be treated according to the Rules we laid down for the Treatment of Wounds of the Abdomen; the reſt muſt be left to God's Providence, and the Strength of the Patient's Conſtitution: during the Cure the Patient muſt be conſtantly kept ſtill and low: if he is of a plethoric Habit of Body it will be proper to bleed him, to prevent Inflammation, and freſh Effuſions of Blood; preſcribing him alſo vulnerary Potions, and giving him daily two or three Doſes of *Lucatellus's* or *Meibomius's Balſam*, for theſe Balſams are of great Efficacy in healing internal Wounds: in hidden Wounds of the Viſcera, that are not to be diſcovered by the Eye or by Feeling, all you can do is to take proper Care of the external Wounds, daily injecting a vulnerary Decoction, and keeping open a free Paſſage for the Evacuation of grumous Blood and Matter from within, ordering the ſame Regimen to be obſerved, both with regard to Medicine and Diet, which we adviſed above, and leaving the reſt to Nature, for Art can give no further Aſſiſtance.

EXPLANATION *of the* FIFTH PLATE.

Fig. 1. Deſcribes a blunt Iron Needle, to paſs a fine Rag or Skein of Silk, well ſaturated with proper Balſams or Ointments, through *Gun-ſhot*, or other *pervious Wounds*, after the Manner of a *Seton*.

Fig. 2. An Inſtrument to ſtop the Blood in Wounds of the large Arteries, deſcribed in *Ch.* II. § VIII. from our Amendment.

AA, A braſs Plate ſomewhat bent.

BB, A ſtrong braſs Screw.

C, A round Plate of a Thumb's Breadth to be fixed upon the Wound.



B
Fig. 8.
Fig
E
E
F
F
E
A
Fig. 9.
A
A

Plate 5.




Fig. 4.



Fig. 5.






Fig. 3.




Fig. 6.




Fig. 7.







Fig. 9.



D, The Button which turns the Screw, and preſſes the Plate C ſtrongly upon the Wound.

E E, A ſtrong leather Belt to ſurround the wounded Part.

F F, Part of the Belt pierced with ſeveral Holes, by which it may be fixed upon the Hooks G G, and lengthened or ſhortened according to the Size of the Limb.

Fig. 3. A crooked Knife, with a round blunt Point, to enlarge Wounds of the Thorax or Abdomen, where that Operation is required.

Fig. 4. A ſtraight Knife with a Button on the Point.

Fig. 5. A crooked Knife with a blunt Point.

Fig. 6. A wooden *Tournequet* in its proper Size, to ſtop Effuſions of Blood, after our Amendment deſcribed above at *Chap.* II, § XII.

A A, The upper Part.

B B, The lower Part.

C, The great Screw.

D, Two ſmall Iron Screws, to which a leather or ſilk Belt is to be fixed.

E, Hooks to faſten the other End of the Belt on, when it is brought round the Limb.

F F, The Ends of the upper and lower Part of the Inſtrument hollowed to receive the Belt, and to keep it ſteady in its Situation.

Fig. 7. Another kind of *Tournequet* made of Iron; the Deſcription is leſs by half than the proper Size of the Inſtrument. *See Chap.* II, § XIV. where it is largely treated of.

Fig. 8. A broad Bandage, called the *uniting Bandage*; this is perforated in the Middle, and rolls up with two Heads; it is uſed in dreſſing Wounds of the Abdomen, which are made lengthways.

Fig. 9. A flexible ſilver Pipe, uſeful to diſcharge the Matter which is collected in Wounds of the Thorax, or in the Empyema.

A, The Openings at the Extremities, and on both Sides.

B B, The Plate round it, with two Holes to paſs a Thread through.

C, The Paſſage that goes through the Pipe to A.

CHAP. X.

Of WOUNDS *of the* THORAX.

Wounds of the Breaſt of three Sorts.

I. WOUNDS of the *Thorax*, as of the Abdomen, are divided into three ſorts: the Wound is inflicted either upon the external Parts of the *Thorax* only; or elſe it penetrates into the Cavity of the *Thorax*, without injuring any of its Contents; or laſtly, the Contents of the *Thorax* alſo partake of the Wound.

How to diſcover whether the Wound is terminated in the external Parts.

II. You may diſcover that the Wound terminates in the external Parts, and does not penetrate into the Cavity of the *Thorax*, by ſeveral Methods: 1. by the Sight: 2. by the Senſe of Hearing, by which you will diſcover whether any Sound proceeds from the Wound at the Time of Inſpiration: 3. by Feeling, when

when your Finger or the Probe meets with Reſiſtance, if you attempt to paſs it into the Cavity of the *Thorax*. 4. by injecting warm Water, which in this Caſe will return ſtrongly upon you: 5. by the Abſence of bad Symptoms, ſuch as Difficulty of Breathing, Fainting, ſick Fits, *&c.* which always attend a Wound that penetrates: when by theſe Methods of examining you are fully ſatisfied that the Wound does not penetrate, you may dreſs it with a digeſtive Ointment, or ſome vulnerary Balſam, and treat it according to the Methods which we have adviſed above for the Cure of ſlight Wounds.

What deep and oblique external Wounds of the *Thorax* occaſion.

III. It ſometimes happens that external Wounds run very deep and obliquely between the Muſcles and the Ribs, and are thereby rendered very difficult to be cleanſed from grumous Blood and Matter. The confined Matter in this Caſe frequently deſtroys the neighbouring Parts, and produces Ulcers and incurable Fiſtulæ: nay, ſometimes it makes its Way through the *Pleura* into the Cavity of the *Thorax*, and forms an *Empyema*, or brings on a *Phthiſis*, or Death itſelf.

How they are to be treated.

IV. The Surgeon's chief Buſineſs in this Caſe is to clear the Sinuſes from the Blood and Matter confined in them: this is to be done either by Preſſure, or by ordering the Wound to be ſucked by an healthy Perſon; by drawing it out with a Syphon, or by making further Openings with the Knife: the reſt of the Cure is to be performed after the ſame Manner which we deſcribed above, *N.* II. The moſt proper Bandage for ſecuring the Dreſſings is the Scapulary with theGirdle. *See Plate* III, *Fig.* 1. which Bandage muſt be eaſy, that the Blood, or Matter, confined in the Wound, may have the freer Vent.

How to empty the Woundwith a Syringe.

V. The *Syringes* that are uſed in this Caſe are of very different Shapes and Sizes; ſome are ſtrait, others crooked. Some Surgeons uſe a Tin Syringe, reſembling that which we have deſcribed at *Plate* VI, *Fig.* 8. but twice as large: The Mouth of it is larger than the reſt of the Syringe, and is of a triangular, round, or oval Figure. *Fig.* 9. repreſents the true Size of it. When you apply this Inſtrument, you muſt clap the Mouth of it to the Wound, and, by drawing back the Handle, endeavour to fill it with Blood. The Inſtrument ſhould have ſeveral Heads of different Sizes and Figures, that it may correſpond with any Sort of Wound: but, concerning the Excellency and Uſe of theſe Syringes, it will be worth your while to conſult ANELLE, in his Treatiſe called *l'Art de ſuccer les Playes.*

How to diſcover if th Wound penetrate the Cavity.

VI. You will diſcover the Wound to penetrate into the Cavity of the *Thorax*; 1. by the Sight, when you can plainly ſee into the Cavity: 2. by the Senſe of Feeling, when you can paſs your Fingers or Probe into the Cavity: 3. by the Hearing, if the Patient makes a particular ſort of Noiſe in drawing his Breath: 4. from the Action of the Air of the Lungs upon the Flame of a Candle, or Feathers, when they are held near the Mouth of the Wound: 5. by warm Water meeting with no Reſiſtance, when it is injected into the Wound: 6: laſtly, from the ſudden Appearance of violent Symptoms, ſuch as Difficulty of Breathing, Sickneſs, Fainting, *&c.* which are brought on by the Preſſure which the Lungs are ſenſible of from the external Air, from a Collection of Blood in the *Thorax*, or from both Cauſes together.

VII. When

What proceeds from a Collection of Blood in the *Thorax*.

VII. When a large Quantity of Blood is ſpilt, and falls into the Cavity of the *Thorax* (which muſt ſometimes be the Caſe) the Expanſion of the Lungs, the Office of Reſpiration, and the Courſe of the Blood through the Lungs, will certainly be impeded; and the Blood by frequent Delays and Obſtructions being entirely inſpiſſated in the Lungs, Life can no longer be ſupported: but where the Quantity of extravaſated Blood is not large enough to obſtruct the Lungs in their Office, the chief Danger that the Patient labours under is, that the extravaſated Blood ſhould putrify by Degrees, and corrupt the Diaphragm, Pleura, or Lungs; which will bring on very bad Symptoms, and in a ſhort Time Death.

Symptoms of extravaſated Blood in the *Thorax*.

VIII. The following Symptoms diſcover an Extravaſation of Blood in the *Thorax:* if, 1. there is a great Difficulty of Breathing, except when the Patient is placed in an erect Poſture: 2. if the Patient lies eaſieſt upon his Back or wounded Side, but finds any other Poſture exceeding troubleſome, or ſometimes impracticable: 3. if he feels a Weight upon the Diaphragm: 4. if he perceives the Undulation of a Fluid upon turning the Body round: and, 5. laſtly, if there has been little or no Diſcharge of Blood from the Wound.

How to get the Blood out of the *Thorax*.

IX. When it appears by theſe Symptoms that there is a Collection of Blood in the *Thorax*, we muſt uſe our utmoſt Diligence to get it out, leſt it ſhould lay a Foundation for great Miſchief: therefore, 1. when the Wound is inflicted upon the middle, or lower Part of the *Thorax*, and has not a very narrow Opening, it will be convenient to lay the Patient upon the [a] wounded Side, adviſing him to fetch his Breath as deep as he can, or to cough: if the Current of Blood is obſtructed by any thick grumous Parts, which will ſometimes ſtop up the Orifice of the Wound, they muſt be removed with your Finger, or with the Probe, or drawn out with a Syringe, or by Suction: 2. if you are called ſo late that the Blood is become too thick to flow out of the Wound, you will be obliged to uſe an attenuating Injection; which may be made of a Decoction of Barley, with the Addition of ſome common Honey, or Honey of Roſes, and a ſmall Quantity of Soap; this is to be injected, not over-warm, into the Cavity of the *Thorax*, and then the Patient is to be ſo ſituated as to let it run out again: this Operation is to be repeated till it appears that all the grumous Blood is waſhed away: the Syringe, which you will ſee deſcribed in *Plate* VI. *Fig.* 8. with the Pipes, *Fig.* 10, 11. will execute this Intention very properly: 3. but if the Wound is ſo narrow or oblique that this Method cannot be proſecuted, it ſhould be enlarged, either with the common Inciſion Knife and Director, or with one of the Knives deſcribed at *Plate* V, *Fig.* 3, 4, 5. This Caution is always to be obſerved, that is, to be very careful not to fatigue the Patient too much, by endeavouring to diſcharge all the extravaſated Blood at one Time: if the Patient is very weak, it is better to do it at proper Intervals, eſpecially if you diſcover any Tendency in him to Swoonings. It will be neceſſary in the mean Time to keep the Wound open by the introducing a leaden or ſilver Pipe into the Wound, ſuch as are deſcribed at *Plate* II, *Lett.* Q, R, S, or rather that flexible one at *Plate* V, *Fig.* 9. though ſome, inſtead of a Pipe, uſe a Tent with a long String at the End of it, dreſſing up with pro-

[a] DIONIS, in his *Surgery*, relates a Caſe of this Kind, where he left his Patient all Night inclined upon the Wound without dreſſing him, and he afterwards recovered him. DE LA MOTTE confirms this by an Inſtance he gives us of the ſame Kind, that occurred to him in his Practice. *See his Obſervationes Chirurgicæ.*

per Plaſters and Compreſſes, ſecuring the whole with the Scapulary, repeating this Method of dreſſing till the Diſcharge ſhall entirely ceaſe, and the external Wound can be conveniently healed.

How the Blood is to be diſcharged when the Wound is in the upper Part of the *Thorax*.

X. When a Wound is made in the upper Part of the Breaſt, or between the upper Ribs, then the Method we have preſcribed of turning the Patient upon the wounded Side, will be of very little Service in diſcharging the extravaſated Blood; for no Poſture will ſatisfy this Intention in this Caſe but ſtanding upon the Head. In this Caſe, if no Relief is to be expected from the Syringe or from Suction, an Opening ſhould be made in the lower Part of the *Thorax*, which Operation the Surgeons call the *Paracenteſis*: the Opening muſt be between the ſecond and third Rib, counting upwards, if it is on the left Side; but on the right Side, between the third and fourth, about a Hand's Breadth from the Spine: the Place where you intend to make the Opening ſhould be marked with Ink; the Inſtrument that is generally uſed upon this Occaſion is called a *Trocar*; it ſhould be driven above the Rib into the *Thorax*, with great Caution and Gentleneſs: after it has penetrated, draw out the ſteel Inſtrument, leaving in the Pipe through which it was conveyed, as a Channel for the Blood to paſs off by; but, if it does not readily paſs, its Evacuation may be forwarded by Suction, or a Syringe: but, as the Lungs are very liable to be wounded by paſſing this Inſtrument forcibly into the Cavity of the *Thorax*, it is beſt, in my Opinion, to divide the common Integuments, the intercoſtal Muſcles and Pleura, with an Inciſion Knife, carefully avoiding the Lungs, which are very apt to adhere to the Pleura, in this Part: when the Perforation is properly made, it is to be kept open in the Manner we have already ſhewn, and the Wound above is to be healed as ſoon as poſſible.

What is to be done when the Lungs adhere.

XI. As the Lungs frequently adhere to the Pleura, the Perforation of the *Thorax* requires great Circumſpection in the Surgeon: the Pleura ſhould be divided with all poſſible Tenderneſs; and when that is done, the Surgeon ſhould examine whether the Adheſion of the Lungs may not ſafely be removed with his Fingers or the Probe: when the Adheſion is very firm, the Pains we have taken to perforate the *Thorax*, and to diſcharge the extravaſated Blood, all prove fruitleſs: we muſt in this Caſe make an Inciſion in another Part, either before, or on one Side; and proceed as above.

How Wounds of this kind are to be treated.

XII. The Cavity of the *Thorax* being thus cleanſed, the Wound is to be dreſſed but once every Day: each Dreſſing ſhould be performed with all poſſible Expedition, and the utmoſt Diligence ſhould be uſed to guard the Contents of the *Thorax* from the external Air: at the Time of dreſſing, a Chafing-diſh of hot Coals ſhould be held near the Wound to warm and thin the Air; and if too great a Quantity of Air is already got into the Cavity of the *Thorax*, it muſt be drawn out with a Syphon. This being rightly performed, the Wound is to be dreſſed up with the utmoſt Expedition.

If any of the Contents of the *Thorax* are wounded.

XIII. When any of the Contents of the *Thorax* are wounded, as the Heart, the Aorta, the Vena cava, the Pulmonary Artery or Vein, the Œſophagus, Thoracic Duct, the Mediaſtinum, or a large Portion of the Lungs (eſpecially if it is a Gun-ſhot Wound,) Death comes too ſuddenly to give the Surgeon Room to exerciſe his Art. On the other Hand, when the Lungs are only ſlightly wounded, that is, when only the ſmaller Ramifications of the Pulmonary Vein or Artery, or the aſpera Arteria, are divided, the Caſe is very dangerous,

but

but not always mortal: though Perſons who recover after Wounds of this Kind are more obliged to the Soundneſs of their own Conſtitutions, than to their Surgeon's Skill.

XIV. We may reaſonably apprehend that the Lungs are wounded, when the Patient voids a great Quantity of frothy Blood by the Mouth, accompanied with a Cough, eſpecially when at the ſame Time the Blood which is voided at the Wound is very florid, and the Patient makes a particular Noiſe when he draws his Breath: the Office of the Surgeon here ſeems to be, to clear the internal Part of the *Thorax* from the extravaſated Blood, and to heal the external Wound, the Methods of doing which, we have already explained: no Application can be made to the internal Wound; that muſt be left to Nature. Whenever the divided Veſſels contract themſelves, and the Blood ſtops of itſelf, the Patient will recover; though Perſons who have recovered from theſe Wounds are remarkably ſubject to Ulcers of the Lungs and Conſumptions. Whenever any of the larger Pulmonary Veſſels are divided, the Violence of the Hæmorrhage either brings preſent Death with it; or, if it ceaſes a little, it returns again, and comes to the ſame End by ſlower Paces: to prevent this as much as poſſible, it will be proper to keep the Patient quite ſtill for ſeveral Days; he ſhould ſcarce ſpeak, he ſhould take cooling and agglutinating Medicines, and avoid all ſharp Things, all that heat the Blood, or provoke Coughing; and, if his Strength will permit it, he ſhould loſe Blood by the Arm. Signs of a Wound in the Lungs.

XV. Sometimes the wounded Part of the Lungs puſhes forward, and ſticks pretty firmly in the Orifice of the external Wound, as FONTANUS, TULPIUS, and RUYSCH have obſerved in their Writings: in this Caſe, if it is forced back again, it will diſcharge a great Quantity of Blood into the Cavity of the *Thorax:* therefore it is better to let it remain in the Situation you ſhall find it, for by this Means it will admit of the immediate Application of proper Dreſſings, and you may ſafely encourage it to adhere to the Lips of the external Wound: and here the Patient muſt be ſtrictly admoniſhed to keep as ſtill as poſſible: but if a [a] wounded Portion of the Lungs ſhould be puſhed out of the *Thorax* beyond the Limits of the external Wound, you ſhould wrap a Piece of fine Linen round this Part, and making a Ligature above the Linen, take off all that is below the Ligature with the Knife, and return the ſound Part of the Lungs into the Body, keeping one End of the Ligature conſtantly hanging out at the external Wound: when you have proceeded in this Manner, keep the Wound open with a Tent, till the Ligature can ſafely be drawn out: how the external Wounds ſhould be treated, we have ſufficiently explained already. Of the puſhing out of the Lungs.

XVI. As to the Medicines which are to be preſcribed for internal Uſe, they conſiſt chiefly, after the Hæmorrhage is over, of vulnerary Decoctions, giving at due Diſtances of Time a Doſe of *Balſamum Lucatelli, vel Meibomii*, obſerving particularly a ſtrict Regulation with regard to Diet: by following theſe Rules, a Surgeon may ſometimes ſave a Patient that has received a Wound of this Kind, at leaſt, where it was impoſſible to perform a Cure, he will have the Satisfaction of having done his Duty. What Internals are to be given.

[a] HILDANUS, *Cent.* II. *Obſ.* 3. relates a Caſe of this Kind, where a Portion of the Lungs forced its Way through a Wound of the *Thorax*; and Part of it appearing black and corrupted, he took it off with a red hot Knife, and then forced the ſound Part back again into the Body the Patient, he tells you, ſurvived this, and recovered a perfect State of Health.

EXPLANATION *of the* SIXTH PLATE.

Fig. 1. A braſs *Tournequet* after PETIT's Manner, but with ſome Alterations: The Uſe of this Inſtrument, and Method of applying it, will eaſily appear, if you compare it with what we have ſaid above in *Chap.* II. *Of Wounds.* § XV. and afterwards in the *Explanation of the fourth Plate*, *Fig.* 2, *and* 6.

Fig. 2. A Handle to fix Needles in when you are to make Sutures: this the *French* call *Portaiguille.*

Fig. 3. Another of the ſame Sort from GARENGEOT.

Fig. 4. PETIT's Handle for Needles.

Fig. 5. A Needle to perform *Gaſtroraphy.*

Fig. 6. Another of a larger Size.

Fig. 7. Another, which is new, to perform the ſame Operation.

Fig. 8. A Syringe for various Uſes, furniſhed with Pipes of different Sorts; by the Help of this you may not only inject Fluids into Wounds of the Abdomen and Thorax, into the Fauces, into Abceſſes, Ulcers, and into the Uterus; but you may alſo by the Aſſiſtance of this Inſtrument draw extravaſated Blood from the Cavity of the *Thorax*, in which Caſe the Syringe ſhould be twice as large; the Mouth of the Pipe A ſhould be triangular, and about two Thumbs Breadth.

Fig. 9. Another Pipe with a round Mouth, intended for the ſame Uſes.

Fig. 10. A ſmaller Pipe, which may be faſtened to the Syringe, *Fig.* 8. for various Uſes.

Fig. 11. Another ſomewhat curved, and perforated on both Sides: this will ſerve to ſuck Blood out of the Cavity of the Thorax, and to throw Injections into that Part, or into the Fauces.

Fig. 12. Another perforated at the End like a Cullender.

Fig. 13. Another like the former, but curved, to throw Injections into the Uterus, and for other Uſes.

Fig. 14. An Iron Inſtrument like an Ear-picker, for various Uſes.

CHAP. XI.

Of WOUNDS *of the* NECK.

Wounds of the Neck of bad Conſequence.


I. WOUNDS of the Neck are no leſs dangerous than thoſe of the Thorax or Abdomen; inſomuch, that I am ſurprized to find ſeveral chirurgical Writers treat of Wounds of this Claſs ſlightly, as if they were ſcarce worthy of their Notice: and I have often wondered, and complained of it in my Anatomy, (Sect. 264.) that in the Diviſion of the Trunk the Neck ſhould be omitted.


How many Kinds of Wounds of the Neck.

II. There are ſeveral Sorts of Wounds in the Neck: ſometimes the Seat of the Wound is only in the common Integuments, and the muſcular Parts; this is attended with very little Danger; but the moſt dangerous, and indeed generally incurable Wounds, are thoſe of the larger Blood-veſſels in theſe Parts; ſuch are thoſe of the Jugular Veins, carotid and vertebral Arteries; or where the Aſpera Arteria is wounded; or the Gula; the Medulla Spinalis; the Nerves that

Plate 6.

Plate 6.




A
C
Fig: 4.
D
B
Fig: 2.
D
A
B
Fig: 8.
A
Fig: 13.
Fig: 14.
Fig: 9.
Fig: 12.
Fig: 11.
Fig: 10.
Fig: 6.
A
Fig: 3.
Fig: 5.

that deſcend by the Neck (ſuch as the Par Vagum, the Intercoſtales, and the Diaphragmatici) or where ſeveral of theſe Parts are wounded at the ſame Time.

III. You will eaſily diſcover with your Eye, or by conſidering the Situation of the Wound, and the Symptoms that attend it, what Parts of the Neck are the Subjects of the Wound: after this Diagnoſtic, the Prognoſtic of thoſe Wounds will eaſily follow: for whoever is thoroughly acquainted with the Condition of a Wound, will find no great Difficulty in determining the Event of it: where the common Integuments and Muſcles alone are wounded, you will have no Reaſon to dread any very ill Conſequences: but where any of the other Parts of the Neck are Partakers of the Injury, you have Reaſon to apprehend the greateſt Danger; becauſe moſt of thoſe Parts are abſolutely neceſſary to Life itſelf: though in this Caſe, where the Wound is ſmall, there are ſome Hopes of a Cure. Diagnoſis and Prognoſis.

IV. Wounds of the Arteries in the Neck are ſcarce ever to be remedied, unleſs the Wound be very ſmall indeed; for in this Caſe the Patient uſually bleeds to Death before a Surgeon can be called to his Aſſiſtance: though, to ſay Truth, if a Surgeon were preſent at the Inſtant ſuch a Wound was inflicted, all his Art and Induſtry would have little or no Effect: for it is extremely difficult to ſtop Blood in this Part, not only from the Largeneſs of the Arteries here ſituated, and from their Vicinity to the Heart; but becauſe it is impoſſible in this Place to make a ſufficient Preſſure upon the wounded Veſſel. Yet the Wounds of the external carotid Arteries, and the Hæmorrhages they occaſion, are more eaſily managed; eſpecially if an experienced Surgeon be applied to in Time. What happens after Wounds of the Arteries in the Neck.

V. A Wound upon the external Jugular Vein is not attended with much Danger, if a Surgeon is called in Time: for only a ſmall Degree of Preſſure is required here, as appears by the frequent Practice of Blood-letting in this Part: but Wounds of the internal Jugulars are extremely dangerous; and this, partly from their Size, which is uſually larger than one of the Fingers; partly becauſe their Situation is ſo deep, that no proper Application can reach them to any Advantage; for theſe Reaſons many Surgeons have determined Wounds of this Kind to be mortal; but I can by no Means admit this as an abſolute Rule, without any Exceptions: on the contrary, I am of Opinion, that where the Wound in the internal Jugular is made by a ſharp Inſtrument, and but ſmall, if a Surgeon is ready at Hand, the Patient may be ſaved: how this is to be effected I ſhall teach below. After Wounds of the Veins.

VI. Wounds of the *Aſpera Arteria* were uſually deemed mortal by chirurgical [a] Writers: I am ſo far from contradicting them in this Sentence, that I ſhall rather endeavour to ſupport it, that is, where the Wind-pipe is entirely divided, or wounded in its lower Part within the Cavity of the [b] Thorax, or joined with a Wound of the carotid Arteries or internal Jugulars, which is frequently the Caſe: but, on the other Hand, if it is only wounded in the fore Part, and the neighbouring Veſſels remain unhurt, it is undoubtedly curable; which Opinion is ſtrongly ſupported by Variety of Examples from my own Experience, and that of other Practitioners: ſee above *Ch.* I. N. XIX. Wounds of the Aſpera Arteria.

VII. There are very little Hopes of Recovery where the Gula is much wounded, or entirely divided; becauſe not only the Office of Deglutition is cut Wounds of the Oeſophagus.

[a] See *de* BOHNIUS, *Vuln. Lethal.* Cap. ii. p. 23.
[b] Ibid. Sect. II. Cap. iii. pag. 121.

off

off, but the Part is ſo ſituated, that it is almoſt impoſſible to wound it without injuring at the ſame Time ſome of the neighbouring Nerves and Blood-veſſels: but, when the Gula is the only Subject of the Wound, and the Opening is very ſmall, the Wound may ſometimes admit of a Cure.

Wounds in the Medulla Spinalis and Nerves.

VIII. Wounds on any Part of the *Medulla Spinalis* are very dangerous, but more particularly ſo when inflicted upon that Part of it which paſſes through the Neck: therefore it is no Wonder that ſcarce any one recovers after a conſiderable Wound of this Kind. The Reaſon of this will immediately appear, when we conſider, that ſeveral Nerves proceed from this Part, which are abſolutely neceſſary to conduct the Œconomy of the Animal; that the vertebral Veins and Arteries will almoſt always be wounded at the ſame Time; and that the Situation of theſe Parts is ſuch, that it is impoſſible to convey the proper Remedies to them: nor are Wounds of the large Nerves of the Neck, ſuch as we mentioned at *N.* II. leſs dangerous than theſe: for if they are divided, the nobler Parts of the Thorax or Abdomen, to which Nature hath determined them, will immediately loſe their Aſſiſtance, and of Conſequence become unequal to the Offices for which they were intended.

How ſlight Wounds of the Neck are to be treated.

IX. The Treatment of Wounds in the Neck is different, according to the different Nature of the Wound: when the common Integuments and muſcular Parts are the ſole Subjects of the Wound, it will require the ſame Method of Treatment which we have adviſed above for all ſlight Wounds, upon what Part ſoever they may be inflicted: where the external Jugular is wounded, the ſame Methods which we uſe after Bleeding in that Vein will be ſufficient.

Cure of a Wound in the Internal Jugular.

X. When the internal Jugular Vein has received a ſmall Wound, the Hæmorrhage will eaſily be ſtopped by filling the Wound well with dry Lint, or ſteeped in *Alcohol. Vini*, or *Spirit. Terebinth.* or any proper ſtyptic Medicine; or with the Fungus called *Crepitus Lupi*; laying over theſe Applications ſquare Bolſters, and ſecuring all with a Bandage, drawn as tight as the Situation of the Part will admit. An Hæmorrhage is much eaſier ſuppreſſed in a Vein than in an Artery. The whole of the Cure depends upon the Degree of Preſſure that you can make upon the wounded Veſſels: ſometimes it happens that the Method of dreſſing which we have juſt adviſed in this Caſe will have no Effect; when this ſhall happen, the Surgeon or his Aſſiſtant muſt keep his Finger conſtantly upon the Wound, or make a Preſſure upon the Part, with ſuch an Inſtrument as we deſcribed in *Plate* V, *Fig.* 2. till the Hæmorrhage is entirely ſtopped: this Preſſure ſhould uſually be continued for a Day or two. The ſame Proceſs ſhould alſo be obſerved in Wounds of the vertebral Veins and Arteries. After the Blood is ſtopped, the Dreſſings ſhould continue upon the Part untouched till the third Day, and then a vulnerary Balſam and Plaſter may be applied to heal the Wound.

How a large Wound of the Internal Jugular is to be treated.

XI. When the internal Jugular Vein has received a large Wound, or is intirely divided, the Patient will preſently die with the Loſs of Blood: but, if a Surgeon ſhould be preſent when ſuch a Wound is received, or ſhould come in inſtantly afterwards, I would adviſe him to make a Preſſure upon the divided Vein with his Finger, and to enlarge the Wound upwards and lengthways, till he can come at enough of the Veſſel to make a ſtrong Ligature upon it by the Aſſiſtance of a crooked Needle, ſuch as I have deſcribed, *Plate* VIII, *Fig.* 4; and then he may fill up the Wound, and treat it as at N. X. By this Means

the

the Life of the Patient may be ſaved, though the Courſe of the Blood through this Veſſel be entirely cut off: I have often tried this Experiment upon a Dog, and he has recovered, and never ſuffered any apparent Inconvenience from it: therefore I think it better to put this doubtful Remedy in Execution, than to leave the Caſe as deſperate.

How a Wound in the Carotid Artery is to be treated.

XII. A Wound in the Carotid Artery is attended with greater Danger than a Wound in the internal Jugular: but, if a Surgeon is preſent when the Wound is received, I think he ſhould make the ſame Attempts to cure it: this is more likely to meet with Succeſs in Wounds of the upper and middle Part of it, than in Wounds of the lower Part: but where the Wound is not in the Trunk of the Artery, but in one of its Branches near the Head, you ſhould fill up the Wound with Lint, dipped in ſome ſtyptic Liquor, if you have it ready: then cover it up with thick Compreſſes, ſecuring all with a tight Bandage, and ordering an Aſſiſtant to make a Preſſure upon the Part for ſome Time with his Hand. *See Part* III. *Ch.* II. *N.* VIII. *and Plate* 37. *Fig.* 8. By theſe Methods I have very ſucceſsfully ſtopped violent Hæmorrhages, that have proceeded from wounded Branches of the Carotid Artery, which I have divided in taking out large ſchirrous, parotid, or ſubmaxillary Glands: in theſe Caſes you ſhould never remove the Dreſſings till the third or fourth Day: nor ſhould the Lint, applied at the firſt Dreſſing be forced out; but remain in the Wound, till it works its own Way; otherwiſe a freſh Hæmorrhage, and that very violent, moſt commonly enſues (I ſpeak from Experience) by which the Patient's Life may be greatly endangered.

How to treat Wounds of the Aſpera Arteria.

XIII. In curing Wounds of the *Aſpera Arteria*, the Surgeon ought, after cleanſing the Wound, to endeavour to unite the divided Parts by the Aſſiſtance of ſticking Plaſters; or, where the Wound is large, by making two Stitches with a crooked Needle, dreſſing them up afterwards with ſome vulnerary Balſam, a ſticking Plaſter, and proper Compreſſes, adviſing the Patient to keep his Head in a prone Situation [a]: the Wound thus treated will eaſily heal, if it is made either by Puncture or by a cutting Inſtrument: but if any Part of the the *Aſpera Arteria* is carried away by a Bullet, the Suture is to no Purpoſe: Wounds of this Kind are more readily healed and filled up by the Uſe of a digeſtive Ointment, or vulnerary Balſam; but this muſt be particularly remembered, that the Head be kept in a prone Situation. If the *Aſpera Arteria* is entirely divided, and the lower Part of it contracts itſelf into the Cavity of the Thorax, ſo that it cannot be laid hold on, and united to the upper Part, the Patient muſt undoubtedly die: if the Artery is not entirely divided, the Surgeon muſt raiſe the lower Part, and unite it to the upper Suture [b].

How to treat Wounds of the Oeſophagus.

XIV. Where the *Œſophagus* is wounded, whatever the Patient attempts to eat or drink paſſes through the Wound, and he is uſually attended with Hiccoughs and Vomiting: where the *Œſophagus* is entirely divided, there is no Poſſibility of curing it; but, where it is only perforated or wounded in Part, you may attempt the Cure by dreſſing the Wound with a vulnerary Balſam, by endeavouring to unite it with ſticking Plaſters, and by adviſing the Patient

[a] Cures of this Kind are to be met with in BARTHOLIN, *in Hiſt. Anatomic. Cent.* V. *Hiſt.* 89. and in TULPIUS, *Obſ.* Lib. i. Cap. 50. and in other Writers; many of whom GARENGEOT has quoted, Tom. ii. C. de Bronchotom.

[b] A remarkable Inſtance of this is relatd by GARENGEOT, *Op. Chir.* Tom. ii. C. de Bronchotom.

to a ſtrict Abſtinence for ſome Days, or at leaſt to take Nouriſhment by the Mouth very ſparingly, at the ſame Time preſcribing nouriſhing Clyſters of Broths or Milk: but when the Neceſſities of Nature require Nouriſhment to be taken by the Mouth, the Wound ſhould conſtantly be diligently cleaned afterwards, leſt any Part of what was taken ſhould ſtick by the Way and putrify, which would bring on very bad Symptoms [c]. After the Wound is cleaned in this Manner, it is to be dreſſed daily with ſome vulnerary Balſam till it heals: but, if the *Œſophagus* be wounded within the Thorax, the Situation is ſuch that Art cannot reach it; the Cure muſt be left entirely to Nature.

How Wounds of the Medulla Spinalis are to be treated.

XV. Wounds of the *Medulla Spinalis* are beſt dreſſed with the *Balſamum Peruvianum, Eſſentia Myrrhæ aut Succini, Spiritus Maſtichis*, or with Medicines of the like Nature, mixed with a ſmall Quantity of *Mel Roſarum* ſpread upon Pledgits, and applied moderately warm; and then covered with a vulnerary Plaſter: the Event muſt be left to God's Providence, and the Strength of the Patient's Conſtitution: ſlight Wounds of theſe Parts ſometimes heal by this Method; but large Wounds here bring certain Death.

How to treat wounded Nerves in the Neck.

XVI. Wounds inflicted upon the large Nerves, which are ſituated in the Neck, are generally mortal; but, where the Wound is very ſmall, the ſame Methods may be attempted which we adviſed in the Wounds of the *Medulla Spinalis*.

CHAP. XII.

Of WOUNDS *of the* HEAD *in general.*

Wounds of the Head very dangerous.

I. NO Wounds are attended with more Danger than thoſe which are inflicted upon the Head; for the ſlighteſt Injury of the Brain will frequently bring on the worſt of Symptoms, and even Death itſelf: nay, Wounds of the Head which do not penetrate into the Cranium, and proceed only from a ſlight Fall or Stroke, even with a blunt Inſtrument, ſometimes occaſion a Rupture of ſome of the internal Blood-veſſels, and an Extravaſation of Blood in the Brain, which is attended with the moſt miſchievous Conſequences: therefore even the ſlighteſt Wounds of the Head require all the Care and Caution that we are Maſters of.

Wounds of the Head are of two kinds.

II. We ought carefully to diſtinguiſh, 1. what Parts of the Head are wounded; and, 2. in what Manner the Wound was made; for ſome Wounds of the Head are made with ſharp Inſtruments, either by ſtabbing or cutting; ſome are made with blunt Inſtruments, which is the Caſe in ſome Blows or Falls, and in Gun-ſhot Wounds: theſe of the laſt Claſs are attended with much greater Danger than thoſe of the former; for they generally give ſuch a violent Shock, as to burſt the finer Veſſels and Nerves of the Brain.

What Parts are wounded.

III. As to the Parts which are wounded, they are either the common Integuments alone, or with theſe the Fleſh of the Face, or the Pericranium, or the Temporal Muſcles, or the Cranium; or ſometimes the internal Parts alſo; next,

[c] The abovementioned Author, in Caſes where the Patient could not ſwallow, recommends the Uſe of nouriſhing Clyſters.

the Dura Mater, Pia Mater, and the Brain, either in its cortical or medullary Part, or in its Ventricles: when the Cranium is wounded, as firſt the inner Lamina of the Cranium, from whence Fragments are often ſplintered, and driven into the Dura Mater or the Brain itſelf; it is either cut, broken, or contuſed: it may not be amiſs to divide Wounds of the Head into two Claſſes; 1. thoſe that affect the Face: 2. thoſe that hurt or wound the Cranium, the Caſtle of the Brain, or ſome of its Integuments.

CHAP. XIII.

Of WOUNDS *of the* FACE.

I. SINCE the Face was intended for Beauty as well as for particular Uſes, two Things are to be remarked; to wit, that we do not leave worſe Injuries upon the Face, and particularly the Eyes, than we were employed to cure; and that we make an even fair Cicatrix: as the Face conſiſts of various Parts, each of which requires a diſtinct Method of Treatment, it will be neceſſary to treat of each of them ſeparately. Of Wounds of the Face in general.

II. In almoſt all Wounds of the Forehead that do not penetrate the Seull, this is principally to be obſerved; that after the Wound is cleaned from grumous Blood, and any foreign Bodies that may have got into it, it ſhould be anointed with ſome vulnerary Balſam, ſuch as the *Balſamum Peruvianum*, *Copaivæ*, or any other of that Kind; the Lips of the Wound are then to be kept together with narrow Slips of ſticking Plaſter, and over this a vulnerary Plaſter is to be laid: where the Wound is large, theſe Plaſters will not be ſufficient to form an even Cicatrix: therefore, to forward this End, it will be proper to ſprinkle the Wound with *Pulvis Sarcocollæ*, *vel Pulvis ex Radice Symphyti*, *Gummi Tragacanth. ac Gummi Arabico præparatus:* you may then apply your Plaſter, Compreſſes, and proper Bandages: the bloody Suture is never to be uſed either in theſe or any other Wounds of the Face, where it can be avoided; for the Stitches increaſe the Number of Scars. If a Wound of the Forehead is made in a ſtrait Line, the *uniting* or *incarning Bandage*, deſcribed in *Plate* II, *Lett. f*, will be of great Service in forming a fine Cicatrix; it is to be applied to the Forehead after the ſame Manner which we adviſed it to be applied to the Abdomen in longitudinal Wounds of that Part. *See Chap.* V, *N.* X. But if the Forehead is wounded tranſverſely, and the Fibres of the frontal Muſcle are divided, it occaſions a great Deformity; for the Power of lifting up the Eye-brows, and of contracting the Skin of the Forehead ceaſes: in this Caſe, after cleaning the Wound, it is beſt to unite it with a Stitch or two, dreſſing it with a vulnerary Balſam or Powder, and laying on ſticking Plaſters, ſecuring all with a proper Bandage, and adviſing the Patient to keep himſelf ſtill. It ſometimes happens in young healthy Perſons, that the divided Fibres of the Muſcles join and unite without any Suppuration, where this Method of Dreſſing is diligently followed: if any great Degree of Hæmorrhage ſhould enſue upon Wounds of this Part, the firſt Intention is to ſtop it with dry Lint, Compreſſes, and a tight Bandage; and at the next Dreſſing, after it has been well cleaned and waſhed with warm Wine, its Of Wounds of the Forehead.

Lips ſhould be brought together as before, with Slips of ſticking Plaſter; or in tranſverſe Wounds with a Stitch or two, if it be neceſſary.

Of Wounds of the Eye-brows.

III. Wounds of the Eye-brows require much the ſame Treatment with Wounds of the Forehead; only in Wounds of the Eye-brows more particular Care muſt be taken to guard againſt Inflammation, leſt the Eyes ſhould partake of the Injury: all ſharp Things ſhould be avoided both in eating and drinking; and if the Patient is of a plethorick Habit of Body, he ſhould loſe Blood in the Arm: the uſual Dreſſings ſhould be covered with Compreſſes, dipped in camphorated Spirit of Wine: if the Wound is large, and the Eye-brows entirely divided, it will be neceſſary to uſe the Suture, and to dreſs them up with a vulnerary Balſam and Plaſter, covering up both Eyes, and keeping them as much as poſſible from Motion: by neglecting this Method, the Situation of the Eyes in this Caſe will have a very frightful Effect; and ſometimes the Patient is deprived of his Sight.

Of Wounds of the Eye-lids.

IV. Wounds of the upper or lower Eye-lid will not readily heal; not ſo much from the Thinneſs of the Parts of which they are compoſed, as from the Quantity of Fluids with which the Eyes are continually moiſtened: at firſt therefore it will be beſt to foment the Eye *cum Decoctо quodam ex Chamomillâ, Hyſſopo, vel Euphraſiâ confectо*, till the Flux of Blood is ſtopped, and the Wound well cleaned: when the Wound is tranſverſe, you may ſtitch it up in the Middle with a fine Needle, ſprinkling it afterwards with the Powder deſcribed at *N.* II. or anointing it *cum Balſamo Copaivæ, de Mecchâ*, or with any other of the ſame Kind, or with *Oleum Ovorum*, laying over it the *Emplaſtrum Diapalmæ*, and tying it up ſo that the Eyes may have very little Power to move: where the Wound is lengthways, you muſt make ſeveral Stitches, and dreſs it up as before.

Of Wounds of the Eyes.

V. Wounds of the Eye are attended with more Danger than any other incident to the Face; not only as the Patient is thereby often deprived of that moſt precious Bleſſing, the Bleſſing of Sight (eſpecially if the *Tunica Cornea* or *Uvea* are wounded, either by themſelves, or conjointly together with the neighbouring Parts,) but as Death itſelf is ſometimes the Conſequence, if the wounding Inſtrument ſhould pierce the Bones of the Orbit, ſo as to injure the Brain or its Nerves. If the Eye is wounded, but not ſo as to let out the vitreous or chryſtalline Humour, the following Method will be of great Service; the Wound ſhould be anointed two or three Times in a Day with a Feather or fine Rag, well dipped in *Unguentum Alabaſtrinum, aut Albumen Ovi, aut Mucilag. Sem. Cydon. & Pſyllii Aq. Roſar. parat.* and afterwards a ſmall Compreſs is to be laid on, being well ſaturated with the following Collyrium. ℞ *Albumin. Ovor. N.* 2. *Aq. Roſar.* ℥iiſs. *Ol. Roſar.* ℥ß. *Camphor. Gr.* iii. *probè conquaſſando.* NUCK gives us a Caſe, where a Man was ſo wounded in the Eye, that Part of the vitreous Humour fell out, nevertheleſs he cured him without leaving any Diſorder in his Sight; his Method of Cure was as follows; he divided the Part of the vitreous Humour that hung out of the Eye from the reſt, and then diligently fomented the Eye with a Collyrium, prepared *ex Albumine, Aquâ Roſarum, Bolo Armenâ & Camphorâ probè conquaſſatis. Gumm. Arabic.* ℈i. *in Aquæ Roſar.* ℥i. *ſolut.* is very ſerviceable in this Caſe; but if it is attended with any great Degree of Inflammation, which is frequently the Caſe, it will be proper to cover the ſmall Compreſs with a larger, dipped in *Spiritu Vini camphorati calido*: the Bowels alſo ſhould be kept looſe for ſome Days with opening and

cooling

cooling Medicines; if there is a plethoric Habit, Blood ſhould be drawn from the Neck or Feet; all warm or ſharp Things ſhould be thrown out of the Patient's Diet, and great Care taken to keep him quiet; by obſerving theſe Regulations, not only the Eye, but the Sight of it alſo may be preſerved: when the chryſtalline Humour, or any Part of it ſticks in the Orifice of the Wound, it ſhould be pulled out, that it may not bring on Deformity, or worſe Miſchief upon the Eye.

Where the Humours are fallen out.

VI. When the vitreous and chryſtalline Humours are fallen out of the Eye, not only the Sight, but Figure of the Eye muſt be entirely deſtroyed: therefore at firſt it ſhould be dreſſed with Compreſſes dipped in warm Wine, or Spirit of Wine, and afterwards with ſome vulnerary Balſam: the Deformity, which the Loſs of Subſtance in the Eye will occaſion, may be avoided by the Help of an artificial Glaſs or Silver Eye. *See Plate* VII, *Fig.* 1: but we ſhall treat more largely of this [a] in another Place.

Sight ſometimes reſtored.

VII. It ſometimes happens when only the *Tunica Albuginea* and *Sclerotica* are ſlightly wounded, the *Cornea* and *Uvea* remaining unhurt, that the Eye recovers itſelf; and though both the vitreous and chryſtalline Humours fall out by the Wound [b], yet they are renewed again by the Benefit of Nature, and the Office of Sight performed as well as before the Injury happened: Dr. SEEGER ſome Time ſince was ſo kind as to communicate the Hiſtory of a Caſe of this Sort to me, whence it appeared that he had reſtored Sight to a Woman after ſhe had loſt the Humours of her Eye: when we have duly conſidered this, we ſhall not altogether reject the Teſtimonies of BURRHUS and KERKRINGIUS, when they affirm to us, that they have acquired the Art of reſtoring the Sight after the Humours are entirely fallen out of the Eye: we may now alſo credit thoſe who [c] affirm that the Sight may be enjoyed without the Aſſiſtance of the chryſtalline Humour, notwithſtanding [d] ſome have ſtrenuouſly maintained the contrary.

Wounds of the Noſe.

VIII. Wounds of the Noſe are generally cured by the dry Suture; but where the Wound divides the Cartilage, and penetrates ſo deep that the Lips of it cannot be kept in Contact by the Application of ſticking Plaſters, the true Suture muſt be made through the Skin on each Side of the Wound: though it ſounds very unlike Truth that any Part of the Noſe ſhould be entirely ſeparated from the reſt, and afterwards united to it again by the Aſſiſtance of Sutures; yet BLEGNY affirms, that this has happened: *See Zodiac. Med. Gall. Edit.* 1680, *pag.* 75. When the naſal Bones are fractured, it is uſual to place ſmall Tubes made of Lead or Silver under them for ſome Time, leſt the Paſſage of the Noſe ſhould be ſtopped by the ſhooting out of the new Fleſh; you will ſee theſe Tubes deſcribed at *Plate* II, P, Q, R. Externally you may uſe ſome Balſam, or *Eſſentia Maſtichis, Succini, vel Myrrhæ*, or ſome glutinous Powder, ſuch as you have ſeen directed at *N.* II: the Lips of the Wound ſhould be kept

[a] Tract. *De Duct. Oculor. Aquoſ.* pag. 126, 127—132.

[b] CREDAT JUDÆUS.

[c] You may find many Inſtances related of Perſons who have enjoyed their Sight after the Loſs of the chryſtalline Humour, in SKENKII *Obſ. Med.* HILDANI *Obſ.* 26. *Cent.* I. *Act. Med. Hafn. Vol.* I. *Obſ.* 69.

[d] See my Treatiſes on the *Cataract. Glaucoma, &c.*

in Contact with each other by the Help of ſticking Plaſters, and of a four-headed Bandage; the Application of which will be explained when we come to treat profeſſedly of Bandages.

Wounds of the Lips.

IX. Wounds of the Lips are made either with ſharp or blunt Inſtruments, or with Bullets: Wounds of the firſt Sort, whether they are made lengthways or tranſverſe, are generally to be cured by the dry Suture; the Patient in this Caſe muſt diligently avoid both Chewing and Talking, his Diet therefore muſt be entirely Spoon-meat; if the Wound is very large, it will require the bloody or true Suture: in Wounds of theſe Parts which are made by blunt Inſtruments, by Falls, or by Bullets, the ſhattered Parts ſhould be brought to Digeſtion, and the Lips of the Wound, after being cleaned, be brought together, either with ſticking Plaſters, or by the Suture, which is uſed for the Hare Lip, which we ſhall deſcribe below.

Wounds of the Cheeks.

X. Wounds of the Cheeks ſhould be treated after the ſame Manner, and with the ſame Circumſpection, which we adviſed for Wounds of the Lips; but if one of STENO's *Salivary Ducts* is wounded in its Paſſage croſs the Cheek from the *Parotid* Gland, the conſtant Diſcharge of Saliva into the Wound will prevent the Cure, till the Duct is perforated in the internal Part of the Cheek, to make a Paſſage for the Saliva into the Mouth: this Method of Cure is propoſed by CHESELDEN, in his *Anatomy*.

Wounds of the Ear.

XI. Wounds of the external Ear are eaſily united by ſticking Plaſters, unleſs the Cartilage is entirely divided, and then it will require the Help of the Needle, and the Application of vulnerary Balſams, with proper Compreſſes and Bandages: when the Ear is wounded in the Neighbourhood of the *Meatus Auditorius*, Care muſt be taken to prevent the Diſcharge of Blood and Matter into that Paſſage, which would do great Miſchief to the *Tympanum*; this may be done by filling the internal Ear with Lint or Cotton.

Wounds of the Tongue.

XII. The Tongue is ſo well guarded by the Jaw-bones and the Teeth, that it is very rarely the Subject of a Cut or Stab, but it is frequently bit in Fits of the Epilepſy, in violent Falls, and it is ſometimes wounded by a Bullet: if the Wound of the Tongue is not very large, it will eaſily heal by the Application of *Ol. Amygdal. dulc. cum Sacch. cand. q. ſ. admiſt. aut Mel Roſar. cum Ol. Myrrhæ per Deliquium.*

How to cure large Wounds of the Tongue.

XIII. Large Wounds of the Tongue will not unite without the Aſſiſtance of the Suture: it is no Wonder therefore that Wounds near the Root of the Tongue always leave a Fiſſure in the Part, ſince their Situation prohibits the Uſe of the Needle: to prevent Loſs of Speech enſuing upon large Wounds of the Fore-part of the Tongue, the divided Parts ſhould be brought together with the Needle, as ſoon and as neatly as poſſible, and afterwards anointed with the Medicines which we preſcribed in the laſt Article, ſince ſticking Plaſters will not take Place here: PURMAN affirms, that he made uſe of ſilver Threads in Sutures upon this Part to great Advantage: *See his Surgery, P.* I. *Chap.* VI. Gun-ſhot Wounds upon the Tongue are to be dreſſed with the Medicines which we recommended above at *N.* XII; for Sutures are of no Service in this Caſe: the Patient ſhould keep from ſpeaking, and live upon Spoon-meats during the Cure, but more particularly when the Wound is juſt beginning to unite.

Wounds of the Palate.

XIV. Wounds of the Palate will heal beſt if you anoint them with *Mel Roſarum* alone, or with the Addition of a ſmall Quantity of *Balſamum Peruvianum*,

or ſometimes *Oleum Myrrhæ per Deliquium:* theſe Remedies alſo have great Efficacy in curing all other Wounds of the Mouth.

CHAP. XIV.

Of the principal WOUNDS *of the* HEAD.

I. WE obſerved above, that Wounds of the *Cranium*, the Seat or Caſtle of the Brain, were to be reckoned under the ſecond Claſs of Wounds in the Head: theſe, by Way of Eminence, are alone called *Wounds of the Head:* they are divided into ſeveral Diſtinctions, according to the different Parts that are wounded, and the different Species of Wounds: theſe we ſhall treat of in the Order we enumerated them in *Chap.* XII, at *N.* III. We ſhall begin with the ſlighteſt, which are thoſe Wounds that are inflicted upon the external Coverings of the *Cranium.* Intent of this Chapter.

II. There are ſeveral Ways of diſcovering that the Wound is terminated in the external Parts of the *Cranium*: 1. by the Eye: 2. by the Probe, which ſhould be uſed very gently here, for Fear of bringing on further Miſchief: 3. by examining the Inſtrument with which the Blow was given, and by conſidering the Degree of Force with which it was impelled: and, 4. laſtly, by the Abſence of violent Symptoms; for a violent Blow upon the Head will always be attended with Vomitings, Vertigo, Blood will be diſcharged by the Noſe, Ears, and Mouth; and the wounded Perſon will loſe his Speech and Senſes: theſe Diſorders will appear ſometimes ſooner, ſometimes later; but always more violent, when the Wound is made by a Fall, or by ſome blunt Inſtrument, in which Caſe the *Cranium* is uſually much ſhattered: the Blood which diſcharges itſelf by the Wound, when it is made with a ſharp Inſtrument, will inſinuate itſelf between the common Integuments and the *Cranium*; in Contuſions that are made with blunt Inſtruments, ſometimes it will lie concealed under the *Cranium*, and by corrupting the *Perioſteum* and *Cranium* will bring on Ulcers and Caries of the Bone; frequently it will occaſion Fever, Convulſions, and Death: but here it muſt be obſerved, that the Symptoms are far from being certain Indications: for ſome, on receiving a Blow, drop down inſtantly, loſe their Speech and their Senſes; yet recover ſurpriſingly; others, at firſt, are ſlightly affected, and afterwards die. Wounds of the external Parts.

III. When the Temporal Muſcles are wounded at the ſame Time, the Patient will be attended with grievous Diſorders; but more eſpecially when this happens by a Blow or a Fall, or by a Bullet; not only becauſe theſe Muſcles are neceſſary for the Offices of dividing the Food, and for forming the Speech; but becauſe they are furniſhed with conſiderable Nerves, Tendons, and Arteries, which will partake of the Injury; and laſtly, becauſe the *Cranium* is thinneſt in this Part. Wounds of the temporal Muſcles.

IV. Wounds that are made on the external Parts of the Head by ſharp Inſtruments, and not attended with any violent Symptoms, are eaſily cured by the ſame Methods which we have before preſcribed for other Wounds, *Chap.* XIII, *N.* II, only: in order to make the proper Applications, it will be neceſſary in the Cure of Wounds on the external Parts of the Head.

the firſt Place to ſhave the Part with a Razor: there will be no Occaſion eve to make Sutures upon theſe Parts, ſince ſticking Plaſters will always anſwe your End: if the Wound be made lengthways on the Integuments of the *Cranium*, after cleaning it well, let the Lips be brought together, and bound with the uniting Bandage: *Chap.* XIII, *N.* II: if the Wound be tranſverſe, ſome Surgeons adviſe the bloody Suture without Exception; but I ſhould rather chuſe, if poſſible, to cloſe it with Plaſters and Bandages, and heal it like other Wounds: unleſs there ſhould be any looſe Pieces of Fleſh or Skin, or the Aperture be ſo wide as not to be united by Plaſters; for in theſe Caſes the true Suture muſt be uſed: you ſhould always endeavour to be as expeditious as poſſible in finiſhing each Dreſſing; the Medicines are to be applied warm, and the Air kept in a moderate Heat with hot Coals: if there ſhould be any great Degree of Hæmorrhage, which will frequently happen from the Number of Veſſels that are liable to be wounded in this Part, it muſt be ſtopped with dry Lint, or, where that is unequal to the Taſk, with the *Alcohol Vini*, *vel Lapi Crepitu*, *vel Pulvere quodam aſtringente*; theſe Applications ſhould be ſecured with a tight Bandage: after the Hæmorrhage is ſtopped, you may dreſs with *Mel Roſarum*, or ſome digeſtive Medicines, till the Wound is well deterged; and then with a vulnerary Balſam, or dry Lint, till it is healed: if the Hæmorrhage be exceeding violent, the Artery muſt be tied up with a Thread: on the other Hand, eſpecially in plethoric Conſtitutions, we ſhould not be too haſty in ſtopping the Blood; for the Diſcharge in this Caſe proves oft-times beneficial, and prevents many bad Symptoms that might otherwiſe enſue.

The Uſe of medicated Bags.

V. It has been frequently the Practice among Phyſicians to order[a] medicated Bags to be applied to the Head, when it has been conſiderably wounded, to prevent or aſſuage the Violence of the Symptoms, ſuch as Tumors, Inflammations, and Pain; theſe Bags are ſtuffed with *Betonicâ*, *Salviâ*, *Majoranâ*, *Serpillo*, *Origano*, *Roriſmarino*, *Floribus Lavendulæ*, *Salviæ*, *Roſarum*, *& ſimilibus*; theſe they boil in Wine, and after having gently preſſed them, they apply them, as warm as the Patient can bear them, to the wounded Part: where the Symptoms are already urgent, they make two Bags, and apply them alternately; by theſe Means the inſpiſſated ſtagnating Blood is rendered fluid, and the Miſchief is frequently removed without having Recourſe to the Trepan: when the Symptoms are too violent to be removed by theſe Applications, we are forced to uſe other Methods, according to the Nature of the Diſorder: of theſe we ſhall treat in the ſubſequent Articles.

How Contuſions are to be treated.

VI. In violent Contuſions of the Head, which will be diſcovered by the Tumor and Softneſs of the Part, by the Separation of the Integuments from the *Cranium*, and by the Collection of ſtagnating Blood which appears to be confined under the Skin; you ſhould endeavour to divide the confined Fluids by attenuating Medicines externally applied, or to diſcharge them by making an Opening with a Knife; or laſtly, to bring them to Suppuration; where the Extravaſation of Fluids is very conſiderable, it is beſt to diſcharge the greateſt Part of them inſtantly by Inciſions, and what remains will be eaſily diſperſed: the Application of the medicated Bags, deſcribed above, will anſwer the Intention of

[a] This Form is entirely laid aſide with us in *England*, and Fomentations made of the ſame Herbs ſubſtituted.

thinning and dividing the ſtagnated Blood; but you may add to the Ingredients, which we mentioned, *Herba Chamædrys*, *Scordium*, *Sabina*, *Abrotanum*, *Abſinthium* *Mentha*, *Ruta*, *Flor. Chamomil. Sambuc. Rad. Bryoniæ*, and Things of the like Intention: the Bags that are ſtuffed with theſe Ingredients may be quilted, that they may be divided into equal Parcels, and not run together in Lumps: where Wine cannot be had to boil them in, you may make uſe of Water, adding a Proportion of Spirits of Wine, or Malt, or Melaſſes, after it has done boiling, and a few Ounces of Soap: but particularly, a Vein ſhould be opened toward the Beginning of the Diſorder, and the Maſs of Blood thinned with proper Infuſions of diluting Herbs, after the Manner of Tea, with all other attenuating Medicines: we ſhall treat more largely upon what is farther to be done in this Caſe, in a following Chpter *upon Contuſions*.

How to bring the Parts to Suppuration.

VII. Where you find it impracticable to effect the Attenuation and Diviſion of the ſtagnating Fluids, it will be proper to attempt the Suppuration of them: in violent Contuſions it will be adviſeable to preſcribe the Application of ſuch Cataplaſms as are directed above at *Chap.* II. *N.* XIII. and below at *Chap.* XV. but in ſlighter Caſes, where there is an Opening, the *Unguentum digeſtivum cum Aloë et Spiritu Vini pauxillo admiſtum* will do the Buſineſs, covering the Part afterwards with a warm Plaſter, ſuch as the *Emplaſtrum de Meliloto*, *Malacticum*, *Diachylon ſimplex vel compoſitum*, *vel Empl. de Galbano*: after the Suppuration is formed, and the Matter diſcharged, the Wound will eaſily heal by the Application of a vulnerary Balſam: but in violent Contuſions, where there is no Opening, or a very ſmall one, by which the Matter cannot be diſcharged, you muſt enlarge the Wound with your Knife, to prevent the neighbouring Parts from being corroded: by this Means the Wound will eaſily be cleaned, and by obſerving the Directions we have frequently laid down above, the Cure will be ſpeedily performed; and you will eaſier diſcover whether the Scull be found or fractured.

Of Wounds in the Pericranium.

VIII. When the *Pericranium* is wounded, but not in ſo great a Degree as to lay the *Cranium* bare, treat the Wound in the Manner we deſcribed above at *N.* IV. *of this Chapter*; omitting the Uſe of the vulnerary Oils there preſcribed, becauſe they would injure the *Cranium*, and ſubſtituting in their Room ſome warm balſamic Medicines, ſuch as the *Balſamum Peruvianum*, *Copaivæ*, *Spir. Terebinth. Eſſentia Myrrhæ*, *Succini*, *Spir. Maſtichis*, and others of that Kind: but where the *Cranium* is left bare, and expoſed to the Air, its internal Lamella, being robbed of its Nouriſhment, by the Deſtruction of the Veſſels by which it was conſtantly ſupplied, will loſe its natural Colour, and become yellow, livid, black, and by degrees ſeparate from the neighbouring Parts, and exfoliate, as we term it, which will greatly protract the Cure of the Wound.

Method of treating the Pericranium and Cranium.

IX. To prevent the Corruption of the *Cranium*, and the Separation of its Lamina, and to expedite the Cure, the Surgeon ſhould immediately cover the denudated Part, by drawing over the Skin, if it has not been too long expoſed to the Air: he ſhould then dreſs it with proper Plaſters and Sutures; by which Means the Cure is commonly effected without Exfoliation: even where the Part has changed Colour, it is not always neceſſary to wait for a Separation of the Lamina; as many are of Opinion it is often ſufficient to apply dry Lint to the naked Bone, and dreſs the Wound with a Digeſtive; by which Method alone it generally heals. In order to haſten the Exfoliation of the *Cranium*, and

forward-

forward the Cure, the Surgeon ought to bore ſeveral [a] Holes through the denudated Part, as deep as the *Diploë*, with an Awl, or with Inſtruments like thoſe deſcribed at *Plate* VIII, *N.* II. *and Fig.* 7. *Lett.* A: this Operation does not only forward the Exfoliation of the Part, but make Way alſo for the ſprouting up of freſh Veſſels, and forming as it were a new *Pericranium*: the Dreſſing, which ought to be performed each Time with Expedition, and not repeated ſo often as in other Caſes, is to be applied in the following Manner: when the Wound is properly cleanſed, Pledgets, firſt dry, afterwards well-ſaturated with *Eſſentia Maſtichis*, *Succini*, or any other mild balſamic Medicine, with the Addition of a ſmall Quantity of *Mel Roſarum*, are to be laid upon the injured Part of the *Cranium*: over theſe you may clap the *Emplaſtrum de Betonica*, and over that the Bolſters and Bandage for the Head (Fr. *Couvre-chef*) deſcribed above at *Plate* III, *Fig.* 1. A: theſe Applications ſhould be continued till the *Cranium* appears to be ſound, and the Wound is in a Condition to heal. When the *Pericranium* is contuſed, but not ſeparated from the *Cranium*, you muſt endeavour to diſperſe the ſtagnated Fluids, by the Application of medicated Bags, deſcribed at *N.* V, VI: if theſe have not the deſired Effect, you may have Recourſe to Scarification, and warm Fomentations.

Of Wounds in the Temporal Muſcles.

X. If the Temporal Muſcles are wounded by a cutting Inſtrument; when the Wound is cleanſed, it muſt be treated in the common Method: ſhould the Artery ſuffer, the Hæmorrhage muſt be ſtopped either by Pledgits, Compreſſes and Bandages, or by a Ligament of Thread. If the Wound be made by Puncture or Contuſion, you muſt have Recourſe to the medicated Bags; and what extravaſated Blood lies beneath, ſhould be drawn off, by Inciſion: when we find that the *Cranium* is fractured under theſe Muſcles, and that there is concreted Blood under the Fracture, then an Inciſion may be made in the Muſcle lengthways, or obliquely, if it be judged neceſſary; that the Wound may be cleared of the ſtagnated Blood and the Fragments of the *Cranium* if there be any, in order to facilitate its Healing.

Of Injuries of the *Cranium*.

There are ſeveral Ways by which the *Cranium* may be hurt; by Falls, Blows Cuts, *&c.* which has occaſioned Authors to divide Injuries of this Part into ſeveral Diſtinctions; to wit, into Contuſions, Depreſſions, Fractures, Fiſſures, and [b] Contra-Fiſſures, that is, where the Fiſſure happens on the Side oppoſite to that which received the Blow.

Diagnoſtic Signs.

XI. There are ſeveral Circumſtances concerned in diſcovering an Injury of the *Cranium*: in the firſt Place, you muſt diligently inſpect the wounded Part, and make Enquiry with what Force the Blow was given that occaſioned it: after this you may ſearch the Wound with a Probe, but very circumſpectly, leſt by puſhing it raſhly forward you ſhould injure the Brain: ſome uſe a Pen in the room of a Probe, when they are ſearching for Fiſſures of the *Cranium*, and if the Pen is pointed at the End like a Tooth-pick, it will eaſily detect any Inequality or Roughneſs of the Bone: but you muſt be very careful not to ſuffer yourſelf to be deceived, as HIPPOCRATES was, by the Sutures: when Fiſſures of

[a] See HILDAN. *Cent.* iv. *Obſ.* 95. and RUYSCH. *Obſ.* 5.

[b] Many Writers have denied this Caſe to be poſſible; but not only HIPPOCRATES in his Book *De Vuln. Capit.* but CELSUS, *Lib.* viii. *C.* 4. and ÆGINETUS, *Lib.* vi. *C.* 90. have plainly deſcribed this Caſe; but amongſt the Moderns D. WAGNER, in a Treatiſe *De Contra-fiſſurâ* and LE-MAIRE, *De Reſonitu*, have put this Matter out of all Doubt.

the

the *Cranium* are ſo very fine, that they eſcape the Eye, and the Touch of the Probe, though the Violence of the Symptoms ſufficiently declare that the Patient has received an Injury of this Kind, it will be neceſſary to lay the Bone bare, and to drop Ink upon the Part of it which you ſuſpect, and wipe it off again immediately with Lint: and if any Part of it is fiſſured, you will find a black Stroke remain, notwithſtanding your Endeavour to wipe the Bone clean: if you are ſtill at a Loſs, put a Key into your Patient's Mouth, and bid him bite hard upon it: if this occaſions a Stridor of the Teeth, and Pain, Surgeons are apt to determine that there is a Fracture or Fiſſure in the *Cranium*: where the Bone has loſt its natural Colour, they will not allow it to be whole. The moſt certain Signs of a fractured *Cranium* are the violent Symptoms that immediately ſucceed the Injury; ſuch as vehement Pains, Vomitings, Vertigo, and Noiſe in the Ears; yet theſe are not always to be depended on: if Blood at the ſame time is diſcharged from the Noſe or Ears, the Senſes and Reaſon entirely loſt, and the Patient is continually ſleeping, the Matter is out of all Doubt. In a few Days after the Wound is received you will have a ſmall Diſcharge of thin fetid Matter: about the ſeventh Day the Integuments ſeparate from the Bone, and the *Cranium* itſelf is ſometimes ſo very foul, that it lets the Matter through to the Membranes of the Brain, which preſently partake of the Diſorder, and occaſion acute Pains, Spaſms, Drowſineſs, Loſs of Motion, or Rigor of the Limbs, Loſs of Speech, Apoplexy, and at length Death: all theſe Miſchiefs may ariſe from a very ſmall Fiſſure of the Scull, Examples of which you will find very frequent amongſt the Writers in Surgery.

XII. This ought to teach us to be very cautious in delivering our Opinions concerning the Event of Wounds in the Head; for we can never promiſe a Cure, though the Wound ſhould at firſt appear to be very ſlight: on the other hand, many who labour at firſt under violent Symptoms, by Bleeding and proper Remedies have been known to recover beyond Expectation: I ſhall here lay down ſome Obſervations which are well worthy of a Surgeon's Attention. It is very difficult to cure a Man who is poxed, or of a ſcorbutic Habit, at the Time he receives a Fiſſure in the *Cranium*: when the temporal Bone is the Subject of the Injury, the Cure is very doubtful; there remain very little Hopes of Recovery where the *Cranium* appears black; they alſo are in extreme Danger who have a black dry Tongue, full of Clefts, and beſet with Puſtules, or are attended with a Diarrhœa or Dyſentery, or where the Water is either quite clear and white, or as turbid as the Urine of Cattle. Prognoſtic.

XIII. The firſt Queſtion to be aſked, when you come to examine a Wound of the Head, is, whether it was made with a ſharp or a blunt Inſtrument? if the Wound was made with a ſharp Inſtrument, and penetrates into the *Cranium*, it muſt be filled at the firſt Dreſſing with dry Lint, in order to ſtop the Blood; but in the following Dreſſings, after the Matter is well wiped away, you may apply the *Eſſentia Succini. Maſtichis, Myrrhæve, cum admixto Roſarum Melle*: theſe Dreſſings are to be repeated as long as the Condition of the Wound ſhall require it: *See above, N.* IV. Where the *Cranium* is very much ſhattered by the Blow, and the Brain wounded, this Caſe is attended with very great Danger, but requires the ſame Method of Treatment with the former, only greater Diligence muſt be obſerved in cleanſing this Wound, and the more Expedition in applying the Dreſſings, to keep it from the Injuries of the Air: If the *Cranium* How Injuries of the *Cranium* are to be treated.

is ſo perforated by a cutting Wound, that it cannot well be cleanſed from the Blood or Splinters; or by a Puncture, that brings on any dangerous Symptoms, the Trepan muſt be applied: if a Piece, quite broke off from the *Cranium*, yet ſticks to the Integuments; that Piece, after cleanſing the Wound, ſhould be reſtored to its Place, the Skin ſtitched together, and the Wound properly dreſſed: this Method generally ſucceeds.

Wounds of the Head with a blunt Inſtrument.

XIV. When a blunt Inſtrument is the Occaſion of an Injury upon the *Cranium*, if the injured Part does not ſufficiently appear of itſelf, we ought to uſe great Induſtry to diſcover it.

How the Wound is to be examined.

XV. You will eaſily diſcover the injured Part, if you divide the common Integuments to the Bone, where they appear tumid and ſoft. In making your Inciſion you ſhould take great Care not to lay too much Streſs upon your Knife, leſt you ſhould force Splinters of the fractured *Cranium* into the Subſtance of the Brain.

How the Inciſion is to be made.

XVI. If you find it neceſſary to make an Inciſion through the Integuments, it may be made in a right Line; but where that is not ſufficient, let it be formed like the Letter X, about an Inch and an half in Length, lifting up the Skin at each Angle, and leaving the Bone bare: the Blood which is ſpilt may be taken up with a Sponge, and dry Lint ſtuffed between the Skin and the *Cranium*: having found out the injured Part of the *Cranium*, you may now apply the *Trepan* if you ſhall think it neceſſary: ſome Surgeons in ſcalping prefer the Figure of the *roman* Letter V, or the *greek* Λ; others prefer a longitudinal Inciſion: in Wounds which are made near the Temples, great Care muſt be taken not to divide the muſcular Fibres: there are Surgeons who contend much for an Inciſion in the Form of a T: but the Situation of the Wound will always determine you with regard to the Figure of the Inciſion which you ſhall make, either for the Diſcovery of a Fiſſure, or to prevent or remove bad Symptoms.

What is to be done after Scalping.

XVII. Having diſcovered the injured Part of the *Cranium*, and cleared away the grumous Blood and Matter with a Sponge, you are next to remove any Splinters of Bone that may come in your Way, with your Fingers or the Forceps; where they hang to the *Pericranium*, you muſt uſe the Sciſſars; where they adhere pretty firmly to the neighbouring Parts of the *Cranium*, it is more adviſeable to replace them, than to endeavour to remove them by Violence: but if there are no Splinters or Fragments of Bones, and the *Pericranium* is bruiſed, inflamed, or bloody, you ſhould then ſcarify the Part, and proceed as above at *N.* VIII.

How a Contuſion is to be treated.

XVIII. But if the *Pericranium* is quite corrupted and ſeparated, cover the Bone with dry Lint; or bore ſeveral ſmall Holes through the external Lamella of the Bone, till you find Blood proceed from the wounded *Diploë:* after this you may dreſs the Part up with balſamic Medicines, (*N.* IX.) If upon repeating the Dreſſings you diſcover freſh, yellow, or black Spots, the Parts ſo diſcoloured are to undergo the ſame Operation; this is the eaſieſt and moſt expeditious Way of remedying this Diſorder.

How Fiſſures are to be treated.

XIX. When you diſcover a Fiſſure in the *Cranium*, attended with no other bad Symptoms, but white, yellow, or brown Spots upon the Face of the Bone, you will find it ſufficient to bore down to the *Diploë*, and dreſs it with warm balſamic Medicines; in the mean time Bleeding and Diluters muſt not be omit-

ted. There is not always Occaſion for *Trepanning* in Fiſſures, as many beſides HIPPOCRATES have declared: but where any violent Symptoms come on, which demonſtrate an Extravaſation of Blood in the Cavity of the *Cranium*, which cannot be evacuated or diſperſed by the Methods abovementioned, the *Trepan* is to be called for without Delay.

XX. The Surgeons amongſt the Antients uſed another Method for the Cure of Fiſſures of the *Cranium*, that were not attended with very bad Symptoms: their Method was to ſcrape away the upper Table of the Bone, 'till they came down to the *Diploe*; for this Purpoſe they uſed *Rugines*, or *raſping Chiſſels*, of different Shapes, ſemi-circular, plain, or acuminated, as you may ſee in *Plate* VII, *Fig.* 3, 4, 5. this Practice is ſtill continued by ſome; but the Method of boring is far leſs troubleſome, and therefore juſtly preferred to it. The Method uſed by the Antients in this Caſe.

Of DEPRESSIONS *of the* CRANIUM.

XXI. The Skull in Infants and Children is ſometimes depreſſed or dented in by a Blow, like Tin or Copper, without any manifeſt Fracture; or at leaſt fractured in ſuch a Manner, that from its Flexibility it does not ſtart out, but ſtill adheres firmly to the neighbouring Bones. But in Adults this Caſe cannot happen; for the Bones in them are become ſo rigid, that it is impoſſible to beat in any Part of the *Cranium* without breaking the Bone to Pieces: theſe Injuries of the *Cranium* are called by the Surgeons *Fractures* or *Depreſſions*; the Brain is frequently injured by theſe Accidents, and the Actions of it diſturbed. Of Depreſſions of the *Cranium*.

XXII. Theſe Accidents are attended with full as bad Conſequences as thoſe we have already deſcribed: according to the Degree of Depreſſion, ſo is it attended with more or leſs Danger: ſometimes it is quite incurable; for in this Caſe the Veſſels of the Brain are very liable to be injured, which frequently produces ſuch an Extravaſation of Blood in thoſe Parts, as muſt neceſſarily bring on grievous Diſorders, and frequently Death itſelf. Diſorders occaſioned by it.

XXIII. You may eaſily diſcover a *Fracture* or *Depreſſion of the* CRANIUM; 1. by your Eye, 2. by the Touch, 3. by conſidering the Cauſe of the Injury, 4. by the Symptoms that ſucceed it; though theſe alone are very uncertain: Depreſſions and Fractures of the *Cranium* are by no Means ſo difficult to diſcover as Fiſſures: that Fractures of the Skull are attended with great Danger, and frequently with Death, nobody will deny, who conſiders well the Structure of the neighbouring Parts. A Fracture in the *Cranium* is eaſily diſcovered.

XXIV. The firſt Thing to be done towards relieving this Diſorder, is to lift up any Part of the Bone that is depreſſed, or beat in upon the Brain, and replace it, if it ſtill adheres to the neighbouring Bones; or to remove any other Body by which that Part is compreſſed: ſometimes a Splinter, which is quite ſeparated from the reſt of the Bone, is driven into the Cavity of the *Cranium*, and lies conſtantly vellicating the Brain and its Membranes with its pointed Parts: this is to be removed without Delay, yet very tenderly, and with the Caution we recommended, *N.* XVIII. How it is to be treated.

XXV. When ſlight Depreſſions are made in the Skulls of Infants, without bringing on any bad Symptoms, you muſt not uſe the forcible Methods of raiſing the depreſſed Part, which we directed above; but call thoſe Medicines into How ſlight Depreſſions in Infants are to be treated.

into Uſe which we adviſed for the Cure of Contuſions, ſuch as the medicated Bags boiled in Wine, or Spirit of Wine camphorated; or, laſtly, apply a Plaſter to the Part, ſuch as the *Emplaſtrum de Meliloto, ſive de Betonicâ:* nor muſt you omit internal attenuating Medicines, *N.* V. Theſe Applications frequently cure ſlight Impreſſions, and prevent the miſchievous Conſequences which might be expected from them.

How a larger Depreſſion is to be treated.

XXVI. But where a greater Degree of Depreſſion happens to Infants, the Elevation or Reſtitution of the Parts is performed in the following Manner: after ſhaving the injured Part, they apply a Plaſter made of very ſticky and gummy Materials, ſpread upon a ſtrong Piece of Leather, to the Middle of which a Cord is faſtened: this Plaſter is laid on pretty warm, and left in its Situation till it is grown cold; the Surgeon then, taking hold of the Cord that is faſtened to it, pulls the Plaſter directly upwards, and with it the depreſſed Part of the *Cranium*: *See Plate* VII, *Fig.* 6. If this does not ſucceed at the firſt Trial, it is to be repeated: the Application of the Cupping-glaſs to the depreſſed Part will ſometimes ſucceed, eſpecially if you ſtop the Patient's Breath at the Noſe and Mouth during the Operation: but if neither the Plaſter nor Cupping prove of any Service, it will be neceſſary to call for the Aſſiſtance of an Inſtrument like an Awger; ſuch an one as you ſee deſcribed at *Plate* VII, *Fig.* 7. *Lett.* B; which is to be applied after the common Integuments and *Perioſteum* are removed: ROHAULT rejects both the Cupping-glaſs and Awger, and adviſes the *Trepan* in their ſtead, where the Symptoms are bad: ſee his Treatiſe above cited, p. 53.

How a fractured *Cranium* is to be treated.

XXVII. But when the *Cranium* is ſo depreſſed, whether in Adults or Infants, as to ſuffer a Fracture, or Diviſion of its Parts, it muſt inſtantly be relieved: the Part depreſſed, which adheres, after cleanſing the Wound, muſt be reſtored to its Place; what is ſeparated muſt be removed, and the extravaſated Blood be drawn off through the Aperture. Some are very high in their Commendations of a ſternutatory Powder for this Purpoſe, aſſerting that the Diſtention of the Brain is ſo violent in the Act of Sneezing, that it will reſtore the depreſſed Parts of the Bone to their former Situation; but the ill Conſequences that may attend this Practice are ſo grievous, that in my Opinion it ought to be rejected. You will find the Elevatories deſcribed at *Plate* VII, *Fig.* 7. *Lett.* C, *and at Fig.* 8. very ſerviceable, if there is a ſmall Foramen to which the Inſtrument can be faſtened: but if there is no Hole already in the Part, you muſt apply the Screw End of the Inſtrument at *Fig.* 7. *Lett.* B, or one of that Kind, by which Application the depreſſed Part may be reſtored: in the mean time an Inciſion ought always to be made through the common Integuments, that they may be drawn back for the Inſtrument to take Place, *N.* XV. and a Foramen ſhould be made with a ſharp-pointed Inſtrument, (*Fig.* 7. *or* 2. *Lett.* A) to admit of the End of the *Trepan.*

A particular Kind of Elevatory with three Feet.

XXVIII. But as the Elevatories at *Fig.* 7. and 8. are ſo contrived, that where the neighbouring Bones are depreſſed or fractured, theſe Inſtruments cannot be applied without Danger of encreaſing the Complaint, it appeared neceſſary to the Surgeons amongſt the Antients to invent another Inſtrument for this Purpoſe, which might be applied with more Safety; this they called, from the Number of its Feet, *Tripes, Tab.* VII, *Fig.* 12. it is near twice as big as the Figure we have given you: the Feet AAA may be placed at farther

ther Diſtances, or brought nearer to each other, as you ſhall ſee Occaſion: the Manner of applying it is this: the Feet of this Inſtrument are applied to the ſound Parts of the Head; and the Screw B, C, by frequently turning round its Handle D D, will preſently lay hold of the depreſſed Part of the *Cranium*, eſpecially if you have before-hand made a ſmall Hole in the Middle of it with the Awl at *Fig.* 2: upon turning the Screw, E E, the *Trepan* is raiſed by Degrees, and with it the depreſſed Part of the *Cranium*: you will conceive this more clearly by examining *Plate* VII, *Fig.* 13: but if any Opening ſhall appear between the fractured Parts of the *Cranium*, it will be better to take off the pointed End of the Inſtrument, and in its room fix the Elevatory G, by the Screw H, about the Part at Letter F of *Fig.* 12. and by the Aſſiſtance of this the depreſſed Part may be raiſed, as we taught above.

HILDA-NUS's Elevatory.

XXIX. HILDANUS deſcribes an Inſtrument for this Intention, which is a much ſimpler Inſtrument than that which we have juſt ſhewn you, and a very convenient one for the Purpoſe, *See* FAB. HILDAN. *Cent.* II. *Obſ.* 4: we have given you a Deſcription of this Inſtrument in *Plate* VII, *at Fig.* 14: you ſhould be provided with the Awger A, and the Hook at *Fig.* 15. through either of which, according as you ſhall ſee neceſſary, the Lever B C may be paſſed, after the Inſtrument is fixed upon the depreſſed Part of the *Cranium:* the Plate D is to be placed upon the ſound Part of the Head, laying Bolſters under it to prevent Pain; then, by raiſing the End of the Lever at B, the depreſſed Part of the *Cranium* will be gently elevated and reſtored to its natural Situation: you will obſerve a Joint at the Extremity of the Lever C, to accommodate the Plate D to the Convexity of the Head in ſome Parts of it, which may be alſo raiſed or depreſſed by the Screw E: if you pleaſe, you may make the Lever longer than it is repreſented here, which will add to its Force: PETIT has deſcribed a new Kind of Lever [a]; which I have given you the Figure of, *Plate* XXXIX, as it may ſometimes be of Service.

A particular Way of removing Splinters.

XXX. But if any Part of the Bone is entirely ſeparated from the reſt, and driven ſo deep into the Cavity of the *Cranium*, that it cannot be elevated or extracted by the Methods which we have already propoſed, you muſt perforate the neighbouring ſound Part with a *Trepan*, and divide the intervening Part with a fine Saw, *Fig.* 9. as deep as you ſhall think you can with Safety: after this you may cut it entirely through with the Chiſſel and leaden Mallet at *Fig.* 10. *and* 11: having made an Opening in this Manner, you will have a full Command of any Splinters or foreign Bodies that are driven into the *Cranium*, and will more eaſily evacuate the extravaſated Blood: Caſes that require this laſt Method of operating are very rare, but they are no leſs neceſſary, though the Operation requires great Pains and Dexterity in the Performance of it.

How to ſecure the Parts in the Situation you have reſtored them to.

XXXI. Having raiſed up the depreſſed Parts of the *Cranium*, and reſtored them to their natural Situation, you muſt take great Care to ſecure them from a freſh Depreſſion; the Patient ſhould lie on the ſound Side of his Head, the fractured or depreſſed Part ſhould be guarded with a braſs or ſteel Plate, and the wounded Part ſhould be treated according to the Rules which we have already laid down.

[a] *Memoires de Chirurgerie*, Tom. i. p. 302.

EXPLANATION

EXPLANATION *of the* SEVENTH PLATE.

Fig. 1. *An artificial Eye* made of Glaſs or Silver, painted after the Life; this may be introduced into the Orbit, and ſupply the Place of the natural Eye, and prevent the Deformity that will enſue upon the entire Loſs of that Organ.

Fig. 2. *An Awl*, or ſharp Inſtrument, to perforate the external Table of the *Cranium*.

Fig. 3, 4, 5. Different Forms of *Rugines*, or *raſping Chiſſels*, to ſcrape the *Cranium*, or other Bones.

Fig. 6. Shews how the *Depreſſion of the Cranium* in an *infantile State* may be relieved by ſticking Plaſters.

Fig. 7. A, a quadrangular, or pointed ſteel Inſtrument, to perforate the external Table of the *Cranium*; B, an Awger; C, an Elevator to raiſe depreſſed Bones of the *Cranium*.

Fig. 8. Another Elevator for the ſame Uſes with the former.

Fig. 9. A ſmall fine Saw; and *Fig.* 10. a ſmall Rugine, which may be uſed with or without the Handle deſcribed to that at *Fig.* 3.

Fig. 11. A wooden Mallet, the Head of which is filled with Lead.

Fig. 12. An Elevator with three Feet: ſee above, *N.* XXVIII.

Fig. 13. Deſcribes the Method of applying this Inſtrument.

Fig. 14. HILDANUS's Elevator: ſee above, *N.* XXIX.

Fig. 15. A Hook belonging to HILDANUS's Elevator.

How extravaſated Blood is to be diſcharged from the CRANIUM.

Diſorders occaſioned by Extravaſation of Blood.

XXXII. In the Injuries of the *Cranium* that we have been deſcribing, that is, in Contuſions, Fiſſures, Depreſſions, and Fractures, one or more of the Blood-veſſels that are diſtributed upon the *Dura Mater* are frequently divided: the Blood that is diſcharged by this Accident greatly oppreſſes the Brain, and diſturbs its Offices; this frequently brings on violent Pains, Deprivation of Senſes, and other Miſchiefs, and at length Death itſelf, unleſs the Patient be timely relieved: if the Quantity of extravaſated Blood be ever ſo ſmall, it will certainly corrupt, and affect the *Meninges* and the Brain itſelf with the ſame Diſorder; from hence will proceed violent Inflammation, Delirium, Ulcers, and what not? even Death itſelf, ſooner or later: and this will frequently be the Caſe after a violent Blow upon the *Cranium*, when a Vein or Artery is wounded, though the Bone ſhould eſcape without any Injury.

Where the Blood is ſpilt.

XXXIII. In theſe Injuries of the Head, the Blood is ſpilt either between the *Cranium* and *Dura Mater*, or between the *Dura* and *Pia Mater*, or between the *Pia Mater* and the *Brain*; or laſtly, into the *Sinuſes of the Brain*: each of theſe Caſes are attended with great Danger, but the deeper the Extravaſation happens, and the more copious the Diſcharge, ſo much the greater will the Danger be.


How to diſcover an Extravaſation of Blood in the *Cranium*.


XXXIV. You may ſuſpect that Blood is extravaſated in the Cavity of the *Cranium* from the Violence of the Symptoms which ſucceed; if the Patient lies ſtill


Plate 7.



Plate 7.




Fig: 14.
A
B
C
E
D
Fig: 15.
B
A
Fig: 7
Fig: 8.
A
B
Fig: 6.
C
D
C
D
Fig: 12.
E
E
A
F
B
A
H
G
A
Fig: 13.



ſtill without Senſe or Motion, if Blood flows from the Mouth, Ears, or Noſe, if the Eyes are much inflamed and ſwelled; if Vomiting ſucceds, when upon the Remiſſion of theſe Symptoms the Patient complains of a remarkable Heavineſs of Head, a Sleepineſs, Vertigo, Blindneſs, Spaſms, and Diſorders of this Kind: when the Quantity of extravaſated Blood is very conſiderable, and oppreſſes the *Cerebellum*, the Patient dies upon the Spot; but when the Extravaſation is not in a very large Quantity, or at leaſt does not affect the *Cerebellum*, Life ſtill remains, but the Symptoms related above come on: ſometimes theſe Symptoms come on very ſlowly, and great Numbers of Perſons, who have appeared at firſt to have been but ſlightly wounded, have died in this Manner, after ſome Time, contrary to all Expectation: therefore I cannot help again admoniſhing the Surgeon, that after violent Blows of the Head, though no violent Symptoms ſhould immediately urge, yet he ſhould be very cautious in delivering his Prognoſtic, and not be too haſty in his Opinion, nor, by treating the Caſe as ſlight and indifferent, endanger the Life of his Patient: but when violent Symptoms immediatly enſue, you may always be ſure that there is an Extravaſation of Blood though no great Injury appears upon the external Part of the Head.

How to diſcover the Part in which the Blood is extravaſated, thoughthere appears no external Wound.

XXXV. If you can find no Fracture, Fiſſure, or Contra-fiſſure, in the *Cranium*, nor even any external Injury upon the Integuments of the Head after a violent Blow, and the Patient is deprived of his Senſes, you will find it difficult to determine in what Part of the Head an Extravaſation is ſeated: it will be proper therefore, 1. to ſhave the Head all over, that, you may be the better able to examine it: for if any Part is ſofter than ordinary, or enlarged, or red from a Stagnation of Blood, it is plain that this is the Part which received the Injury: you may alſo examine Perſons who were preſent at the Accident, from whom you may frequently get Light into the Affair: but, if you are ſtill left in the Dark, 2. cover the whole Head, after it is cloſe ſhaved, with an emollient Plaſter, laying over it medicated Bags well-heated; this Application will, in a few Hours, produce Tumor and Softneſs upon the injured Part: 3. ſometimes the Patient, though he lies ſpeechleſs, and to all Appearance ſenſleſs, will be continually clapping his Hand to the aggrieved Part: 4. if either Side of the Patient has loſt Senſe and Motion, and is become paralytic, it is an apparent Sign, whatever ſome may think to the contrary, that the Injury was received on the contrary, or ſound Side: *See* MORGAGNI *Adverſaria Anatomica* VI. *et Diſſert. de Reſonitu, Argentorat.* 1722. *Edit. Pag.* 23: If you diſcover any Wound in the Skin, you ſhould enlarge it with the Knife, till you come at the Injury in the *Cranium*, whether Depreſſion, Fiſſure, Contra-fiſſure, or Fracture.

How the injured Part is to be treated.

XXXVI. When you have diſcovered the Seat of the Injury, the firſt Intention is to diſcharge the extravaſated Blood, which muſt otherwiſe endanger the Patient's Life; and then to clean the Wound, and remove all Splinters or extraneous Bodies. Many Writers in Surgery adviſe the inſtant Uſe of the Trepan, to make Way for a Diſcharge of the extravaſated Blood; but ſince this is a difficult and dangerous Operation, and many have recovered without having Recourſe to it, I ſee no Reaſon for attempting it, unleſs we are driven to it by abſolute Neceſſity: therefore I think it is beſt to try firſt the Force of attenuating and dividing Medicines in this Caſe.

XXXVII. With

How inſpiſſated Blood ſhould be attenuated.

XXXVII. With this Intention, 1. open a Vein, and draw away as much Blood as the Strength of your Patient will admit; this will take off the Impetus of the Veſſels, and prevent the Extravaſation of more Blood: 2. preſcribe a pretty briſk Purge, to leſſen the Quantity of Fluids, for which Purpoſe you may alſo give ſharp Clyſters: 3. foment the Head with medicated Bags, and apply a Melilot Plaſter to it: 4. endeavour to rouſe the Patient by volatile Applications to his Noſtrils, ſuch as *Sal volatile oleoſum*, *Spiritus Salis Ammoniaci*, *vel Spiritus Cornu Cervi per ſe:* laſtly, 5. give frequently attenuating Fluids warm, ſuch as Infuſions prepared *ex Theâ*, *Betonicâ*, *Salvia*, *Roriſmarino*, *Lavendulæ Floribus*, *Ligno Saſſafras*, and the like: this Method will contribute greatly to the thinning and diluting the Blood.

XXXVIII. Yet this does not immediately procure the deſired Effect, therefore it muſt be continued for ſome Time, and the Preſcriptions frequently repeated; and more particularly when the Symptoms ſeem by Degrees to abate: the Repetition of Bleeding in this Caſe may ſeem ſtrange to ſome, but it muſt be to thoſe who are ignorant of the good Effects it produces, by leſſening the Quantity of Fluids, and by reſtoring the Courſe of the ſtagnating Blood: if the Patient finds a little Relief from the firſt Bleeding, it will be proper to repeat the Operation a ſecond and a third Time, eſpecially if he is young and athletic, and to apply the Remedies which we have recommended above in the Intervals, till the Diſorder is entirely removed.

Sometimes the Trepan is neceſſary.

XXXIX. But when you find, notwithſtanding theſe Applications, that the Symptoms rather encreaſe than abate, you will be obliged to make a Perforation in the *Cranium* with the *Trepan*, near the Seat of the Wound, that there may be a Paſſage for the Diſcharge of the confined grumous Blood: but there ſhould be great Caution always uſed in this Operation: if the extravaſated Blood or Matter be collected under the *Dura* or even the *Pia Mater*, an Inciſion muſt be made in theſe Membranes, without Reſerve, that the Enemy may be removed; the Wound is then to be cleanſed, and afterwards healed by proper Applications: when you cannot diſcover the Part of the Head which is principally affected, and the Symptoms are ſtill as violent, or rather aggravated, you muſt perforate the Skull in ſeveral Places, till you hit upon the right: for, if this Method does not always produce the Effect deſired; yet, with CELSUS, it is far better to try a doubtful Remedy, than none at all: I ſhall teach the Manner of performing this Operation, and the Methods of healing the Wound, in another Part of this Work, which treats profeſſedly of *Chirurgical Operations*.

Principal Writers on Wounds of the Head.

XL. If you deſire to ſee Hiſtories of Cures of Wounds of the Head, conſult HIPPOCRATES *De Capitis Vulneribus*, *cum* ARANTII & PAAWII *Commentariis*, and CELSUS on the ſame Subject: add to theſe, BERENGARIUS *De Fractura Cranii*, ARCÆUS *de Vulneribus*, SCULTETUS *in Obſervationibus*, 1 *ad* 23. HILDANI *Obſervationes variæ*, TULP. *Obſ.* L. i. C. 14. SCHULTZIUS, *De Capite læſo*, BELLOSTIUS *in Chirurgo Noſocom.* WOYTIUS and WALTHERUS *De Capitis Vulneribus*, and ſeveral others; but particularly amongſt the modern Writers, ROHAULT's Book on Wounds of the Head, called *Traité des Playes de Tete*, 4*to*, 1720, and LE DRAN, *in his Chirurgical Obſervations*.

CHAP.

CHAP. XV.

Of CONTUSIONS.

I. A CONTUSION is any Hurt of the Body that is inflicted by a blunt Inſtrument; and ſince, in this Caſe, an infinite Number of ſmall Veſſels and Fibres are injured and broken, a Contuſion may properly be ſaid to be a Congeries of an infinite Number of exceeding ſmall Wounds: it is well enough called by the GREEK PHYSICIANS *Ecchymoſis*, and by CELSUS, *Vulnus Colliſum*, *Lib.* V. *Cap.* 26. A Contuſion what.

II. Contuſions may be diſtinguiſhed into ſeveral Sorts: 1. ſome may be called *ſimple Contuſions*, that is, when only the ſoft external Parts are injured; ſome are *compound*, when the internal or bony Parts alſo partake of the Injury: 2. ſo ſome Contuſions are ſlight, ſome of great Conſequence; others prove mortal, and in ſome Caſes immediately; this depends upon the Cauſe of the Injury, and the Nature of the Part injured: 3. laſtly, ſome Contuſions are ſo circumſtanced, which is very wonderful, that the internal Parts ſhould be violently affected, whilſt the external Parts remain whole and unhurt; for we are experimentally taught, that a Man may receive a Blow with a blunt Weapon, or even with a naked Hand, upon the Head, Breaſt, or Belly, which ſhall occaſion inſtant Death, though there ſhall appear no external Signs of Injury: *See* BOHNIUS *De Vulner. lethal. Sect.* I. *Cap.* I. Differences.

III. Contuſions are uſually occaſioned, 1. by violent Blows given with blunt Weapons, ſuch as Staves, Bludgeons, or Stones, or a Bullet almoſt ſpent: 2. the ſame will happen from a Fall upon the Stones, or any other hard Body: 3. Contuſions are occaſioned by the Body being preſſed between two Doors, by Preſſes, Screws, Mills, Wheels, and ſuch like Machines; for, by Accidents of this Kind, the Veſſels are either entirely broken, or the Blood is violently ſqueezed out of them. Cauſes of Contuſions.

IV. When the ſmall Veſſels and Fibres have been broken by a Contuſion, the Fluids that were contained in them will be forced out; hence will proceed Obſtructions, Corruption, Inflammation, and Ulcers, or even Gangrene, and ſeveral other fatal Miſchiefs, in Proportion to the Violence of the Cauſe, and the Nature of the affected Part. When the external Parts are contuſed, the Skin at the ſame Time remaining whole, the Blood will ſtagnate under it, and occaſion red, black, and livid Spots, which we call a *Sugillation*; from whence ariſe ſeveral other Miſchiefs; and if this happens near a Bone, a Caries, or a Fracture What ſucceeds a Contuſion of the ſoft Parts.

V. When a bony Part is the Subject of a Contuſion, then, 1. the ſame Miſchiefs will enſue from the Injury inflicted upon the *Perioſteum*, which we have already deſcribed as happening to the *Pericranium* in Wounds of the Head: but when this Diſorder, 2. is accompanied with a Fracture, the ſame Miſchiefs will enſue, which uſually attend fractured Bones, and theſe always increaſe in Proportion to the Force of the Blow; on which Account the Contuſions from Bullets, *&c.* are generally attended with the worſt Conſequences. If the Injury is in the Bones of the *Cranium*, the *Thorax*, or the *Vertebræ*, you may expect all the Miſchiefs, of which we have largely ſpoken above in the preceding Chapters: laſtly, 3. when the medullary Juice of the Bones is affected, you Of the Bones.

may expect every violent Diſorder, whether the Bones are fractured or not; for the Blood which is diſcharged out of the Veſſels that are ſent to the *Medulla*, will preſently corrupt and produce a Gangrene; or, by corroding the Bones, bring on a Caries, Ulcers, and incurable Fiſtulæ; which will make it neceſſary to take off the Limb, to ſave the Life of the Patient; for the medullary Juice is in the ſame Condition in theſe Caſes with the Brain in Fractures or Contuſions of the *Cranium*.

Of the Joints and Muſcles.

VI. Contuſions of the Joints uſually bring on violent Pains and Inflammations, Convulſions, Gangrene, Sphacelus, Rigidity of the Limbs, and Caries; the ſame will ſometimes happen from Contuſions of the muſcular Parts. When the internal Parts are contuſed, great Miſchiefs uſually enſue, but that depends entirely upon the Nature of the injured Part, and the Degree of the Injury; ſometimes Inflammations, Rupture of the Veſſels, Varices, Aneuriſms, Hæmorrhages, Stagnation of the Fluids, Corruption, Gangrene, Suppuration; and ſometimes, as a neceſſary Attendant upon theſe, Death. When the Head receives a conſiderable Contuſion, the Senſes are then taken away, the Limbs become either convulſed or rigid, and Death preſently follows, in the Manner we have already explained, treating upon Wounds of the Head. If a violent Contuſion falls upon the Thorax, a Difficulty of Breathing follows, with Spitting of Blood, fainting Fits, Inflammation, and Ulcers of the Lungs, which uſher in Death. After Contuſions of the Abdomen, you may expect Vomiting of Blood, Inflammations, Suppurations, or Gangrene of the Viſcera, and at length Death[a]: if any internal Veſſel is burſt by the Violence of a Blow, it is no Wonder if the Patient dies upon the Spot, though there be no Mark of Violence left upon the external Parts: laſtly, if the Eye is contuſed, Tumor and Inflammation will ſucceed, and frequently Loſs of Sight.

What ſtill remains to be done.

VII. Contuſions may be examined, 1. by the Eye, when they are inflicted upon the external Parts of the Body; Tumors are formed, the injured Parts are diſcoloured, at firſt becoming red or black, then livid, yellow, green, and at laſt black again; if the Contuſion is not very conſiderable, the Parts will of themſelves recover their natural Colour: 2. when the Contuſion is not within the Reach of the Eye, you muſt feel for it; an unnatural Softneſs of the Limb, or a Fluctuation of the extravaſated Blood under your Fingers, will pretty clearly point out the injured Part to you: 3. Pains and Rigidity of the contuſed Part will make the ſame Diſcovery: laſtly, 4. you may form ſome Judgement of the Degree of the Injury received, from conſidering the Manner in which it was given, and the Size and Nature of the inflicting Inſtrument: you will judge what internal Parts are injured, by the Symptoms which ſucceed, and by obſerving which of the Functions of the Body are diſturbed or deſtroyed: if a Fracture attend the Contuſion, it will eaſily be diſcovered by the Eye, the Touch, and the Ear.

[a] An Inſtance of this Kind happened in the Year 1726, at a Village near *Helmſtadt*; a Schoolmaſter there beat one of the Children very ſmartly, with a Stick of no great Size, but the Boy died in a few Days afterwards; upon opening him, the Viſcera of the Abdomen appeared grievouſly bruiſed and lacerated: I opened another Boy ſoon afterwards, who was killed by a Blow, and found his Liver divided quite through the Middle, though there appeared no external Injury: Conſult here the Quotation from BOHN, in the preceding Page: In the Year 1738, a Boy's Spleen was torn by the Kick of a Horſe, and the Cavity of the Belly found full of Blood.

Prognoſis.

VIII. What we have ſaid above, concerning the Nature and neceſſary Effects of Contuſions of each parcicular Part, will give the Surgeon great Light in forming his Prognoſtic; nevertheleſs, it will not be improper to ſubjoin a Rule or two in this Place. Slight Contuſions are attended with little or no Inconvenience or Danger, beſides diſcolouring the Skin; and even that Deformity is of a very ſhort Date, for the ſtagnating Blood is preſently licked up again, and the Spots vaniſh: but in larger Contuſions, where there is a great Collection of ſtagnating Blood in the muſcular Parts, an Abceſs, Gangrene, or Sphacelus, will eaſily follow. Contuſions of the internal Parts are extremely dangerous; and the Degree of Danger encreaſes in Proportion to the Violence of the Contuſion, and the Conſequence of the Part in performing the neceſſary Offices of Life. If inſtant Death does not happen in this Caſe, yet it is uſually attended wirh ſuch dangerous Inflammations, that the Patient conſumes away by Degrees, and very rarely eſcapes. Contuſions of the Bones, particularly of their *Medulla*, and of the Joints or Ligaments, are very dangerous, eſpecially thoſe which are inflicted by Gun-ſhot; but the Contuſion or Fracture of the *Cranium* from the Vicinity of the Brain, and of the Bones of the Thorax from their near Relation to the Heart and Lungs, exceeds the reſt in the miſchievous Conſequences which attend it, as we have largely enough explained above.

Cure of ſlight Contuſions.

IX. Your principal Care in the Cure of Contuſions ought to be, to divide the inſpiſſated Fluids, and at the ſame Time to prevent the Parts from ſuppurating, and being affected with Gangrene. There are ſeveral Methods ſucceſsfully uſed for the Cure of ſlight Contuſions; for Example, when a Tumor ariſes in the Forehead from a Fall, which very frequently happens to Children, it will eaſily be cured by fomenting it *cum Vino calido, Spiritu Vini vel ſolo vel camphorato, Aqua Reginæ Hungariæ*; or by applying cold Water or Vinegar mixed with Salt to the Part; or by clapping a broad Piece of Money, or a Plate of milled Lead, upon the Tumor, and faſtening it on with a very tight Bandage. Perſons of ſlender Circumſtances may find eaſier and cheaper Remedies; nor will they be baulked in their Expectations, if they apply linen Rags dipped in freſh warm Urine to Tumors of this Kind.

Of larger Contuſions.

X. Larger Contuſions may be dreſſed with Decoctions *ex Scordio, Sabina, Abrotano, vel ſeorſim vel junctim, in Vino, vel Aqua ſalſa*, repeating them warm with linen Cloths, and with the medicated Bags: you will find great Benefit by applying a Sponge dipped *in Decocto Saponis Veneti in Urina recenti*: your End alſo will be ſufficiently anſwered by Applications of *Spiritus Frumenti*, or *Aqua Calcis, cum admixto Spiritn Vini camphorato, vel Acetum Lithargyriſatum, item Acetum cum ſemine Carui coctum*: theſe Remedies are all to be applied warm.

Of violent Contuſions.

XI. When the Contuſion is ſo violent, that it is apparently impoſſible to divide the ſtagnating Fluids, and return them into the Circulation; and the Parts are haſtening to become gangrenous, you muſt ſcarify them without Delay, carefully avoiding the larger Trunks of the Veſſels: by this Means you will ſet the ſtagnating Fluids at Liberty, and prevent dangerous Conſequences, as Tumors and Inflammations, Suppuration and Gangrene; and the Cure will be eaſily effected.

What is farther to be done.

XII. Having done this, you are in the next Place to apply proper Fomentations, or medicated Bags, made in the Manner we directed in *Chap.* XIV.

N. 10. or according to the following Preſcription: ℞: *Rad. Bryoniæ* ʒij *vel* ʒiij. *Herbæ Sabinæ—Scordii—Abrotani, Arboris Vitæ, ſive Thuyæ vel Abſinthii ana,* Mij. *Singula iſta minutim diſſecantur, affuſiſque Vini circiter Libris duabus, per Horæ quadrantem probe decocta, per Panniculum laneum precolantur. Debinc Saponis Veneti vel Hiſpani aliquot Unciæ huic Decocto probe calido admiſcentur, complicatique panniculi lanei ex eodem expreſſi per ſingulas fere Horas læſæ Corporis parti calide ſuperinjiciantur*: rub the Tumor well with hot Cloths before you foment it, which will keep the Blood in its fluid State; or, if it is already concreted, it will divide it, and make it fit to return into the Veſſels, or at leaſt to eſcape through the inviſible Pores of the Skin. If you cannot be ſupplied with Wine to make your Fomentation, you muſt uſe ſalt Water; which, if you are not near the Sea, you may make of common Water two Pints with the Addition of a Handful of Salt: if any one is better pleaſed with the Form of a Cataplaſm, he may prepare a very cheap, and no leſs uſeful one in the following Manner: ℞. *Pulver. Radic. Bryoniæ, Saponis Veneti ana* ʒiij. *coq. in Aquæ recentis vel Aquæ ſalſæ q. ſ. ad Conſiſtentiam Cataplaſmatis:* this will have ſtill greater Efficacy if you add *Gummi Galbani vel Ammoniaci* ʒj. *in Vitell. Ov. q. ſ. ſolut.*

Of internal Remedies and a proper Diet.

XIII. Where the Contuſion is of any Conſequence, you ſhould never neglect the Adminiſtration of internal Medicines; and here your Intention is to promote the Diſcharge of Sweat and Urine, by preſcribing dividing and attenuating Decoctions and Infuſions to be drank plentifully: theſe may be prepared *ex Theâ, Betonicâ, Veronicâ, Salviâ, Roriſmarino, Ligno Saſſafras, Herbâ Arnicâ, vel Petroſelini Radicibus:* the Efficacy of theſe Medicines in dividing inſpiſſated Fluids is ſcarcely to be conceived, eſpecially if you now and then add to a Draught of one of theſe Infuſions a Drachm of *Venice* Soap: you will find no leſs Aſſiſtance from the *Pulvis ad Caſum Auguſtanorum*, or from *Sperma Ceti, vel ſolum vel cum admixtis Sanguine Hirci, Mumia, Cancrorum Lapidibus, in Pulverem redact.* Theſe may be given to a Drachm at a Doſe, in a Draught of any of the former Infuſions: in plethoric Habits you ſhould never forget to open a Vein, and repeat it as often as you are threatened with an approaching Abceſs or Gangrene; the Patient muſt abſtain from Fleſh and ſtrong Liquors, living wholly upon Broths and thin Spoon-meat.

What ſtill remains to be done.

XIV. The Fluids that were collected together by the Contuſion being pretty well diſperſed by the Methods we have recommended above, the remaining Part of the Cure that principally regards the Wound (which frequently accompanies this Caſe) is eaſily performed, by filling it up with Pledgits ſpread with a digeſtive Medicine, and laying on a warm Plaſter over the Dreſſings; which will ſave the Surgeon the Trouble of preparing Cataplaſms and Fomentations for this Purpoſe, and anſwer his End as well: the *Emplaſtra Diaſaponis, Diachylum, de Meliloto, de Spermate Ceti, de Galbano*, all anſwer this Intention; or if you pleaſe you may uſe the following ℞. *Empl. de Meliloto* ʒiv. *Galban. puri ſolut.* ʒij. *Farin. Rad. Bryon.* ʒj. *Flor. Sulphur. Æthiop. min. ana* ʒß. *Ol. Chamæmel. q. ſ. M. f. Emplaſtrum:* in the mean time, the Regimen which we directed above, both with regard to Medicine and Diet, ſhould be ſtrictly obſerved: the moſt dangerous Contuſions are cured in this Manner much eaſier than by Suppuration or Scarification. Having anſwered the Intention of diſperſing the ſtagnating Fluids, and cleanſing the Wound, nothing remains but

to

to forward the Union of it by Applications of the vulnerary Balſam, and at laſt dry Lint, as we have already adviſed for healing other Wounds.

How the Parts are to be treated when they ſuppurate.

XV. It ſometimes happens, when the contuſed Parts lie very deep, or the Surgeon is ignorant of his Buſineſs, or the Patient refuſes to ſubmit to proper Treatment, that the ſtagnating Fluids will corrupt and ſuppurate: when the Suppuration is begun, it muſt be forwarded, 1. by emollient Cataplaſms prepared *ex Rad. Malv. Althææ, Liliorum alborum, Herbis Malvæ, Althææ, Parietariæ, Mercurialis, Brancæ Urſinæ, Meliloti, Verbaſci, Ficubus, Lini Semine, Fœnogræci, Farinis Variis, Micis Panis cum affuſo Aqua vel Lacte coctis ad Pulticulam, Butyroque, Adipe, Oleiſve emollientibus, Lini ſcilicet, Chamæmelæ, Liliorumque Oleis dilut.*; theſe are to be applied to the Part as hot as they can be well born: 2. ſometimes, in this Caſe, it will be proper to mix warm Medicines with Emollients, ſuch as *Cepæ ſub Cineribus toſtæ, Fermentum Panis, varia Gummata, Galbanum ſcilicet, Ammoniacum, Bdellium, Opoponax in Vitell. Ovor. ſoluta*; theſe are to be mixed with the emollient Ingredients which we enumerated above: for Example, ℞ *Herbæ Malvæ, Althææ, Parietariæ, Meliloti ana M* i. *conciſa coquantur in Aqua ſimplici q. ſ. ad Conſiſtentiam Cataplaſmatis. Adde Ceparum ſub Cineribus toſtarum,* ℥iv. *Galbani, Vitell. Ov. ſolut.* ʒii, *Ol. Lilior. albor.* ℥iſs, *Farinæ Sem. Lini q. ſ. ad Conſiſtentiam*: theſe Applications are to be repeated till the Suppuration is thoroughly formed: in ſmall Contuſions, the *Emplaſtrum Diachylon cum Gumm.* will ſufficiently anſwer this Intention.

How the Matter is to be diſcharged.

XVI. When the Whiteneſs and Softneſs of the Tumor evidently diſcover that the Matter is thoroughly formed, and fit to be diſcharged, you may lay open the Part with your Knife, and afterwards digeſt and heal the Wound in the ſame Manner as we have frequently directed above; if it breaks of itſelf, it ſhould be treated in the ſame Manner: where the Aperture is too ſmall, it muſt be enlarged with your Knife, that it may be eaſier cleanſed, and more conveniently healed.

How a Gangrene and Sphacelus are to be treated.

XVII. Large Contuſions are ſometimes attended with violent Inflammation or Gangrene; in this Caſe make frequent and deep Inciſions upon the Part, and dreſs the Wounds *cum Theriaca Spiritu Vini Camphorato dilut.* applying warm Fomentations externally, not omitting the internal Medicines preſcribed at *N.* XIII. (but I ſhall treat more largely upon this Head in *a Chapter upon Gangrene and Sphacelus*): when a Sphacelus is begun, if it is only in the common Integuments, you muſt apply Scarification, a digeſtive Ointment, Medicines that reſiſt Gangrene, and Suppuration; but, if the whole is ſphacelated, that is, entirely corrupted and mortified, the Limb muſt be entirely taken off, in the Manner we ſhall ſhew you when we come to deſcribe *Chirurgical Operations.*

Cure of internal Contuſions.

XVIII. When the internal Parts are contuſed, unleſs the Patient has immediate Aſſiſtance, Inflammations, Suppuration, and Gangrene inſtantly enſue; which ſoon terminate in Death: therefore, in theſe Caſes, the Surgeon ſhould endeavour to diſperſe and attenuate the concreted Blood with the utmoſt Expedition, by frequent Blood-letting, by gentle opening Medicines and Clyſters, *Chap.* XIII, *N.* XXXVII, by preſcribing the warm Decoctions and Infuſions which we directed above at *N.* XIII: if the Diſeaſe is curable, theſe Methods will prevent Suppuration or Mortification. Theſe Parts do not admit of Inciſion, and the Uſe of abſorbent Powders, ſuch as *Lapides Cancrorum, Sanguis Hirci, Cornu Cervi*

Cervi, *Pulvis ad Caſum*, and the like, is trifling in this Caſe. We have already ſufficiently explained how Contuſions of the Head in particular ought to be treated, in the preceding Chapter; but, in Contuſions of the Breaſt, or the Belly, you can direct nothing better externally than a Compreſs, ſteeped in Spirit of Wine camphorated, or a Bladder filled with warm Milk, in which the Flowers of Chamomile or Alder have been boiled; apply theſe continually warm to the Part affected: for further Applications, conſult the Method laid down for the Treatment of Wounds in theſe Parts.

Contuſions of the Eye.

XIX. When the Eye, that moſt noble and uſeful Organ, is contuſed by any Accident, it will ſwell immediately, and be entirely deprived of Sight, except the Contuſion is very ſmall, and proper Remedies are inſtantly and carefully applied: if the Eye, therefore has received a ſlight Contuſion, you may waſh it frequently, for the firſt Day, with cold Spring Water, covering it with linen Rags, wet with the ſame: on the next Day, rub it externally *cum Spiritu Vini camphorato*, covering it with Stuphs wrung out of vinous Decoctions *ex Euphraſiâ*, *Veronicâ*, *Hyſſopo*, *Salviâ*, *Florib. Chamæmel. & Semin. Fœnicul.* If you cannot get theſe Herbs, you may apply Bolſters, dipped in *Vino calido*, renewing them often; if the Contuſion is large, or the Patient of a plethoric Habit, you ſhould open a Vein. *On Bloodſhot Eyes, ſee* CELSUS, *pag.* 369.

Of violent Contuſions of the Eye.

XX. If the Contuſion of the Eye is ſo violent, that you can plainly ſee the extravaſated Blood through the *Cornea*, and all Objects appear red to the Patient, open a Vein either in the Foot or Neck, as you ſhall think moſt convenient, and repeat it, if neceſſary; foment the Eye with Stuphs wrung out of the Decoctions which we preſcribed above, and order him to bathe his Feet in warm Water two or three Times in a Day, adviſing him alſo to obſerve the ſame Regimen with regard to Diet and internal Medicine, which we deſcribed at *N.* XIII. By the ſtrict Obſervation of theſe Rules, he will recover his Sight, if the Diſorder be not become deſperate, eſpecially if you frequently drop warm Pigeon's Blood into the Eye. If theſe Attempts to diſperſe the ſtagnating Blood are fruſtrated, you may very probably ſucceed by making an Opening in the *Cornea* with your Lancet; the Manner of doing this to Advantage you will find deſcribed in *Chap.* LX, and LXI. of the *Second Part* of this Work, which treats profeſſedly of *Operations*.

CHAP. XVI.

Of VENOMOUS WOUNDS, *and thoſe that are made by the* BITES OF ANIMALS.

Poiſonous Wounds are diſtinguiſh.

I. WE are informed by antient Tradition, that the *Indians*, and the barbarous Nations all over *Africa*, uſed to poiſon their warlike Weapons, to aggravate the Wound, and deſtroy their Enemies with greater Certainty; which ſome of them continue to this Day; this Cuſtom has long ago been laid aſide by the *Europeans*, as inhuman. Wounds that are inflicted by Weapons of this Kind are attended with extreme Danger: for, as this Sort of Miſchief is in a great Meaſure concealed and unexpected, there is no Room to make,

make Uſe of proper Precautions to prevent or remedy the Evils that will enſue from it.

II. For though ſeveral *Phyſicians* and *Surgeons* have aſſerted, that you may diſtinguiſh Wounds made by a poiſonous Weapon, not only by the filthy Stench of the wounded Parts, and the unuſual Colour of the Diſcharge that proceeds from them, to wit, yellow, green, livid, and black; but particularly by the Increaſe of Pain, by the extravagant Degree of Tumor and Inflammation that they are attended with; Palpitation of the Heart, Swooning, Spaſms, Diſtortion, or Rigidity of the Limbs, cold Sweats and Shiverings, with which the Patient is conſtantly afflicted in this Caſe: nevertheleſs, if I may be allowed to judge, I muſt determine theſe Symptoms to be altogether doubtful and uncertain; for what Surgeon does not know, that all theſe Symptoms may be brought on either by the bad Habit of the Patient, or from the Nature of the wounded Part, if it is nervous or tendinous: or, in a Word, from an hundred other Cauſes, where Poiſon is no ways concerned? For the Marks of them are very uncertain.

III. You have much greater Certainty of a Mixture of Venom in the Wound, when it is made by the Bite of a *venomous* or *mad Animal* (for there is ſcarce any Species of Animals but what is at ſome times ſubject to Madneſs), eſpecially of a Dog, a Cat, a Wolf, an Ape, a Man, a Serpent, a Scorpion, or of any other venomous Inſect: but, ſince the Coldneſs of our Climate renders us very rarely ſubject to Injuries from the Bites of venomous Serpents, or indeed of any other venomous Animal but a *mad Dog*, it will be moſt to our Purpoſe to treat chiefly of that Subject; at the ſame time not entirely neglecting the Deſcription of other Wounds inflicted by Biting: and firſt we ſhall ſpeak of the Bites of Animals which are not mad. Of Bites.

IV. Bites of enraged Animals are attended with very grievous Conſequences, though they are not afflicted with Madneſs: [a] CELSUS has long ago taught us, that the Bites of a [b] Man, an Ape, a Cat, a Dog, or of any wild Beaſt or other Animal, frequently bring on terrible Miſchief: in that Paſſage of his where he ſays, *Omnis* [c] *fere Morſus quoddam Virus habet*, "almoſt all Bites whatever have "ſomething poiſonous in them;" he is not to be underſtood as if he had aſſerted that all Wounds, made by Bites, have actually ſome Particles of Poiſon, properly ſo called, inſtilled into them; but rather as ſpeaking of the bad Symptoms which muſt neceſſarily enſue from the violent Laceration and Contuſion of the Muſcles, Nerves, Tendons, Ligaments, and Bones, by the Bite of a large Dog, a Horſe, a Wolf, or a Bear, or any other large Animal. If the Wound is ſlight, encourage the Diſcharge of Blood from the Part, by preſſing it with your Bites of Animals which are not mad.

[a] *Lib.* v. *Cap.* 27. *N.* 1.

[b] PANAROLI *Penec.* 2. *Obſ.* 42. HILDANI *Cent.* I. *Obſ.* 84. & 85. *ac de morſu Equi*, ibid. *Cent.* II. *Obſ.* 86. SEREN SUMMONIC. *Cap. De hominis & ſimiæ morſu.*

[c] In ſeveral Editions of CELSUS you will find FERÆ for FERE, *Omnis autem* FERÆ *morſus quoddam virus habet*; but I think the other Reading preferable to this; for CELSUS does not treat in this Place of the Bites of wild Beaſts alone, for they are very uncommon Caſes, but of the Bites of a Man, an Ape, and particularly of a Dog (which Animals he manifeſtly diſtinguiſhes in this Place from wild Beaſts), which Bites he deſcribes as bringing on violent Miſchiefs, eſpecially if the Animal is much enraged: therefore CELSUS very properly, in an extenſive Senſe, declares, *Omnem* FERE *Morſum habere quoddam Virus, ſive Venenum*; which Opinion is not applicable to wild Beaſts alone, but to all Animals whatever, for Reaſons which we ſhall preſently lay down: MORGAGNI is of the ſame Opinion with me concerning the Interpretation of this Paſſage, which he explains, according to his uſual Cuſtom, with great Learning and Perſpicuity. *In Epiſtol. in* CELSUM, *ſag.* 126.

Fingers,

Fingers, ſucking it with your Mouth, or by the Application of Cupping-glaſſes, or by enlarging the Wound with a Lancet: waſh it afterwards with warm Wine, or camphorated Spirits of Wine, and apply Bolſters to it dipped in the ame Liquor, repeating it every three or four Hours till all Danger of Inflammation is gone off. CELSUS recommends Salt, as the beſt Remedy for the Bite of a Dog, if it is applied dry, and well rubbed in; it is afterwards to be healed with the Oil of Turpentine, or ſome vulnerary Balſam. If the Wound is very conſiderable, it will be abſolutely neceſſary to enlarge it with the Knife, unleſs the Opening is already very large; the Diſcharge of Blood alſo ſhould be encouraged in this Caſe, by the ſame Methods which we adviſed in the foregoing; and you ſhould open a Vein to prevent a dangerous Inflammation. I lately ſaw the bad Effects of a Neglect of this Practice, in the Caſe of a Boy, who was bit by a Dog near the Knee, and was ſeized with a violent Inflammation over the whole Leg and Thigh, for Want of a proper Evacuation of Blood. The Wound ſhould be diligently waſhed with Wine, warm Spirits of Wine, or ſalt Water, dreſſing it up with Lint and linen Bolſters wet with the ſame Liquors: theſe Dreſſings are to be repeated frequently every Day, to prevent a violent Inflammation: you may dreſs afterwards with Honey, or a digeſtive Ointment, and heal with a vulnerary Balſam, as in other Wounds.

How to know a mad Dog.

V. In order to know whether your Patient has been bit by a mad Dog, it is neceſſary that we ſhould firſt ſettle the Marks by which a mad Dog is diſtinguiſhable from other Dogs: when a Dog is mad, he foams at the Mouth, and lolls out his Tongue, claps his Tail betwixt his Legs, and runs up and down without ceaſing, as if he was purſued; he makes a hoarſe Noiſe when he barks, and is afraid of all Animals that come in his Way, ſnapping at every thing he meets, even at his own Maſter, upon whom he uſed to fawn: other Dogs are afraid of him, and avoid him.

Miſchiefs that enſue from the Bite of a mad Dog.

VI. Men that are bit by a mad Dog are uſually afflicted with grievous Diſorders, ſometimes ſooner, ſometimes later, in Proportion to the Malignity of the Poiſon that is imbibed by the Wound, and to the Patient's Habit of Body at the Time he receives the Bite: when once the Poiſon begins to exert itſelf, the Patient is ſeized with great Anguiſh, continual Groanings, Sighing, acute Pains, and Fevers.

Prognoſis.

VII. If nothing is done to relieve this Diſorder, he is ſeized with a *Hydrophobia* about the ninth Day, a miſerable Circumſtance! ſince he is continually afflicted with Thirſt, and at the ſame time labours under ſuch a Dread of all Fluids, that he durſt not ſatisfy it, but rages and foams like a Dog, till, being quite ſpent, he [a] expires: therefore, in this Caſe, it well behoves us to be early in our Applications to Wounds of this Kind; for, when the *Hydrophobia* appears, nothing is to be looked for but certain Death [b].

How the Wound is to be treated.

VIII. Where ſhall we find a Remedy for this dreadful Diſorder? Many are of Opinion, that to puſh a Man unawares into a Pond or River is a certain Cure;

[a] There have been ſeveral Inſtances where the Poiſon has lain dormant in the Blood for one, two, nay for ſeveral Years, and has at length broke out, and carried off the Patient after the uſual Manner; WEBSTER has given us ſeveral ſurpriſing Relations of this Kind in his Book *De Magiâ*.

[b] It will be worth your while to conſult VERDRIES upon this Subject, *in Lib. De æquilibrio ment. & corpor. circa finem*; and MARESCOTTUS, *De Variolis, pag.* 57. where he treats of the *Hydrophobia*.

this was a common Remedy in the Times of CELSUS, *Lib.* V. *Chap.* 27: Some think it ſufficient to bathe the wounded Limb frequently in cold Water for ſeveral Days together, and to dreſs the Wound with ſome of the mad Dog's Hair: they imagine this laſt Method cures a Man by *Sympathy*, as they affect to call it: others, as CELSUS adviſes, throw the Patient into Water when the *Hydrophobia* is coming on, and endeavour to force him to drink againſt his Will; for by this Means, they aſſert, that they take off his Thirſt, and the Dread of Water at the ſame Time: on the other Hand, almoſt all the moſt experienced Surgeons recommend the following Method, as the ſafeſt and moſt worthy to be tried: to enlarge the Wound with the Knife, to promote the Flux of Blood, to cleanſe it *cum Aquâ ſalſâ, vel cum Aceto, vel Spiritu Vini & Theriaca*, and to endeavour to draw out the Poiſon by Cupping-glaſſes: and, laſtly, if the Texture of the Part will permit it, that is, when only the common Integuments or fleſhy Parts are wounded, they apply the actual Cautery to the Wound, and dreſs it afterwards like other Burns: but if the Part affected will not admit of the Cautery, the Wound muſt be enlarged by Inciſion, then dreſſed with Vinegar and Treacle, and covered with a Bolſter dipped in the ſame Ingredients: and to prevent Inflammations and a conſequent Fever, a Vein ſhould be opened (eſpecially in plethoric Habits) agreeable to the Practice of the Antients. *See* CELSUS, *Lib.* V. *Cap.* 27. *N.* 2. AQUAPENDENS, *in Operat. Chirurg. pag.* 331. adviſes this Method to be uſed to all Wounds that are infected with Poiſon: but in theſe laſt Caſes you ſhould firſt diligently enquire, whether the Arrow, or other Weapon, by which the Wound was inflicted, was poiſoned or not, or whether the Violence of the Symptoms give you ſufficient Reaſon for ſuch Conjecture; for where it remains doubtful whether the Weapon was poiſoned or not, you ſhould deal more tenderly with your Patient, and not proceed to the Uſe of the actual Cautery, but treat the Wound after the Method juſt deſcribed.

Kæmpfer's Method of curing the Bites of Serpents.

IX. KOEMPFER, who was one of the chief Phyſicians in the *Eaſtern* Countries, and well verſed in the Nature of the venomous Serpents, with which that Part of the World abounds, tells us in his *Amœnitat. Exotic. pag.* 581. and in his *Itiner. in Chinam & Japan*, that he has frequently cured the Bites of theſe Animals without the Help of the Cautery, by making a Ligature upon the Limb above the injured Part, and ſcarifying the Wound, anointing it well afterwards *cum Theriaca*, and covering it with a Cataplaſm made of the ſame Medicine, giving alſo a Doſe of it frequently by way of Sudorific: he declares, that he never loſt a Patient, where he had an Opportunity to treat him in this Method. As this is a ſimple, eaſy Method, and proves by Experience to be a very ſafe one, I ſee no Reaſon why we ſhould not prefer it to one attended with great Cruelty and Pain in the Operation; yet this Method may be found leſs effectual in curing the Bites of *European* Serpents.

Other Methods.

X. Some anoint the Wound with the *Oleum Nucis Moſchatæ* inſtead of the *Theriaca:* others apply *a Toad* to the Part, either alive, or dried and ſoftened with Vinegar, imagining that this Animal has a ſpecific Virtue in extracting Poiſon from a Wound: others again are extremely fond of the *Ophites*, or ſerpentine Stone, called *Pedro del Cobra*, which they are told is found in ſome Species of Serpents in the *Indies:* they affirm, that if you lay this Stone upon a Wound made by the Teeth of a venomous Serpent, or Viper, it will imbibe all the Poiſon, and if you afterwards ſoak it in Milk, it will depoſit it in that Fluid: com-

pare with this Place, KOEMPFER, *in Amœnitat. Exotic. pag.* 57, *& ſeq.* though he only adviſes it *contra Serpentum Ictus*; but the celebrated VALLISNERIUS, in his Book *De Generatione*, *p.* 141. denies that it is equal to the Cure of a Bite from an *Italian* Viper; therefore, I think, very little Credit is to be given to it. The following Cataplaſm is in great Reputation with ſome for this Intention, ℞. *Cepæ ſub Cineribus aſſatæ, & Allii Bulbum unum, Theriacæ, Fermenti Panis Valentiſſimi ana* ʒ j. *Sinapi* ʒ ß *quæ ſingula infuſo Aceto calido in formam Cataplaſmatis probe conteruntur, Vulnerique ſuperimponuntur:* Dr. MEAD recommends Viper Fat as a certain Remedy for the Bite of a Viper: and ſince that, Olive Oil has been in the higheſt Eſteem.

The reſt of the Cure. XI. In a Day or two after your Patient has been bit by a mad Dog, the Wound ſhould be dreſſed *cum Melle vel Unguento digeſtivo admiſt. Ung. Ægyptiaco, vel Mercurio præcipitato rubro bis quotidie:* it may be kept open with theſe Dreſſings for ſome Weeks, or for about forty Days, till the Poiſon is thoroughly diſcharged. You ſhould always be very careful not to heal Wounds of this Kind too ſoon, eſpecially where they have not been cauteriſed; for the principal Part of the Cure in theſe Wounds conſiſts in keeping the Part open, and encouraging a Diſcharge: wherefore CELSUS always recommends very ſtimulating Medicines.

The internal Method. XII. Beſides the external Remedies that we have adviſed, it will be proper to preſcribe ſtrengthening Medicines and Sudorifics to be given internally, according to the Strength of the Patient: ſome of the Antients, according to CELSUS, put the Patient into a warm Bath, and ſweated him there as long as he could bear it, with the Wound uncovered, that the Poiſon might diſtil out in greater Quantities, waſhing it well afterwards with Wine, which is an Enemy to all Poiſons; when they had repeated this Proceſs for three Days, they thought him out of all Danger: it would be very convenient in this Caſe to give him now and then a Glaſs of Wine inwardly, and a Spoonful or two of good Wine-Vinegar, in which ſome Sage had been boiled, with a Drachm of *Theriaca* in it; and between whiles to adminiſter Draughts of *Infuſum Scordii vel Salviæ in Aquâ calidâ*, putting the Patient into a warm Bed, or into a Bath, to encourage him to ſweat largely: this ſhould be done for ſeveral Days ſucceſſively. You may give, for ſeveral Mornings, *Valerianæ Radicis* ʒ j. in the room of *Theriaca*, which I find is much the Practice in *Italy*; or *Radix Gentianæ*, in the ſame Quantity, with a Draught of one of the Infuſions which we juſt now preſcribed: ſome, after the Example of GALEN and BOYLE, inſtead of *Theriaca*, give *Sal volatile Viperarum, vel ex Cancro Fluviatili combuſto paratum*, which they have ſo great an Opinion of, that they venture to affirm it to be *an infallible Specific* in this Caſe.

XIII. Several amongſt the Moderns recommend the *Scarabæus Maialis Melle conditus & tritus, vel Scarabæi Succus*, which they ſuppoſe to have very great Efficacy in deſtroying Poiſon, and preventing its bad Effects, if it be repeated for ſome Days: others have no leſs Opinion of the Virtues of the Heart, Liver, or Brain of a mad Dog or Wolf, which they affirm to have very ſalutary Effects, if given to the Patient in Time; but for many Reaſons I think this by no means a juſtifiable Practice. PARÆUS directs *Garlic* to be given frequently; but I think the moderate Uſe of ſome generous Wine, and the Juice of Citrons and mild acid Fruits, or Wine-Vinegar, either ſimple, or mixed with Honey, will be

of

of great Service, not only in ſtrengthening the Patient, but in deſtroying the Poiſon.

XIV. The ſame Methods of Cure which we adviſed above, *N.* VIII. *and* XIII. will be ſerviceable againſt the Stings of *Scorpions*, or other venomous Animals: the Scorpion affords an eaſy Remedy againſt his own Sting; for ſome bruiſe him and lay him upon the Wound; others drink him in a Glaſs of Wine. *See* CELSUS, *Lib.* V. *Cap.* 27. *N.* 3. where he ſays, *Venenum Serpentis non Guſtu, ſed in Vulnere nocet.* Some dreſs the injured Part with Oil of Scorpions, which they eſteem a ſure Method of Cure: others do nothing but draw Blood from the Arm. The Antients in this Caſe hired Men to ſuck the Blood and Poiſon out of the Wound, which they did, ſpitting it out again, without injuring themſelves in the leaſt. See the above cited Paſſage from CELSUS: but the Patient at the ſame Time did not neglect the Uſe of the Methods which we preſcribed above, both with regard to internal and external Medicines and Applications. The beſt Cure for the *Sting of Bees* or *Waſps* is, *Acetum cum Theriaca*, or *Theriaca cum Spiritu Vini*, or *Bolus Armena cum Aceto*. The Method of curing a Gangrene ariſing from the Bite of a Horſe, may be ſeen in HILDANUS, *Cent.* ii. *Obſ.* 86.

INSTITUTIONS OF SURGERY.

PART II. BOOK II.

Of FRACTURES.

CHAP. I.

Of FRACTURES *in general.*

Fractures, what.

I. UNDER the Name of *Fracture*, ſpeaking in general Terms, we conceive every Solution of Continuity in the Bone, either from an internal or an external Cauſe, whether the external Cauſe was a ſharp or blunt Inſtrument: but as we uſually call thoſe Injuries of the Bone that are brought on by acute Inſtruments *Wounds of the Bone*, ſo we properly call thoſe *Fractures of the Bone*, where the Bone is broken by the Force of a blunt Inſtrument: therefore Fractures generally happen when any Part of the Body where a Bone is ſituated receives a violent Shock, either by a Fall, by Jumping, or a Blow with a Piece of Timber, a Stone, or by a Shot from a Gun: there are alſo Inſtances where this Accident has happened from an internal Diſorder, to wit, from the Scurvy, a Caries, or the venereal Diſeaſe, which have rendered the Subſtance of a Bone ſo brittle, that it has been fractured without any apparent external Accident. *See* CELSUS, *L.* viii. *C.* 1. MARCELL. DONATUS *Hiſt. Med. L.* iv. *C.* 5. *ex* PANDOLPHINO, *p.* 272. CONNOR, *Diſſ. Med. Phyſ. de ſtupendo Oſſium Coalitu, pag.* 11. *Fractura Oſſium a Cauſâ internâ mirabilis.* SAVIARD, *Obſ.* LXII. HEYNE *De Oſſium Morbis, N.* XXIX.

Different Species of Fractures.

II. We may diſtinguiſh Fractures into ſeveral Claſſes or Species: firſt, every Fracture is either *ſimple*, that is, when no other Parts beſide the Bone are injured; or *compound*, that is, when you have at the ſame Time a Wound, a Diſlocation, Hæmorrhage, Inflammation, Fever, Caries, or Contuſion of the Bone; or where the Bone appears to be fractured in ſeveral Places, or more than one Bone, at the ſame Time: other Differences ariſe with regard to the Situation of the Fracture: ſometimes it happens in the Cranium, Ribs, Clavicles, Vertebræ; ſometimes in the upper or lower Limbs; ſometimes in the Middle of the Bone; ſometimes in either of the Extremities: again, ſome

Fractures

Fractures are tranſverſe, others oblique; in which Caſe it frequently happens that the Points of the Bones wound the neighbouring Parts, puſhing quite through the Muſcles and common Integuments, or at leaſt pricking them grievouſly, and bringing on Pain, Inflammation, Tumor and Spaſms: violent Contuſions alſo may be claſſed under the Head of Fractures; for the Bones in this Caſe are frequently broke into Splinters, by the Falling of any heavy Body upon the Part, or by Fire-arms, or the Preſſure of Mill-wheels, or the Wheels of Carriages.

Of Fiſſures in the Bones.

III. To Fractures of the Bones we may properly add *Fiſſures*, when the Bones are divided either tranſverſely or longitudinally, not quite through, but cracked after the Manner of Glaſs, by any external Force; for although moſt Surgeons have looked upon the Mention of Fiſſures as an idle Jeſt, eſpecially of thoſe that are ſaid to be made in a longitudinal Direction, and others have paſſed them over ſilently in their Writings, or, where they have by Chance been mentioned, no Method of Cure has been directed for them; yet there is not one of them that I know of, who was ever able to demonſtrate the Impoſſibility of theſe Fiſſures; ſince they often happen in the *Cranium*, and indeed in other Bones: all they can pretend to alledge is, that they have never fallen under their Obſervation; but I find Inſtances of this Kind of Diſorder, with a Method of Cure deſcribed for it, in Authors of undoubted Credit: *ſee* HEYNE *De Morbis Oſſium, N.* XXIX. and particularly that famous *German* Surgeon FELIX WURTZIUS, *in Chirurg. Part* II. *Cap.* 28. which makes me ſo far from calling the Fact in Queſtion, that I think it ought rather to be a Spur to a young Surgeon to conſider well the Marks that WURTZIUS has deſcribed, and to make a more diligent Search after Caſes of this Kind, than has hitherto been made: we ſhall ſpeak more largely to this below.

How Fractures are to be diſcovered.

IV. It is no difficult Matter to examine *Fractures of the Bones*, 1. *by the Eye*, when the Pieces appear through the Skin, when the injured Part is apparently ſhorter than the ſound, or when you ſee that the Patient cannot make Uſe of it: 2. *By the Touch*, when you perceive a præternatural Inequality of the Bone, or that it bends in a Part where Nature never intended it ſhould; and here, by the way, we muſt recommend it to the Surgeon, if it be poſſible, to fix the Patient immediately, at the firſt Searching of the Fracture, where he is to lie during the Courſe of the Cure: 3. *by the Ear*, when we hear the Ends of the broken Bones cruſh againſt each other upon moving the Limb: but, 4. we may ſtrongly ſuſpect a Fracture of the Part, when it has received a Blow with great Violence from a heavy Body: and, 5. we ſhould not neglect to obſerve, that the Parts are more ſubject to this Injury in Winter, than in Summer; laſtly, 6. ſometimes, particularly in Fractures that are made in a tranſverſe Direction, the broken Parts of the Bone will immediately of themſelves recover their natural Situation, and leave very little room to ſuſpect the Diſorder: therefore it is neceſſary to be very cautious and prudent in forming your Judgment in Caſes of this Kind. If your Patient has entirely loſt the Power of moving any Limb, or puts it in Action with the greateſt Difficulty, after having received a violent Blow upon that Part; or if he feels violent Pain when you handle it, or move it for him, this affords great Reaſon to ſuſpect a Fracture: but to make yourſelf more certain in this Caſe, it will be proper to take hold of the injured Limb with both your Hands, and ordering an Aſſiſtant at the

ſame Time to move it about, attend diligently whether you cannot hear the broken Ends of the Bone rub againſt each other, and obſerve whether you cannot diſcover a præternatural Dent or Sinking in any Part of the Limb: the Motion that your Aſſiſtant makes ſhould be done with great Care and Tenderneſs.

How to diſcover Fiſſures.

V. *Fiſſures in the Bones* are not eaſily detected; ſince neither your Senſes of Seeing, Feeling, or Hearing can give you Light enough to determine any thing with Certainty in this Caſe: and this ſeems to be the Reaſon why moſt Surgeons are deceived in this Caſe, as GOVEIUS well obſerves in his *Chirurgie veritable, pag.* 79. If we will believe thoſe Authors, who declare to us, upon their own Experience, that theſe Caſes ſometimes happen, we ſhall find there are ſufficient Signs to diſcover a Fiſſure of the Bone: they always ſuppoſe a Fiſſure when you have ſuch violent Pains, after any external Violence, that the injured Part will not bear Handling, and cannot ſupport the Parts above it; when you have more than ordinary Tumor, and theſe Symptoms do not yield to the uſual Applications: after this you are to expect violent Inflammations, Suppurations, Fiſtulæ, and Caries. Theſe Authors are of Opinion, that aged Perſons are moſt ſubject to this Diſorder, becauſe their Bones are very brittle. When we conſider the Nature of a Fiſſure, we ſhall not be long in gueſſing whence all the bad Symptoms attending it can ariſe: for the Bone being once cracked, the Blood and Sanies which fill up the Vacancy will preſently putrify and corrupt the Medulla, the neighbouring Parts, and at laſt the Bone itſelf, which will eaſily produce the Miſchiefs we have deſcribed.

Diſorders attending a fractured Bone.

VI. Great Variety of Miſchiefs attend a fractured Bone; which differ, 1. with regard to the injured Part, and the Nature and Diſpoſition of the neighbouring Parts: 2. with regard to the Manner in which the Fracture is made; for oblique Fractures, and thoſe whoſe Splinters or Points wound and vellicate the neighbouring Parts, are much more painful and dangerous than tranſverſe Fractures. Fiſſures are attended with more or leſs Danger in Proportion to the Size, as appears from what we have delivered above: but, 3. we may judge of the Miſchief that is likely to attend a Fracture, from the Number of Pieces into which the Bone is broken: and, 4. by obſerving whether the Fracture happens in the Middle of the Bone, or at either of its Extremities. The principal Inconveniencies that attend a Fracture, are theſe; the Patient loſes the Uſe of the Limb, the lower Part of the Limb will be contracted by the Muſcles, which will make it appear diſtorted and deformed; the Laceration of the *Periоſteum*, and the Veſſels of the *Medulla* bring on great Danger of Fiſtulæ and Caries. When the Nerves are pricked and irritated by Splinters or Points of the broken Bone, the Patient ſuffers great Pain, Convulſions, Inflammation, and Fever: if any Veſſels ſuffer Preſſure, the Circulation of the Blood is retarded: therefore no wonder if Inflammations, Abceſſes, Gangrene, and Death, are the Conſequences. If the Preſſure is upon a Nerve, the Part to which it was determined becomes paralytic, loſing both Senſe and Motion, and by Degrees uſually waſtes: ſometimes, whilſt the Bone is uniting, the broken Parts are ſupplied in too plentiful a Manner with Juices, and the Callus is formed irregularly, which occaſions Deformity in the Limb. When you have a Wound in the fleſhy Parts in Conjunction with a Fracture in the Bone, you will moſt likely be troubled with a violent Hæmorrhage: when the Blood-veſſels are injured, the Skin remaining ſound and unhurt, great Effuſions of Blood, and dange-

dangerous Swelling and Suppurations, are the general and almoſt inevitable Conſequence.

VII. The Surgeon ought to be very cautious in delivering his Prognoſtic concerning Fractures: he ſhould avoid being too haſty in promiſing a quick, eaſy, and certain Cure, leſt his Art ſhould be overcome by accidental Diſorders, and he be accuſed of Knavery or Ignorance; for I don't know how it happens, the moſt unſkilful Perſons in Surgery ſpeak of fractured Bones, as Caſes of the leaſt Importance, and make nothing of promiſing a Cure: whereas it is undoubtedly true, that it is ſometimes impoſſible to reſtore a broken Limb to its former Shape and Strength, though your Surgeon is perfectly Maſter of his Art: therefore, ſince Fractures are ſometimes cured eaſily, but at other Times are attended with the worſt of Conſequences, it will be an Argument of Diſcretion in a Surgeon to deliver his Prognoſtic in ſuch a Manner, that it may not regard the fractured Part alone, but may give Warning alſo of the Accidents that are likely to happen to the neighbouring Parts, or which may be occaſioned by the Age, or the bad Habit of Body of the Patient, or by any other Circumſtances: and in this he ſhould always take Care not to be over haſty. Prognoſis in general.

VIII. I would recommend the following Obſervations to the Surgeon: 1. Simple Fractures, when you are called ſoon after the Accident, are much more eaſily cured than Fractures that are complicate with an external Wound, a Diſlocation, a great Contuſion, an Hæmorrhage, a Caries, or with any other grievous Diſorders: 2. Fractures are more eaſy or difficult of Cure, according to the Part on which they happen: thus ſmall Bones, ſuch as the Clavicles or Ribs, or the Bones of the Fingers, are uſually cured in twenty Days; the Radius, or Cubitus, in thirty; whereas the Os Humeri, or the Tibia, require from thirty to fifty Days: 3. Men of good Conſtitutions, and in the Prime of Life, are cured ſooner, with leſs Trouble, than Perſons of a bad Habit of Body, or advanced in Years. In particular.

IX. Where the Situation of the Bone is not altered by the Fracture, or the broken Parts ſtart very little, they are much eaſier replaced, than where they are entirely ſeparated from each other, and a great Space intervenes between them: tranſverſe Fractures admit of an eaſier Cure than oblique Ones: Fractures near the Articulations are attended with worſe Conſequences than thoſe which are made above the Middle of the Bone: for where the Fracture happens near either Extremity of the Bone, the Joint frequently ſuffers, which occaſions Loſs of Motion in the Part; the Ligaments alſo and Tendons are uſually bruiſed in this Caſe: from whence ariſe violent Pains, Inflammations, and Convulſions, and ſometimes even Gangrene, and Death itſelf; or at beſt, the Patient muſt ſubmit to an Amputation.

X. When two Bones of the ſame Limb are fractured, the Cure is more difficult than when this Accident happens only to one of them: when the Bone is broken into ſeveral Pieces, the Patient will ſeldom eſcape Abceſſes, or even Gangrene or Sphacelus; that either the Limb muſt be taken off, or Death will enſue: at leaſt the Cure will require a great deal of Time, and the Limb will never entirely recover its Shape: therefore, where a Surgeon ſees this, he ought always to forewarn the Patient, or his Relations, what Danger he apprehends.

XI. Where the broken Bones are inſtantly reduced, your Cure will be performed with greater Eaſe, than where they have been for ſome Time ſeparated: therefore

therefore where the Surgeon is called a conſiderable Time after the Fracture has been made, he cannot promiſe to reduce the Bones eaſily, or to make a ſpeedy Cure.

XII. When any Parts of great Conſequence to the animal Œconomy are ſituated in the Neighbourhood of the Fracture, the Caſe will certainly be attended with great Danger, if not with Death: ſuch are Fractures of the Cranium, from the Vicinity of the Brain; of the Vertebræ, from the Medulla Spinalis; of the Ribs or Sternum, the Oſſa Ilei and Pubis, from the Situation of the Viſcera of the Thorax, and Abdomen: Fractures alſo of the Bones, to which the larger Arteries or Veins are connected, are very dangerous; more particularly when any Splinter or Point of the broken Bone vellicates or wounds a large Veſſel; for very violent, if not mortal Hæmorrhages muſt neceſſarily enſue, eſpecially when this happens in the Axilla or Groin, which is often the Caſe.

XIII. When the Ends of the fractured Bone break through the Muſcles and common Integuments, you will find great Difficulty in reducing the Bone to its proper Situation, from the great Number of Muſcles, Nerves, and Blood-veſſels that lie in the Way; the Laceration of which will bring on great Miſchiefs, and frequently Deformity and Weakneſs upon the Limbs, eſpecially if it is the Os Humeri, Tibia, or Femur, ſo as to render the Amputation of it neceſſary.

XIV. The moſt temperate Air and Seaſon of the Year is moſt convenient for the Cure of this, as well as all other Diſorders: the Cure alſo ſucceeds more happily in Children and young Perſons than with aged Perſons: when Fractures happen to big-bellied Women, they are ſeldom cured till they have got rid of their Burthen.

XV. When the Bone is broken into Fragments, the Conſequences are generally Inflammations, Suppurations, or Fiſtulæ, which will not admit of any Remedy till the Splinters are all removed. If the Fracture is occaſioned by an internal Diſorder, ſuch as a Caries of the Bone, you will find it much more difficult to cure, than when it proceeds from any external Violence: nay, it is frequently an incurable Caſe, unleſs the Occaſion of it, to wit, a ſcorbutic or dropſical Habit of Body, or a venereal Taint, be removed.

XVI. When a large Piece of Bone is driven away by a Piſtol or Muſquet Ball, it is better to cut off the lower Part of the Limb, ſince the two Ends of the Bone are never likely to unite, than to deceive the Patient with the fruitleſs Hopes of a Cure, and weaken him to the laſt Degree, with the Attempt[a]: but when only a ſmall Piece of the Bone is carried off in this Manner, you may ſafely enough attempt the Union of the Parts, but the Limb will be ever ſhorter than the other; and if the Injury is in the Foot, he will be always lame.

XVII. When the Blood inſinuates itſelf through a Fiſſure into the internal Part of the Bone, by corrupting there, it produces a Caries, or Spina Ventoſa, incurable Fiſtula, Tabes, and Sphacelus, which always require Amputation of the Limb, and frequently deſtroy the Patient: the ſame Accidents will happen in Fractures of any Kind, when the extravaſated Blood mixes with the Medulla, and corrupts it.

XVIII. Fractures of the lower Limbs are much more inconvenient than thoſe of the Arm: though Diſorders of the upper Limbs are eaſier concealed,

[a] HORSTIUS, in *Obſerv. Med. P.* II. *Lib.* IV. *Obſ.* 10. gives us an Account of a Man who ſuffered a Loſs of Subſtance in the Bone of his Foot, of the Size of three Fingers Breadth; but he cured his Patient without leaving any Lameneſs: if the Story is true, it is very extraordinary.

whereas

whereas thoſe of the lower Limbs appear preſently, eſpecially in Men, from the Lameneſs and Deformity which they occaſion, which require great Care in the Treatment of them: on the other Hand, the Deformity is more conſpicuous in the Arms of Women; in theſe Fractures therefore, more Care is required, and a nicer Operation, that the Callus may be formed as ſmooth as poſſible: but where there is a Fracture of the Bones from Gun-ſhot Wounds, eſpecially on the Joints, as the Tarſus, the Knee, Elbow, Shoulder, or Thigh, it is attended with great Danger, and often with Death, if not prevented by immediate Amputation.

Cure of FRACTURES.

XIX. The Surgeon's principal Care in Fractures is to unite the broken Bone, to which three Things are neceſſary: 1. that the Bone be reſtored to its natural Situation, which is to be done by extending it and replacing it: 2. that after the Bone has recovered its natural Situation, it be kept there by giving it Reſt, and applying proper Bandages: laſtly, 3. you are to uſe proper Means to prevent, or remedy, the Diſorders that uſually attend this Accident. The Knowledge of Anatomy is neceſſary to perform theſe Intentions; for, 1. the Surgeon muſt be acquainted with the Situation and Structure of the Bones, that he may know whether the injured Limb is ſupported by one or more Bones, whether they are large or ſmall, whether they are firm or ſpongy, whether they are even or uneven, whether one or more Bones are broken at the ſame Time: 2. what Muſcles there are in the Neighbourhood of the Bone, their Situation and Office: laſtly, whether any conſiderable Nerves or Blood-veſſels are near the fractured Part: all which Things are abſolutely neceſſary to be known by any one, who expects to ſucceed in the Cure of theſe Diſorders. Cure.

XX. When the fractured Bones maintain their natural Situation, you are under no Neceſſity of extending or replacing the Limb, but of applying a proper Bandage: but when the fractured Parts recede from each other, ſome Degree of Extenſion is neceſſary, which muſt be always ſuited to the Diſtortion of the Limb. The greater Diſtance there is between the Extremities of the divided Parts of the Bone, ſo much ſhorter will the Limb be, from the Contraction of the Muſcles; therefore the Extenſion in this Caſe ought to be in Proportion ſo much the greater; but to prevent the Patient from ſuffering any Violence, every Thing ought to be done tenderly, and with great Care. Of Extenſion.

XXI. The Extenſion of fractured Limbs ought to be performed in the following Manner: 1. the Patient is to be kept firm and ſteady; the Poſture of Body, to be obſerved at this Time, differs according to the Circumſtances of the Caſe; ſometimes the Patient ſhould ſit, either upon a Stool, or upon the Floor; ſometimes it will be better for him to lie upon a Table or a Bed: 2. an Aſſiſtant ſhould ſupport the Limb with his Hand, both above and below the fractured Part: 3. the Aſſiſtant who holds the lower Parts of the Limb ſhould extend it ſtrongly, till you can replace the fractured Part of the Bone; if his Hands alone are not ſufficient to make the required Extenſion, he muſt uſe a Cord, or rather a Napkin; if one Man has not Strength enough for this Office, you muſt employ two or more: you muſt be very careful not to How the Extenſion is to be made.

uſe too great Roughneſs in this Operation, leſt you ſhould give your Patient unneceſſary Pain.

Means uſed by the Antients for Extenſion.

XXII. The Surgeons amongſt the Antients, when they found that neither Hands nor Napkins were ſufficient to make a proper Extenſion, (which was indeed a very rare Caſe) contrived ſeveral mechanical Inſtruments to anſwer this End: for this Purpoſe you will ſee ſeveral Pullies with Ropes deſcribed; the *Scamnum Hippocratis*, and ſeveral Machines of this Kind, which you will find in the Works of the principal Surgeons, ſuch as ORIBASIUS, PARÆUS, ANDREAS A CRUCE, SCULTETUS, and others: but if we attend to the Obſervations of ſeveral modern Surgeons, which are made with great Accuracy, we ſhall find that Machines of this Kind do not act ſufficiently equal in all Parts at the ſame Time, and that you will find great Difficulty in applying them: beſides, they are not always at hand in Times of War, and upon many other Occaſions: therefore it is no Wonder that you ſcarce ever ſee or hear of theſe Inſtruments amongſt the Surgeons of the preſent Times; eſpecially ſince you will always find your Hands or the Napkin ſufficient for any Extenſion that can be required.

What is to be done where you have Tumor and Inflammation.

XXIII. There remains one Obſervation to be made with relation to the Extenſion of the Limb: when the Surgeon is called at ſome Diſtance of Time from the Accident, when a Tumor and Inflammation are come on, it is beſt to defer the Extenſion of the Parts till theſe Symptoms are removed: for it is impoſſible to make a proper Extenſion whilſt the Parts are affected in this Manner, without bringing on the moſt acute Pains, Convulſions, and Danger of Sphacelus: but if the Symptoms of this Kind appear but in a ſmall Degree, it is better to attempt the Extenſion of the Parts inſtantly, before the Inflammation increaſes.

How the Inflammation is to be treated.

XXIV. Where the Inflammation is already arrived at ſo great a Height as to forbid the Extenſion of the Parts, the Surgeon's principal Care ſhould be directed to aſſuage this Symptom: the ſame Methods which we propoſed for diſperſing Contuſions, (*Chap.* XV, *Book* I, *N.* X, *&c.*) Blood-letting, looſening the Bowels, adviſing the Patient to drink large Quantities of aqueous Fluids, preſcribing ſuch internal Medicines as are known to abate Inflammations, and fomenting the Parts with warm diſperſing Fomentations, will anſwer this Intention: theſe Applications will uſually remove the Inflammation in four and twenty Hours, in ſuch a Manner that you may ſafely undertake the Extenſion of the Limb: inſtead of the foregoing Fomentations you may uſe the following, which very powerfully anſwers the Intention it is preſcribed for: ℞ *Herb. Scordii M* ij. *vel* iij. *Aq. ſimplicis* ℔ j. *Spiritus Vini* ℥ vij. *quæ ſimul per Horæ quadrantem probè decoct. admixtiſque Salis Culinaris* ℥ j. *it. Nitri* ℥ ß *identidem cum Linimentis calidè ſupra fractum membrum deligantur:* where the Inflammation is ſo violent, that it will not yield ſufficiently in the Time above-mentioned, to admit of the Extenſion of the Limb, and the Patient is plethoric, you muſt open a Vein, and repeat theſe Applications till they take Place, and the Symptoms diſappear.

Of Splinters.

XXV. Sometimes you will be troubled with Splinters of the Bone in your Way, which vellicate and prick the neighbouring Parts, and will render the Reduction of the Bone very difficult: if the Splinters are free, and have no Connexion with the Bone, you muſt remove them carefully; if they hang by a Portion

tion of the Periosteum, divide them with your Scissars, for you will never find that they will easily unite again with the rest of the Bone, but will always give you great Uneasiness and Trouble in your Cure: if the Splinters adhere to the neighbouring Parts, and do not much impede the Reduction of the Bone, it will be best to replace the Bone, and to leave the Splinters either to loosen and come away by the Suppuration of the Parts, when they may be taken out without giving the Patient great Pain; or sometimes they will grow again to the rest of the Bone: when they adhere very firmly to the principal Parts of the Bone, we should be so far from attempting to extract them by Force, that we should endeavour to replace them with the greatest Exactness: when this is performed with Accuracy, they will frequently unite to the rest of the Bones: but where that is not to be expected, we must get them out by Degrees in the best Manner we can.

Of Splinters sticking through the Flesh.

XXVI. Where Points of the broken Bones or Splinters stick so far out, that they are an Hindrance to the Reduction of the Bone, you should diligently consider whether you can by any Means contrive their Re-union to the Bone; which you may judge of by observing at what Distance they are removed from some large Bone, and whether there is a large Quantity of Flesh intervening: where they cannot be reduced or re-united to the Bone, they may be removed by a Pair of strong-pointed Forceps, *See Plate* VIII, *Fig.* 1: or if they stick very firm you may use a fine Saw, *Plate* VII, *Fig.* 9: when you have removed the Splinters, you are in the next Place to make your Extension, and reduce the Bone. Till they are removed, the Reduction and Re-union of the Bone are generally impracticable.

Of Splinters that are concealed under the Skin.

XXVII. If the Splinters are concealed under the Skin, and you cannot lay hold on them with your Hands, you must first try if you can reduce them to their natural Situation: if this cannot be done, make an Incision through the Skin, and take them out.

Of Extension.

XXVIII. To make a proper Extension of the Limb, two Assistants should be employed, in the Manner we described above at *N.* XXI, and the Surgeon should take hold of the extended Part, and direct it with his Hands, sometimes a little outwards, sometimes a little inwards; now upwards, then downwards; putting it into different Positions, as the Circumstances of the Case shall require, till the Parts have recovered their natural Situation.

How to discover when the broken Bones are replaced.

XXIX. You may know that the Bones have regained their natural Situation, by the Remission or Absence of Pain, and by observing that the fractured Limb is of the same Figure and Length with the sound Limb: if these Signs of Recovery are wanting, you have good Reason to suppose that the Operation is as yet ineffectual, and the Extension is to be repeated or continued in the Manner we have described, till the Bone is replaced.

They are to be kept in that Situation.

XXX. The Bones being properly replaced, the next Thing to be done is to secure them in their Situation, that they may unite to the best Advantage.

How Fractures are to be bound up.

XXXI. Two Things are chiefly required to answer this End: 1. to bind it up properly; and, 2. to lay the Limb in a convenient Posture: the Apparatus for securing the Situation of the Limb is composed of *Bandages*, *Bolsters*, and *Splints*[a],

[a] The famous PETIT of *Paris* forbids the Use of *Splints*, and supplies their Place with *Bolsters*, which I think will by no Means answer.

which are to be made of thick Paper, of Wood, or, if the Surgeon ſhall think proper of thin Plates of Copper, Braſs, Steel, Tin, or Lead: *See Plate* VIII, *Fig.* 7: but I think the beſt are thoſe made of Wood or Paper. The Manner of Dreſſing the Limb is as follows: in the firſt Place, a Roller is to be paſſed round the fractured Limb; upon this are to be placed Bolſters, and over them Splints, which are to be ſecured by a tight Bandage over all. In ſome Caſes other Inſtruments are neceſſary, ſuch as Boxes made of Paſte-board, Wood, or Metal, to fix the fractured Limb in: *See Plate* IX, *Fig.* 9: other Inſtruments are alſo neceſſary in this Caſe; the particular Manner of applying which, to the Arms, Legs, and other Parts, we ſhall deſcribe below, when we treat particularly of Fractures; and there you will find that ſimple and compound Fractures require different Bandages: this Apparatus of Inſtruments is required only to ſecure the Bones in their Situation, and to forward their Union: it is no Wonder therefore that Fractures are ill cured, where the Surgeon is ignorant of the proper Methods of applying the Bandage, or the Patient is unruly, and will not give the Limb proper Reſt.

The Uſe of Plaſters in this Caſe.

XXXII. Although great Numbers of Surgeons at this Time make it their conſtant Practice to apply a Plaſter to the fractured Part of the Limb before they make the Bandage, yet the moſt prudent and ſkilful Surgeons amongſt the Moderns entirely reject Applications of this Kind, as not only uſeleſs, but injurious to the Patient: for theſe Plaſters can do no Service without the Bandage; but the Bandage alone, if it is dexterouſly made, is ſufficient to keep the Limb firm: and the Plaſter carries this Inconvenience with it, that it ſtops up the Pores of the Skin, and produces Tumors, and moſt violent Itchings: for my own Part, I am entirely of Opinion, that all Kinds of Fractures may be very happily cured without the Uſe of Plaſters, and I am confirmed in this Opinion by long Experience: but if, notwithſtanding this, any one ſhould be bigoted to the Uſe of Plaſters, I would adviſe him to be cautious not to make them of too great Length: they ſhould not entirely ſurround the Limb; but a Thumb's Breadth of it at leaſt ſhould be left bare, leſt the Blood ſhould be obſtructed in its Courſe, which would bring on Tumors, Gangrene, and Sphacelus.

Of the Dreſſings.

XXXIII. Before we treat more particularly of Fractures, it will be proper to ſay ſomething briefly of the Apparatus of Dreſſings required in Caſes of this Kind: and ſince the chief Help ſeems to be expected from *Bandages*, we ſhould principally contrive that, beſides having the general Properties of due Length and Breadth, they ſhould alſo be accurately adapted to the Shape of the broken Limb: in Fractures that are not attended with a Wound, you ſhould apply two ſingle-headed Bandages, each of which ſhould take its Beginning upon the injured Part; one aſcending, when it has gone thrice round the Limb, and the other deſcending in a contrary Direction, and then aſcending again: the ſame Windings and Circumvolutions may be made with a ſingle Bandage, if it be long enough; as will appear from our Treatiſe of Bandages.

The Bandage ſhould be neither too tight nor too looſe.

XXXIV. In order to keep the Parts in their natural Situation, the Bandage ſhould be made pretty firm: but if you tighten it too much, you will interrupt the Circulation of the Blood, and excite Tumors, Inflammation, and Gangrene: on the other Hand, if the Bandage is made too looſe, it will eaſily come off, and ſet the diſunited Parts at Liberty; the middle Way therefore is moſt eligible: you will diſcover the Mean between theſe two Extremities by obſerving a ſlight

a ſlight Degree of Tumor below the Bandage, after it has been applied for ſome Time: if the Tumor encreaſes to a violent Degree, you muſt looſen the Bandage; if the Parts do not enlarge at all, you muſt bind it ſtill tighter.

Of Bolſters and Splints.

XXXV. *Bolſters* and *Splints* are to be prepared in Proportion to the Size of the fractured Limb: where the Limb is of an unequal Size in different Parts of it, you muſt fold up the Bolſters in the Manner we have deſcribed at *Plate* IX, *Fig.* 13: you will by this Means be able to apply the Splints to greater Advantage: the Splints ſhould be tied on with three Tapes, the Middle of which is to be faſtened firſt, and then the others.

How the fractured Limb is to be placed.

XXXVI. In Fractures of the lower Arm, after you have applied you Dreſſings and Bandage, you may ſuſpend it in a *Scarf* or *Sling* (by the *French* called *Eſcarpe*) which is to hang from the Neck: *ſee Plate* XXXVIII, *Fig.* 17: in Fractures of the Leg you may reſt the Limb upon Pillows, *Plate* IX, *Fig.* 5, or in Boxes, *Plate* IX, *Fig.* 9, placing Cuſhions or Pillows under it: theſe Machines alſo are to be faſtened to the Limb with Tapes, that it may remain fixed and immoveable: ſome Surgeons faſten a Pillow under the Limb, after the Application of the Bandage, in Imitation of SOLINGIUS: *See the* Amſterdam *Edition, printed in* 1698, *Plate* XV, *Fig.* 9: others uſe wooden Boxes, ſuch as you will find deſcribed by SOLINGIUS and SCULTETUS: but the moſt prudent Surgeons prefer Cuſhions or Pillows; for this is not only more uſeful than any other Method, but it is alſo very handy and eaſy to come at: we uſe in this Place a Sort of a *Sole*, *Fig.* 6, made of thick Paper or Wood, which keeps the Foot ſteady: this ſhould be lined with a ſoft Bolſter, to keep it from galling or fretting the Foot: *ſee Fig.* 7: it is to be faſtened to the Pillows by the Tapes *a a a*, *Fig.* 6: a Piece of Linen, in the Shape of a Ring, is to be ſewed to the lower Part of this Bolſter, and faſtened on with the Strings, *b b*, *Fig.* 8: this is a Contrivance to ſuſpend the Heel, to prevent Inflammation, Pain, and other Miſchiefs that are frequently brought on by lying upon it too long. The two-headed Bandage has its Uſe in this Caſe; for the Heel may be put into this, and the two Heads of it being ſewed to it, will be kept faſt on: the two Heads of the Bandage are to be placed one under the internal, and the other under the external *Malleolus*, to prevent too great Stricture upon the *Tendo Achillis*, which would bring on acute Pains and Inflammations: you may make an Arch over the Foot with a Piece of Hoop, which will keep the Bed-cloaths from being troubleſome, and at the ſame Time not prevent the Application of warm Napkins, or Fomentations to the Part: *ſee Plate* IX, *Fig.* 10.

How the Patient is to be treated.

XXXVII. The Patient ſhould lie upon his Back, with his Head and the fractured Limb ſomewhat higher than the reſt of his Body; he ſhould have a Rope with a Handle at the End of it hung from his Bed's Teſter, that he may be able to take hold of it, and raiſe himſelf up when there is Occaſion: if he is of a plethoric Habit of Body, you will do well to bleed him in the Arm, to prevent Inflammation: the Surgeon ſhould be very frequent in his Viſits at the Beginning of this Diſorder, and very diligent in examining whether the Bandage, and other Applications, remain ſufficiently firm or not: if any Thing is out of Order, he is to correct it; the Regimen with regard to Diet ſhould be the ſame which we adviſed above, when we treated of Wounds: *ſee Chap.* I, § XLIII, *and the following:* he muſt not attempt to riſe on any Account, but be ſupplied with Bed-pans, *&c.*

XXXVIII. The-

When the Dreſſings are to be opened.

XXXVIII. The firſt Dreſſings ſhould be opened and renewed ſooner or later, in Proportion to the Nature and Number of the Symptoms that accompany the Fracture: when the Bandage remains ſufficiently tight, and no bad Symptom appears, you ſhould not looſen it, till the fifth or eighth Day: but where you have Inflammations, Tumors, Pains, and violent Itchings; or where the Bandage is too looſe or too tight, which is frequently the Caſe, you muſt inſtantly take off the Dreſſings, and change them: the ſecond and third Dreſſings muſt be performed in the ſame Manner with the firſt; with this only Difference, that at the third Dreſſing, if you perceive no Tumor, you may make the Bandage tighter than before, and by this Means prevent the luxurious Growth of the Callus, which would occaſion Deformity.

Cure of Fiſſures.

XXXIX. When you have Reaſon to judge, by the Symptoms related above at *N.* III, *and* V, that your Caſe is a *Fiſſure*, you may follow FELIX WURTZIUS's Inſtructions upon the Head: he always laid his [a] Plaſter, which he made Uſe of in Fractures, upon the diſordered Part, and upon that he placed Splints, and adviſed the Patient to reſt for ſome Days, and the Tumor will quickly diſappear: when you find the Tumor advanced in Size, and ſoft, it is a plain Indication that it contains a Fluid, which is to be let out by Inciſion: when you have evacuated the corrupted Fluids, you ſhould put a Tent into the Wound, dipt in the *Unguentum fuſcum Wurtzii*, uſing afterwards the Bandage which is applied to Fractures accompanied with a Wound: if we liſten to WURTZIUS, Ointments, Cataplaſms, Fomentations, and Baths, are of no Service in this Caſe, but are prejudicial: for collected Fluids putrifying, corrupt the neighbouring Parts, and the Bones, and bring on Caries, and other grievous Diſorders: the Symptoms that ariſe from Fiſſures are frequently attributed to Defluxions, or to the Gout: whoever deſires to be more fully informed of the Nature of this Caſe, I would adviſe him to conſult WURTZIUS, *Part* II, *Chap.* XXVIII, *p.* 381, *edit. Baſil. ann.* 1687: GOUEIUS aſſerts, that Fiſſures, when they are juſt made, may be cured by the Application of Bandages without the Aſſiſtance of other Remedies: *ſee Veritable Chirurgie, pag.* 86.

CHAP. II.

Of DISORDERS *accompanying* FRACTURES.

Of a Fracture accompanied with Wouud.

I. IF a Fracture is accompanied with a Wound, after you have reduced the fractured Bones, you muſt treat the Wound in the ſame Manner with other lacerated Wounds: firſt, the Wound is to be well cleanſed with warm Wine, Spirits of Wine, or ſalt Water; in the next Place, it is to be filled with dry Lint, to ſtop the Hæmorrhage; then to be dreſſed with digeſtive Ointment [b]; laſtly,

[a] The Plaſter is made in the following Manner: ℞ *Reſinæ pur. & candidæ* ℔ii. *Terebinthin. vulg.* ℔ſſ. *leni igne liquefiant, injectoque demum Radic. Ulmariæ Pulv.* ℥iv. *tantiſper bene ſubigantur, donec modicè frigeant:* when you have a Mind to ſpread it upon Linen or Leather, throw it into hot Water: the Author is very high in his Commendation of this Plaſter at *pag.* 320, of his Surgery.

[b] If any Hæmorrhage ſhould happen, you muſt obſerve the Methods we deſcribed when we were treating of Wounds.

it is to be anointed with ſome vulnerary Balſam till it is thoroughly healed. Since it is neceſſary to open the Dreſſings every Day, in order to cleanſe the Wound, but at the ſame Time it would be of very bad Conſequence to move the Limb, therefore a great Length of Bandage in this Caſe would be very wrong, eſpecially in the *Femur* or *Tibia:* for it would be troubleſome to lift the Foot up to roll on a long Bandage, which would diſturb the fractured Bones, and throw them out of their natural Situation: for this Reaſon the beſt Surgeons neglect the Uſe of long Bandages in this Caſe, and apply the Bandage of eighteen Heads, *Plate* IX, *Fig.* 4. which may be looſened at Pleaſure: when the Wound is healed, which happens frequently before the Bones are united, you ſhould lay aſide the Bandage of eighteen Heads, and bind up the Limb with long narrow Rollers till the Cure is thoroughly perfected: but we ſhall explain this more largely below, when we come to treat profeſſedly of Bandages.

A Fracture attended with an Ulcer;

II. When a Fracture is attended with an Ulcer without a Caries, which frequently happens in the Leg or Thigh; it is to be dreſſed every Day, after the ſame Manner as we directed for a Wound in the ſame Circumſtances. Having firſt replaced the Bone, the Limb is to be bound up with the Bandage of eighteen Heads, till the Ulcer is healed: but when the Ulcer is healed, and the fractured Parts of the Bone not ſufficiently united, you muſt lay aſide the Uſe of the eighteen-headed Bandage, and apply long narrow Rollers, as we adviſed above for a Fracture attended with a Wound.

With a Caries.

III. Sometimes the Fracture happens upon a Part that has been long troubled with an Ulcer and Caries; this Caſe is very difficult of Cure, nay frequently it admits of no Cure at all: very few Writers in Surgery have laid down any Directions, by which we may be guided in this Caſe: PETIT indeed deſcribes the Caſe of a *fractured Tibia attended with a Caries:* but as he has related the Caſe of the Tibia alone, neglecting to deſcribe it as happening to other Parts, he has in my Opinion, by no means ſatisfied the Subject: however, this may ſerve as an Example to be imitated in ſimilar Caſes, till we ſhall be furniſhed with more perfect ones: a young Man, who had been for ſome time troubled with an Ulcer and Caries, about the Middle of the Tibia, had the Misfortune to break the Bone in the very Part, the Fibula remaining at the ſame time whole; therefore no Extenſion was required in the Cure of this Fracture: PETIT, in the firſt Place, took off all the vitiated Fleſh that was ſituated near the fractured Part, with his Knife, and reduced the Ends of the Bone into their proper Situation with his Fingers, and then filled up the Ulcer with dry Lint, and covered all with the eighteen-headed Bandage, as above: after ſome Days, when the Fever was quieted, he cauteriſed the Extremities of the fractured Bone that were affected with Caries, and afterwards took off the carious Parts with the Trepan, which the *French* call *Trepan exfoliatif:* Having done this, he applied Lint to the naked Bone, well ſaturated *cum Tinctura Aloës:* but he dreſſed the fleſhy Parts firſt *cum Unguento digeſtivo*, and afterwards *cum Unguento fuſco*, to keep down the Luxuriancy of the hard Fleſh, which is very prejudicial in this Caſe: and this Method of dreſſing he continued for fifty Days, till the diſordered Parts of the Bone ſeparated from the ſound: he then began to encourage the Growth of new Fleſh, by applying vulnerary Balſams, and healed both the Bone and Ulcer after the uſual Method.

IV. But

A fractured Thigh with a Caries.

IV. But the Caſe is attended with far greater Difficulties when the Fracture happens upon an ulcerated Part, attended with [a] Caries in the Thigh; which Caſe I find entirely neglected in PETIT's Book of Fractures: I knew a Student of about twenty Years of Age, who had been troubled for many Years with an Ulcer and Caries, in the middle and internal Part of his Thigh, near the Situation of the crural Artery: the Fleſh in this Part was ſo thick, that the Caries did not appear, and the Vicinity of the great Artery prevented us from enlarging the Ulcer with the Knife, or from cauteriſing the Bone, ſo that all the Medicines which were applied had no Effect; at length, as he was walking about, the Thigh broke in this diſordered Part, without the Aſſiſtance of any external Force: what ſhould we do now? we were prevented from enlarging the Wound, or cauteriſing the Bone, by the Reaſons I juſt mentioned; and though we replaced the Bone, and applied a proper Bandage, yet it would never unite, but the Patient dragged on a miſerable Life: therefore it is worth our ſerious Conſideration, what is the beſt Method of Cure for Fractures of this Kind, when they happen in the Thigh, Arm, or other Parts where the Bones be concealed and cannot be laid bare with Safety: but this is rather to be wiſhed for than expected.

Of Callus.

V. The Surgeon has done his Duty in the Treatment of a Fracture, when he has diligently replaced the Bones, and taken Care to preſerve them in that Situation: for Nature has provided for the reſt, by ſupplying the divided Parts with a Callus; a Sort of Gelly, or liquid viſcous Matter, that ſweats out from the ſmall Arteries and bony Fibres of the divided Parts, and fills up the Chinks or Cavities between them; this firſt appears glewy, then of a cartilaginous Subſtance, but at length becomes quite bony, and joins the fractured Parts ſo firmly together, that the Limb will often make greater Reſiſtance to any external Violence with this Part, than with thoſe which were never broken, in the ſame Manner as we frequently ſee it happens to Pieces of Wood well glewed.

Its Growth.

VI. But as the new Fleſh in Wounds will frequently ſprout out too faſt, ſo will the Callus in Fractures, and by this Means render the Limb uneven and deformed, eſpecially in Fractures attended with a Wound: where this is the Caſe, and you ſee plainly that you cannot prevent it, you had beſt inform your Patient of it in Time, leſt he ſhould blame his Surgeon as the Author of his Deformity: for it cannot always be prevented or remedied; nor can you take off the Luxuriancy of a Callus as you can of the Fleſh, for ſeveral Reaſons of Conſequence: therefore, when once it is formed, it remains without Cure.

How to prevent the Luxuriancy of the Callus.

VII. But ſome Meaſures may be taken to prevent the Callus from exceeding its due Bounds, by making the Bandage ſomewhat tighter than ordinary, and wetting it firſt with the beſt rectified Spirits of Wine: this will not only keep the viſcous Matter within its Bounds, but will alſo forward its Induration: which may be obſerved in the Tibiæ of Men and the Arms of Women, as thoſe Parts are more frequently expoſed to View: but in Fractures attended with a Wound, as they will not admit of too tight a Stricture; eſpecially if you uſe the eighteen-headed Bandage, it is extreme difficult to prevent a Luxuriancy of Callus; when once the Callus is indurated, we have no Medicine that will take it down or deſtroy it: nevertheleſs, there are ſome who pretend that it is

[a] *Lib. De Morb. Oſſ.* Tom. II. p. 270, &c.

to

to be diſperſed by the *Emplaſtrum de Ranis Vigon. cum Mercurio*, tying a Plate of Lead over it; the Callus grows ſometimes faſter, ſometimes ſlower, according to the Size of the fractured Bone, the Habit of the Patient's Body, the Temperament of the Air, and laſtly, in Proportion to the Patient's Age: when it comes on but ſlowly, ſome Surgeons place great Confidence in the Patient's taking *Oſteocolla*, half a Dram at a Doſe.

VIII. Violent Itching is beſt prevented by removing oily fat Remedies, and therefore the Plaſters themſelves, from the Limb; for they are compoſed of ſuch Particles that they ſtop up the inſenſible Pores of the Skin: if the Itching remains after the Removal of theſe Applications, you may waſh the Part with warm Wine, Oxycrate, or Spirit of Wine, covering it up with ſoft, fine Linen: if Bliſters riſe upon the Part, they ſhould be ſnipt with the Sciſſars. To prevent violent Itching.

IX. Inflammations are to be treated in the Manner we adviſed above in *Book*, I. *Chap.* XV. but to remove Pains and Convulſions, you ſhould diligently attend to what we laid down in deſcribing the Cure of Wounds; but above all, you ſhould be very accurate in replacing the fractured Bones, and in obſerving whether they maintain the Situation which you reſtored them to; and if you obſerve any Splinters quite free from the neighbouring Parts, you ſhould inſtantly remove them, and endeavour to lay the Limb in an eaſy Poſture: in theſe Circumſtances you ſhould not neglect to open a Vein, and to apply emollient and diſperſing Cataplaſms and Fomentations, preſcribing at the ſame time Medicines to be given internally with this Intention, and adviſing the Patient to obſerve a proper Regimen with regard to his Diet: without obſerving theſe Rules, violent Inflammations, Sphacelus, and Death itſelf, will frequently enſue. Of Inflammation, Pain, and Convulſion.

X. If the Inflammation is ſo violent as to threaten a *Gangrene* of the Part, you muſt bleed inſtantly, lay aſide the long narrow Bandages, and apply the Bandage of eighteen Heads, uſe Fomentations prepared *ex Aquâ Calcis & Spiritu Vini camphorato cum Eſſentiâ Aloes & Myrrhæ; vel ex Spiritu Vini camphorato & Sale Ammoniaco*, or the Remedies we recommended above, treating of Fractures, *C.* I. *N.* XXIV. and in the Chapter of *Contuſions*; but if the Part is already affected with *Gangrene*, you muſt make frequent and deep Scarifications, to ſet the ſtagnating Fluids at Liberty, not neglecting at the ſame time to apply externally the Fomentations we recommended above, and to give the Bark inwardly. When the *Gangrene* has penetrated ſo deep into the parts, that it is beyond the Reach of Fomentations, and begins to be ſphacelated, you muſt take off the Limb, to ſave the Life of the Patient. How a Gangrene is to be treated.

XI. If the Fracture is attended with a conſiderable Diſcharge of Blood, you ſhould diligently examine whether the Hæmorrhage proceeds from a Vein or an Artery; whether the Flux of Blood is to be ſtopped by Preſſure, by the Help of dry Lint, Bolſters, and Bandages; or by ſtyptic Medicines, or by making a Ligature upon the injured Veſſels; or laſtly, by the actual Cautery, as we have taught above at *Chap.* II. on *the Cure of Wounds*. After the Blood is ſtopped, the Bones are to be replaced, extraneous Bodies are to be removed, and the Limb bound up. Of Hæmorrhage.

XII. If a Relaxation of the Nerves, or Waſting of the Limb, ſucceed a Fracture, there are very little Hopes of Help: however it will be adviſeable, (1) to rub the Limb well with hot Cloths; (2) with ſpirituous Medicines, ſuch as Of Palſy and Waſting of the Limb.

Spirit. Formicar. Lubricor. Matricalis, C. C. Sal Ammoniac, Eſſentia Euphorbii, Caſtor, and others. (3) to foment the Limb with warm Fomentations and Baths made *ex Vino Herbiſque corroborantibus, aromaticis ac nervinis, vel Thermis naturalibus:* (4) laſtly, the beſt Remedy, in my Opinion, is, to wrap the tabid Limb up in the Skin of an Animal that is juſt killed, and remains in its natural Heat: for by this Means the Flux of the Blood and nervous Juices to the Part, is very much excited; and more particularly ſo when you preſcribe at the ſame time nervous and ſtrengthening Medicines to be given internally.

Stiffneſs of the Joint.

XIII. When the Joint is become rigid and inflexible, which Diſorder the *Greeks* call an *Anchyloſis*, if it is occaſioned by a Diſcharge of the Juices of the broken Bone into the Joint, which concrete there inſtead of forming a Callus in the fractured Part, this Caſe will turn out very difficult to cure: but if this Diſorder is occaſioned by having kept the Joint for a long Time without Action, or from a Concretion of the Juices that are ſecreted in theſe Parts to make them ſlippery and eaſy to move; it will be very proper to foment the rigid Part with emollient Fomentations and Baths; to rub it frequently with Oils and Fat of Animals, or with emollient Ointments; and to move it backwards and forwards frequently with your Hands, till it ſhall recover its natural Faculty of moving [a].

Fracture with Diſlocation.

XIV. You have frequently a Diſlocation as well as Fracture of the Bone, in one and the ſame Limb: when this is the Caſe, the Luxation muſt be remedied in the firſt Place; and then the fractured Parts may be reſtored to their natural Situation; each of them muſt be dreſſed with a proper Bandage: ſometimes the Fracture happens ſo near the Head or Articulation of the Bone, that it is impoſſible to fix your Hands or Inſtruments to make a proper Extenſion: in this Caſe, the Fracture is firſt to be attended to; which muſt be cured, before you can attempt to remedy the Luxation; though you ſhould be very careful, during the Cure of the Fracture, to foment the luxated Limb *cum Spiritu Vini, vel ſolo, vel camphorato, vel & Aceto calefacto:* this Method may keep the Part free from Inflammation and Tumor. I will not pretend to affirm, that this Method of Cure is always to be depended upon; for it frequently happens that the luxated Parts are to be reduced by no Art: but as this is the only probable Method of relieving the Patient, and as there are frequent Inſtances of its being attended with Succeſs, even where the Luxation has been of ſome Months, or even a Year's ſtanding, I think it ought by no Means to be rejected [b].

In what Manner Limbs are to be broken again, when they have been ill ſet.

XV. If a fractured Limb appears crooked and deformed after the Cure has been performed, which Accident happens either from the Negligence of the Surgeon, or from the imprudent and reſtleſs Behaviour of the Patient; I know of no other probable Method of reſtoring the Limb to its former Shape and Beauty, than by making a ſtrong Extenſion of it, and breaking it in the Part where it is juſt united; by this Means the Parts may be replaced in a more proper Manner: great Care and Circumſpection is required in the Treatment of the ſecond Fracture. When the Deformity complained of is but ſmall, and the Callus intirely indurated, or where the Patient is in Years and infirm, I ſhould not adviſe this Method of Cure to be attempted; ſince it is not only attended with great Pain, but with great Danger alſo: on the other hand, when the Callus is tender, and the Patient young and vigorous, I think this Operation

[a] For the Cure of an *Anchyloſis* ſee LE DRAN, *Obſ.* 93, 94. and BOERHAAVE's *Aph. Pract. N.* 556.
[b] See PETIT, on *Diſeaſes of the Bones.*

may

may be fairly attempted: in the mean Time, it is neceſſary to obſerve here, that, before you undertake this Cure, you muſt endeavour to ſoften the Callus, by uſing emollient Baths, Fomentations, and Ointments, for ſeveral Days.

CHAP. III.

Of FRACTURES *in particular.*

The Connection and Argument of this Chapter.

I. SINCE we have already treated of Fractures of the Bones in general, it, remains now that we ſpeak to particular Fractures: and firſt, in this Chapter, we ſhall treat of thoſe that happen in the Head: we ſpoke largely enough above in *Chap.* XIV. of Fractures of the Cranium; therefore we ſhall now proceed to deſcribe other Kinds of Fractures.

FRACTURE *of the* NOSE.

Fracture of the Noſe.

II. In the Noſe, both Bone and Cartilage are the Subjects of Fracture, which happen ſometimes on either Side; ſometimes in the Middle, chiefly by a Blow or Fall; this is eaſily to be diſtinguiſhed by the Sight or Touch: if either of the Bones in the Front of the Noſe are fractured, it produces a Flatneſs in the Noſe, and the Air meets with Obſtruction in its Paſſage through the Noſtrils; if the Bone on either Side is fractured, the Part becomes hollow; when the Cartilage is diſturbed, the Noſe inclines too much to one Side: *See* CELSUS *upon this Head, Lib.* VIII, *Cap.* 5. Sometimes the Fracture happens without a Wound, but is much oftner attended with a Wound of the common Integuments: *See* SAVIARD, *Obſ.* 107. If the Injury of the Noſe is very violent, the Fracture cannot be ſo perfectly cured, but ſome Deformity will ſtill remain: the Vicinity of this Part to the Brain, which is frequently injured at the ſame Time, renders Caſes of this Kind frequently very dangerous: a *Caries* alſo, *Ozæna*, and *Polypus*, are no uncommon Attendants upon this Diſorder; by which Means the Senſe of Smelling, the Faculty of Speech, and the Actions of Inſpiration and Exſpiration, are very much diſturbed.

After what Manner Bones of the Noſe are to be replaced.

III. In order to reſtore the fractured Bones of the Noſe to their natural Situation, the Patient is to be placed in a Seat oppoſite to the Light, and his Head held back by an Aſſiſtant; the Surgeon is to raiſe the depreſſed Parts with a Spatula, Probe, or a Quill, applying externally the Thumb of one Hand, and the Fore-Finger of the other: if the Bones of the Noſe are fractured on both Sides, they are to be raiſed on each Side after this Manner, and the Cavity of the Noſtrils is to be filled up with long Doſſils, to prevent the Bones from collapſing; covering the Part alſo, for this End, with ſome Plaſter, having firſt applied ſuch Dreſſings as are ordinarily uſed to recent Wounds: if the Bone is fractured into ſeveral Splinters, they are to be forced into their proper Places by the Fingers; but if a Splinter is ſo entirely ſeparated from the Bone that it will not eaſily unite with it again; you muſt remove it with your Forceps.

How a Fracture with a Wound ſhould be treated, and Bandage be performed.

IV. When the Fracture of this Part is accompanied with an external Wound after you have replaced the Bones, you ſhould dreſs the Wound (at firſt) with dry Lint, covering it with a vulnerary Plaſter; afterwards you muſt uſe balſamic Medicines; ſuch as *Ung. digeſtiv. Eſſent. Aloes, Myrrhæ, Succin. Maſtich* All greaſy and oily Medicines are to be diligently avoided here, and in all

other Caſes where the Bone is injured; becauſe they are very hurtful in theſe Caſes: but, where you have no external Wound, it will be ſufficient to apply a ſticking Plaſter to the Part, to ſecure the Bones in their Situation; and by this Means you will find they will unite in about fourteen Days; if no Abceſs or Caries ſupervene. If the Bone ſhould require a ſtronger Support than what we have hitherto mentioned, you may make one of ſingle or double Cap-paper, which may be adapted to each Side of the Noſe, and ſupported with Bolſters: *See Plate* VIII, *Fig.* 8: the whole muſt be ſupported with a Bandage of four Heads, which muſt not be bound on too tight; which will appear to you more clearly, when you conſult what we ſhall ſay below, where we are to treat profeſſedly of Bandages: before the Plaſters and Bandages are applied, ſome introduce a ſilver or leaden Pipe, or Quill, into each Noſtril, to render the Faculty of Breathing eaſier: *See Plate* II, *Lett.* P *and* Q: in order to ſecure theſe Pipes and the Bones of the Noſe in their proper Situation, they uſe the four-headed Bandage. Some amongſt the modern Surgeons intirely reject the Uſe of all this Apparatus, except the Bolſters, Bandage, and Plaſter; for they are of Opinion that it does more Harm than Good, and that the Introduction of Pipes, or even Tents, into the Noſtrils, will occaſion ſo great a Degree of Irritation, and ſuch a Difficulty of Breathing, as is not to be borne: beſides, when once the Bones of the Part are properly replaced, they are not ſo eaſily diſturbed as is commonly imagined: in theſe and all other Caſes, where there is no Neceſſity for a more laborious and complicated Treatment, the ſimpleſt and eaſieſt Method of Cure is always to be preferred.

CHAP. IV.

Of a FRACTURE *of the* JAW.

Of a Fracture of the Jaw.

I. THO' both Jaws are liable to Fractures, the upper is leſs ſo than the lower and even that, than the reſt of the Bones: when they happen in the upper Jaw, the divided Parts muſt be replaced, as near as poſſible; and then covered with a Plaſter, as in the Noſe: when in the lower Jaw, it is broke either on one Side or on both; and the divided Parts in this Caſe do not recede any conſiderable Diſtance from each other; for the Muſcles of this Part are ſo ſituated, that the Bones are not much ſeparated from each other by their Action. but the Degree of Injury depends upon the Violence of the Blow received.

By what Signs a Fracture of the Jaw is known.

II. That Kind of Fracture in this Part is ſooneſt diſcovered, where the Bones are ſeparated from each other; for not only your Eye, and often your Ear, but eſpecially the Touch, will ſpeedily and evidently demonſtrate what is diſplaced in the Jaw; and whether the natural Poſition of the Teeth be diſturbed: beſides which, the Patient's ſuffering violent Pains, and ſometimes Convulſions, is uſually a pretty certain Sign that the Jaw is fractured; but, if the Pieces of the Bone are not ſeparated, the Fracture is diſcovered with much more Difficulty.

How the Bones of the Jaw are to be ſet or replaced.

III. A Fracture of the Jaw being thus diſcovered, our next Intention is to reſtore the broken Bones to their proper and natural Poſition: the Patient is therefore to be commodiouſly ſeated againſt the Light; and his Head to be held firm

firm by an Aſſiſtant: the Surgeon is then to introduce his Thumb or Fore-Finger of one Hand into his Mouth, applying his other Hand externally; and by this Means he is to preſs the Fragments of the Jaw on each Side, till they have regained their former Situation; which may be known by the regular Diſpoſition of the Teeth: but if any of the Teeth be found looſe or ſlipped out, it may not be improper, if nothing hinders, to reſtore them afterwards to their Places[a], and to faſten them by gold or ſilver Wire, or with Cerate, to ſuch as are next them; for by this Means they have been frequently held firm: if the Jaw ſhould happen to be broke on both Sides, they muſt be reſtored one after the other by the ſame Method as before: but then the Operation is uſually more or leſs ſucceſsful in Proportion to the Surgeon's Skill in the Anatomy of this Part: if there ſhould be a Piece not moved out of its Place, there will be no Occaſion to reſtore it.

What is to be done after the Jaw Bone is ſet.

IV. After the Bones are properly reduced, they muſt be covered with, firſt, a Plaſter, and then a Compreſs, dipped in *Sp. Vini*, and applied internally; and another Compreſs ſewed to a Piece of Paſte-board in the Form of a half Jaw, is to be laid on externally. *See Fig.* 9. *Tab.* VIII. Theſe are to be kept on by the Bandage with four Heads, perforated in the Middle, to let in the Chin; or elſe it muſt be very carefully bound up with the particular Bandage for this Caſe, which we ſhall deſcribe when we come to treat profeſſedly of Bandages: but whenever the Jaw is found to be fractured on both Sides, it is uſual to introduce and apply internally, after the Compreſs dipped in *Sp. Vini*, another made of thin Paſte-board, perforated in its Middle, and accommodated to the Figure of the Chin, as at *Fig.* 10. In this Manner its Middle (*a*) that is perforated, is to be applied to the Chin; and its two Extremities (*bb*) toward the Ears: but Fractures of this Part may be well enough cured without Plaſters and Splints, where we can commodiouſly apply a Bandage; for the Bones are not very eaſily diſplaced, when they are once reduced: in what Manner this Part is to be bound up, we ſhall make pretty evident, when we come to treat of Bandages in particular.

How the Patient ſhould behave himſelf under the Cure.

V. To forward the Agglutination of the fractured Jaw, after Phlebotomy, the Patient ſhould be reconciled to reſt as much as poſſible; but above all he ſhould ſtrenuouſly avoid, particularly for the firſt Days, all Talk and Eating: it ſeems therefore to be much the ſafeſt Way to live upon, till the Jaw is grown firm, only fluid Aliments, ſuch as Broths and Soops, poached Eggs, Jellies, and the like, taking Care not to lie on the Back, and ſtrictly to avoid turning on the Face, or either Cheek: by which Means the Fracture will be well in about twenty or thirty Days; eſpecially if the internal Parts of the Mouth that are injured, be frequently moiſtened with a little *Mel Roſarum*.

VI. If the Fracture be attended with a Wound, it muſt be undone every Day, and treated as we have taught in *Chap.* IV, *N.* VI. till it be healed. An Example of a Fracture in both Jaws may be ſeen in LE DRAN, *Obſ. Chirurg.* 3, *Tom.* I. but of the lower Jaw only, in *Obſ.* 8.

[a] GOUEUS indeed diſſuades us from this Method, thinking that the Bones will by this Means be again diſplaced; but TURNER (and ſome others) in his *Surgery*, gives an Inſtance where it ſucceeded; and ſo does LE DRAN, *Obſ.* 3. Tom. I.

CHAP. V.

Of a FRACTURE *of the* CLAVICLES, STERNUM, *and* HUMERUS.

Of a fractured Clavicle.

I. THE Clavicle[a] is extremely ſubject to be fractured both from its tranſverſe Poſition, and from its Smallneſs; which happens either in its Middle, near the *Humerus*, or near the *Sternum*: but in which-ever of theſe Parts it is broke, that End next the *Humerus* always deſcends lower than the other, next the *Sternum*; from the Weight of the Arm, which was before ſuſtained by the Clavicle and Head of the *Sternum*: and notwithſtanding that Part of it next the *Sternum* remains immoveable, by the Deſcent of its other End. it can ſcarce happen but they will, in ſome Meaſure, collapſe one over the other.

How a Fracture of the Clavicle is to be diſcovered.

II. It is no great Difficulty to know when this Part is fractured: for (1) it will be hardly poſſible for the Patient to lift up his Arm: (2) his Arm will hang inclining towards his Breaſt, whereas before it was ſtraight, or tended rather backward: (3) and laſtly, as the Clavicles are covered with ſcarce any Muſcles, the Fracture will be clearly evident both to the Touch, the Eye, and the Ear; eſpecially upon any ſmall Motion of the Part.

The *Prognoſis* of a fractured Clavicle.

III. When the neighbouring Parts are not affected, this Fracture is attended with no bad Conſequences; but if the adjacent Veins, or Arteries, or even Nerves are injured, there is generally great Danger: the Reduction of a broken Clavicle is not very hard to be effected, eſpecially when the Fracture is tranſverſe; nor is it uſual for the *Humerus*, with the Fragment of the Clavicle, to be ſo far diſtorted as not to be eaſily replaced with the Fingers: but the Difficulty is much greater to keep the Bone in its Place, when the Fracture is once reduced, eſpecially if the Bone was broken obliquely; for which there are two Reaſons: *viz.* the circular Bandages, with which the Bones of the Arms and other Extremities are uſually held very firm, cannot be applied here, by reaſon of the Form and Situation of the diſordered Part; and then the Weight of the depending Arm itſelf ſoon pulls aſunder what the Surgeon has been replacing: it is no Wonder, therefore, if the Juncture of the Clavicle be often found either uneven or unfirm after its Agglutination: yet we do not want Examples where fractured Clavicles have been very happily and firmly cured, eſpecially when the Patient keeps himſelf quite free from Motion.

How the broken Clavicle is to be reduced.

IV. A Fracture of the Clavicle is to be reduced in the following Manner. The Patient muſt be placed on a low Seat, and an Aſſiſtant is to thruſt his Knee againſt the Middle of the Patient's Back, between his two Shoulders; then laying hold of the Patient's Shoulders with each Hand, he muſt pull them gently and gradually backwards; by which Means the Clavicles will be properly extended: while this is doing, the Surgeon muſt ſtand before, and endeavour to replace the Bone with both his Hands, ordering the Aſſiſtant to hold the Bone in that Poſition: he is then (1) to apply the narrow and thick Compreſs (*Tab.* IX. *Fig.* 13.) folded up at each End, ſo as to fill up the Cavities above and below the Clavicle: upon theſe (2) he is to lay two more narrow

[a] A Fracture of the Clavicle is by CELSUS (*Lib.* VIII, *Cap.* 8.) called *Jugulum fractum*; but all the modern Surgeons and Anatomiſts give the Name of *Clavicle* to this Bone, and attribute a quite different Signification to the Word *Jugulum*.

Compreſſes,

Compreſſes, made in the Form of the *Letter* X. Over all theſe, he is (3) to to apply a Piece of Paſte-board (*Tab.* VIII, *Fig.* 12.) accommodated to the Shoulder and Neck, and firſt ſteeped in *Sp. Vin.* or Oxycrate: then he muſt (4) place a Ball under the Arm, or bind it ſeveral Times with a thick Roller, to prevent the *Humerus* from ſubſiding: and laſtly (5.) the whole is to be diſcreetly bound up, and the Arm ſuſpended in a Saſh or Sling, that is put about the Neck: the Plaſters that were uſed to be frequently applied in this Caſe, have been generally found uſeleſs.

V. As it is ſometimes very difficult to keep the Arms from puſhing inwards, which would diſturb the Agglutination; it will be of Service to uſe a wooden or iron Inſtrument (*Tab.* VIII, *Fig.* 13.) in the Form of a T, ſo contrived as to keep back the Shoulders: the Sides of this Inſtrument are about the Breadth of three Fingers, and lined with Cloth or Leather. It is to be applied thus: *viz.* the two Arms AA, are to be placed againſt each Shoulder, and the perpendicular Part B, is to go againſt the Middle of the Back: through the Aperture C, is paſſed a double Ligature to faſten it to the Body, the two Arms being firſt put through the Rings AA, which may be widened or narrowed at Pleaſure: the tighter the perpendicular Part B is faſtened to the Body, the more the Shoulders are by that Means drawn backward: but if they cannot be this Way drawn tight enough, a Compreſs, folded lengthways, is to be firſt placed between the Back and the Inſtrument; by which Means the Shoulders will be drawn more ſtrongly backwards: the Rings AA, may be made of Iron or Leather, ſo as to be taken in, or let out, as there may be Occaſion.

VI. Whenever there are any looſe Splinters of the Bone that are intirely ſeparated, which not only wound and hurt the Fleſh, but obſtruct the Meeting of the Clavicle; it ſeems altogether requiſite to open the Skin and remove them, before the Reduction of the Bone, treating the Wound as uſual: but if there ſhould be any Splinters which ſtill adhere to the Bone, and prick the adjacent Parts, or impede the Reduction; they muſt be alſo either taken off with the *Forceps*, (*Plate* VIII, *Fig.* 1.) or elſe forced into their Places, whereby they may be again united to the Bone: but to divide the Parts, and remove the Fragments, requires great Caution; leſt ſome of the large ſubclavian Veins or Arteries be wounded in the Operation, and a fatal Hæmorrhage be thereby produced. What is to be done in caſe of looſe Splinters.

VII. The *Scapula* is uſually fractured, either near its *Acromion* or Head, where it joins with the Clavicle, or in ſome other Part; which will be diſtinguiſhed by the Eye or Touch. If in its *Proceſſus Acromium*, the Reduction may be eaſily made, by lifting up the Arm to relax the Deltoide Muſcle; or by puſhing the Arm evenly upwards, and drawing the fractured Parts together with the Fingers; but then they eaſily ſlip away again, by any ſlight Cauſe, and ſo are with Difficulty agglutinated; more eſpecially they are eaſily ſeparated by the Weight and Motion of the Arm, and the Contraction of the Deltoide Muſcle; inſomuch that there is ſcarce any body that ever cures a fractured *Acromion* ſo as to admit afterwards of a free Motion of the Arm upwards[a]. In the mean Time all Means muſt be uſed to retain the replaced Bones in their right Situation: a Compreſs wet with *Sp. Vin.* is to be applied to the Fracture; a Ball is to be put under the Arm-pit to to ſupport it; the whole is to be bound up with Of a Fracture of the *Scapula*.

[a] Such is the Opinion of CHESELDEN, treating of this Bone, in his Anatomy.

the Bandage commonly called *Stica*, and the Arm is to be ſuſpended in a Saſh or Sling, hung about the Neck; and the Patient muſt reſt himſelf without Intermiſſion: but if the Neck of the *Scapula*, which lies under the *Acromion*, or its *Acetabulum* ſhould be fractured, (which is a Caſe that ſeldoms happens as it is difficult to diſcover, by reaſon of its thick Covering), it is a hundred to one, but from the Vicinity of the Articulation, the Tendons, Muſcles, Ligaments, Nerves, and large Veins and Arteries, there will follow a Stiffneſs and Loſs of Motion in the Joint, a violent Inflammation, Swelling, and Abceſs, with the worſt of Symptoms, and even Death itſelf; as happened in a Caſe I ſaw, of a certain Profeſſor at *Helmſtadt*[a]: but when the Fracture falls on ſome other Part of the *Scapula*, the Symptoms are generally much milder.

How the fractured *Scapula* is to be reduced.

VII. That the fractured *Scapula* may be ſet with the greater Readineſs, an Aſſiſtant is to extend the Arm gently forwards; the Surgeon in the mean Time dextrouſly replacing the Fracture with his Hands, ſhould apply afterwards the proper Compreſſes, and Slips of Paſte-board, ſuitable to the *Scapula*, and firſt wet with *Sp. Vin.* or Oxycrate; which are then to be firmly bound on with the *Stellate*, or four-headed Bandage, as we ſhall direct at large in the third and laſt Part of this Treatiſe.

FRACTURE *in the* STERNUM.

Of a fractured *Sternum*.

IX. The *Sternum* is equally ſubject to Depreſſions and Fracture, from Falls or Blows, with the reſt of the Bones: when either of theſe happen, the Part is not only uneven and painful, but the ſubjacent Arteries and Veins are alſo contuſed or ruptured; whence ariſe Pains in the Breaſt, Difficulty of Breathing, violent Coughs, ſpitting of Blood, or elſe Extravaſations of it in the *Præcordia*, or between the Duplicature of the *Mediaſtinum*, with many bad Symptoms of the like Nature.

The Signs of a fractured *Sternum*.

X. The Signs therefore of a fractured *Sternum*, will be, in my Opinion, ſufficiently evident, from what follows: namely, its Depreſſion or Fracture will appear not only from the Symptoms beforementioned (*N.* IX.) but frequently alſo from the *Sternum's* being unequal or moveable to the Touch; eſpecially when one Part grates againſt the other: the Depreſſion of the *Sternum* will be alſo apparent, not only from the Symptoms of the preceding Section, but alſo from the Cavity or Inequality made in this Part, which is a Sign peculiar to this Diſorder.

How the *Sternum* is to be replaced.

XI. In order to ſet the Fracture, if any Part of the Breaſt-bone be diſplaced, it will be very ſerviceable to lay the Patient on his Back, upon a Bed, or rather a Table, putting a hard Pillow, a large Piece of Cloth rolled up, a Drum, or other Cylinder under his Back, preſſing down his Shoulders, by which Means the *Sternum* will be elevated and extended: and to facilitate the Reduction, the Surgeon muſt preſs the Sides of the Breaſt together, and ſhake them pretty ſtrongly: by this Means you not only puſh the Ribs forwards, but at the ſame Time force what is depreſſed in the *Sternum* into its natural Situation: but when this Method is impracticable, or not proper, the Skin muſt be divided, and the depreſſed Part of the *Sternum* lifted up into its Place, by means of a Lever,

[a] The ſame has been obſerved by CHESELDEN *(loc. citat.)* and by DOUGLAS.

commonly

commonly called an Elevator; or elſe by a Screw, gently wormed into the Part, and pulled upwards. Notwithſtanding this Way of Cure is more operoſe and difficult than the former, it is preferred by GOUEY (in his *Chirurgie veritable*) and PETIT (*de Morb. Oſſ.*) as the beſt and readieſt Method: as for the fitteſt Method of retaining the *Sternum* after its Reduction, we ſhall treat of that more at large, when we come to the Doctrine of Bandages: but if, as it ſometimes happens after the Reduction, violent Pains continue under the *Sternum*, and if Blood ſhould gather and ſuppurate internally between the Duplicature of the *Mediaſtinum*, it will not be improper to trepan the lower Part of the *Sternum* (as PETIT adviſes) after the Manner we do the *Cranium*: and when the putrid Matter is diſcharged, and the Cavity cleanſed, it ſhould be carefully treated with ſome vulnerary Balſam; laſtly, if any Blood ſhould be found extravaſated in the Cavities of the *Thorax*, the Cure ſeems to depend intirely upon diſcharging this by the *Paracenteſis*, in the Manner we have deſcribed under Wounds of the *Thorax*. As to the Buſineſs of Dreſſing, after the Application of Compreſſes dipped in warm Wine, or *Sp. Vin.* we muſt go on with that Kind of Bandage called the *Napkin-and-Scapulary*.

CHAP. VI.

Of FRACTURES *in the* Ribs, Vertebræ, Os Sacrum *and* Innominatum.

Of fractured Ribs.

I. SOMETIMES the Ribs are fractured, or only fiſſured, in ſuch a Manner, that barely ſome external or internal Part of them are hurt, and not moved out of the natural Place: which Caſe is uſually attended with no bad Symptoms, and is often ſcarce diſcoverable, the Bone growing together again of itſelf: but if the whole Rib be fractured, and ſome Part of it moved out of its Place, it is a more dangerous Caſe: for the coſtal Muſcles, and the *Pleura* that lines the internal Cavity of the *Thorax*, will be very much diſturbed, inflamed, or torn by the ſeparated Fragments of the Bone. When a Rib is fractured, it projects either externally or internally, much in the ſame Manner as if it was a broken Arch: when it projects externally, the Symptoms are uſually much the milder [a]; but when it is driven inwards, the Caſe is much worſe, eſpecially if any of the intercoſtal Veins or Arteries be divided ſo as to let Blood run into the *Thorax*: in Conſequence of which, we need not wonder if violent Prickings, Inflammation, difficult Reſpiration, Cough, Fever, Spitting of Blood, Suppuration, Extravaſation of Blood in the *Thorax*, or cellular Interſtice of the *Mediaſtinum*, and other bad Symptoms, ſhould follow in Courſe; eſpecially if the neighbouring *Viſcera* [b] be wounded, or more Ribs be broken at the ſame Time: if theſe be not timely remedied, they produce violent Fevers, Inflammation, and Ulceration of the Breaſt and Lungs, *Empyemas*, incurable *Fiſtulæ*, and *Caries* of the Bones; and ſometimes Death itſelf will be the End,

[a] Indeed GOUEY denies that broken Ribs are ever drove outwards; but PETIT (*Lib. de Morb. Oſſ.*) witneſſes that there may be ſuch a Kind of Fracture.

[b] All Fractures of different Ribs at the ſame Time are ranked by BOHN amongſt mortal Wounds: *Lib. de Vuln. leth. C.* 3. and I myſelf have been Witneſs to ſuch a Fracture, where the Patient died in a few Hours.

particularly in a Fracture of more than one Rib at the ſame Time: it frequently happens, unleſs the Fracture be a ſimple one, that the ſoft Parts are punctured, and an external Wound made, by ſome ſharp Piece of the Bone: If the Parts are wounded, it occaſions ſometimes a very profuſe Hæmorrhage, often very difficult to ſtop, as one cannot eaſily get at the Arteries beneath the Ribs: and if the Blood ſhould run into the *Thorax*, it can ſcarce be diſcharged from thence but by the *Paracenteſis*, or elſe by dilating the Wound, when it happens between the baſtard Ribs. If by any external Force the Cartilages ſhould be ſeparated from the Ribs, we term it a Fracture, and treat it in the ſame Method with other Fractures in this Part, which we are going to deſcribe.

The Method of diſcovering a Fracture of the Ribs.

II. When the fractured Parts of a Rib keep in their natural Situation, they continue even and ſmooth to the Eye, and are unaccompanied with any conſiderable Pain; it is therefore difficult to diſcover the Fracture: but yet, upon ſlightly moving the ſame, it will be attended with ſome Pain, though it will the more readily grow together: but when the fractured Parts recede from each other, the Deformity will be apparent both to the Eye and Touch, and a Noiſe will be heard upon moving them. If a ſharp Piece of the Bone ſhould moleſt the *Viſcera* internally, it will occaſion the greater Part of the Symptoms mentioned at *N.* I, and from the Intenſity and Malignity of thoſe, we judge the Fracture to be more or leſs dangerous: but it alſo frequently happens, that a Fracture of the Ribs occaſions a windy Tumor, called by the Greeks, *Emphyſema*; formed by the Air inſinuating itſelf, by a ſmall Wound, between the Skin and Muſcles, into the Subſtance of the cellular or adipoſe Membrane; ſpreading itſelf afterwards up to the Neck, Head, Belly, and other Parts, much after the Manner in which Butchers blow up their Veal.

How a ſlight Fracture of the Ribs is to be ſet.

III. In order to replace fractured Ribs, it is always previouſly neceſſary to inquire whether the Splinters project externally or internally: when the firſt is the Caſe, the Patient is to be placed on a high Table, and the ſeparated Bones muſt be gently forced by the Fingers into their Places, the proper Compreſſes dipped in *Spir. Vin.* muſt be laid on, and then covered with Slips of Paſte-board or Splints; and laſtly, the circular Bandage, or elſe the *Napkin-and-Scapulary*: but when the latter is the Caſe, while the Patient retains a deep Breath, the Surgeon carefully compreſſes both Sides of the Rib with his Hands, agitating till they are properly fixed. What is farther neceſſary to be done in this Caſe, will come under the Head of Bandage; unleſs that the Paſte-board is to be here omitted, and the Napkin not drawn very tight; but the Dreſſing need not be undone, unleſs it be over looſe, and ſome Symptoms or the Patient's erect Poſture require it: by theſe Means, Fractures of this Kind are uſually cured in about three or four Weeks Time: through the whole Courſe of the Cure, as CELSUS (*Lib.* VIII, *Cap.* 9) adviſes, the Patient muſt carefully avoid all Talk and Clamour, Paſſions and Anger, violent Motions of the Body, Smoke, Duſt, and every thing that will occaſion Sneezing or Coughing. But if the Reduction cannot be effected by the Means hitherto delivered, it may not be improper to try by ſome ſticking Plaſter, as in a Depreſſion of the *Cranium* at *Book* I, *Chap.* XIV, *N.* XXIV.

What is to be done in the more dangerous Fractures of the Ribs.

IV. If any ſharp Pieces of the Ribs ſhould pierce the *Pleura*, it will occaſion moſt violent Pains, a Difficulty of Breathing, a Cough, Spitting of Blood, Inflammation,

ſlammation, Fever, and other ſuch grievous Symptoms: therefore it will be proper to open the Skin, and extract the Fragments which ſtick in the Fleſh with the Fingers, Plyers, Hooks, or other proper Inſtruments: unleſs this Method be followed, the Patient will be in great Danger; to prevent which, Phlebotomy, Clyſters, cooling and anodyne Medicines are to be uſed, and a thin Diet muſt be followed: this Method of Inciſion is alſo more particularly neceſſary when the ſticking Plaſter, and other Means adviſed, prove inſufficient to reduce the Fracture.

Of the Veins and Arteries hurt.

V. When there happens to be a Wound of any of the Veins or Arteries which run under the Ribs ſo as to let their Blood flow internally, the Caſe will be much the ſame with the Wounds mentioned in *Book* I, *Chap.* X, and it ſeems then neceſſary to open the *Thorax* near the fractured Part, ſufficient to admit the Finger, anointed with ſome Liniment, and dipped in ſome ſtyptic Medicine, which is to be held upon the Veſſels till the Blood ſtops: but when the Finger proves ineffectual, the divided Veſſel muſt be diſcovered, and cloſed either with a Ligature or an actual Cautery, properly applied: and in order to diſcharge what is lodged in the *Thorax*, when the Wound itſelf is in the lower Part thereof, the Surgeon muſt dilate and keep it open with Lint, as was adviſed before in Wounds of the *Thorax:* but when the Height of its Situation in the Breaſt, near the upper true Ribs, will not admit of a convenient Diſcharge by that, a freſh and more convenient Opening or *Paracenteſis* muſt be made in the lower Part of the *Thorax*; and the Wound in the ſuperior Part muſt be cloſed: ſee *Book* I, *Chap.* X, *N.* X.

How the *Emphyſema* and Contuſion are to be treated.

VI. When an *Emphyſema* happens, it will be very proper to inlarge the Opening in the Skin, if too narrow, with the Scalpel; and to bring down the Tumor with Frictions and Bandage, carrying the Compreſſion gradually towards the Opening, ſo as to expel the included Air by Degrees: but if there ſhould be a Contuſion alſo, it muſt be treated in the Method which we have already laid down, in the Chapter (XV, *Book* I) of *Contuſions:* if a violent Cough or Inflammation follow, it muſt be remedied by Bleeding, and other proper Medicines: ſee an Example in LE DRAN, *Obſ.* 29, *Tom.* I.

Fractures of the *Vertebræ*.

VII. When any of the *Vertebræ* are fractured, either by a Fall, Blow, or any other Cauſe, without hurting the ſpinal Marrow; we may reaſonably ſuppoſe that the Fracture is confined to ſome of the oblique or ſpinal Proceſſes; and therefore the Patient will be in no great Danger: but when the Body of the *Vertebra* is either broke or ſplit by ſome external Force[a], and the contiguous ſpinal Marrow bruiſed or compreſſed; all Parts of the Limbs and *Viſcera* beneath that Vertebra become immoveable and rigid: no Wonder then, if a ſpeedy or ſlow-paced Death often follows, in Proportion to the Degree of Damage; here it may be alſo proper to recall to Mind, what has been ſaid in the preceding Book, on Wounds of the *Medulla ſpinalis:* and laſtly, if the tranſverſe Proceſſes of the *Vertebra* are broke, which incline towards the Cavity of the *Thorax*, it is ſcarce poſſible that the Heads of the Ribs which are there connected, ſhould eſcape being fractured alſo; which makes the Caſe very deplorable.

[a] GOUEY thinks the Body of the *Vertebræ* cannot be fractured, unleſs by a Buſſet; but I have ſeen them from a violent Fall off a high Place, and the Patient died ſoon after, from bruiſing the *ſpinal Marrow*; as they generally do in this Caſe.

The Signs of Fractures in the *Vertebræ*.

VIII. Fractures in the *Vertebræ* may be judged to exiſt from (1) conſidering the Nature of the external Violence, whether it be a great Fall, Blow, or the like; but more eſpecially (2) from the Pains ſeated about the affected *Vertebræ*; and laſtly (3) from the Manifeſtation thereof, to the Touch, Eye, and Ear.

How to replace the fractured Proceſſes or *Apophyſes*.

IX. When any the Proceſſes of the *Vertebræ* are found broken, it will be much the beſt Way to force them into their Places with the Fingers, placing narrow Compreſſes dipped in warm Spirit of Wine on each Side the *Vertebræ*, and over them Slips of thick Paſte-board, to be kept on by the *Napkin-and-Scapulary:* for by this Means, the Bones of the *Vertebræ*, which are very ſoft and ſpungy, will quickly and eaſily grow together again.

How the worſt Kind of Fractures in the *Vertebræ* are to be treated.

X. If in any Caſe the ſpinal Marrow ſhould be divided, Death will be generally an inevitable Conſequence: but to offer the Patient no Aſſiſtance becauſe we deſpair, would ſeem cruel and uncharitable; therefore we muſt try our Skill, though our Attempt ſhould be in vain; in order to which, the Surgeon muſt lay bare the fractured *Vertebra* with a Scalpel, and replace or elſe remove ſuch Fragments as injured the ſpinal Marrow: the Wound is to be afterwards gently cleanſed as uſual, and dreſſed with the Balſams mentioned *Book* I, *Chap.* II, *N.* XV, clapping over them a Compreſs dipped in warm Spirit of Wine, or Lime-water, and Spirit of Wine camphorated, to be held on with the *Napkin-and-Scapulary*, till the Wound ſhall terminate either in a perfect Cure or Death.

Fracture of the *Os Sacrum*.

XI. It ſometimes alſo happens, that by a Fall or a Blow, the *Os Sacrum* becomes in like Manner fractured; which may be diſcerned to be broken, from conſidering the external Violence, the Pains, by the Touch, *&c.* as is uſual in other Fractures.

How to ſet a Fracture of the *Os Sacrum*.

XII. As ſoon therefore as the *Os Sacrum* is found to be fractured, its Fragments are to be forced into their Places with the Fingers: but if any Part of it be depreſſed inwards, it may be convenient to introduce a Finger (that has firſt had its Nail cut cloſe and been dipped in Oil or Butter) up the *Anus*, in order to thruſt the depreſſed Fragment into its proper Place, to which it is to be directed externally by the other Hand: this being performed, we muſt apply ſome Plaſter ſuitable for Fractures, with Compreſſes dipped in *Sp. Vin.* over it, to be kept on by the T Bandage; or the Plaſter may be omitted and only the Compreſs and Bandage retained: and laſtly, to facilitate the Agglutination, the Patient ſhould keep his Bed quietly on his Sides for about a Fortnight: or if he muſt needs ſit at Times, let it be in a Chair without a Bottom, to avoid diſplacing of the Bone, from touching the Seat.

How the *Os Innominatum* is to be replaced.

XIII. When the *Os Innominatum* is broke, which ſeldom happens, it is readily diſcovered by the Injury and Symptoms in the neighbouring Parts, and is more particularly dangerous when the Patient diſcharges a brown and bloody Matter: in reſtoring this Bone, the Patient muſt lay down on his ſound Side; the Bone is to be replaced with the Hands, covered with Compreſſes, dipped in *Sp. Vin.* and bound up with the Bandage *Spica:* afterwards Bleeding, with cooling and relaxing Medicines, muſt be uſed, and a thin Diet obſerved; and let the Patient lie either on the ſound Side, or on his Back[a].

[a] See ROONHUYS, *Obſ.* p. 142, *Edit. Belg.*

CHAP.

CHAP. VII.

Of FRACTURES *in the Bones of the* Humerus, Cubitus, *and* Hands.

Fracture of the Humerus or Fore-Arm.

I. THE *Os Humeri* is broke either in its Middle, which is the leaſt dangerous; or elſe near its upper or lower Head, which is much worſe, as being more difficult to cure, and producing more violent Symptoms, acute Pains, Tumors, and Inflammation: indeed Fractures of this Part are uſually very obvious to the Senſes, being expoſed to the Eyes and Hands; but then they require a different Treatment, according to the particular Part injured: it ſometimes alſo happens, that the fractured Parts of this Bone keep their Places; but it more frequently falls out, that they ſlip one over the other, by which Means the fractured Limb becomes ſhorter than the ſound one: but it will ſometimes, though ſeldom, happen, that the two Parts of the Bone ſhall recede much from each other, by reaſon of the Weight of the Arm, which they ſuſtain: if the firſt be the Caſe, the Fragments are uſually more eaſily and readily replaced; but in the latter, there is required more Force and Skill to reduce the Bones to their Places, from whence they were removed; eſpecially if the Patient has tenſe Nerves and large Muſcles, as is uſually obſerved in ſtrong Men.

How a broken Arm is to be ſet.

II. In a Fracture of the *Os Humeri*, the Arm may be readily extended in the following Manner: let the Patient be ſeated on a high Stool, and an Aſſiſtant lay firm hold of his Arm above the Fracture, keeping his Elbow gently bended; then the lower Part of the Arm, beneath the Fracture, is in like Manner to be taken hold of, and the Arm is to be gently extended forward, by endeavouring to remove eaſily each Part from the other in a right Line: then the Surgeon himſelf lays hold of the fractured Part of the Arm with both his Hands, and ſtrives to replace the Bones, held in a due Extenſion by the Aſſiſtant, into their proper Situations; judiciouſly rolling up the Part with proper Bandages, agreeable to what has been ſaid of them in general in the Introduction, and what we ſhall explain more at large in the particular Doctrine of Bandages alone. If one Aſſiſtant be not able ſufficiently to extend the Arm of a robuſt Patient, the Office may be undertaken by two; or elſe thin Napkins or other linen Bandages may be wound round each Articulation of the Arm, and given to ſeveral Aſſiſtants, to be pulled in oppoſite Directions, till the Limb be ſtretched a little longer than it naturally ought; and then the Surgeon is to replace the Bones; but if the Surgeon's Hands and Napkin prove inſufficient, which is ſeldom the Caſe, HILDANUS's Girt, with the Sling (as deſcribed, *Plate* VIII, *Fig.* 17), muſt be applied to the Arm above the *Cubitus*; by which you may extend it, and reduce the Bone to its proper Situation.

Of Fractures in the Cubitus, or Lower Arm.

III. The lower Part of the Arm, called by Anatomiſts *Cubitus*, contains two Bones; the *Radius* and the *Ulna*: a Fracture in this Part may therefore happen to only one, or to both of theſe Bones; and that, either in their Middle or Extremities: but when they are both broke together, the Bones are not only very eaſily diſtorted from each other, but are alſo replaced and joined together again with much more Difficulty: but if one only ſhould be broke, whilſt the other:

other remains whole, the fractured Parts do not much recede out of their Places, nor are they very difficult to reduce and retain: for the ſound Bone is found to be a better Direction and Support in this Caſe, than either Splints or Bandages: when the Fracture happens towards the lower Head, near the *Pronator quadratus* Muſcle, the fractured Part is ſtrongly drawn (by that Muſcle, and the intervening Ligament that is ſpread between the *Radius* and *Ulna*) towards the ſound Bone, which makes it more difficult to replace: this is therefore a very material Circumſtance to be conſidered in the *Prognoſis* and Cure of this Fracture.

The Signs of a Fracture in the *Cubitus*.

IV. A Fracture in theſe Bones of the Arm may be will enough diſcovered by the Signs common to Fractures in general: but whether one or both be broke, and which of them is the Bone and its particular Part fractured; theſe may be known by the Sight and Touch, and by properly moving the Joint in or out, as may be neceſſary: it is however much eaſier to diſcover a Fracture in the *Ulna*, from its Inability to ſupport the Joint, as uſual, than that of the *Radius:* the Ear will alſo frequently aſſiſt the Sight, in the Search after this Fracture; for there will be generally perceived a Grating of the Bones, upon moving the Patient's Hand in and out, whilſt the upper Part of the *Cubitus* is held firm.

In what Manner the *Radius* is to be ſet.

V. If the *Radius* is to be ſet or replaced, whoſe Fragment is contracted towards the *Ulna*, an Aſſiſtant muſt hold the Arm whilſt the Surgeon inclines the Patient's Hand towards the *Ulna*, to draw back the contracted Part of the *Radius:* when this is done, he muſt carefully reduce them by Compreſſion on each Side with the Palms of both his Hands, ſo as to reſtore the compreſſed Muſcles, between the *Radius* and *Ulna*, and Fragments of the *Radius*, to their proper Places: the Arm is to be then bound up in the Method we ſhall hereafter deliver: and the Limb is to be put into a ſort of Caſe, (*Tab.* VIII, *Fig.* 14), made of Paſteboard or light Wood, to be ſuſpended in a Sling put about the Neck [a].

How the *Ulna* is to be replaced.

VI. In ſetting a Fracture of the *Ulna*, the ſame Method is to be obſerved with that of reducing the *Radius* as before, binding and ſuſpending it in the ſame Manner; but there is this Difference neceſſary to be obſerved, that in the Extenſion the Hand muſt be bent towards the Thumb and *Radius*, before the diſtorted Part of the *Ulna* can be compreſſed into its Place.

How we are to treat a Fracture of both the Bones.

VII. When both Bones of the *Cubitus* are broke, the Method of Cure will be much the ſame with that uſed to each of the Bones, when broke ſingly; unleſs that there is required more Strength and Circumſpection in replacing and retaining them, and the Bandage muſt be applied with greater Caution. And let the Surgeon, with the Palms of his Hands, compreſs the Patient's Fleſh on each Side of the Arm; by which Means the two Bones will mutually accord in returning to their natural Situation: we muſt be alſo careful to obſerve, that, while the Arm continues a good while without Motion, the Mucilage of the Joint does not harden, or the Ligament become ſtiff, and the Arm or *Cubitus* be thereby rendered immoveable: it will be therefore not improper to unbind the Part every other or third Day, and to move it carefully and gently, a little backward and forward, and ſometimes to foment it with warm Oil or Water; for by this Means its natural Motion will be eaſily preſerved.

[a] *Vide* SCULTET. *Tab.* LVI, *& infra Tab.* XXXVIII, *Fig.* 17.

VIII. The

VIII. The Bones of the Wriſt are ſeldom the Subject of Fracture, on account of their Smallneſs: but it ſometimes happens to them, from the Stroke or Compreſſure of ſome hard or heavy Body: when this is the Caſe, there uſually remain but little Hopes of effecting a Cure: for the Ligaments and Tendons are here ſo numerous, and the Bones themſelves are ſo very ſmall, that it ſeems ſcarce poſſible to reduce them into their Places, or make them grow together again: and on this Account, the Joint of the Hand generally becomes ſtiff and immoveable: or elſe violent Inflammations, Abceſſes, Suppurations, *Fiſtulæ*, and *Caries* of the Bones do thence ariſe; which, on account of the Softneſs of the Bones, and the Difficulty of diſcharging the Matter, are ſeldom remedied but by amputating the Hand: agreeable with this, RUYSCH (*Obſ. Anat. Chirurg. page* 10) among others, inſtances a Fracture of this Kind, which, after three Years Treatment, remained ſtill uncured. Fracture of the Wriſt.

IX. But that the Surgeon may not ſeem to be altogether negligent on his Part, he is rather to try what he can do in the Caſe, than to leave the Patient deſtitute of Help: it will be therefore moſt proper for an Aſſiſtant to lay hold of the Hand and Arm, above the fractured Wriſt, and to extend them as much as is ſufficient, in oppoſite Directions: while this is doing, the Surgeon muſt uſe all his Endeavours to reſtore the Fragments to their proper Places, with his Hands: and after he has very carefully reduced the Fracture, it is to be bound up with a ſuitable Bandage. How a Fracture of the Wriſt is to be treated.

X. As the *Metacarpus* is much more ſubject to Fractures than the Wriſt, becauſe its Bones are larger; upon the ſame Account it is alſo more eaſily replaced and cured: there can be hardly a better Method of reducing this Fracture, than that of ſpreading the Hand upon a ſmooth Table by an Aſſiſtant, the Surgeon carefully uſing all his Endeavours to replace the Bones with his Fingers, ſecuring them with a proper Bandage: an Inſtance of a Fracture in the Wriſt with a Wound, may be ſeen in LE DRAN's *Obſ.* 56, *Tom.* I. Fracture of the Hand, or *Metacarpus*.

XI. When one or more of the Bones in the Fingers are broke, the Surgeon's principal Buſineſs is, to carefully replace what has been removed, and to roll up the Finger a little Way with a narrow Bandage, and then to bind it firmly to the next ſound Finger: if the Thumb is fractured, it muſt be ſupported with ſmall Splints, and ſecured with a narrow Bandage: the Method of commodiouſly applying the Bandage when ſeveral of the Fingers are broke at once, will be declared hereafter in the Doctrine of Bandages: but when the Hand or a Finger is ſo violently maſhed as to have no Room to expect a Cure, it is more adviſeable to cut it intirely off, than to conſtantly torment the Patient, and perhaps put him in Danger of his Life. Fractures of the Fingers.

CHAP. VIII.

Of a fractured THIGH.

I. THE Thigh-bone, though the largeſt and ſtouteſt in the whole Body, is frequently broke after ſeveral different Manners; and that either in its Middle, or towards its Head and Articulation: but more frequently near that Part which Anatomiſts call its Neck, near its Articulation with the Hip-bone: which, whenever it happens, is very difficult to ſet, and more difficult to retain in its Place. When the Bone is broke in two Places at once, the Fractures of the Thigh.

Danger is ſtill greater: and if the Patient ſhould eſcape Death, which they uſually do not, it is a common Caſe for him to be ever afterwards lame: ſometimes the Bone is broke tranſverſely, ſometimes obliquely; and at other Times the Ends ſlip one over the other, which makes it a very bad Caſe: for the Muſcles of this Part being very robuſt, and ſtrongly contracted, draw the lower End of the Bone with a conſiderable Force upward, ſo as to make it require a conſiderable Strength to extend and replace it. The oblique Fracture more frequently ſlips out of its Place again than the tranſverſe, and generally leaves the Thigh ſomewhat ſhorter than the other, notwithſtanding the Surgeon has performed his Duty with Exactneſs: it is therefore neceſſary to uſe, in theſe Caſes, beſides the Means to be hereafter mentioned, a more ſtrict Bandage than in the tranſverſe Fracture, to prevent the replaced Bones from being eaſily moved.

How a Fracture of the Thigh is to be ſet.

II. In reducing a fractured Thigh, we are to conſider whether the Bone be broke near its Neck, or in ſome other Part; which Conſideration is always very neceſſary for the better replacing and binding up the Limb: whenever, then, a Fracture of the Thigh-bone happens, either in the Middle or towards its lower Head, it is to be extended and replaced with the Hands, like other Fractures; excepting that the extending Force here required, eſpecially in robuſt Patients, muſt be much greater: therefore more and ſtronger Aſſiſtants are to be here employed, who are ſufficiently to extend the Limb with their Hands; or, where their Hands will not ſuffice, Slings, Napkins, or linen Bandages may be bound round each Head of the Thigh, whereby the fractured Bone may be extended both Ways, while the Surgeon cautiouſly reduces the Fracture with his Hands, and treats it with a proper Dreſſing.

The Girt or Belt of HILDANUS.

III. But when the Extenſion cannot be performed effectually by the Hands, Slings, nor Bandages, which is a Caſe that ſeldom happens, we muſt then have Recourſe to the Belt or Girt of HILDANUS, *Tab.* VIII, *Fig.* 17, which is to be drawn and buckled very tight above the Knee, being firſt introduced through the Eyes of the Hooks A A, upon which is to be faſtened a ſtrong and ſmall Rope B B; about the Middle, C, whereof are to be applied the Hands of the Aſſiſtants, or Napkins, *&c.* by which Means a ſufficient Extenſion may be made, in order to replace the Fragments in their former Situations: nor is this Contrivance reſtrained to the lower Limbs only; for it may be applied upon Occaſion, to extend Fractures of the *Humerus* and *Cubitus:* if a fractured *Cubitus* is to be extended, the Girt is to be faſtened above the Hand; if the *Humerus*, above the Elbow.

Of the compound Pulley, or *Polyſpaſton.*

IV. If the laſt Method of Extenſion ſhall prove ineffectual by itſelf, it ſeems every Way neceſſary to try if any Thing can be done more to the Purpoſe by the Pullies of *Tab.* VIII, *Fig.* 15: the Hook A, of one Pulley, is to be faſtened upon the Rope of *Fig.* 17, at its Part, C; the Hook of the other Pulley B, is to be hung upon the Ring A, of the Hand-ſcrew B, of *Fig.* 16; which is to be firſt ſcrewed tight into ſome Beam or Rafter: then, the Patient being held firm, about the other Head of the fractured Limb, by Means of Slings, Napkins, or other ſtrong and long linen Bandages, to prevent his giving way to the Extenſion; the Rope C, put through the Pullies D and E, of *Fig.* 18, muſt now be drawn through, till the Thigh-bone be ſufficiently extended, ſo as to admit of a convenient Reduction thereof by the Surgeon: here it is to be obſerved, that the more Wheels the Rope paſſes round in the Pullies D, and E,

of

of *Fig.* 18. the more eaſily and gradually will the Extenſion be performed, inſomuch that by this Inſtrument one Man may draw more than ten without it.

A Fracture in the Neck of the Thigh-bone.

V. When the Neck itſelf of the Thigh-bone is broke, to which, from its oblique or tranſverſe Direction, and ſpungy or brittle Subſtance, it is very ſubject; it makes a Fracture not only very difficult to reduce, but ſuch a one alſo as can be ſeldom cured without leaving the Limb lame or ſhorter than the other, as HILDANUS (*Cent.* V. *Obſ.* 86.), RUYSCHIUS, and others teſtify. Now the Reaſons for this Calamity are more than one: for (1) the Fragments cannot, but with great Difficulty, be preſſed into their right Places, by Reaſon of the great Thickneſs and Strength of the Muſcles which cover them: (2.) it ſeldom happens that the Bones can be retained in their natural Poſition, after they have been very well ſet; becauſe the Muſcles, which paſs over and are inſerted a little below the Neck of this Bone, draw its lower Part upwards; and both theſe happen generally the more eaſily, (3.) becauſe of the oblique Poſition of the Neck of this Bone, which is inſerted into its Head in a Direction not perpendicular nor parallel, but as it were ſloping on one Side of the ſame, as will evidently appear upon viewing this Bone in a Skeleton; ſo that we have hence none of us any Occaſion to wonder, if Lameneſs and other bad Accidents follow as Conſequences of this Kind of Fracture.

The Difficulty of diſcovering a Fracture in the Neck of the Thigh-bone.

VI. To the foregoing Reaſons we may add, (4) that it is very difficult to diſcover when the Neck of the Thigh-bone is fractured, the Caſe being almoſt always taken for the Head of the ſame Bone being ſlipped out of its *Acetabulum* or Socket: till firſt PAREY (*Lib.* XIV. *Cap.* 21.) then SCHENCKIUS (*Obſ.* XI, *Lib.* 5.) after them the celebrated RUYSCH [a] (when the Obſervations of the two former were forgot) and, ſince him, ſeveral other eminent Surgeons and Phyſicians [b] have made it very evident that the ſpongy Neck of the Thigh-bone is, and may be oftener broke in two, than its Head, defended by very ſtrong Ligements, be puſhed out of its deep Socket by any external Violence. Of this conſiderable Obſervation, the Phyſicians and Surgeons of not only the former, but even the laſt Age, were ſo generally ignorant, that they never in the leaſt ſuſpected the Caſe to be a Fracture, but treated the Patient as if the Thigh had been luxated, tormenting and miſerably diſtorting the Member with the Machines uſed in that Caſe: ſince, therefore, this Method of treating the Patient has been found by Experience to be not only fruitleſs, but barbarous; it is highly neceſſary we ſhould recommend another Practice, and ſuch as might prevent thoſe acute Pains, violent Inflammations, and many dangerous Symptoms which might otherwiſe enſue.

How this Kind of Fracture is to be diſcovered and cured.

VII. When we think the external Force to have been ſufficient to produce a Fracture; when the Patient cannot bear any Streſs upon the Limb by ſetting his Foot to the Ground; when very acute Pains are felt about the Articulation itſelf; and when we find the affected Limb ſhorter than the ſound one, it being an eaſy Matter to turn the Foot almoſt round from one Side to the other, and perceive any Cracking or Grating of the Bones in that Motion, we may then reaſonably ſuppoſe that the Neck of the Thigh-bone is fractured, we may then reaſonably ſuppoſe that the Neck of the Thigh-bone is fractured: we muſt

[a] In *Theſaur. Anat.* VIII. *Tab.* III. *Fig.* 1. and *Theſ.* IX. *Tab.* I. *Fig.* 1.

[b] CHESELDEN, *Anatom.* upon the Bones of the lower Extremities, and in *Tab.* VI. G. H. DOUGLAS, *Philoſoph. Tranſact. N* CCCLXXXI. Ann. 1716; and PETIT, on *Diſeaſes of the Bones.* SALTZMAN, *Diſſert. de Fractura Femoris frequentiori*, and others.

then carefully avoid the violent Extenſion of the Limb, which was uſed formerly under the Notion of a Luxation, by the Inſtruments contrived by SCULTETUS, and others, for that Purpoſe: our Buſineſs here is, to extend the Limb very gently and gradually, till the diſordered Limb be of the ſame Length with the ſound one; and this by means of a Napkin, proper Slings, or the Hands of a ſtout Aſſiſtant faſtened round the Foot, or elſe by the preceding Girt and Pulley; in a Manner by which we may be able to rejoin, in ſome Meaſure, if not perfectly, the Neck of the Thigh-bone with its Head ſtill firmly adhering in its Socket: and though a Shortneſs of the Limb, or Lameneſs, is generally left behind after this Fracture; yet becauſe there are ſome cured without thoſe Attendants, I muſt approve, as very uſeful, ſuch a ſtrict Bandage as may apply and retain the Neck to the Head of the Bone, ſo as that they may gradually grow together again: for which Purpoſe, we uſually apply the Bandage called *Spica inguinalis*, in this Caſe; then a large and broad linen Cloth or Napkin is placed between the Thighs, to keep the Body of it from ſubſiding; and laſtly, Ligatures are put about the Knee and Ancle, whereby the Foot is faſtened to the lower End of the Bed, with a little Pad of Straw, to prevent the Limb from being contracted upwards: but we ſhall deſcribe all this more at large, when we come to the Doctrine of Bandages: indeed PETIT teaches, that this Kind of Fracture is to be bound up ſimply in the ſame Method with other Fractures of the Thigh; but that this is not reaſonable, the experienced herein will readily allow: having proceeded thus far regularly, and placed the Patient in as convenient a Poſture as poſſible, we muſt all along obſerve, with a ſtrict Eye, whether the afflicted Member be either equal or ſhorter than the ſound one. If it ſhould be found to become ſhorter, there will be great room to ſuſpect that the Neck of the Thigh-bone is ſlipped out of its Place again: and therefore it muſt be gently extended again, after unbinding of it, till it becomes of the ſame Length with the ſound one as before: but when the Foot of this continues of the ſame Length with that of the ſound Limb, there is great room to hope that the Patient will be happily cured; if continued Reſt and a proper Diet be regularly obſerved: what remains, is to be left to Nature.

How ſuch a fractured Thigh is to be retained in its proper Situation.

VIII. If we had an Inſtrument that would keep the fractured Thigh properly extended, and of the ſame Length with the ſound one, for about fourteen Days, or till the Cure was perfect, we could go on with much more Certainty and Succeſs, in the Cure of Fractures in the Neck of the Thigh-bone, than we do. He therefore would be Author of a no ſmall but important Advantage that ſhould contrive a Machine fit for this Purpoſe: for though HILDANUS has deſcribed (*Cent.* V. *Obſ.* 86.) an Inſtrument proper for extending Thighs which are obliquely fractured; there is yet great room to doubt of its Fitneſs for this Kind of Fracture: for he does not, that I know of, ſupply us with any Inſtances of Extenſions or happy Cures that have been made by this Inſtrument: but till we have a more proper Machine contrived, and when the other Means are not found of themſelves ſufficient, it will not be amiſs to uſe the forementioned Inſtrument of HILDANUS; or, when that is alſo of itſelf inſufficient, to add the Straw-pad, the large four-headed Bandage, and the reſt of the Apparatus deſcribed by HILDANUS; or to bind two long Napkins about each Groin, faſtening them by Nails or Rings to the Head-bed-poſts or Sides, ſo as to retain the Patient's Body ſufficiently firm from deſcending: but that the lower

Part

Part of the Limb may not give Way upwards, a Ligature or Bandage is to be put round the Knee and Ancle, to be faſtened to the Bed's Feet, as we obſerved at § VII. by which Means the Limb may be retained in its proper Poſture till the broken Neck of the Thigh-bone be joined firmly together: the ſame Method of binding and retaining may be alſo uſeful in other Fractures of the Thighs, but it is found not only uſeful, but really neceſſary in oblique Fractures of this Limb: but to prevent the Napkins or Ligatures from galling the Groins, it may be ſometimes proper to interpoſe ſoft Compreſſes or Lint; and for Advice concerning the proper Poſture in which a broken Thigh is to be retained, beſides what has been briefly ſaid at *Chap.* I. *N.* XXXVI. we ſhall be more full and particular in the Doctrine of Bandages [a].

A Fracture of the Thigh with a Wound.

IX. If a Fracture of the Thigh be accompanied with a Wound, it makes the Caſe very dangerous and difficult to cure: and if theſe Accidents ſhould happen to be inflicted on the neighbouring Joint, Death is generally the Conſequence; more eſpecially when any of the large Blood-veſſels are wounded, as muſt be evident from the great Hæmorrhage. So alſo is the Fracture dangerous, when the Wound is ſeated in the back Part of the Thigh; becauſe it is with great Difficulty to be cleanſed and dreſſed.

Cure.

X. In theſe Fractures with a Wound, the eighteen-headed Bandage, *Tab.* IX. *Fig.* 4. is to be uſed for the Dreſſing: this is deſcribed at large in our third Part, upon Bandages: but, if the wounded Part be very much contuſed, ſo that extravaſated Blood be lodged under the Skin and about its Interſtices, it is to be carefully opened by ſeveral Inciſions of a ſufficient Depth, that the extravaſated Blood, which would, in a ſhort Time putrify, may be by this means diſcharged: the injured Parts are to be afterwards waſhed with *Aq. Calcis* mixed with a fourth Part of *Sp. Vin. Camph.* or ſome ſuch reſolving Liquor, till the contuſed Parts are digeſted.

When accompanied with Hæmorrhage.

XI. When this Kind of Fracture is accompanied with Loſs of Blood, which is not very violent, nor the Bone near, the Wound is to be dreſſed with dry ſcraped Lint, properly folded, ſo as to fill the Wound; more and larger Compreſſes are to be laid over theſe, and the whole is to be retained with a proper Bandage, as is uſual in Hæmorrhages: but if the Flux be greater, we recommend the Uſe of aſtringent Liquors, ſuch as are uſed to ſtop the Hæmorrhages of Wounds, eſpecially the moſt highly rectified Spirit of Wine, which is here found to be extremely ſerviceable and effectual; but if it run ſtill more vehemently, from an Artery, the Veſſel is to be firſt diſcovered by the *Tournequet*, and afterwards ſecured by Ligature: when this Kind of Fracture is attended with very great Hæmorrhage, and a violent Splintering of the Bone from Gun Shot, ſo as to indicate the crural Artery to be lacerated; if our Deſire is ſincerely to ſave the Life of the Patient, our beſt Method will generally be, to amputate the Thigh and tye up the Artery in Time: for by this means the Patient will be more eaſily preſerved, than if we ſtrive to ſave the lower Parts of his Limb; for the crural Artery is ſo large that it ſeldom grows together, and if it does, the lower Parts are ſoon ſeized with a Gangrene: after the Blood is ſtopped and the Wound cleanſed, the Fragments of the Bone may

[a] In ſimple Fractures of the *Femur* or *Tibia*, the eighteen-headed Bandage may be properly applied.

be replaced, and the Limb carefully bound up with Compreſſes, Splints, and the Bandage with eighteen Heads, defending it in a Caſe of Straw, by the *French* called *Fanons*. The Wound is to be afterwards unbound every Day, cleanſed from its Matter, and dreſſed with ſome digeſtive Ointment or vulnerary Balſam, till it be healed: Inſtances of Fractures of the Thigh with a Wound may be ſeen in SCULTETUS, *Obſ.* 77 and 78. and PURMAN, *Obſ.* 63.

An EXPLANATION *of the* EIGHTH TABLE.

Tab. VIII. explained.

Fig. 1. Is a Sort of large and ſharp *Forceps*, proper to cut off the Splinters or Fragments of Bones, which ſtick out: but to make them cut the eaſier, the Handles ſhould be two or three Inches longer than the Figure.

Fig. 2. Is a ſingle Hook.

Fig. 3. Is a double Hook, ſerving for various Purpoſes in Surgery and Anatomy.

Fig. 4. Is a Needle for taking up Arteries with a Ligature in Hæmorrhages, and many other Caſes: A, is its blunt Point, B, its Eye tranſmitting the Thread, C, its little Head.

Fig. 5. Is a Caſe to hold the ſubſequent Inſtrument, which is uſed to hold and apply the *Lapis infernalis*, or Cauſtic Stone.

Fig. 6. The Inſtrument itſelf, made of Steel, for holding and conducting the ſaid Stone: *a*, the Nippers which lay Hold of the Stone, *b*, the little Ring which ſhuts and holds them faſt upon the Stone, *c*, the other End of the Inſtrument uſed as a ſticking Quill to ſupport the Lips of Wounds.

Fig. 7. Exhibits the Figure of a Splint, made of thin Wood or Paſte-board, to be uſed in Fractures of the Arms and Feet: its Breadth ſhould be about three or four Fingers, and its Length ſuitable to the Size of the Limb.

Fig. 8. Is a Paſte-board Splint, ſuch as is ſometimes uſed in Fractures of the Noſe: its Size is to correſpond to that of the Noſe.

Fig. 9. Is a Splint of Cap-paper, ſuited to the lower Jaw, when fractured only on one Side.

Fig. 10. Is a double Splint of the ſame Kind, for the lower Jaw, when fractured on both Sides: it is to be applied ſo that the Aperture (*a*) in the Middle may let in the Chin: but its two Extremities or Wings (*bb*) which may be folded together in the Middle (*a*), are to be applied towards the Ears.

Fig. 11. Is a Compreſs, in Form of an X, to be uſed in Fractures of the Clavicle.

Fig. 12. Is a Paſte-board Splint, to be laid over the former Compreſs, in the ſame Fracture.

Fig. 13. Is an iron or ſteel Inſtrument in the Form of a T, uſeful to retain the Shoulders in a proper Poſture, in Fractures of the Clavicle: AA, its tranſverſe Part, to which are faſtened iron Rings, to retain and keep back the Shoulders: B, its perpendicular Part going down the Back: C, an Aperture in its lower End by which it is to be faſtened with a Ligature round the Waiſt, to be tied before on the Belly. *See Chap.* V. § V. foregoing.

Fig. 14. Is a Paſte-board Caſe, in which a fractured Arm is to be lodged after it has been ſet and dreſſed: its Size is to be anſwerable to the Arm.

Plate 8. Vol. I. P. 148.




Fig: 1.

Plate 8.




Fig: 5.



Fig: 6.



Fig: 7.




Fig: 8.




Fig: 9.



Fig: 10.



Fig: 13.




Fig: 14.



Fig: 15.






Fig: 16.




Fig: 11.



Fig: 12.



Fig: 17.



Fig. 15. Is a *Polyſpaſton*, or compound Pulley, uſed to extend Fractures, deſcribed before at *Chap.* VIII. § IV. A and B, are two Hooks, by which the Inſtrument is faſtened on both Sides. C, the Rope, by drawing which, an Extenſion is made upon the broken Limb. D, and E, are the two Pullies, conſiſting of ſeveral Wheels, by which the Force of the Drawer is very much increaſed.

Fig. 16. Is a ſtrong iron Screw, whoſe Worm or Thread B, is to be forced by the two Handles, into ſome Beam or Rafter; and upon its Ring A, is to be hung the Pulley E, foregoing.

Gig. 17. Is the Girt of HILDANUS, ſometimes neceſſary to make Extenſions upon the upper and lower Limbs: AA, two Hooks, upon which is hung the Sling or Rope BB; C being the Place where the extending Force is to be applied. See above *Chap.* VIII. *N.* XIII. The Girt ſhould be three or four Fingers wide, and a Foot and a half long.

CHAP. IX.

Of a FRACTURE *of the* PATELLA, ROTULA, *or* KNEE-PAN.

I. IN order the better to underſtand and cure a Fracture of the *Patella*, it is previouſly neceſſary to learn from Anatomy, the Manner in which it adheres by Means of Ligaments and Tendons to both the Leg and Thigh; where we may alſo obſerve, its Aſcenſion with the contracting Muſcles upwards in extending the Foot, its Deſcenſion upon bending the ſame, and the great Force it ſuſtains both Ways in violent Motions of the Body: when a Fracture of this Bone happens, from a Fall, Blow, or any other external Violence; the Courſe of the Fracture is either longitudinal, tranſverſe, or in ſeveral Directions at the ſame Time; but of all, the tranſverſe Fracture is moſt frequent: the longitudinal happens much ſeldomer, and is more readily cured; becauſe the Fragments in this Caſe, generally keep in their right Places [a]: but when the Bone is broke tranſverſely, and into many Pieces, the Caſe is uſually much more dangerous; for though the lower Part of the Bone keep in its Place, as being not annexed to any Muſcles, yet the ſuperior Part of the Bone is drawn upwards, by the very ſtrong Muſcles to which it is joined, which makes it very difficult to reduce and retain. The Nature of this Kind of Fracture.

II. The Diſcovery of this Kind of Fracture is uſually Matter of no great Difficulty; for it may be eaſily perceived, by the Eye and the Touch, whether the *Patella* be ſound or divided; and alſo, when it is divided, whether it be broken tranſverſely, longitudinally, or into many Pieces: whether the Fragments adhere to each other, or are ſeparated at ſome Diſtance. In examining this Fracture, forcible Flexures of the Knee are to be avoided as of no Service, but very painful and pernicious; becauſe by this Means, the Fragments are pulled further from each other, and PETIT gives an Inſtance of Death occaſioned A Fracture of the *Patella* eaſy to diſcover.

[a] Indeed, GARENGEOT (*Lib. de Inſtrum. Tom.* II. *pag.* 310.) thinks, that this Bone cannot be broke longitudinally; but that this Caſe ſometimes happens, has been ſhewed by PETIT, among many others, in his Chapter of a fractured *Patella*.

thereby:

thereby: but it ſometimes happens, through the Obeſity of the Patient, and the little or no Separation of the fractured Parts, that this Caſe is not ſo ſoon to be diſcovered as is otherwiſe common; but then the Fracture is alſo leſs dangerous; for the Juice of the Bone, of which the *Callus* is formed, cannot ſo eaſily inſinuate itſelf into the Articulation, whereby the Knee would become rigid and immoveable, which frequently happens in ſome Fractures of the Bone.

Prognoſis. III. It is generally a very difficult Matter to make a perfect Cure of this Fracture, as thoſe experienced herein have often found: for if we may believe Practitioners, the Joint is generally left either rigid, or at beſt its Motions are performed with Difficulty: for, beſides the Inſinuation of the oſſific Juice, which was deſtined to the Formation of *Callus*, into the Receſſes of the Articulation; the Mucilage alſo, which lubricates the Joint itſelf, mixes and indurates with it: ſo that the Bones of the Leg and Thigh being joined together like two Pieces of Wood with the ſtrongeſt Glew, the Joint becomes ſtiff, the Bones grow together and become like one: and this happens the more readily becauſe of the long continued Inactivity of the Joint till the Bone is united, which is extremely neceſſary in theſe, and eſpecially in tranſverſe Fractures; by which long Inactivity, the lubricating *Mucus* of the Joint generally grows thick and hard: but it alſo uſually happens, that the Tendon which ſuſtains the *Patella*, and chiefly directs the Motion of the Joint, is violently contuſed at the ſame Time, and from the ſame Cauſe with the Fracture of the *Patella*: upon which Account, alſo, the Motion of the Knee is greatly impeded or wholly deſtroyed. We therefore need not wonder that thoſe who have once broke one of their Knee-pans, ſhould be ſo ſubject to frequent Falls, and in Conſequence of them break the other; ſince the violent Contuſion of this Tendon always leaves an incurable Weakneſs in the Joint.

Cure. IV. With regard to the Cure of a fractured *Patella*, it muſt be attempted in the following Method: In a longitudinal or perpendicular Fracture, the Patient muſt be laid upon his Back, and extending his Foot, the Surgeon in the mean Time replaces the Fragments on each Side with both his Hands, binding them up carefully with the uniting Bandage; which muſt be applied here in the ſame Manner with that uſed in large Wounds of the Belly and Fore-head, which we have before taken Notice of, and ſhall deſcribe more largely in the Doctrine of Bandages. But when the *Patella* is broken tranſverſely, or into ſeveral Pieces, the Patient being put in the ſame Poſture and extending his Foot as before: the Surgeon is then carefully to endeavour to bring together, compreſs, and replace the Fragments of the Bone in their natural Situations, with the Palms of his Hands, Thumbs, and Fingers, retaining them firm with the Application of a Plaſter in Form of a half Moon (*Tab.* IX. *Fig.* 2.) or perforated (as at *Fig.* 3.) and then the Foot of the afflicted Member is to be bound up and placed ſo that it cannot be eaſily bent or otherwiſe diſturbed. We intend to be more particular on the whole Buſineſs of the Cure, in the Doctrine of Bandages: but notwithſtanding there are to be found ſeveral particular Machines invented by Surgeons for retaining this Kind of Fracture;

* SOLINGEN recommends an Inſtrument of this Kind in his *Surgery*, in the Chapter of a broken *Patella*: and in *Tab.* XV. *Fig.* 26. *Edit. Amſtel.* 1698, we find the Machine delineated.

GARENGEOT (*Lib. de Inſtit. Chirurg.*) has alſo deſcribed another; and we are acquainted with ſtill more.

they

they all ſeem to be much of ſuch a Make as to fall vaſtly ſhort of being ſufficient for the preſent Deſign: but to prevent the replaced Bone from being diſturbed or broken a-freſh, which is an Accident we find often happens; it muſt be carefully obſerved that the Patient do not any way exerciſe his Leg till after the Expiration of the ninth or tenth Week; for a Fracture of the Knee-pan is ſeldom ſufficiently united before that Time: and ſuch as uſe their Legs before that Time, generally halt in Walking, as RUYSCH (*Obſ.* 3.) obſerves. Further, upon this Kind of Fracture, the Obſervations which PURMANNUS has collected in his Surgery (*P.* iii. *C.* 21.) deſerve to be conſulted.

CHAP. X.

Of FRACTURES *in the Bones of the* LEG *and* FOOT.

I. THERE is but little new to be ſaid on Fractures of the Leg and its two Bones, the *Tibia* and *Fibula*, which has not been before obſerved here; ſo that there is no Occaſion for more than the general Directions, which we have before laid down, to be obſerved in the Cure of every Kind of Fracture: *viz.* that the broken Bones are to be properly extended by the Hands or Slings, and then accurately replaced; to be afterwards properly bound up, and retained in the moſt ſuitable Poſture: this I have further to obſerve, that ſometimes both the Bones, and at other Times one of them only are broken: if both, it ſeldom happens that each of them are broke directly in the ſame Place, but one of them a little higher than the other: if the *Tibia* alone be broke, it is eaſily diſcovered, it being placed ſo near the Skin: but if the *Fibula* alone, which is buried under ſo many Muſcles, the Fracture is not ſo eaſy to be diſcerned: and when only the *Fibula* is broke, the Patient is generally under much leſs Diſorder, in ſuch a Manner, that it frequently permits them to walk: but to obtain a proper Knowledge of the Diſpoſition of this Bone when it is fractured, the Calf of the Leg is to be graſped by one Hand, whilſt the other Hand moves the Foot; and in the mean Time the Hand which holds the Leg will perceive whether and where it is fractured. Fracture of the Leg.

II. If, as it frequently happens, a Fracture of the *Tibia* ſhould be accompanied with an external Wound of the Skin; this muſt be firſt well cleanſed, and the Splinters of Bone, with all foreign Bodies, removed: then the broken Bone, after a proper Extenſion, may be reduced into its right Place, the Hæmorrhage, if there be any, may be afterwards ſtopped, (as we ſhewed at *Chap.* VII. § XI.) and the Limb then be bound up firmly with the eighteen-headed Bandage, cut ſomewhat in the Form of a Book, as at *Tab.* IX. *Fig.* 4. which we ſhall demonſtrate more fully hereafter in *Chap.* VIII. of Bandages: but if any Fragments of the Bone ſhould ſtick out ſo as to obſtruct its Reduction, they ſhould be firſt removed by a Pair of ſharp *Forceps*, or a fine Saw, before any Attempt be made to reduce or bind up the Fracture: having proceeded rightly ſo far, the laſt Step is, to place the Limb in a Straw Caſe, or elſe in a braſs Frame (*Tab.* IX. *Fig.* 9.) purpoſely accommodated to retain Fractures of the *Tibia*; renewing the Dreſſing and Bandage daily, 'till the Wound be healed. Sometimes little Pieces of the Bone will be ſet at Liberty and expoſed to Sight

the Suppuration, in the Courſe of the Cure; which are to be then laid hold of, removed, and the Cure continued as before: an Example of a fractured *Tibia* with a Wound may be ſeen in SCULTETUS, *Obſ.* 82, and 8.

PETIT's Machine for theſe Practices.

III. A very uſeful and proper Machine or wooden Caſe for retaining the preceding Fracture has been alſo contrived and deſcribed by Monſ. PETIT, a celebrated Surgeon at *Paris*, firſt in the *Act. Acad. Reg. Pariſ. Ann.* 1718. and afterwards in his Treatiſe of *Diſeaſes of the Bones*, from whence GARENGEOT transferred it into his Book of *Chirurgical Inſtruments:* we chuſe to exhibit the Machine rather from the *Act. Reg. Pariſ.*[a] than from the Inventor's *Book on the Bones*, or GARENGEOT's of *Inſtruments*; becauſe in the two latter, the Inſtrument is repreſented only put together, and therefore may not be intelligible to ſome, as if exhibited in a double Light, according to the other: you have it therefore firſt whole or put together, in *Tab.* IX. *Fig.* 11. and then ſeparated into its component Parts at *Fig.* 12: the Baſis or principal Part of the Machine AA (*Tab.* IX. *Fig.* 12.) is to be gently put under the broken Leg (after it has been firſt ſet, the Wound properly dreſſed, the whole bound up with the Bandage of eighteen Heads, and defended with Splints tied on with three Strings, as is uſual.) The two lateral Parts of the Caſe BB, and its Front C, which ſerves as a Sole to the Foot, are faſtened together by the Hinges D D, and kept ſhut by the Hooks EE, as may be ſeen at *Fig.* 11. by which Means the Foot cannot ſlip or ſhake, but is held firm and eaſy to the Patient. FF is the lower Part or Foot of the Machine, ſerving as a Foundation to the reſt: at its End G G, it is joined by Hinges to the preceding Floor A A, whoſe ſloping Part ſlides under the Thigh: over the Floor A A, Pieces of ſtrong Tape or Ticken are to be nailed tight to the Sides, upon which the Limb reſts eaſier than upon the Plank or Board: the other Parts of this Caſe ſeeming to be very obvious from the Figure, we ſhall, for Brevity, omit any Explanation of them, and only obſerve that its Size is to agree with that of the Limb; but by reaſon of the vaſt Numbers of Fractures which happen in a War, and the great Scarcity and Cumberſomeneſs of theſe Machines at ſuch Times, the Camp Surgeons are generally obliged to ſubſtitute Caſes of Straw in the room of them. At every Dreſſing of the Limb, if PETIT's Machine be uſed, the Hooks EE are to be undone, and the three Sides opened; but when the Wound and Fracture are dreſſed and bound up, the Foot muſt be exactly placed and the Caſe faſtened as before.

Fractures of the Bones of the Foot.

IV. Laſtly, the Bones of the Foot, which compoſe the *Tarſus*, *Metatarſus*, and Toes, are equally liable to Fractures in the ſame Way with thoſe of the Hands: but by reaſon of the great Complication of Nerves, Tendons, Ligaments, and Membranes, Fractures in this Part are uſually attended with Wounds and the worſt of Symptoms, as Inflammations and Gangrenes: the Bones are to be replaced, and the Cure carried on much in the ſame Manner alſo; except the Difference of Bandage, which we ſhall explain when we come to the particular Doctrine of them[b]: This we may alſo obſerve in the general, that Fractures of the Feet, like thoſe in the Hands and Ankles, can ſeldom be ſo

[a] Tho' it is a great Pity that the Author has not there ſubjoined a particular Explication of his Figures by annexed Letters or Numbers; becauſe it is probable that ſome Parts will not be rightly underſtood by many.

[b] See LE DRAN's *Chirurgical Obſervations*, 108.

per-

perfectly cured as to leave no Stiffneſs nor Want of Motion behind, if they ſhould eſcape the Company of an Ulcer, *Caries*, or incurable *Fiſtula*; which laſt bad Symptoms are often to be remedied by no Means but that of amputating the Member, nor will even that always preſerve the Patient from Death; and the injured Part ſhould be carefully guarded againſt Inflammations and Gangrenes, by proper Medicines; particularly, Fomentations of Lime-water and Spirit of Wine camphorated: nor ſhould you by any Means omit Bleeding and the additional Application of internal Remedies: it is one's Intereſt, therefore, in violent Fractures and Contuſions of this Part, to give timely Intimation of the Danger to the Patient, or at leaſt to his Friends: leaſt the miſerable Condition of the Patient ſhould be afterwards raſhly attributed to ſome Miſconduct in the Surgeon, as they too often are: but if any body be deſirous of a larger Acquaintance with Fractures of the Bones, I muſt recommend to him the diligent Peruſal of the celebrated PETIT's Treatiſe on *Diſeaſes of the Bones*.

CHAP. XI.

Of Bones broken by ſharp pointed Inſtruments, which may be termed WOUNDS *of the* BONES.

I. HITHERTO we have been treating of Fractures of the Bones, occaſioned by blunt Inſtruments: it remains now that we conſider ſuch as are produced by ſharp ones, as Darts, Swords, Spears, *&c.* which may not improperly be called Wounds of the Bones; for which Reaſon few Writers have treated of them ſeparately: for theſe Weapons do not only cut aſunder and ſeparate the ſoft and fleſhy Parts, but do alſo the ſame to the hard Bones, which they divide ſometimes ſlightly, ſometimes greatly, and often they make a Solution equal to a Fracture: but theſe Wounds cannot be inflicted upon the Bones without being attended with a great Variety of Symptoms, which are often very grievous, according to the Size and Depth of the Wound, the Nature of the Part, and the Force with which it was inflicted; as whether the Violence be received in the Head, Noſe, Jaws, Fingers, Hands, Arms, Shoulders, Legs, or Thighs: as therefore the Knowledge of theſe Accidents is of great Importance, and as they require a ſomewhat different Method of Treatment from other Fractures, it was here proper to ſay ſomething in particular of the beſt Method to be taken for their Cure. Wounds of the Bones.

II. But before we proceed to the Method of Cure, it muſt be firſt obſerved, that ſuch ſlight Wounds as do not penetrate deep into the Bone, are generally not ſo very dangerous: eſpecially if we proceed regularly in the Cure, keeping the Bone covered as much as poſſible with its Integuments from the Action of the Air, and wholly reject the Uſe of fat or oily Medicines, as very prejudicial to the Bones: but when they penetrate deep, wholly divide the Bone and its adjacent Parts, or violently affect any Organ more directly neceſſary to Life, in the Head, Neck, *Spina Dorſi*, and Breaſt, with a Puncture or Diviſion of the larger Veins, Arteries, Nerves, and Tendons of the upper or lower Limbs: the Danger is then much greater, and the Cure more difficult, Death being often the Conſequence. Their *Prognoſis*.

The Method of Cure.

III. In the Cure of these Fractures by sharp Instruments, PETIT inadvertently advises, in his Treatise on *Diseases of the Bones*, though in other Respects a very good Surgeon, "That in this Kind of Accidents in the Bones, if the Solu-"tion be inflicted lengthways, the Lips of the Wound are to be closed to-"gether, and cured with the uniting Bandage; but such as are inflicted very "obliquely, or wholly transversely, are to be joined together by Suture and "the Bandage that has eighteen Heads:" but as this Method is unsuccessful in many Wounds of this Kind, and so might lead young Practitioners out of the Way, it will not be improper here to expound this Matter more fully, and set it in a clearer Light: indeed in the first Kind of these Wounds I do almost agree with him; especially when they are slight, as when the Skull is not wholly nor deeply penetrated, and without Contusion, nor the Brain much hurt, as we have observed in Wounds of the Head, *Chap.* XIV. *N.* II. But when the contrary of these obtains, we must proceed more cautiously, and in a Method very different, keeping the Wound open with Lint, cleansing it, and when cleansed, healing it with Balsams, as we have observed in treating of Wounds: for by a too speedy Closure of such Wounds, the most violent Symptoms, and often Death itself, have been frequently brought on. So also in the slighter Wounds of this Kind, which are inflicted obliquely or transversely, I do not approve, with PETIT, of using promiscuously the Suture and eighteen-headed Bandage: but on the contrary, instead of a general Use, I think them the most seldom necessary: for I have seen cured by others, and have often cured myself, many of those Wounds in the Bones without the Use of that Bandage or Suture. To make the Thing more apparent by Example; in oblique Wounds of the Head, Forehead, and *Cranium*, which are none of the violent Kind, the Parts may be retained and closed much easier by a Plaster and common Bandage, than by Sutures made with Needles and Thread, as PETIT seems here to direct; and still much less Occasion is there for the Bandage with eighteen Heads: but as I have said in the Chapter of *Wounds in the Head*, these are generally more easy to cure by agglutinative Powders, Balsams, and Plasters, whether the Bones wounded be the Jaws, Clavicles, Shoulder-blades, or in the upper or lower Extremities; but when the divided Part hangs down, so as not to be kept rightly rejoined to its opposite by these Means, the Suture then seems altogether necessary.

Wounds of the Finger Bones.

IV. If the Bones of the Fingers should be thus wounded, or wholly divided by a Sword, so as only to hang by the Skin and Flesh; I have happily cured them, without the Suture and eighteen-headed Bandage, in the following Manner: I first accurately replaced the divided Bone, and retained them joined together in that Posture by winding round a Slip of Plaster, then applied a Compress dipped in *Sp. Vin.* laying over little Splints of Paste-board for the Retention of the broken Bones in their right and natural Posture; and, lastly, I bound up the whole firm with a proper, long, and narrow Bandage, suspending the Hand in a Sling, hung about the Neck for that Purpose: this I left so for several Days, ordering nothing more than for the Patient to keep up to a proper Diet and Rest: at length I carefully undid the Bandage, and tenderly removed the Compress, but not the Plaster, still supporting the Finger in its right Situation; and after cleansing the Wound as well as it would admit, I dropped in some vulnerary Essence, and applying a fresh Compress dipped in

Sp.

Sp. Vin. bound it up again as before: thus it was again left for ſeveral Days more, and in about every three Days it was dreſſed in the ſame Method, till after the Space of about a Month it was quite firm and well.

Wounds of the Arm and Leg Bones.

V. If either of the Bones of the *Cubitus* is divided, it generally happens to be the *Ulna*, that being moſt expoſed to the Weapon in fighting; nor does it then require either the forementioned Suture or the Bandage: but the Wound being cleanſed, is to be treated with ſome vulnerary Eſſence or Balſam, and with Lint dipped in the ſame Eſſence; after which, are to be laid on in order the Plaſter, Compreſs, and Paſte-board Splints wetted with *Sp. Vin.* which are to be bound round the thick Part of the *Cubitus* near the Wound with a long Bandage, that, as they dry they may accommodate themſelves the better to the Figure of the Part: and laſtly, the Arm is to be ſuſpended in a Sling hung as uſual about the Neck: and thus dreſſing the Wound every other Day, or, in Proportion to the Diſcharge, every Day, a Cure may be brought about without any Suture, which I here judge to be pernicious: but when either of the Bones of the Leg are broke, I do then indeed uſe the Bandage with eighteen Heads, as in other Fractures of the Leg and Thigh; but ſcarce ever the Suture: becauſe there is ſeldom or never Occaſion for it in Fractures of the *Tibia* alone, which is covered with ſcarce any Thing more than the Skin: and it is extremely rare that it is required in Fractures of the *Fibula*, unleſs ſome of its large Muſcles are divided: for we ſhould refrain from the Uſe of Sutures as much as poſſible, becauſe they generally excite Inflammation, Pain, Convulſion, and other bad Symptoms; ſo that we cannot approve of their Uſe, but in the greateſt Neceſſity where we perceive the Cure of the Wound cannot be effected without.

Wounds in the Arm and Thigh Bone.

VI. If the Thigh Bone ſhould be cut by a Sword, then the better to cloſe and retain thoſe ſtrong Muſcles, a Suture made with Needles and Thread, as in ſome other Wounds (*Book* I. *Chap.* I. § XXXIII. and XXXIV.) will certainly be of Service: the Wound is to be treated in the Method we have there taught, bound up with the eighteen-headed Bandage, and the Limb is to be placed carefully in a Caſe of Straw, as in other Fractures; ſo alſo if the Bone of the *Humerus* or Arm, ſhould be penetrated by a Sword, it ſhould, for the ſame Reaſon, be treated with the Suture as before; yet not dreſſed with the eighteen-headed Bandage, but a long and narrow one, as in other Fractures of the Arm: the Arm is afterwards to be ſupported by a ſhort Napkin, faſtened about the Neck; by which Means the Muſcles will be brought to a more ready Union, and the Cure ſooner and eaſier perfected: when we find the fleſhy Parts are united, the Threads muſt be cut, and drawn out, as in other Sutures: for the reſt, we proceed, as in all other Fractures of this Kind.

Of both the Bones of the *Cubitus* and Leg divided together.

VII. If it ſhould at any Time happen that both the Bones of the *Cubitus* or *Crus* are divided, ſo as to leave the Member hanging only by the Fleſh, Skin, and Blood-veſſels, (which is an Accident that very rarely happens without wholly amputating the Limb), then alſo the Suture with the eighteen-headed Bandage is to be applied; but the Suture can be of no Service when the Part is wholly or ſo far cut off as to hang by the Skin, its Nerves and Blood-veſſels being divided; eſpecially when the Part is ſo conſiderable as the Leg or Arm; for in that Caſe it is much the beſt to take the Limb quite off, to ſtop the

violent Hæmorrhage of the Veſſels, as in other Amputations, and to dreſs the Member in the ſame Manner.

Wounds of the Jaw-Bone, Clavicles, and *Scapulæ*.

VIII. When the lower Jaw is ſo cut by a Sword, that the Piece ſeparates much, and cannot be otherwiſe properly retained, then alſo the Suture muſt be brought into Uſe; adding a proper Balſam, Plaſter, Compreſs, and ſuitable Bandage: if the Clavicle, or *Acromion Scapulæ*, ſhould be in like Manner wounded by ſome ſharp Inſtrument, the Treatment and Bandage are to be performed in much the ſame Manner; gently unbinding, cleanſing, and dreſſing every other, or every Day, as we have obſerved in the reſt of theſe Accidents, till the Cure is perfected.

IX. The Hæmorrhage, which in theſe Injuries is often very large, muſt be ſtopped by Compreſſes, Aſtringents, or Ligature upon the Veſſels, according as which may ſeem moſt ſuitable to the Caſe: Gun-ſhot Wounds of the Bones are to be treated in the ſame Manner; or like Fractures. See further on this Subject, *Ch.* III. of *Gun-ſhot Wounds*; and in my Treatiſe of *Wounds of the Bones*.

An EXPLANATION *of the* NINTH TABLE.

Tab. IX.

Fig. 1. Is a Compreſs folded together by Degrees, called by the *French Compreſſe gruduée*, to be applied in Fractures of the Thigh, to make its ſmall Part towards the Knee of the ſame Thickneſs with its other, that the Splints may act more equally upon it by the Bandage.

Fig. 2. Are two lunar Plaſters, to include and hold firm the fractured Knee-pan after it has been ſet.

Fig. 3. A perforated Plaſter for the ſame Uſe.

Fig. 4. Is a Fracture of the Leg, with an external Wound A, to be bound up with the Bandage of eighteen Heads BBBB; which commodious Kind of Bandage ſeems to have been unknown to the Antients.

Fig. 5. Is a ſtraw Couch or Caſe for a broken Thigh, called by the *French Fanons*, the Letters AAAA denote two Sticks covered with Straw, bound on with ſtrong Packthread: to both Sides of theſe is alſo faſtened a ſtrong Cloth BB, of about two Feet broad, and three long: this Couch is uſually made twice the Length of the Thigh, ſo as to reach from the Groin and *Os Ilium* to the End of the Foot.

Fig. 6. Is a Sole of thick Paſte-board or Wood, fitted to the Size of the Patient's Foot: it is to be applied to the Bottom of the fractured Foot, and bound on by the three Tapes *a a a*, to retain or ſtay the Foot in its proper Poſture; whence CELSUS calls it *Mora*.


Fig. 7. Is a quilted Compreſs to be applied between the Foot and the Stay, to be ſoft, and defend it from any rough Action of Paſte-board or Wood.

Fig. 8. Is a ſoft linen Ring joined to the foregoing Compreſs, to let in and hold the Heel: it is to be faſtened to the Foot by the two Tapes *b b*.

Fig. 9. Is a braſs Trunk for ſecurely retaining a broken Leg: it conſiſts of three Parts ABC, which are joined by the Hinges 1, 2, 3, 4, 5, 6. The middle Part B is the Baſis or Chief of the Machine, which, like an hollow Pipe, receives the bound-up Limb: the outer Parts A and C are as moveable Lids

or

Fig. 11.






Fig. 12.






Plate 9.

Fig. 1.




Fig. 2.



Fig.




Fig. 4.






Fig. 5.



Fig. 13.






Fig. 6.

Fig. 7.




Fig. 8.



Fig. 10.

Fig. 3.



Fig. 9.





Fig. 10.



Fig. 11.








Fig. 12.







or Wings, which may be turned back or folded together: to each of these Lids AC, are joined three almoſt ſquare Loops EEE, through which are paſſed Tapes, to draw them tight together, and keep them firm upon the fractured Leg: its Size muſt agree with the Leg.

Fig. 10. Is a wooden Arch to put over a broken Leg, to keep it from being diſturbed by the Bed-cloaths, *&c.*

Fig. 11. Is PETIT's new Machine Caſe, (in *French Boete*) or a Box, for retaining a broken Leg, after it has been ſet and dreſſed as uſual: it is deſcribed fully at *Chap.* X. § III. of *Fractures.*

Fig. 12. Is the ſame in Pieces, to ſhew its Structure the better: the Letter M denotes the perforated Bracket, which receives the wooden Axle or Hinge II, that it may be elevated or depreſſed: the reſt are ſufficiently explained above at *Chap.* X. § III.

Fig. 13. Is a Compreſs folded at one End, to fill up the Small of the Leg, that the Splints may compreſs the more equally and firmly.

INSTI-

INSTITUTIONS OF SURGERY.

PART I. BOOK III.

CONCERNING LUXATIONS, or LUXATED BONES.

CHAP. I.

Of LUXATIONS *of the* BONES *in general.*

What a Luxation is.

I. HITHERTO we have treated of Fractures of the Bones: we ſhall now proceed to Luxations, or, according to CELSUS, Diſlocations; we ſay a Luxation or Diſlocation has happened when any Bone is moved out of its Place or Articulation, ſo as to impede or deſtroy its proper Motion and Office: ſo, for Example, we judge there is a Luxation when the *Acromion* is looſed from the Clavicle, when the Head of the *Humerus* is ſlipped out of the glenoeide Cavity of the *Scapula*, or the Head of the Thigh-bone puſhed out of its *Acetabulum* by ſome Violence, *&c.* So that it hence appears that Luxations are proper only to Bones that have moveable Joints or Articulations; but in a common Way of Speaking, People term it a Luxation when the Bones of the Noſe are diſplaced, or when *Epiphyſes* are ſeparated from their Bones in Infants, whereby they loſe their natural Uſe; and, indeed, CELSUS has ranked theſe Receſſes of the Bones from each other under the Claſs of Luxations.

What is required previous to their Diſcovery and Cure.

II. From what has been ſaid of Fractures, it may be eaſily concluded what is neceſſary to be done by thoſe who deſire to be happily verſed in the Knowledge and Cure of Diſlocations; firſt, that they ſhould have a clear Idea and Remembrance of the Form of each Articulation, with their Ligaments and Muſcles: which may be in ſome Meaſure obtained from accurate Figures in anatomical Books, but rather from a frequent and diligent Inſpection of the Skeleton and recent Bodies: for the Ligaments and Cartilages which are abſent in the bare Skeleton, may be fully obſerved, in their natural State, in a recent Subject.

III. Luxations

III. Luxations are generally diſtinguiſhed by Phyſicians into *perfect*, and *imperfect*. The imperfect conſiſt chiefly in this, that the Bones are here diſlocated or removed out of their Places but in Part, yet ſo as that they cannot perform their Office. Some are for diſtinguiſhing this Kind of Injury by the Name of *Subluxation* or *Diſtortion:* but the perfect Luxation is, when moveable Bones are wholly ſeparated or diſplaced from their Articulation with each other: as when the *Humerus* or Thigh-bone is removed quite out of its Socket: in both theſe Kinds of Luxations the Bone may ſlip out in ſeveral Directions; as externally or internally, behind or before, and above or below: another conſiderable Diviſion of Luxations is, into *ſimple* or *compound:* the latter, when, beſides the Diſlocation, there is ſome other bad Symptom, as a Wound, Fracture, Weakneſs or Straining of the Ligaments, Contuſions, violent Inflammation, or the like; but in the firſt there are none of theſe: the laſt Diviſion of theſe Injuries is into *recent*, or juſt inflicted, and *inveterate*, or of ſome Standing. The more free and moveable the Bone is in its Articulation, the more ſubject and eaſy to be diſlocated: ſo that it is no Wonder if the Bones of the Arm are oftner diſplaced from their Articulations with the *Scapulæ* than thoſe of the *Cubitus* and Wriſt, and the *Vertebræ* of the Neck and Loins oftener than thoſe of the Back. Of the ſeveral Kinds of Luxations.

IV. What we have been ſaying is in common to all Diſlocations: but it remains that we deſcribe every particular Kind of Luxation, beginning with the Head. We may ſuppoſe the Head to be luxated when (1.) the Bones of the Noſe gape; or (2.) when the lower Jaw ſtands in or out further than the upper; but it cannot be eaſily ſhoved out backward, becauſe hindered by a Protuberance of the *Os Petroſum*; or (3.) when the Head with the *Vertebræ* of the Neck are diſtorted to one Side, as it may have been ſometimes obſerved by the Surgeon; or, laſtly, (4.) when the Bones of the *Cranium* are forced apart by violent Pains, Fever, or Dropſy in this Part. Luxation of the Head.

V. Tho' all the *Vertebræ* which compoſe the Spine have a proper Motion, they are none of them eaſily removed wholly out of their Places, ſo as to make a perfect Luxation: but the *Vertebræ* of the Neck are much eaſier diſplaced than the reſt, becauſe ſmaller and more moveable; tho' theſe are generally connected very cloſely and ſtrongly to each other and the larger *Vertebræ:* ſo alſo the *Vertebræ* of the Loins are extremely difficult to diſlocate, tho' more moveable than thoſe of the Back, being ſeparated by thicker Cartilages, and without Sinuſes. Laſtly, the *Os Coccygis* may be ſhoved outwards in hard Births, and is ſometimes diſplaced and bent inward by a Fall, or the Force of ſome other hard Body; by which means it preſſes on the *Rectum*, and very bad Symptoms follow. Luxation in the Spine.

VI. As the Breaſt is made up of various Bones, ſo it is alſo ſubject to various Luxations: thus the Ribs may by ſome violent Blow or Fall be ſhoved from their Articulation with the *Vertebræ* into the *Thorax*, to the great Damage of the Breaſt and Lungs: ſometimes it happens that the enſiform Cartilage, at the Bottom of the *Sternum*, is depreſſed or thruſt inward by ſome Violence, ſo as to greatly afflict the Stomach [a]. The Clavicles are alſo ſometimes diſlocated Luxations of the Breaſt.

[a] See CODRONCHIUS, *of Diſeaſes reigning at Imola*, &c. An. 1602. and a new Diſtemper, called, *The Depreſſion of the enſiform Cartilage. Bonon.* 1603.

at

at one or both their articulated Heads, those joined to the *Sternum*, and those to the *Scapulæ*, but mostly the first: which, whenever it happens, the Arm hangs down unsupported, and its Motion is obstructed.

Luxation of the *Humerus.*

VII. If any one Bone is to be easily dislocated, it is that of the *Humerus*, partly because its Head is not lodged in any deep *Sinus*, and partly from its very ample and free Motion: it may be forced out either before, behind, or downwards; but never upwards, without breaking the *Acromion* or coracoide Process; for these confine down the Head of the *Humerus* very firmly above.

Luxation of the *Cubitus.*

Though the *Cubitus* does, at Times, undergo various Luxations, it can seldom happen, unless the Violence be great: and then it usually suffers only an imperfect Luxation; from the Shortness of the Ligaments, the Deepness of the Articulation, and its external as well as internal Defence with Ligaments: for the *Cubitus* to be luxated forwards is hardly ever the Case, being prevented by the Protuberance, *Olecranon:* but then it easily and frequently slips out backwards; as from duly considering the Articulation will be very apparent.

Luxation of the Hand.

VIII. The Wrist is very seldom dislocated from the Bones of the *Cubitus*, and hardly ever suffers more than an imperfect Luxation, from the Shortness and Strength of its Ligaments: but if it should be luxated, it will much easier slip out backward and forward, than inward and outward: the Reason of which is not difficult; for there is a bony Process on each Side the *Carpus*, where it is articulated to the *Radius* and *Ulna*, which defends it from being easily displaced: sometimes the small Bones of the *Carpus* are subluxated among themselves, whence generally arises an Extension and Stiffness in the Hand: in like Manner may the Bones of the Fingers be displaced; but then they are more easily reduced and cured.

Luxation of the Thigh.

IX. Among Luxations of the lower Extremities, that of the Thigh-bone comes first to be considered. The Head of the Thigh-bone may be forced out either upwards, downwards, forwards, or backwards: but which of these Ways it happens to be displaced, may be determined generally from the Touch, and from the different Direction and Length of the Limb: what we have before taken Notice of (*Book* II. *Chap.* VIII. *N.* VI.) is also here worth fresh Observation; *viz.* that the Head of this Bone is not near so often pushed out of the *Acetabulum* by some external Violence, especially in grown Persons, as is commonly suspected: for the modern Surgeons, contrary to their strong Opinion of a Luxation, have generally found a Fracture in the Neck of the Thigh-bone: nor is this to be wondered at; since the Head of this Bone is articulated into so deep a Socket, and secured by such strong Ligaments, that it cannot be dislocated in a dead Subject by the strongest Man, or other Violence: whereas, on the contrary, the Neck of this Bone is found to be very small, infirm, and brittle: so that it will be much easier for the Neck thereof to be broken, than its large Head to be forced out of its Socket: the Reason why this Fracture has been so commonly taken for, and treated as a Luxation, seems to be owing to the close Concealment of this Part by so many thick Muscles; and on that Account the *Trochanter Major* has been frequently mistaken for the Head of the Thigh-bone.

A Luxation of the Thigh-bone usually happens from an internal Cause.

X. From what has been said, we may perceive the Reason why the antient Surgeons had generally such bad Success in reducing this their supposed Luxation, scarce ever making a Cure without laming the Patient: to say nothing

of the Torture and bad Conſequences of their improper Extenſions by Machines: they thought their not being able to reduce theſe Luxations was, becauſe they could not make an Extenſion ſtrong enough to overcome the robuſt Muſcles of this Part: upon which Account, they invented all Sorts of Pullies and ſtrong drawing Machines, whereby they might extend and draw with the greateſt Force: Figures of which may be ſeen in SCULTETUS's *Armamentarium:* but as the Bone was not diſlocated, but fractured, all the Good they did the Patient was little elſe than exciting violent Pain, Convulſion, Inflammation, Abceſs, and other grievous Symptoms: for nothing is more certain than that a true Luxation of this Bone from external Violence was ſcarce ever at the Bottom of any of their Caſes, which they, as ſome now do, ſuſpected to be ſuch: for it is ſcarce poſſible the Head of this Bone ſhould ſlip out of its Socket; unleſs ſome great Weakneſs or Relaxation of its Ligaments, and a Congeſtion of morbid Humours between the Joint has happened ſome Time before, by which Means this otherwiſe very ſtrong Ligament may, by Degrees, be ſo elongated and relaxed, as eaſily to give Way to ſome future external Force, which is obſerved to happen in Children rather than Adults: in theſe young Patients the Head of the Thigh-bone generally falls inwards toward the great Aperture of the *Os Pubis*, and can ſeldom be replaced: it moſt commonly adheres to that Bone; and the Children who labour under this Complaint, if properly ſupported by ſtrengthening Medicines, are yet able to walk, though not without limping.

Luxation of the *Patella* and Knee.

XI. A Diſlocation of the Knee-pan is ſeldom diſcoverable by an unſkilful Surgeon, eſpecially when the Motion of the Bone, from its natural Seat, is very eaſy and large: for if he be deſtitute of anatomical Skill in the Joint, there is great Danger of his treating it for a Diſlocation of the Knee, tormenting the Patient with Pain from an uſeleſs Extenſion: but ſuch as have before duly conſidered the natural Diſpoſition of theſe Bones, will readily perceive whether the Diſlocation be of the *Patella*, or of the Knee; for the Knee-pan is always puſhed either without or within-ſide the Joint: but for the Knee itſelf, though the Head of the *Tibia* may be forced on either Side that of the Thigh-bone; yet, as the Articulation is very broad and grooved, being defended and held faſt by exceeding ſtrong Ligaments, it never happens to be perfectly luxated.

Luxation of the Foot.

XII. The Foot, indeed, is not exempt from being puſhed out before or behind from the *Sinus* of the *Tibia:* but it cannot be diſlocated on either Side, becauſe prevented by the two Heads of the Bones which form the Ankle, unleſs they ſhould chance to be broke at the ſame Time: the lower Head of the *Tibia* may be ſometimes ſeparated by a great Force from that of the *Fibula*, and the Foot may at the ſame Time be diſlocated outwards, as we read in ſome Obſervations: CELSUS has treated of this Species of Luxation, *Book* VIII. *Cap.* 11. The Bones of the *Tarſus* are connected to each other by very ſtrong Ligaments, and ſo cannot be eaſily diſlocated: but they are ſometimes ſo violently ſtrained, as to occaſion moſt ſharp Pain, Convulſion, and *Sphacelus*, unleſs prevented by timely Aſſiſtance. Laſtly, the Toes are ſeldom luxated; but if they ſhould, they muſt be treated like the Fingers.

The Cauſes of Luxations.

XIII. The *Cauſes* of Luxations are either external or internal: the *external* are Falls, Blows, Leaps, Strugglings, and ſuch like: the *internal* are preternatural Congeſtions in the Articulations: as when morbid Humours gather and

relax the Ligaments, ſo as to make the Joint diſlocate of itſelf, or by a Force not much greater, as riſing up, walking, leaping, *&c.* a ſad Inſtance whereof, I ſaw in a Student at *Altorf:* and another Inſtance of the ſame Kind I remember in the Manager of a conſiderable Farm, who frequently diſlocated his Shoulder-bone from a very ſlight Motion of his Arm: the weaker Men are, the more ſubject to this Sort of Luxation: hence it is that the Bones in the Limbs of Infants are ſo eaſily diſtorted, and wholly ſeparated from their *Epiphyſes*, upon a Fall, or rough Handling: it is alſo worth obſerving, that ZWINGER (*Theat. Pract.* II. *pag.* 109.) knew a lame Woman that bore three lame Sons.

The Signs of Luxations.

XIV. Many and various are the Signs of Luxations of the Bones: as from (1.) the Want of Motion in the Joint: (2.) the Change of Figure or natural Poſture of the Limb: (3.) an unuſual Hollowneſs of Protuberance, there being always a Tumor on that Side where the Bone is out, and a Cavity on the other where it came from: (4.) from the Difference of Length in the Limb, which is uſually ſhorter when the Bone is diſlocated upwards, and longer, when downwards; or laſtly, (5.) from the Pains excited by the violent Diſtortion of the Ligaments: for unleſs the Diſlocation be ſpeedily and rightly reduced, it is ſcarce poſſible but there muſt follow violent Convulſions, Inflammations, *Sphacelus*, and Death itſelf, merely from the vehement Diſtention of the Ligaments: but when the Bone is gradually thruſt out of its Place from internal Cauſes, then there is ſcarce any Uneaſineſs perceived: in the mean Time, to make a more ready Diſcovery of Diſlocations in general, it may be very proper to have in Readineſs an univerſal Rule; *viz. That whenever the Head of any Bone is removed out of its Place, its other End will be diſtorted in an oppoſite Direction:* when the upper End of a Bone is thruſt inward, its lower one will ſtand outwards; and when the firſt is outwards, the latter will be bent inwards.

The Signs proper to particular Luxations.

XV. Though theſe common Signs of Luxations, with a Knowledge in the Mode of each Articulation, may be generally ſufficient to diſcover moſt Diſlocations (as, for Inſtance, where there is a Luxation, you will feel a Cavity and a *Sinus* by preſſing your Finger upon the Place; and, unleſs it be ſoon reduced, a Swelling and Inflammation will enſue); yet we ought not to be ignorant of ſeveral other Signs which are proper to ſome Luxations only: thus in a Diſlocation of the Lower Jaw, the Mouth gapes open and cannot be ſhut by the Patient: when one *Vertebra* is puſhed over another, all the Parts beneath it are deprived of Senſe and Motion: for none of the *Vertebræ* can be diſlocated in any Manner, without compreſſing or wounding the *Medulla*, which is tranſmitted through their Middle, in Conſequence of which, the Courſe of the Spirits through it and its Nerves to the lower Parts, will be either diſturbed or wholly intercepted: when one of the Ribs is diſlocated, the Breath is very difficult to be drawn, and other bad Symptoms of the like Kind ariſe: but to open at large the peculiar Signs of every other Luxation, is not the Buſineſs of this Place: eſpecially as they may be readily deduced from the Action of each particular Part where they happen.

Signs of imperfect Luxations.

XVI. A Subluxation or Strain may be diſcovered, when the Patient has ſuffered under ſome great external Violence, and the particular Joint is afflicted with Immobility and violent Pains, the natural Figure or Poſition of the ſame being little or nothing changed: but however, upon a more ſtrict Examination

tion of the Part affected, there may be almoſt always found ſome little Inequality in the Articulation or Limb.

XVII. Laſtly, Luxations which proceed from internal Cauſes may be known from the following evident Signs: (1.) the Limb is ſo much relaxed as to be eaſily turned about in any Direction: (2.) there will be a Cavity about the Place of the Articulation, and the Fingers will perceive a Hollowneſs upon preſſing them between the Bones, and a preternatural Swelling will appear in another Place: (3.) the Bone that has ſlipped out may be eaſily replaced, but then it ſoon falls out again of itſelf; ſo great is the Weakneſs of the Ligaments and Muſcles, that they are not able to keep the Bone in its right Place: hence, (4.) the diſlocated Limb will be longer than the ſound one: it is alſo (5.) generally not accompanied with any Pain, Inflammation, or Convulſion, as is uſual in other Luxations: laſtly, (6.) from the Seat of this Luxation, being generally in the upper Joint of the Thigh or Arm, and ſometimes in the Articulation of the Foot with the *Tibia*. Signs of Luxations from internal Cauſes.

XVIII. If any Surgeon deſires to be well ſkilled in the *Diagnoſis* and *Prognoſis* of Luxations, I adviſe him to be well verſed in the Structure and Difference of the Parts affected, as well as to compare the Caſe carefully with the ſeveral Cauſes and other Circumſtances of Luxations; for thus we find that *imperfect* and *ſimple* Luxations are reduced with much more Eaſe, and treated with much greater Succeſs, than ſuch as are attended with Wounds, Fractures, Convulſions, Inflammations, or the like: the Reduction is not only more difficult in Proportion to the Number of Accidents or Symptoms, but alſo as the Bones are more or leſs diſtant and ſeparated from each other; inſomuch, that the Bones cannot often be replaced, by reaſon of the Fracture and great Inflammation; or, if they are once reduced, it is very difficult to retain them in their Places, and perfect the Cure without Lameneſs, from the great Weakneſs of the Ligaments: which laſt is uſually the more certain in Luxations from internal Cauſes: but in Luxations that happen from internal Cauſes in very young Subjects, the lower Part of the Limb generally waſtes, and becomes altogether weak and flaccid: Luxations that have juſt happened are in the general much eaſier and ſooner cured than thoſe of long ſtanding: for in the latter there generally ariſes a Tumor with Inflammation, and the Juices gather in great Quantity, by which Means the Ligaments are extremely relaxed, or the Articulation ſo glewed up and obſtructed, that it cannot receive the Head of the Bone as before: nor is it unuſual for the Head of the diſlocated Bone, in an inveterate Luxation, to lodge itſelf in ſome new *Sinus*, on one Side its natural one; by which Means the Head of the Thigh-bone has adhered and grown to the external Part of thoſe of the Hips, or elſe to its *Acetabulum*; that Cavity itſelf being filled up with ſome preternatural and tenacious Juice: it may be obſerved in general, that Luxations are very ſeldom mortal, except in the Head, and in the *Vertebræ*. The *Prognoſis* of Luxations.

XIX. If any Bone be diſlocated in Infants, or ſeparated from ſome *Epiphyſis*, the Caſe is very dangerous, and uſually attended with very bad Conſequences: for (1.) the Head of the very ſoft and cartilaginous Bone is ſo diſtorted as to be ſeldom if ever reducible to its natural Figure: (2.) theſe Kinds of Luxations are uſually concealed by Maids and Nurſes, ſo that they do not come under the Care of the Parents or Surgeon, till it is too late: (3.) the Caſe of Infants Luxations in Infants.

rarely admits of a regular Extenſion and Reduction of the Part, from the Extremity of Pain, which they are incapable of bearing: (4.) when the Luxation in younger Patients has been of any ſtanding, the Head of the Bone is ſo enlarged, and the *Sinus* or Socket ſo contracted, that it is often impoſſible to reduce it: (5.) it may happen that the Surgeon, ignorant of the true Cauſe, will take it to be and treat it as proceeding from a Flux of Humors, often too violently extending thoſe ſoft and now cartilaginous Parts, and throwing them into ſome very bad Poſture: laſtly, (6.) Want of Skill in the Surgeon may be an Occaſion of the Bones not being happily replaced in Infants; for Nothing is more improper than the violent Diſtenſion ſome Surgeons uſe in theſe Caſes, whereby they ſeparate thoſe ſoft Bones and their *Epiphyſes* more from each other, and occaſion many bad Symptoms.

If attended with a Wound.

XX. Diſlocations attended with a Wound, eſpecially of the Shoulder or Thigh-bone, are of very bad Conſequence, and often endanger the Life of the Patient; in CELSUS's *Opinion*, *Book* VIII. *Chap.* 25. whether the Bones be replaced or not, there is generally great Danger; and ſo much the more, the nearer the Wound is to the Joint: HIPPOCRATES has declared that no Bones can be reduced with Security, beſide thoſe of the Hands and Feet: *Vectiar.* 19. 5.—See more on this Subject, in that Paſſage of CELSUS juſt now quoted: though I by no Means recommend the following him implicitly.

CHAP. II.

Concerning the CURE *of* LUXATED BONES.

The Cure of Luxations.

I. THE Method of treating Luxations of the Bones does pretty much agree with, and is in a great Meaſure the ſame with, that uſed in Fractures: for in Diſlocations, as in Fractures, the whole Deſign of the Surgeon is, (1.) to reſtore the luxated Bone to its Place, firſt by Extenſion, and then by Reduction with his Hands: (2.) to preſerve and retain what is ſo replaced in its natural Poſition: and laſtly, (3.) to prevent and cure the ſeveral Symptoms which uſually attend: the Reduction is uſed to be commodiouſly performed by placing the Patient on a Stool, Table, Bed, or the Ground, as the Surgeon ſhall think moſt ſuitable to the Caſe: it is however to be obſerved here, that thoſe Luxations are moſt readily reduced on a Stool, which happen in the Jaw, Clavicle, Arm, or Hand: on a Table, ſuch as happen in the *Vertebræ* or Thighs: on a Bed, ſuch as happen in the Legs or Feet: and laſtly, thoſe Diſlocations are moſt commodiouſly reduced on the Floor which happen on the Shoulders or *Vertebræ* of the Neck.

Of (1.) the Extenſion.

II. The Extenſion, as we obſerved, in diſlocated Bones, is to be made much after the ſame Manner as in Fractures: *viz.* the outer or lower Part of the diſlocated Limb is to be extended by an Aſſiſtant, till the Head of the diſordered Bone be found to correſpond exactly with the *Sinus* from whence it was luxated: this may be done by the Hands, but if they are not ſo convenient, the Extenſion ſeldom fails of being made ſo well by a Napkin, as to render the Machinery delineated in ſuch an ample Manner by ORIBASIUS, PAREY, ANDREAS A CRUCE,

CRUCE, SCULTETUS, and others, generally unneceſſary: ſince they can effect ſcarce any thing more, unleſs it be to terrify and diſcourage the Patient in the Extenſion, by their formidable Shew.

Of (2.) the Réduction.

III. To replace the luxated Bone again in its natural Seat, the Surgeon muſt regulate the Aſſiſtant's Extenſion, by ordering it to be ſtrong enough, and in a right Direction; in the mean Time he is to compreſs the Articulation gently with his Hands and Fingers, till he find the elapſed Bone recover its right Place.

How to know when the Bones are rightly replaced.

IV. An accurate Reduction of a Luxation is known to have been effected by the ſame Signs which have been before mentioned in the Doctrine of Fractures: it is a good Sign (1.) if the Bone be heard to ſnap or crack in its Reduction: (2.) when the diſordered Limb is found to be of the ſame Length with the ſound one: (3.) when the Pains grow leſs: or laſtly, (4.) when the Limb can perform its uſual Motion.

The Reduction is often to be delayed.

V. But as Fractures are often prevented from being directly ſet by being attended with Inflammation, Hæmorrhage, or Tumor; ſo alſo Luxations often cannot be ſafely reduced before thoſe impeding Symptoms are firſt removed, or at leaſt much abated, by a proper Treatment: (ſee *Book* II. *Chap.* II. § XI.) In ſuch Caſes alſo where the Luxation is accompanied with a Fracture, the Reduction muſt be put off till that is firſt ſet and joined: for the Extenſion cannot be ſafely attempted till the Fracture be well joined by a firm *Callus*.

How the luxated Limb is to be treated after Reduction.

VI. After the Bones have been puſhed into their Places from whence they were forced out, the next Buſineſs is carefully to retain them there: but Bones that are intire are much eaſier retained than thoſe that have been broken: for the latter cannot be contained in their right Poſture without ſtrict Bandage and Reſt; whereas there is in the firſt Caſe ſeldom much Occaſion for Bandage, or any great Reſt: for thus in freſh Diſlocations of the Jaw, Bones of the Fingers, Hands, *Cubitus*, and *Humerus*, the Bone may be immediately reduced without further Bandage or Reſt; becauſe they are generally held firm enough by their proper Ligaments and Muſcles: it ſeems rather more neceſſary to bend, extend, and gently move the Limb ſometimes, than to endanger its becoming ſtiff and immoveable by a long Inactivity: but when the Luxation happens in the lower Extremities, it ſeems better to let the Patient reſt a few Days in his Bed, moving the Limb gently as ſoon as he finds it capable, and afterwards he may riſe and walk cautiouſly with it.

Of an inveterate Luxation.

VII. On the other Hand, when the Ligaments have been much ſtretched by a violent and long-continued Diſtenſion, or have been rendered infirm by any other Means, it ſeems altogether neceſſary to make uſe of ſome proper Bandage, and to recommend Reſt to the Patient, till the Ligaments have regained their former Strength: but here it muſt be alſo carefully obſerved, to let the diſordered Articulation ſometimes have a little gentle Motion, by an eaſy Flexion and Extenſion of the Limb, to prevent any Stiffneſs or other bad Conſequence from ſuch a continued Reſt: in the mean Time, it may not be improper to moiſten the Bandages and bathe the Part well with *Sp. Vin. Aq. Hungar.* or ſome other warm and ſtrengthening Spirit, by which Means the Ligaments are uſed to become very firm and ſtrong: the Bandages themſelves ſhould be neither too tight nor too looſe: the Reaſon for which, we have given in *Book* II.

Chap.

Chap. I. § XXXIV. treating on Fractures. As for the Application of Plaſters, which has been ſuch a prevailing Cuſtom in theſe Caſes, they may be altogether omitted here, as in Fractures, without any Danger: they ſeem even to do more Service by their Abſence than Preſence.

Of the Symptoms of Luxations.

VIII. The Inflammations, Tumors, Pains, Convulſions, Hæmorrhages, and other ſuch Symptoms which happen before or after the Reduction of a Luxation, are to be treated and cured in the ſame Method with that we preſcribed before in the Cure of Wounds and Fractures, *Book* I. *Chap.* II. § XVII, XVIII. *Book* II. *Chap.* II. § I. But as ſoon as the Bones are replaced, the forementioned Symptoms generally vaniſh, by Degrees, of themſelves. When the Ligaments are very much weakened, it is extremely uſeful to bathe the Part, after it has been firſt well rubbed with hot Linen Cloths, with highly rectified Spirit of Wine ſet on Fire, uſing plentifully afterwards ſome ſtrengthening Spirit, (as at *Book* II. *Chap.* II. § IX.) and then binding it up with a proper Bandage: but if violent Pains ſhould remain, notwithſtanding the Luxation be reduced, there is Reaſon to fear that there is a Fracture along with it: we muſt therefore endeavour to be ſatisfied with regard to this Certainty; and if we find a Fracture, we muſt uſe our Endeavours to ſet it: but if you find no Fracture, you muſt perſevere in applying ſtrengthening Fomentations and highly rectified Spirits; for after conſiderable Diſlocations, (eſpecially in the Knee or Foot, which bear the Streſs of the Body) if they are not immediately reduced, the Pains often prove obſtinate, are of long Continuance, and require great Patience. If a ſlight Fever ſhould attend, Bleeding, a thin Diet, and cooling Medicines are to be uſed. If a Gangrene ſhould appear, which may ſometimes happen, it muſt be treated not only with the Medicines which we have before recommended, but alſo with Fomentations and digeſtive Cataplaſms, binding up with the eighteen-headed Bandage: for the reſt of the Symptoms, they may be treated as we propoſed, *Book* II. *Chap.* II. always taking Care to let the Diſlocation be reduced firſt. If a Luxation ſhould be attended with a Wound, we muſt make uſe of the eighteen-headed Bandage, and proceed with the reſt as we have directed in Hæmorrhages, *Book* II. *Chap.* II. in the Doctrine of Wounds. If in Diſlocations attended with a Wound, an Hæmorrhage enſue, you muſt proceed at the firſt Dreſſing in the Method above deſcribed, where we treated of Wounds, *Book* I. *Chap.* II. and the Wound muſt be healed with balſamic Medicines. If an Abſceſs ſhould be formed, it will be much the beſt to open it as ſoon as ever we find it to be ripe: for elſe there will be Danger leſt by the long Stay of Matter, it ſhould corrode the Articulation and Bones, and produce the worſt Kind of *Fiſtulæ*, which are often to be remedied by no Means, but that of amputating the Limb. When the Bones are diſlocated with ſo much Violence as to break and deſtroy the Ligaments, Tendons, and adjacent Skin; the Caſe is then, as HIPPOCRATES has obſerved, altogether incurable: for the more we ſtrive to replace them, the leſs Inclination have they to join again firmly; and by exciting Convulſions and a Gangrene, take off the Patient: therefore, whenever Luxations are attended with ſuch grievous Accidents as are certainly deſperate, if we would preſerve the Life of the Patient, we muſt of Neceſſity ſpeedily take the Member intirely off: if the Luxation is attended with a Fracture, then the Luxation muſt be reduced firſt, if poſſible,

and the Fracture is to be ſet afterwards: but, when this cannot be done, it will be proper to have Recourſe to what we have before obſerved on Fractures, *Book* II. *Chap.* II. § XI: if any Joint ſhould become ſtiff and immoveable, it will be proper to treat it in the Manner mentioned near the Place now cited: laſtly, if the Luxation be inveterate, and the diſlocated Bone, after the Uſe of Baths and emollient Fomentations cannot yet be reduced, it is much better to abſtain from violent Extenſions, and give it up, than to torture the Patient with exceſſive Pains, which might probably excite the moſt grievous Symptoms.

CHAP. III.

Of LUXATIONS *in particular; and on thoſe of the* HEAD *and* NOSE.

Diſlocation of the Head.

I. HAVING treated of Luxations in general, it remains that we conſider each particular Luxation by itſelf: we ſhall therefore begin firſt, with thoſe of the Head, and then deſcend to the reſt, as we did in expounding the Doctrine of Fractures: there are not wanting ſome who deem it a Luxation of the Head, when the Bones of the *Cranium* are ſeparated any Diſtance from each other; whether it proceeds from an *Hydrocephalus* in Infants, or from violent Head-achs, or ardent Fevers in Adults: but there is no room for us here to treat more largely on theſe Luxations: the Method of treating the firſt, we ſhall deliver when we come to conſider the *Hydrocephalus:* but as the other very ſeldom, if ever, happens, it ſeems to be curable by no other Method than that of Bandage and Compreſſion.

Luxation of the Noſe.

II. It ſometimes, though not often, happens, that the Bones of the Noſe are ſeparated from each other, or diſtorted out of their natural Places, by ſome violent Blow or Fall: when ſuch an Accident happens, it is ſeveral Ways diſcoverable: as (1.) by the Sight, when we behold the deformed Poſition of the Noſe: or (2.) by Feeling: or laſtly, (3.) by the Ear, when we perceive with what Difficulty the Patient draws his Breath through his Noſtrils: but as we before obſerved, theſe Luxations do but ſeldom happen: for the Bones of the Noſe are ſo firmly connected to the *Os Frontis* and other Bones, that they will ſooner break than ſeparate from each other.

Cure of a luxated Noſe.

III. When this Caſe happens, the Patient is to be ſpeedily placed in a high Chair, that an Aſſiſtant may ſtand behind and hold his Head firm, in a proper Poſture: the Surgeon is then to introduce with one Hand, a thick Probe, a Gooſe-quill, (as in the Caſe of a Fracture, *Book* I. *Chap.* III.) or little Stick ſhaped for the Purpoſe, up the Noſtril internally, by which Means the depreſſed Parts of the Noſe may be thruſt into their Places: in the mean Time he applies his other Hand externally, to guide and direct the Parts which are moved from within: this being done, there is ſcarce any thing elſe required but to let a bit of Sticking Plaſter lie upon the Noſe at the ſame Time: but if any Thing ſhould occaſion a Wound in the Noſe at the ſame Time, the Cure muſt be carried on in the Way which we propoſed before under a Fracture of the Noſe.

IV. If

IV. If a Tooth be removed from its Place by a Fracture of the Jaw, or any other Accident, it may be properly termed a Luxation: but this, when replaced, from the Teſtimony of many Writers, generally ſettles again in its Socket, and adheres firmly to the Gums.

CHAP. IV.

Of a DISLOCATION *of the* LOWER JAW.

How the Jaw may be luxated.

I. THE lower Jaw is indeed ſeldom luxated, becauſe it is held ſo firm by ſtrong Ligaments and Muſcles, by whoſe Aſſiſtance it is retained in two *Sinuſes* in the Baſis of the *Cranium:* but when it is by Accident forced out from thence, it may chance to be on one Side only, or elſe on both, it being then thruſt directly forwards: and this happens moſt frequently from opening the Mouth too wide in yawning: though it has ſometimes been occaſioned by a violent Blow or Fall: if it be luxated on both Sides, the Chin will incline downward, and the Jaw will be thruſt very forward: but if only on one Side, the Chin will be inclined toward the oppoſite Side; the elapſed little Head of the Jaw not being capable of Diſlocation but forward and inward [a]; for the Proceſſes of the Bones of the *Cranium* prevent the Jaw from being diſlocated backwards: hence it ſeems ſtrange that any one ſhould aſſert, contrary to the common Obſervations and Writings of the beſt Practitioners, that the lower Jaw may be luxated backwards as well as forwards: this is ſo inconſiſtent, that though he ſhould confirm his Opinion by Examples and Obſervations, it muſt be looked upon as the Conſequence of ſome Difference in the Articulation from what is uſual in Nature.

How to diſcover a Luxation of the lower Jaw.

II. The lower Jaw is chiefly known to be luxated on one Side, when the Chin is diſtorted on the oppoſite Side: for that Part to which the Chin inclines, is the ſound: but that from whence it recedes, is the luxated one: the Mouth in this Caſe gapes wider than uſual, ſo that the Patient cannot ſhut it, nor eat with his Teeth; the lower Range of Teeth being projected beyond, and on one Side the upper: but when the Jaw is luxated on both Sides, then the Mouth not only gapes wide and open, but the Chin alſo hangs down, and is thrown directly forwards: ſo that it is no Wonder if the Patient cannot ſhut his Mouth, ſpeak diſtinctly, or even ſwallow any Thing without much Difficulty.

Prognoſis. III. When the Jaw is out only on one Side, and the Caſe recent, the Cure is uſually not ſo very difficult; but when both Heads are diſlocated, and not preſently reſtored to their Places, it always occaſions the worſt of Symptoms, as Pains, Inflammations, Convulſions, Fevers, Vomitings, and at length, as HIPPOCRATES obſerves, Death itſelf comes on: and theſe Symptoms are the more violent, as the adjacent Nerves, Tendons, and Ligaments ſuffer a greater Extenſion: but if an expert Surgeon comes in Time, the Luxation is not very difficult to reduce.

Cure. IV. When this Kind of Luxation happens, the Patient is to be directly ſeated on a low Stool, ſo that an Aſſiſtant may hold his Head firm back againſt his Breaſt. Then the Surgeon is to thruſt his two Thumbs as far back into the Patient's Mouth as he well can: but they are to be firſt wrapped round in a Handker-

[a] See MONRO's Learned Diſſertation on this Subject. *Act. Edinb.* Vol. I. Art. II.

chief,

chief, to prevent them from ſlipping or being hurt; and his other Fingers are to be applied to the Jaw externally: when he has got firm Hold of the Jaw, it is to be ſtrongly preſſed, firſt downwards, then backwards, and laſtly upwards, but ſo as that they may be all done in one Inſtant: by which Means the elapſed Heads of the Jaw may be very eaſily ſhoved into their former Cavities: but the Surgeon ought to be always careful to ſnatch his Thumbs quickly out of the Patient's Mouth, leſt they ſhould be compreſſed, bruiſed, or bit, by reducing the Jaw into its Place.

Of the Jaw luxated on one Side.

V. If the Jaw be out on one Side only, every Thing muſt be done in the ſame Manner: but the luxated Side of the Jaw muſt be forced more ſtrongly downward and backward than the ſound one. Some ſay this Luxation may be ſometimes very readily reduced by a violent Stroke on the oppoſite Side of the Jaw: but this is a Method too pleaſant to be uſed with Safety in moſt Patients. As for Bandages, there ſeems to be no great Occaſion for them in this Caſe, unleſs the Luxation has remained ſome Time before it was reduced; for then it may not be improper to apply for ſeveral Days the four-headed Bandage, with ſome ſtrengthening Spirit, which may be taken off when the Patient intends to eat.

CHAP. V.

Of LUXATIONS *of the* HEAD *and* SPINE.

How the *Vertebræ* may be luxated.

I. THE Luxations which happen in the *Spine* and *Vertebræ* of the Back are generally imperfect ones. For it appears from an accurate Conſideration of the Structure and Articulation of theſe Bones, that none of the *Vertebræ* can be entirely diſplaced without being fractured, and alſo compreſſing or wounding the Spinal Marrow, which muſt produce Danger of inſtant Death. Even the imperfect Luxations of theſe Bones are very dangerous: which happen either between the two ſuperior *Vertebræ* of the Neck and the Head, or elſe between the reſt of the *Vertebræ*, when they are forced from each other.

Luxation of the Head.

II. Such as have a Luxation between the Head and upper *Vertebra*, ſeldom eſcape being carried off by a ſpeedy and ſudden Death: for in this Caſe the tender *Medulla* which joins immediately with the Brain, and is lodged in the Spine, the Brain itſelf, and the Nerves which ariſe beneath the *Occiput*, are too much diſtended, compreſſed, or lacerated; the two condyloide Proceſſes of the *Occiput* uſually ſlip out of their glenoide Sinus's in the firſt *Vertebra* of the Neck, when a Perſon falls headlong from a high Place, from off a Ladder, from on Horſeback, or when he receives a violent Blow upon his Neck: they dying very ſuddenly in this Caſe, are vulgarly ſaid to have *broke their Neck*, tho' there is generally no more than a Luxation: yet it ſometimes happens that the *Vertebræ* of the Neck are really fractured: if Life ſhould remain after ſuch a Luxation, which very rarely happens, the Patient's Head is commonly diſtorted with his Chin cloſe down to his Breaſt, ſo that he can neither ſwallow any Thing, nor ſpeak, nor even move any Part that is below his Neck: therefore, if ſpeedy Aſſiſtance be not had, Death enſues, from the Compreſſure or Hurt of the *Medulla*.

How the Head is to placed.

III. But to repulſe this unwelcome Meſſenger, the Patient is to be immediately laid flat upon the Ground or Floor: then the Surgeon kneeling down with his Knees againſt the Patient's Shoulders, is to bring them together ſo as to contain the Patient's Neck between them: this done, he quickly lays hold of the Patient's Head with both his Hands, and ſtrongly pulling or extending it, he gently moves it from one Side to the other; till he finds, by a Noiſe, the natural Poſture of the Neck, and the Remiſſion of the Symptoms, that the Diſlocation is properly reduced: by this Method the Surgeon retains the Patient firm between his Knees, and performs the Extenſion and Reduction between his Hands.

Another Method of Reduction.

IV. The ſame may be effected by another Method much like the former: as when the Patient ſits upon the Ground, his Shoulders being preſſed down, and his Head laid hold of under the Ears, and pulled ſtrongly but cautiouſly upwards, inclining it a little to each Side, till the Signs enumerated before (at § III.) demonſtrate it to be reſtored to its natural Place. If any of the other *Vertebræ* of the Neck ſhould be diſlocated, the Reduction is to be made in the ſame Manner: therefore there is no Occaſion to give them here a ſeparate Treatment.

PETIT's Method of Reduction.

V. But M. PETIT (*Lib. de Morb. Oſſ.*) rejecting the former Methods, has taught us another Way of reſtoring a Luxation of the Head, though he does not mention that he ever uſed it: he forms two Slings, having a large Opening about their Middle, as is delineated in *Tab.* X. *Fig.* 1, 2: The Patient lying on his Back, he takes the Sling, *Fig.* 1, and puts his Head through the Opening AB, which is made purpoſely large enough, and proportionable to the Size of the Head: the Part of the Sling A comes under the Patient's Chin, the Part B is placed under the *Occiput*, and the two Extremities of the Loop CC, come up over his Ears, the Ends D and E being the Parts by which the Extenſion is made: but to hold the Patient firm, he recommends another Sling, *Fig.* 2, through the Opening F, the Head is tranſmitted ſo as to make the Part of the Sling G come down his Back, and the Part H to come over his Breaſt, the two Extremities of the Sling I I, are to be joined together between the Thighs, and by this Means the Body is to be held from giving Way to the Extenſion made by the other: while the Head and *Vertebræ* of the Neck are kept ſufficiently extended by pulling theſe Slings in oppoſite Directions, the Surgeon endeavours to replace the luxated Bones: but, to ſay Truth, the preceding Methods ſeem to me to have the Preference: partly becauſe they are more ſimple, and performed without any Aſſiſtants or other Inſtruments than the Hands, which former are not always to be had; and partly becauſe the Patient may be relieved much ſooner by theſe Means; for while the Machinery is fetching or adapting, the Patient will, in all Probability, be dead. PETIT lays down no other Method of reducing this Luxation, throughout his whole Book, than by his Slings, not even how to aſſiſt the Patient in ſuch Caſes: whereas the Accident may happen very often in the Country, where ſuch Slings and Aſſiſtants cannot be had to help the Patient: in the mean Time, a Napkin, or long Slip of Linen of two or three Hands breadth, ſlit to let the Patient's Head through, will make a good Subſtitute for theſe Slings when they are not at hand.

What to be further done.

VI. But after any of the *Vertebræ* are replaced by any Method, it will be proper, in order to prevent a Tumor, and reſtore the ſtretched Ligaments of

the Neck to their former Vigour, to bathe it with *Aq. Hungar. Sp. Vin. camph.* or ſome other ſtrengthening Spirit applied warm, as alſo Compreſſes dipped in the ſame: the Patient ſhould bleed, and reſt gently for ſome Days, till the Neck be found ſufficiently ſtrong and well: as for Bandages, there ſeems to be little Occaſion for them here, unleſs it be ſuch as are deſigned to keep on the Compreſſes, dipped in ſome ſtrengthening Spirit.

Of Luxations in the other *Vertebræ* of the Back.

VII. With reſpect to the reſt of the *Vertebræ* of the Back, they are ſeldom moved quite out of their Places, unleſs they are fractured, they being retained, for the greateſt Part, by adhering to the adjacent Ligaments and Muſcles: therefore the Luxations which happen among them are uſually imperfect; no more being diſplaced than their two upper or lower Proceſſes, and they often but on one Side: and this happens ſometimes to one of the ſpinal *Vertebræ*, and ſometimes to more: but it is here to be briefly obſerved, that it is uſual to include among the Number of luxated *Vertebræ*, that which is ſound and firm, but intercepted by others which are not ſo: thus whenever the upper *Vertebræ* of the Loins from the laſt of the Back, and lowermoſt *Vertebræ* of the Loins next the *Os Sacrum* are luxated, we commonly ſay and reckon there are five *Vertebræ* out of their Places: when, ſtrictly ſpeaking, only the two outermoſt or the uppermoſt and lowermoſt of thoſe *Vertebræ* are diſturbed; the three middle Ones retaining their natural Situation and Connection.

How Luxations of the Spinal *Vertebræ* can happen.

VIII. If any one cloſely conſiders the natural Structure and Connection of theſe Bones, it will pretty evidently appear, that the ſpinal *Vertebræ* are not to be luxated but by ſome very conſiderable Violence: for, beſides their being moſt cloſely joined to each other by means of *Proceſſes* or *Apophyſes*, they are tied together and connected very firmly by exceeding ſtrong Ligaments and Cartilages: and this is the Reaſon why the ſpinal *Vertebræ* are not luxated, unleſs thoſe Cartilages and Ligaments ſhould break, in violently bending the Back, or receiving ſome great Blow or Fall thereon: for theſe Accidents are generally ſo far from ſeparating them, that they drive them more cloſely together: but if this ſhould happen from ſome very great Violence, it ſhatters the ſpinal *Vertebræ* and their *Medulla*, and quickly kills the Patient, as I myſelf have ſometimes ſeen: therefore, whenever a *Vertebra* is luxated without being fractured, the Body muſt, of Neceſſity, incline ſtrongly forwards, or on one Side: for in this Caſe, the ſuperior Proceſſes of the *Vertebræ*, by which they are faſtened to each other, will be ſeparated from the inferior Proceſſes, by which means the *Vertebræ* will be diſpoſed to be eaſily removed from each other: and they will incline towards the right Side, when the Hurt is on the left, and the contrary.

The common Signs of Luxations in the *Vertebræ*.

IX. The Signs common to Luxations in the *Spina Dorſi* are chiefly the following: the Back itſelf is found to be crooked or unequal, after the external Violence has been inflicted: the Patient can neither ſtand nor walk, and his whole Body ſeems to be paralytic: the Parts which are beneath the luxated *Vertebra* are nearly without all Senſe and Motion: the Excrements and Urine cannot be diſcharged, or elſe they are ſometimes emitted involuntarily; the lower Extremities grow dead by Degrees; and, at length, Death itſelf follows: but theſe Symptoms vary in Proportion to the Degree of Violence in the Luxation: for the more Diſorder the *Spina Dorſi* undergoes, the more grievous and dangerous will be the conſequent Symptoms.

The particular or proper Signs.

X. But what Number of the Spinal *Vertebræ* are luxated, muſt be judged of by the Degree of that preternatural Incurvation[a]: for where there is but one *Vertebra* luxated, the Curvature is gibbous, making a Sort of Angle: if the Proceſſes of the *Vertebræ* are diſplaced forwards, then the *Spina Dorſi* will ſeem to bend inwards; and the Patient will always have violent Pains upon bending his Body: on the contrary, when he lies upon his Back, the Pains will be more gentle: if the *Vertebra* is luxated on the right Side, the Body may be obſerved to incline towards the Left, and it will be eaſier bent on the right than left Side: if the *Vertebra* be luxated on the left Side, the contrary of all theſe Appearances uſually follows.

Prognoſis.

XI. If any one be deſirous to preſage the dubious Events of Luxations in the *Vertebræ*, I would have him remember that theſe Caſes are generally very dangerous and uncertain; and that, even when the *Medulla* is neither contuſed nor wounded, but from the Difficulty of reducing the luxated *Vertebræ*: and the more the *Vertebræ* are diſplaced, the more will the *Medulla* be injured, the worſe will be the Symptons that ariſe, and the more precipitate will be the Patient's End: the nearer the luxated *Vertebra* is to the Head, the greater and more extenſive is the conſequent Danger: for as Injuries are the eaſieſt to be inflicted upon the *Medulla* in thoſe Parts, ſo they are always of the worſe Conſequence: therefore Luxations in the Neck are always more pernicious than thoſe which happen in the Back; and thoſe in the Back are much worſe than thoſe which happen in the Loins: and what may ſeem wonderful is, that the Symptoms appear much milder in Caſes where ſeveral *Vertebræ* are luxated, than they do when there is only one; and ſtill much milder, when the Proceſſes on both Sides are diſplaced, than when only one of them are luxated: for in the latter of theſe Caſes, the *Medulla* is more compreſſed upon a leſs Space, as will appear evident to ſuch as carefully conſider the Structure of the *Spina Dorſi*: but then in ſlight Luxations the *Vertebræ* may be more eaſily replaced, and therefore Men may be often in leſs Danger of Death on that Account.

Cure.

XII. To make the Caſe no better than it is, Luxations of the Spinal *Vertebræ* are, in general, very difficult to reduce: the Artifices uſed by the Ancients were ſo foreign and inadequate to the Caſe, that they ſeem to have been uſed to no Purpoſe, proving rather a Torture than a Remedy. The following ſeems to be the moſt ſuitable Method of reducing Luxations of the *Vertebræ*: when the *Apophyſes* of the *Vertebræ* are diſlocated on both Sides, the Patient is to be laid leaning upon his Belly over a Caſk, Drum, or ſome other gibbous Body: then two Aſſiſtants are ſtrongly to preſs down both the Ends of the luxated Spine, on each Side; by which Means the Bones of the Spine will be ſet free from each other, lifted or puſhed up in the Form of an Arch, and ſo gradually extended: this done, the Surgeon preſſes down the luxated *Vertebræ*, and at the ſame Time nimbly puſhes the ſuperior Part of the Body upwards: and by this Means the luxated *Vertebræ* are ſometimes commodiouſly reduced into their right Places: but, if Succeſs ſhould not attend the firſt Time, the Method ſhould be repeated two or three Times more.

XIII. PETIT lays a thick Cloth rolled up like a Cylinder acroſs upon the Bed, and placing the Patient over it, treats him in the ſame Method which we juſt now propoſed: when the *Vertebra* comes out on one Side, the Patient is then

[a] See SAVIARD's *Obſ.* of Luxations of the *Vertebræ Dorſi*.

to

to be placed inclining in the prone Poſture now mentioned; but ſo that, when the left *Apophyſis* is diſplaced, one Aſſiſtant may preſs the lower *Vertebræ* inwards to the right, and another Aſſiſtant may depreſs the right *Humerus*, & *vice verſa*. For if there be any convenient Method of reducing the ſpinal *Vertebræ* when luxated, there can ſcarce be any more commodious than that here propoſed: and from hence I ſee it will appear evident, that the generality of thoſe Slings, Bandages, Pullies, Leavers, and other Inſtruments, which the antient Surgeons uſed to faſten about the Patient's Hips, Shoulders, and Breaſt, and are to be ſeen figured and deſcribed in ORIBASIUS, PAREY, and SCULTETUS, muſt be on every hand allowed to be ſo far from ſuitable for reducing theſe Luxations, that they muſt be generally pernicious.

XIV. For the Remainder, it ſeems proper, after the *Vertebræ* are reduced, to bathe the Spine with *Sp. Vin.* or to lay on Compreſſes dipped in *Sp. Vin. camph.* and to bind the Parts up with the *Napkin-and-Scapulary*. Afterwards the Patient is to be laid in a ſoft and even Bed: Bleeding, and bathing the weak Parts with ſtrengthening Spirits, are to be uſed as there may be Occaſion. The Bandage muſt be very ſeldom taken off, and all the Symptoms which happen in theſe Luxations are to be palliated as uſual, till the Cure is perfected.

CHAP. VI.

Of LUXATIONS *of the* Os COCCYX, *Ribs, and Clavicles.*

I. THE *Os Coccyx* may be thruſt inwards by a violent Fall or Blow, and it is often puſhed outwards in hard Birth. When this happens, it is uſually attended by violent Pain and Inflammation about the lower Part of the Spine, Abceſſes form in the *Inteſtinum Rectum*, and the *Fæces* are conſtipated or ſuppreſſed: to diſcover the Luxation of this Bone the more readily, we have Recourſe to the Uſe of our Hands and Eyes, as well as to the Knowledge of the forementioned Symptoms; nor is the replacing this Bone very difficult, if attempted by a careful and expert Surgeon [a]; for if it be thruſt outwards, it muſt be depreſſed into its right Place by the Thumb, after which may be applied Compreſſes dipped in warm Wine, or its Spirit, made broad above, and narrow below, to fill up the poſterior *Sinus* of the *Nates*: and theſe may be held on by the T Bandage of HELIODORUS, *Tab.* II. *Fig.* h: but that Part of this Bandage which comes between the Thighs, ſhould be ſlit and placed ſo that the Patient may go to Stool without undoing the Bandage, and to prevent the Bone from being by that Means diſplaced again. A Luxation of the *Os Coccyx* outwards.

II. When the *Os Coccyx* happens to be luxated inwards, the firſt Finger is to be introduced into the *Anus*: after it has had its Nail cut and been dipped in Oil, it muſt be thruſt as far as poſſible, that it may the more readily drive out the depreſſed Bone; the other Fingers being applied externally, are to conduct the Bone into its right Poſture: when this has been done, it will be proper for the Patient to reſt ſome Time upon the Bed; and when he ſits up, it ſhould be in a Luxation of the *Os Coccyx* inwards.

[a] PETIT acquaints us, that, from neglecting the Aſſiſtance of a Surgeon in this Caſe, an Abceſs enſued, a Caries, a ſlow Fever, and which terminated in Death.

[b] See his Treatiſe on *Diſeaſes of the Bones*, T. I. C. 5.

Chair

Chair with a Hole in its Bottom, leſt the affected Part ſhould be otherwiſe compreſſed or diſturbed.

Luxations of the Ribs.

III. The Ribs are indeed ſometimes, though but ſeldom, diſlocated: for upon the Aſſault of ſome external Violence, it is not uncommon for them to be diſplaced either upwards, downwards, inwards, or outwards: they cannot be eaſily luxated outwards, becauſe prevented by the vertebral Proceſſes, and reſiſted by very thick and ſtrong Muſcles: but when they are drove into the Cavity of the *Thorax*, they not only lacerate the *Pleura*, or Membrane, which lines the Cavity of the *Thorax*, but do generally great Injury to the contained Parts: in Conſequence whereof ariſe moſt ſharp Pains, Inflammations, Difficulty of Breathing, Cough, Ulcers, Immobility, and many other dangerous Symptoms of the like Nature: but by what Signs ſuch Diſlocations of the Ribs are to be diſcovered, there is no Occaſion to conſider here at large; ſince the external Form and Poſture of the Side, with the troubleſome Symptoms now enumerated, generally afford evident Demonſtration whether any of the Ribs are luxated, and on which Side.

How the Ribs are to be reduced, when luxated upward or downward.

IV. The more numerous and grievous the conſequent Symptoms are, the greater is the Danger, and the more ſpeedily ſhould the Luxation be reduced: when the Rib is diſlocated either upwards or downwards, in order to replace it conveniently, the Patient is to be laid on his Belly upon a Table, and the Surgeon muſt ſtrive to reduce the luxated Rib into its right Place with his Hands: or, the Arm of the diſordered Side may be ſuſpended over a Gate or Ladder, as is ſhewn by Figures in PAREY and SCULTETUS; and while the Ribs are thus ſtretched up from each other, the Heads of ſuch as are luxated may be puſhed into their former Seat.

How the Ribs are to be reduced, when luxated internally.

V. But thoſe Luxations, wherein the Heads of the Ribs are forced into the *Thorax*, are generally found to be much the moſt difficult to reduce; ſince neither the Hand, nor any other Inſtrument, can be applied internally to direct the luxated Heads of the Ribs: but, notwithſtanding there are many eminent Surgeons who pronounce this Caſe to be wholly incurable; yet, in my Opinion, we ought not to deſpair of being frequently ſucceſsful: in this Caſe it ſeems proper to lay the Patient on his Belly over ſome gibbous or cylindric Body, and to move the the Fore-part of the Rib inwards towards the Back, ſhaking it ſometimes, for thus generally the Head of the luxated Rib ſlips into its former Place: but if this Method of Cure will avail nothing, and the deplorable Condition of the Patient requires ſpeedy Help, we have no Remedy left but Inciſion, and endeavouring to replace the luxated Head of the Rib with the Fingers, Plyers, or little Hooks, after the ſame Manner which we propoſed before in Fractures of the Ribs, *Book* I. *Chap.* X. § VIII, *& ſeq.* In the mean time, where the Symptoms are not very urgent, and the Heads of the Ribs but little diſplaced, it is adviſeable neither to cut the Fleſh, nor violently force the Ribs; becauſe there are ſeveral Inſtances where the luxated Ribs have retained their diſlocated Stations without any Hurt: but above all, Care muſt be taken to lay on a Compreſs dipped in warm *Sp. Vin.* or *Sp. Vin. camph.* to be retained on the afflicted Part of the Side by the *Napkin-and-Scapulary*.

Luxations of the Clavicles.

VI. Though the Clavicles are ſometimes diſplaced, it is but ſeldom, by reaſon of their ſtrong Ligaments: they may be diſlocated either from the Top of the *Sternum*,

Sternum, or *Proceſſus Acromion* of the *Scapula*, to which they are connected, by ſome external Violence, as a Fall, Blow, the lifting ſome great Weight, or the like. With regard to the Cure, the ſooner Aſſiſtance is had to the Patient, the more eaſily may the Reduction of the Clavicle be performed: but when the firſt is delayed, the latter will be more difficult, inſomuch that inveterate Luxations of the Clavicles are generally found incurable.

VII. The Clavicles may be diſlocated in two Manners; from the *Sternum*, either internally towards the *Larynx*, or externally upon the Breaſt: when the firſt Caſe happens, a Cavity may be generally obſerved upon the Part affected, and the *Trachea*, with the Carotid Arteries, the Jugular Vein, Nerves, and *Oeſophagus*, which are all together, will be very much diſturbed and compreſſed: on the contrary, when it is luxated forwards upon the Breaſt, it ſhews itſelf by a preternatural Tumor inſtead of a Cavity, upon that Part. (1.) near the *Sternum*.

VIII. In what Manner the luxated Clavicles may and ought to be extended and reduced again into their natural Places, has no Buſineſs to be inſerted again in this Place: becauſe every thing is to be obſerved the ſame as we propoſed in reducing Fractures of the Clavicles, *Book* II. *Chap.* V. *N.* 4. but this muſt be particularly regarded, to carefully remove the Injuries of the Neck, as ſoon as the Bones are replaced: if any Kind of Luxation requires an accurate Retention by Bandage, it muſt certainly be this of the Clavicle: eſpecially when the Luxation has happened ſome time before its Reduction: for, beſides that the Clavicles have ſcarce any Muſcles to ſupport them, their Ligaments are generally ſo much ſtretched and weakened in this Caſe, that they are in no wiſe ſufficient to ſuſtain the Weight of the Arms: it will therefore be proper to apply ſuch a Bandage to the Neck, as we ſhall deſcribe at large in the Doctrine of Bandages. How the Clavicles are to be replaced.

IX. Such Luxations of the Clavicles as happen near the *Proceſſus Acromion*, are generally much the more difficult to diſcover; ſo obſcure, that as HIPPOCRATES (*Lib. de Articulis*, *N.* 62.) and PAREY witneſs, abundance of the beſt Phyſicians, and Surgeons not a few, have been deceived in the *Diagnoſis* hereof, taking it to be a Luxation of the *Humerus*, and ſo have miſerably tortured the Patient to no Purpoſe: whenever this Luxation happens, as PAREY obſerves, the ſuperior Part of the *Scapula* ſticks up; but in the Place where the Clavicles are ſeparated from the *Acromion* Proceſs, a Cavity may be obſerved: moſt acute Pains ariſe, and the Arm itſelf cannot be moved or lifted up: if therefore the luxated Clavicles are not timely reduced, it is no Wonder that we meet with ſome People, who, from neglecting the Caſe, entirely loſe the Uſe of their Arms afterwards, ſo as that they cannot lift them up to their Head or Mouth: GALEN himſelf ſays, (*in Comment. in* HIPPOCRAT. *Lib.* I. *de Articulis*, N. 62.) "I "myſelf had once, in ſtruggling, my Clavicle ſo vaſtly ſeparated from the "*Acromion*, that there appeared a *Sinus* between the Bones, of near three "Finger's Breath:" in the mean time, a ſtrict Bandage continued about the Parts for forty Days, to make the diſunited Bones again coaleſce, will be found very ſerviceable. (2.) near the *Acromion*.

CHAP.

CHAP. VII.

Of a LUXATION *of the* HUMERUS.

How the *Humerus* may be diſlocated.

I. THE *Humerus*, from the Length and Laxity of its Ligaments, the Largeneſs of its Motion, and the Shallowneſs of the Cavity in the *Scapula*, into which it is articulated, is thereby rendered of all Bones the moſt ſubject and eaſy to be luxated. The Head of this Bone may often be diſlocated under the Arm-pit, ſometimes forwards, ſometimes backwards, and even below the *Scapula*; but ſeldom perpendicularly downwards, and never directly upwards, unleſs the *Acromion* and *Coracoide* Proceſſes of the *Scapula* ſhould chance to be fractured at the ſame Time: beſides, as long as the ſtrong *Deltoide* and *Bicipital* Muſcles of the *Humerus* remain intire, they greatly reſiſt and keep down the *Humerus* from being luxated upwards.

Signs of a luxated *Humerus*.

II. When the *Humerus is luxated downwards*, (1.) there ſuddenly appears a Cavity, and upon preſſing with the Fingers, you will perceive a *Sinus*; but under the Arm there muſt be a Tumor, becauſe the Head of the Bone is thruſt there: (2.) the *Proceſſus Acromion* will ſeem to ſtick out further than uſual, becauſe of the adjacent *Sinus:* (3.) the luxated Arm will be longer than the other, and it cannot be lifted up towards the Head without violent Pain, and ſometimes it cannot be lifted up at all, or even extended: but when the *Humerus is luxated forwards* as well as downwards, there will be obſerved the ſame *Sinus* under the *Proceſſus Acromion* as before, and a Tumor will appear from the Head of the *Humerus* projecting towards the Breaſt, under the *Axilla:* the Arm itſelf alſo cannot be moved without exciting the moſt acute Pain: laſtly, when the *Humerus is luxated backwards*, the *Cubitus* is thrown forwards towards the *Præcordia*, and the Head of the Bone makes a Protuberance in the Shoulder: the Arm itſelf cannot be bent nor extended, nor even pulled outwards from the Breaſt, without occaſioning the moſt violent Pains: but no Luxation of this Limb is attended with ſuch dangerous Symptoms, as when it is diſlocated forwards or inwards; becauſe the luxated Head of the *Humerus* cannot avoid injuring the large Arteries and Nerves of the Arm; in Conſequence of which, various Symptoms will ariſe.

Prognoſis.

III. If Aſſiſtance be had to theſe Luxations ſoon after they have been inflicted, before the bad Symptoms come on, the Reduction of them into their natural Places again may be effected without much Difficulty: more eſpecially, if the Bone be luxated directly downward or backward, it may be very eaſily reduced; but very difficultly when luxated inward, under the Pectoral Muſcle: ſo it may be eaſily replaced, when the Arm retains its natural Length: but if it be ſhorter, and the Accident has been done ſome Time, or accompanied with Tumor, Inflammation, or a Fracture of the *Proceſſus Acromion*, it is then a very difficult Matter to reſtore the Limb to its former Strength and Motion; but when the Head of the *Humerus* grows faſt to ſome of the adjacent Parts under the Arm, it can ſeldom be reſtored by any Means whatever: the Reduction is alſo more difficult in People that are ſtrong or fat, than in ſuch as are lean or weak.

IV. As

How a Luxation of the *Humerus* is to be reduced.

IV. As ſoon therefore as the Luxation is diſcovered in the *Humerus*, the ſafeſt Way will be to ſeat the Patient on the Floor, or on a low Stool, as at *Tab.* X. *Fig.* 3. A. two ſtrong Aſſiſtants are to be placed on each Side the Patient, one of which, B, ſhould ſecure his Body, and, if poſſible, the *Scapula* too [a], that it may not give way to the Extenſion: while the other, C, lays firm Hold of the luxated Arm with both his Hands, a little above the *Cubitus*, gradually and ſtrongly extending it. But before that Extenſion be made, the Surgeon himſelf, D, ſhould have a large Napkin, of a ſufficient Length, tied at the Ends, and hung about his Neck ſo that the Knot may be behind; but the other Part of the Napkin, E, muſt hang over his Breaſt: then the Patient's Arm muſt be put through the Napkin up to the Shoulder, and the Surgeon at the ſame Time lays hold of the Head of the *Humerus* with both his Hands: this done, he orders the Aſſiſtant to ſufficiently extend the Limb, and in the mean Time he elevates himſelf the Head of the Patient's *Humerus* by the Napkin about his Neck, directing it with his Hands, till it ſlip into its former Cavity in the *Scapula*. But I would adviſe the Surgeon to move the Head of the *Humerus* one way and the other, according to the Manner in which it is luxated; which muſt be left entirely to his Diſcretion: and by this means I have happily reduced a great many recent, though not inveterate Luxations of this Joint, particularly three in one Month, and that by no other Aſſiſtance or Machinery.

The bare Hands are ſometimes inſufficient for the Extenſion.

V. Though the Method now deſcribed for reducing this Luxation ſeems to be the moſt ſafe, ready, and commodious of any hitherto invented for that Purpoſe; yet it is found, that the Extenſion cannot, by this Means, be made ſufficiently ſtrong in ſome Caſes; and this particularly when the Patient is very robuſt, or when the Caſe has been delayed ſome Time, without any Aſſiſtance: therefore, when one or two Aſſiſtants are not able to retain the Patient, and ſufficiently extend his Arm, it is much the beſt Way to uſe a long Napkin with more Hands; or to apply the Girt of HILDANUS (*Tab.* VIII. *Fig.* 17.) about the *Humerus*, a little above the *Cubitus*, and to make the Extenſion by a Rope put through the two Hooks, and by another Rope faſtened to the Middle of that, letting as many Aſſiſtants pull as may be ſufficient, according to the Circumſtances of the Caſe. But when the Extenſion is made with a great Force, it requires to be antagoniſed by a ſtill greater Force, to keep the Patient ſteady: therefore it is proper to retain the Patient, by two Aſſiſtants; and if they are not ſufficient, to uſe a long Napkin or Piece of ſtrong Linen, ſlit and made in Form of the Slings at *Tab.* X. *Fig.* 1, 2. that the luxated *Humerus* may be put through the Slit up to the *Scapula*: the one Half of this Linen Sling ſhould come over the Breaſt, the other Half behind the Back, and both to meet afterwards together in a Knot: this is to be faſtened upon a Hook, or given into the Hands of ſeveral Aſſiſtants, or elſe it may be faſtened to a Beam, or ſome other fixed Point, ſo as to keep the Patient from being moved out of his Place. While this is performing, the Surgeon's immediate Buſineſs is, to accurately lift up, agitate, and reſtore the luxated Bone to its right Place, as we before directed: but when this Method alſo alone is inſufficient to extend the *Humerus*, it will be proper to apply to it the Pulley, *Tab.* VIII. *Fig.* 15. and keeping the Patient firm, to make a prudent

[a] A particular Stay might be contrived, for retaining the *Scapula*, by a long Napkin perforated in the Middle.

Extenſion of the *Humerus*, much as we propoſed before in a Fracture of the Thigh, *Book* II. *Chap.* VIII. *N.* III.

The *Ambe* of HIPPOCRATES.

VI. In theſe Kinds of Luxations, when the Hands are inſufficient for Extenſion, the Antients, and particularly HIPPOCRATES, made uſe of a Machine which they called (ἀμβὴ) *Ambe*, which may be ſeen delineated in *Tab.* X. *Fig.* 4 and 5. It conſiſts of a Pillar, or *Fulcrum* A A, and the moveable Lever B C, which is placed under and bound to the *Humerus* in the Manner of *Fig.* 5. by the Ligatures E E E. When this is done, the End of the Lever B is carefully and gradually preſſed downward; by which Means the other End of the Lever C is moved upward, and thus the luxated Arm is both extended, and its Head replaced at the ſame time. This was frequently uſed with ſo much Succeſs by them, that the Machine got a great Name, and is to this Day called the *Ambe* of HIPPOCRATES [a]. Notwithſtanding it was very ſucceſsful, and may be ſtill in ſuch Caſes, where the Head of the *Humerus* was luxated directly downward; yet, when the Head of the *Humerus* is luxated on one Side, or beneath the Neck of the *Scapula*, as generally happens, the Inſtrument elevating only directly upwards could not reduce the Luxation, but contuſed or lacerated the adjacent Parts, or elſe threw up and preſſed againſt the Neck of the *Scapula*, often exciting violent Pains, in ſuch Manner that (to ſay nothing now of its other Defects) it has been generally neglected by moſt for this long while, and is now wholly rejected.

Of other Artifices for this Purpoſe.

VII. To proceed, we muſt not omit taking Notice here, that there are ſeveral other Methods and Contrivances invented, not only by the Antients, but alſo many of the modern Phyſicians and Surgeons, for reducing a Luxation of the *Humerus*. Thoſe of the Antients are delineated by ORIBASIUS (*Lib. de Machinamentis*,) PAREY (in his *Surgery*, *Book* XV.) GERSDORFF, BRUNSWIG, SCULTETUS (in their *Chirurgical Writings*) and other eminent Surgeons. As for the modern Contrivances, two of their Machines are publiſhed in the *Acta Eruditor. Ann.* 1683, *pag.* 37. another in JUNGKENII *Chirurgica Germanica*, *pag.* 168. where he treats of Luxations; another in PURMANNI *Chirurg. Curioſ. Tab.* XIV. *pag.* 692; and ſtill another in PETIT's *Treatiſe on Diſeaſes of the Bones*: and though theſe latter ſeem to be each in great Eſteem with their own Authors, every one thinking he had mended the Defects of his Predeceſſors; yet there are ſome of the *French* Surgeons who eſteem and publickly declare them to be either unneceſſary, or leſs ſuitable than the *Ambe* of HIPPOCRATES [b]. There are even ſome who look upon all Machines as unneceſſary in this Caſe, but the Hands, and Napkins, or Slings; as GOUE, a *Frenchman* too, in his *Surgery* [c].

Of PETIT's Machine.

VIII. But becauſe PETIT is an ingenious Surgeon, and well verſed in his Profeſſion, I thought it would be worth while to exhibit here the Machine which he ſo vaſtly commends, and to give a ſhort Deſcription thereof. See *Plate* X. *Fig.* 7: but ſuch as deſire a more full Account, may conſult the Author's Book of Inſtruments itſelf [d]. PETIT made it his Buſineſs to contrive his Machine ſo as not only to make a ſufficient Extenſion of the Limb, which

[a] See a Book intituled, *Diſſertation en forme de Lettre.* [b] See ditto.

[c] DOUGLAS too is of Opinion, that Writers in Surgery have been highly blameable in recommending uſeleſs and ridiculous Machines in a Luxation of the *Humerus*. *Op. Chirurg. Syll. pag.* 47.

[d] As PETIT's Figures are from wooden Plates, and but very indifferent, I have endeavoured to repreſent them as clearly as poſſible, that they may be better underſtood.

others

others had invented Means to anſwer very well before, but alſo to make a counter Extenſion or Reſiſtance at the ſame time, to retain the Patient, and particularly his *Scapula*, ſufficiently firm from giving way to the Extenſion of the Limb made by the Inſtrument: with this View he made a ſort of Buttreſs or Supporter *(l'Archoutant)* of Ticken, a Foot long, of ſufficient Strength, and lined with Leather, as at *Tab.* X. *Fig.* 7. The Arm is to be firſt put through the Opening A, ſo as to make one End B come over the Breaſt, and the other End C to go croſs the Back. Its two Holes D D, let in the two Horns or Legs of the Machine, *Fig.* 6. *a a*, the other End of which, B, is lodged upon the Ground. In this Machine there are ſeveral little Pullies *c c*, *c c*, as in the *Polyſpaſton* of *Tab.* VIII. *Fig.* 16. round which paſſes the Rope *d d d:* there is alſo a moveable Handle E, by which the Rope is wound up through the Pullies, and the luxated Arm by that means extended. But that the Arm may be the better extended, he uſes a peculiar Sling A A, *Fig.* 8. made of ſoft and double Leather, fourteen Inches long: this he faſtens ſtrongly round the lower Part of the *Os Humeri*, a little above the Elbow; the Skin being firſt pulled upwards, is to be kept firm upon the Limb by means of a Silk Cord, three Quarters of an Ell long, ſewed in a particular Manner to the Leather of the Sling, and to be faſtened by a Knot at the two Ends *b b:* to this Silk Cord is faſtened another Sling *c d e*, by two moveable Loops *f f*, to which is to be annexed the Rope *d d d*, which paſſes round the Pullies of the Machine. The Apparatus being all rightly fitted, he orders his Aſſiſtant to wind up the Rope by the Handle E, *Fig.* 6; the Rope becomes by that means ſtretched, and the Arm to which it is faſtened is gradually extended. In the mean time the Surgeon directs the Head of the *Humerus* with his Hands, that it may again obtain its natural Place, which it very often does of its own accord, without the Direction of the Surgeon[a].

Of other Methods leſs in uſe.

IX. But to give my Opinion impartially concerning the Uſe of Machines for reducing a Luxation of the *Humerus*, I muſt needs ſay, that the Surgeon's Hands and a Napkin, with ſtrong and dextrous Aſſiſtants to make the Extenſion, and hold the Patient firm, will, of themſelves, be generally ſufficient for the Buſineſs: but if any one be willing to uſe other Methods, he may pitch upon thoſe as the beſt, which ſufficiently extend the Bones, and equally ſtretch the Muſcles every way alike, that the Head of the *Humerus* may be aptly replaced. Upon this Principle we may readily judge whether the *Ambe* of HIPPOCRATES be ſufficiently proper or no to be applied in this Caſe: or the ſtill more uncertain Method of pulling and extending the luxated Arm over a Gate, Ladder, or Beam, by a Couple of tall and ſtrong Aſſiſtants, in ſuch Manner as to liſt the Patient off his Legs: or when a luſty and ſtrong Aſſiſt ſits down on the Floor, and preſently laying hold of the Patient's Arm, ſuddenly raiſes himſelf up thereby: or laſtly, when the Patient is ſeated on the Ground; and by placing the Foot under the Head of the *Humerus*, it is violently pulled upward, or any other Way extended: all which Methods are handled at large by PAREY, in his *Surgery*, *Book* XV. But here it muſt be cautiouſly obſerved over and over, that

a This Machine of PETIT, and his Treatment of a Luxation of the *Humerus*, is ſmartly cenſured in a *French* Journal *(des Scavans)* M. *March*, 1734 (which the Author of the late quoted Diſſertation has ſeconded) and many notorious Errors are there diſcovered.

the Nerves, Veins, Arteries, Muſcles, and the Bones themſelves, be not contuſed or broke, by the too great Strength and Suddenneſs of the Extenſion. That ſuch Accidents as theſe may readily happen in a rough Extenſion of the luxated Arm over a Gate, *&c.* where the Patient is ſuſpended by it, we ſhall find no Room to doubt, eſpecially if we conſider the Reaſons and Inſtances cited by PETIT in his Chapter on a Luxation of the *Humerus*, and others. And ſince this is the Caſe, the Surgeon's principal Care and Buſineſs in the Extenſion will be, to let the Arm be ſtretched out with a Force ſtrong enough, but equable, before he ſtrives to replace the luxated Head of the Bone: otherwiſe he acts to no Purpoſe, or to a very bad one, by bruiſing and lacerating the neighbouring Parts.

A new Machine.

X. There is ſtill another new and very conſiderable Machine with a Pulley, which I received not long ago from a very eminent Surgeon, deſigned for the Reduction of an obſtinate and inveterate Luxation of the *Humerus*; the great Advantages of which he very much praiſed and recommended to me: but becauſe I have not yet had Opportunity to uſe it, and ſo could not experience its Effects, I muſt refer the Deſcription thereof to another Opportunity.

CHAP. VIII.

Of a LUXATION *of the* CUBITUS.

In what manner the *Cubitus* may be luxated.

I. THE *Cubitus* conſiſting of two Bones, the *Ulna* and the *Radius*, is articulated by *Ginglymus*, which the *French* call *Charniere*, as is evident from what is ſaid of theſe Bones in the Writings of Anatomiſts. The Connection of theſe Bones is ſuch, that the *Ulna* or *Cubitus*, as being the largeſt Bone, and ſeated in the inferior Part of the Arm, does, of itſelf, perform the whole Flexion and Extenſion of the Arm, yet it cannot perform that Motion without carrying the *Radius* along with it: ſo that the *Radius* always follows the *Ulna* in Flexion and Extenſion: but on the other hand, the *Radius* may be turned along with the Hand both inward and outward, without at all moving or bending the *Ulna*: as when the Pronation and Supination of the Hand is made thereby. Both theſe Bones of the *Cubitus* are ſo articulated with the lower Head of the *Os Humeri*, that large Protuberances are received into deep Cavities or Grooves, and the whole inveſted and faſtened with exceeding ſtrong Ligaments; ſo that notwithſtanding the *Cubitus* may be luxated in all four Directions, outward, inward, forward, and backward, yet it is but ſeldom that it ſuffers a perfect or entire Diſlocation, unleſs the upper Part of the *Ulna*, called *Olecranon*, be broken, or the Ligaments of the *Cubitus* much weakened, by ſome very great external Violence.

How to diſcover a Luxation of the *Cubitus*.

II. If the *Cubitus be luxated backward*, which is the moſt frequent of all others, then the Arm becomes crooked and ſhorter, and it cannot be extended. In the inward Part of the Bend of the Arm, the Head of the *Humerus* may be obſerved to ſtick out; in the back Part of the ſame, the Head of the *Ulna* or *Olecranon* will be protuberant, and between both Bones will appear a *Sinus* or Cavity. But it very ſeldom happens that the *Cubitus is luxated forward*, from the Largeneſs of the *Olecranon*; unleſs that be fractured at the ſame Time.

Time. But if this ſhould happen, the Head of the *Humerus* will ſtick out behind, and that of the *Cubitus* before; and there will be a Cavity more or leſs in Proportion to the Degree of the Luxation. When the *Cubitus is luxated externally*, the Protuberance appears on the Outſide of the *Cubitus*; and the contrary when luxated inwards. To conclude, unleſs the Ligaments and Muſcles of the *Cubitus* are quite broken in two, it is ſo far from being capable of ſuffering a perfect Diſlocation, that no more can happen to it than a Subluxation, *i. e.* it can recede but a very little Way out of its right Place: but whatever of this Kind happens, the Caſe may be very eaſily underſtood, by feeling and inſpecting the Part, if there be no Tumor: but if the Joint be much ſwelled, it is very difficult to be diſcovered.

Prognoſis. III. Since, in the more violent Kind of theſe Luxations of the *Cubitus*, the Tendons and Ligaments muſt be very much ſtrained; it is no wonder (if theſe be not ſpeedily helped) that there ſhould follow grievous Pains, Tumors, Inflammations, Convulſions, Vomiting, Fever, and at length Gangrene and Death: an ample Witneſs wherof is PAREY, in *Book* XIV. *Chap.* 18. and *Book* XVIII. *Chap.* 33. and to make no Diſſimulation in the Caſe, when the *Cubitus* is diſlocated, it is a very difficult Matter to replace it again, by Reaſon of its Inequalities and ſtrong Ligaments: and this more eſpecially when the Luxation is very violent or inveterate; for the ſlighter and more recent the Luxation, the more eaſy will be the Reduction.

How a Luxation of the Cubitus is to be replaced. IV. Be the Luxation however more or leſs, the Patient muſt be ſpeedily placed in a Chair, and both Parts of the Limb, the *Humerus* and the *Cubitus*, muſt be extended in oppoſite Directions, by two ſtout Aſſiſtants, till the Muſcles are found pretty tight, with a free Space between the Bones: then the luxated Bone muſt be replaced with the Surgeon's bare Hands, or together with Bandages: and that the Proceſſes may fall into their *Sinuſes*, the *Cubitus* muſt be afterwards ſuddenly bent. But if the Tendons and Ligaments are ſo violently ſtrained, that they can ſcarce perform their Office; it will not be improper to anoint them well with emollient Oils, Ointments, or the Fat of Animals, or to apply emollient Fomentations and Cataplaſms. Where the bare Hands are not ſufficient to make a proper Extenſion upon the Limb in this Caſe, it will be very proper to uſe the Means and Inſtruments which we before propoſed in *Book* II. *Chap.* VIII. *N.* 3. and 4.

How the Cubitus is to be treated after Reduction. V. As ſoon as the Reduction has been by theſe Means effected, the Articulation muſt be bound up with a proper Bandage, and the Arm is to be afterwards ſuſpended in a Napkin or Sling about the Neck: but Care muſt be taken, as HIPPOCRATES himſelf adviſes, that the Bandage be not ſuffered to be on too long, nor the Arm to be kept all the Time ſtill, without ſome gentle Motion: for thus there would be Danger of the Mucilage of the Joint becoming inſpiſſated, whereby the Articulation might become ſtiff, and quite loſe its Motion: but happily to prevent this, it is very neceſſary to undo the Bandage every, or every other Day, and to gently bend and extend the Limb: afterwards, Compreſſes dipped in burnt Wine, may be applied hot, and held on firm with a Bandage, till the Swelling and Inflammation, if there be any, ſubſide, and the Ligaments and Articulation regain their former Strength.

CHAP. IX.

On LUXATIONS *of the* HAND, CARPUS, METACARPUS. *and* FINGERS.

Luxation of the Hand.

I. NOtwithſtanding the Hand is very accurately connected to the two preceding Bones, and particularly to the *Radius*, by means of the *Carpus* and ſtrong Ligaments, yet it ſometimes ſuffers Luxation in all four Directions: but it is generally not ſo eaſy to be luxated on either Side, as forward or backward, becauſe of the two Proceſſes of the *Radius* and *Ulna*, which guard it on each Side. The Hand is ſaid to be *luxated forwards* or *inwards*, when it recedes from the Muſcles which bend the Fingers; to be *luxated backward* when it departs from the Muſcles which extend the Fingers: much alſo in the ſame manner, the Hand is judged to be *luxated outward*, when the *Carpus* makes a Tumor near the Thumb, and a Cavity near the little Finger: to be *luxated inward*, when the contrary happens. This being rightly conſidered, it will not appear difficult to diſtinguiſh the Signs by which we are to diſcover a Luxation of the Hand.

Symptoms and *Prognoſis*.

II. For, if a Luxation of this Kind ſhould happen, it can hardly avoid being accompanied with violent Pains, on account of the Ligaments (though ſtrong) being too vehemently ſtrained: the Fingers alſo cannot be bent nor extended, from the violent Compreſſion of their Tendons: upon which Account, it is no Wonder if there follow grievous Inflammation, Tumor, Abceſs, Stiffneſs of the Joint, Gangrene and *Sphacelus*, or *Caries* of the ſpongy Bones in the *Carpus*; which Evils are ſeldom remedied but by amputating the Limb: but when the Luxation is but ſlight and recent, the Cure may be effected with much more Eaſe, and the Diſlocation will not be attended with ſuch grievous Symptoms.

How a Luxation of the Hand is to be reduced.

III. It therefore ſeems to be the ſafeſt Way immediately to reduce what is diſplaced: and that this may ſucceed the better, two Things are to be chiefly regarded: (1.) that the luxated Hand be ſufficiently extended by two Aſſiſtants, one of which is to lay hold of the Hand, and the other of the *Humerus*, pulling in oppoſite Directions: (2.) that the Part of the extended Hand, where the *Sinus* is, be placed on a Table, or ſome other flat Body, that whatever ſticks up may be depreſſed: by which Method the Hand, in whatever Part luxated, may be very readily reduced into its natural Seat.

Luxation of the Carpal Bones.

IV. It alſo ſometimes happens, that one or two of the eight little Bones of the *Carpus* are luxated and diſtorted from their natural Seat by ſome external Violence. When this happens, there will be perceived a Tumor in one Part, and a Cavity in another, which may be alſo felt by the Fingers; beſides, violent Pains will be felt by the Patient: for the reſt, as this kind of Luxation is very eaſily diſcovered, partly by the Sight, and partly alſo by Feeling; ſo, when it is recent, it is almoſt as readily cured, letting the Hand be extended in the Manner we before propoſed (at *N.* III.) and the diſlocated Bone be afterwards forced into its Place.

Luxation of the *Metacarpus*.

V. The four ſmall Bones, which are found in the *Metacarpus*, or Palm of the Hand, may be ſometimes luxated from the *Carpus* itſelf, to which their upper Parts are connected: which uſually happens from ſome external Violence; notwithſtanding they have a natural Inclination to reſiſt ſuch Luxation: for the

two Bones which are ſeated in the Middle between the two other external ones, cannot be diſlocated to either Side: as the two external ones which ſuſtain the firſt and little Fingers cannot be luxated inwardly, but are more eaſily driven outward: tho' each of them may be luxated on the fore or back Part of the Hand. But which ever of theſe happens, the particular Diſorder may be diſcovered and examined by feeling and inſpecting, and the Cure may be carried on in altogether the ſame Method which we directed before at § IV.

VI. Laſtly, the Bones of the Fingers, to which we join thoſe of the Thumbs, are liable to Luxation at each of their Articulations, and that in ſeveral Directions. But theſe Accidents, if recent, are both very eaſy to diſcover and cure: for the Ligaments being not very robuſt, the Fat and Muſcles thin, and the *Sinuſes* of the Articulations ſhallow, render the Extenſion very eaſy, and the Reduction of them into their former Places may be done very readily. While one Hand of the Surgeon extends the Finger, he ſtrives with his other to replace the Bones in their natural Seat. The Bandage proper to dreſs the Finger after Reduction will be explained more at large in the third Part of this Book, where we treat profeſſedly on Bandages. Luxation of the Fingers.

CHAP. X.

Of a LUXATION *of the* THIGH.

I. VERY rare is it that the Head of the Thigh-bone is diſplaced out of its *Acetabulum*; tho' formerly it was ſuppoſed to be pretty frequent, the Phyſicians taking a Fracture thereof for a Luxation, as we have obſerved in treating on Fractures. See *Book* II. *Chap.* VIII. § VI. and *Book* III. *Chap.* I. § IX. the Reaſon whereof may be taken from the Articulation itſelf. (1.) how very deep is the *Sinus*, called by the Ancients *Sinus Coxæ*, and by the Moderns *Acetabulum*, into which the Head of the Thigh-bone is received. (2.) with what a broad concave Cartilage is almoſt the whole Head of that Bone covered. (3.) how ſtrong are the Ligaments with which it is faſtened. (4.) how greatly it is defended with exceeding ſtout and thick Muſcles. (5.) but how very brittle is the Neck of this Bone beyond any other Part thereof: from all which it appears, that the Neck muſt be far more frequently and eaſily broke eſpecially in Adults, than its Head diſlocated[a]: and tho' ſomething of this Kind may ſometimes happen, ſo as to make the Head of the Thigh-bone ſlip out of its *Acetabulum*; yet that generally proceeds more from internal than external Cauſes[b]: for we find it has been obſerved by very ſkilful Phyſicians, that the Ligaments of the Thigh-Bone, tho' very ſtrong, may be by various Cauſes, and particularly by a Flux of Humours, ſo relaxed and weakened, as to let the Head of that Bone ſlip ſpontaneouſly out of its *Acetabulum*: ſo that it ſhould ſeem no great Wonder if the Thigh ſhould be ſometimes luxated even The Thigh ſeldom luxated.

[a] To theſe we may add that the celebrated CHESELDEN in his Anatomy ſays, that upon opening two Subjects, whoſe caſe every body thought to be a Luxation, the Neck of this Bone was found fractured: and WISEMAN, with other eminent Surgeons, wholly deny any Luxation in this

[b] RUYSCH obſerves that the Head of the Thigh-bone is ſeldom diſlocated by external Force; but frequently by Humours from within: ſee his *Theſaur. Anatom.*

while

while the Patient lies in Bed, without any external Violence, as I have ſometimes ſeen; ſo that when they riſe, one Leg appears longer or ſhorter than the other, and ſeems as if it were unhinged. *Vid.* HIPPOC. *Aph.* 59, 60. § VI. ZWINGER, *Theat. Pract. Part.* II. *pag.* 119. *ſub tit. Luxatio.*

Happens oftener in Infants than Adults.

II. But this Caſe does not happen ſo eaſy in robuſt Adults, as in ſuch as are more young and tender, as we before obſerved. I remember to have ſeveral Times obſerved this Caſe of a ſpontaneous Luxation, though other Phyſicians and Surgeons were of a contrary Opinion, becauſe they could not find that any external Violence had gone before: but though it was preceded by no external Violence, Experience has taught me, that the Head of the Thigh-bone may thus ſlip out of its *Acetabulum*; being the Conſequence of preternatural Humors or ſome other Diſeaſe, whereby the Ligaments and Articulation are rendered infirm.

When the Thigh is luxated, it is generally perfectly ſo.

III. Whenever the ſaid Head of the Thigh-bone is thruſt out, it is almoſt always wholly diſplaced, ſo as to make a perfect Luxation. The exact Roundneſs of this Head, with the great Force of the circumjacent Muſcles, and the Narrowneſs of the Sides of the *Acetabulum*, will not admit the Bone to be diſlocated a little Way only: for as ſoon as the Head of this Bone is thruſt up to the Edge of the *Acetabulum*, it muſt unavoidably either turn quite out, or elſe fall back again into its right Place[a]: yet there are ſome who hold that the Thigh may ſuffer an imperfect Luxation.

The Thigh may be luxated in ſeveral Directions.

IV. The Thigh is uſually luxated four ways; *upward*, *downward*, *backward*, and *forward:* but it is moſt frequently diſlocated downwards and inwards, towards the large *Foramen* in the *Os Pubis*[b]: for beſides that the cartalaginous Defence on the lower Part of the *Acetabulum* is not ſo high as the reſt, the *Ligamentum rotundum* is found to give Way more eaſily in that Part than any other: and laſtly, the adjacent Muſcles are found to be weakeſt in their Reſiſtance on this Part, being inſufficient to keep the Head of this Bone from ſlipping out: and then there is a certain Eminence in this Edge of the *Acetabulum*, which keeps the Head of the *Os Femoris* from falling back again into its right Place: but if the Head of this Bone be diſplaced outwards, it generally ſlips upwards at the ſame time; it being ſcarce poſſible but the very ſtrong Muſcles of the Thigh muſt then draw the Bone upwards, and then there is no Eminence there, in the Edge of the *Acetabulum*, to reſiſt the Head of the Bone in that Paſſage: but ſhould it at any time be luxated by an external Force, there muſt certainly be a Rupture of the round Ligament; as ſome Authors have obſerved.

Sign of the Thigh luxated (1.) downward and forward.

V. When the Thigh is diſlocated forwards and downwards, which is what uſually happens, the Leg hangs ſtraddling outward, and is longer than the other: alſo the Knee and Foot turn outwards: the Head of the Bone itſelf will be felt near the lower Part of the *Inguen* and *Os Pubis*: ſometimes there is a Suppreſſion of Urine in this Caſe; when ſome Nerve, which communicates with the Bladder, is violently compreſſed: in the Buttock may be perceived a Cavity, from the *Trochanter Major* and the reſt of the Bone being diſplaced: and if the Thigh-bone be not timely replaced into its *Acetabulum*, the whole Limb withers

[a] See WISEMAN's Surgery, *pag. m.* 463.
[b] As VERDUC has obſerved, *Operat. Chirurg.* pag. m. 300.

ſhortly

ſhortly afterwards [a]: and this is the Reaſon why the Patient can bear little or no Streſs upon that Limb, but muſt always incline and throw the Weight of his Body upon the other, unleſs he would fall down: in like Manner, when they walk or go forward, the Perſon muſt move that Limb in the Form of a Semicircle; but as for the Body itſelf, it is obliged to be ſupported under the Arms by Aſſiſtants, or elſe by Crutches and Sticks: though there are not wanting particular Caſes, ſome of which I have been Witneſs to, where the Head of the luxated Thigh-bone has grown ſo firmly to the adjacent Parts without the *Acetabulum*, as to become, in Proceſs of Time ſo ſtrong, as to ſupport the Body without Crutches or Sticks, though they always halted in walking.

(2.) upward and outward.

VI. But if the Thigh-bone be diſplaced backward, it is uſually drawn upward alſo at the ſame Time, as we before obſerved: hence there will be perceived a Cavity behind the *Inguen*; but upon the Haunch or Buttock, a Tumor; becauſe the Head and *Trochanter* of this Bone will be thruſt there. The Tumor in the Haunch being thruſt upwards, the reſt of the Limb will become ſhorter than the other, and the Foot will ſeem to turn inwards: the Heel will not touch the Ground, and ſo the Perſon will ſeem to ſtand upon his Toes: and laſtly, the luxated Limb may be bent with more Eaſe than extended: alſo the Body is uſually ſuſtained more firmly by this Limb when luxated backward than forward; becauſe, in the firſt Caſe, the Feet are removed farther from each other: and this is the Reaſon why a great many in Caſes of this Kind, which have been cured by Surgeons without reducing the Bone, are able to ſtand firmly and walk, eſpecially if they have a Shoe with a very high Heel to it: but there generally follows ſomething of a ſlight Withering or Decay in the Limb afterwards, from the Nerves being in ſome Meaſure compreſſed: though this Accident is much ſlighter here than at § V. Laſtly, it is extremely rare that the Thigh is luxated forward or backward, without being alſo drawn upward or downward, as we before obſerved: but if it ſhould ſo happen, it may be evidently diſcovered by what we have been juſt now ſaying, and from conſidering the Structure of the Articulation.

How to diſtinguiſh between a Fracture and a Luxation of the Thigh.

VII. As it is very difficult to diſcover whether the Thigh-bone be diſlocated or fractured, both by feeling and inſpecting, becauſe of the great Thickneſs of the Muſcles and Integuments; it is therefore, in my Opinion, a Matter of ſome Conſequence to propoſe the following Signs, which we recommend for diſcerning one from the other. We do not without Reaſon judge the Thigh to be luxated (1.) when we find the Ligaments of the Bone have been relaxed by ſome preceding Congeſtion of Hūmors, and when no external Violence has been exerted upon it, eſpecially in young Patients: (2.) when neither the Symptoms, Pain, Tumor, or Inflammation follow: and laſtly, (3.) when the whole Limb may be bent and turned about at the *Acetabulum* without any Cruſhing of the Bones, which is otherwiſe common in Fractures. The contrary of theſe Signs are ſtrong Indications that a Fracture is preſent: more particularly if the Foot in grown Perſons be ſhorter, from the Injury of any external Violence, and you hear a Grating of the Bones in moving the Limb.

Prognoſis.

VIII. If it be difficult to diſcover whether the Thigh be fractured or luxated, as we have before made evident, its proper Treatment and Cure will be found much more ſo. See *Book* III. *Chap.* I. § IX. For this Difficulty there are many Reaſons: for (1.) the Force and Thickneſs of the adjacent Muſcles

[a] See HIPPOCRAT. Sect. V. *Aph.* 59, and 60.

themſelves hinder the Thigh from having a ſufficient Extenſion; eſpecially if it be in the ſtronger Sort of Men: hence, (2.) for the ſame Reaſon, the Reduction of the Bone will be very difficult to effect; and when it is effected, it will be a hard Matter to diſcover: and (3.) if the Thigh ſhould happen to be replaced quite home as it ſhould be, yet there is great Danger of its ſlipping out again, from the Laxity of the Ligaments and Slipperineſs of the Parts: to which we may add (4.) that the Ligaments happen to be ſometimes quite broke or lacerated, from the Greatneſs of the external Violence: and we muſt not forget that (5.) the Mucilage of the Joint becomes often ſo inſpiſſated in the *Acetabulum*, as not only to prevent its Reduction, but often alſo to thruſt it out again when once replaced, (6.) in Infants the Head of the luxated Bone increaſes, and at the ſame Time the *Sinus* contracts; that if there be not a ſpeedy Reduction, there can be none: ſo that it is no Wonder if ſuch become halt or lame, as have their Thigh-bone luxated, and reduced not at all, or elſe when it is too late: yet in this Diſlocation in a younger Patient, if it be recent, and the Ligaments are not broke, there are Hopes of reducing it; eſpecially, if you apply proper Remedies: for in this Stage of Life the Muſcles are very pliant, and the Bone is eaſily extended, and replaced: but it is not ſo eaſily retained there, in Conſequence of the Weakneſs and Flaccidity of the Parts.

Cure of the Thigh luxated forward and downward.

IX. The luxated Bone is always to be replaced in a Method agreeable to the Nature and Direction of the Diſlocation. When it is diſplaced forwards and downwards, the Patient is to be laid flat upon his Back on a Table: then a linen Napkin or ſtrong Sling is to be made faſt over the Groin about the Part affected, ſo that one End of the Sling may come over the Belly, and the other over the Nates and Back, to be both tied together in a Knot upon the Spine of the *Os Ilium*, and afterwards faſtened to a Hook fixed in ſome Poſt, or held firm by ſome Aſſiſtants; rather the firſt, if we uſe the *Polyſpaſton* or Pulley, to retain the Patient's Body firm from giving Way in the Extenſion: in like Manner, at the Bottom of the Thigh, a little above the Knee, there muſt be alſo faſtened another Napkin, Sling, or the Girt of HILDANUS at *Tab.* VIII. *Fig.* 17. with a Compreſs between it and the Thigh; or PETIT's Sling, *Plate* X. *Fig.* 8. Both the Slings being drawn tight, the Thigh is to be extended, not vehemently, but only ſo much as is ſufficient to draw the Bone out of its *Sinus*, that it may be replaced into its *Acetabulum* by the Surgeon's Hands: one Hand is to preſs the Head of the Thigh-bone outward, while the other conducts the Knee inwards: or, the Reduction may be made by Napkins, faſtened round the Extremities of the Thigh like Slings, much as in a Luxation of the *Humerus*; which will be more likely to ſucceed, if the Knee be at the ſame Time preſſed inwards by the Hands. When the fore-recited Means are not ſufficient to make the Extenſion, it will be neceſſary to make Uſe of the *Polyſpaſton* or Pulley which we propoſed before in *Book* II. *Chap.* VIII. § IV. As ſoon as the Thigh is found to be ſufficiently extended, the Surgeon muſt take particular Care to reſtore the luxated Head of the Thigh-bone with his Hands from the *Os Pubis* into its former Seat.

X. Whenever the Thigh is luxated backward, the Patient is to be placed flat on a Table, with his Face downward, and the Thigh is to be extended in directly the ſame Manner, but a little more ſtrongly than we juſt now propoſed: and the Reduction is to be effected afterwards by the Surgeon's Hands, an Aſſiſtant

ſtant in the mean Time extending the Limb, and turning it inwards: by this Method the Head of the Thigh-bone generally ſlips very readily again into its *Acetabulum*. This being all rightly effected, the next Buſineſs is to let the diſordered Limb be well bound up, as we ſhall teach in the Doctrine of Bandages, and the Patient is to be cloſely recommended to reſt in his Bed for three or four Weeks, not without the Uſe of ſtrengthening Medicines.

The Uſe of PETIT's Machine.

XI. But in either Caſe, whether the Thigh be luxated forward and downward, or backward and upward, PETIT greatly recommends his Machine before deſcribed in the Chapter on a Luxation of the *Humerus*; becauſe the Hand and other Inſtruments are here very often inſufficient, becauſe of the many ſtrong Muſcles in this Part: but to make uſe thereof, the *Retinaculum* or Stay delineated at *Tab.* X. *Fig.* 7. is required to be not ſo broad, and it may be without the Opening A, (See *Plate* X. *Fig.* 9.) as the Thigh is not to be tranſmitted through it: but the Middle thereof is to be applied to the Tubercle of the *Iſchium*, one End being folded behind, and the other before: the Patient is to be placed on his ſound Side, that the luxated Thigh may lie upwards; but the Machine itſelf is to be placed between the Thighs, the Knee of the diſtorted Side being a little bent: the Sling *Fig.* 8. *Tab.* X. is to be faſtened firmly round the lower Head of the Thigh, above the Knee, the Skin being firſt drawn tight upwards, as we adviſed before in a Luxation of the *Humerus:* it is then to be firmly faſtened to the Rope paſſing round the Pullies of the Machine *Fig.* 6. *d d:* and laſtly, the Legs or Horns of the Machine *a a*, are to be put through the Apertures in the *Retinaculum* D D, *Fig.* 7. and by winding up the Rope by the Hand E, *Fig.* 6. it is to be gradually and carefully extended, till the Surgeon perceives by the Limb that it is ſufficient; this done, the Surgeon ſtrives to reduce the Head of the Thigh-bone into its *Acetabulum*, from the *Sinus* where it was lodged, as we have before directed at § IX.

More particularly in a Luxation forward and downward.

XII. But more particularly, if the Thigh be luxated forward and downward, and ſticks near the large *Foramen* in the *Os Pubis*, the Reduction in this Caſe is often very difficult. PETIT has in this Caſe ſubſtituted for the Legs or Horns of the Machine *a a Fig.* 6. others which are expreſſed at *Fig.* 10. which have at their Ends tranſverſe or lunar Proceſſes: one of theſe A, he applies to the *Os Ilium*, and the other B, to the Middle of the Thigh: he afterwards ties a Napkin about the Thigh, near the *Inguen*, which he makes faſt to the Rope about the Pullies of the Machine: he then makes the Extenſion by turning the Handle of the Machine, by which Means the Inſtrument exerts its Force in three different Places: the Part A retains the Patient firm, and reſiſts the *Os Ilium* as an immoveable *Fulcrum*; the Part B, when the Rope is drawn tight, turns the lower Part of the Thigh inwards; but the Napkin, which is faſtened about the upper Part of the Thigh, does by Means of the Rope draw it outward; all which Motions are neceſſary to be performed, in order to reduce this Luxation: but be cautious againſt too ſtrong an Extenſion, becauſe the Limb is already too long of itſelf: yet the Extenſion ought to be continued till the Surgeon can replace the Bone from the *Sinus*, where it was lodged, into its *Acetabulum:* for if it be let looſe before this is effected, the Extenſion will be found to have been altogether uſeleſs, and muſt be repeated again.

XIII. If it ſhould ſuffer an imperfect Luxation (which yet very ſeldom, if ever happens, as we obſerved at § III.) and if the Head of the Bone ſhould ſtop upon the lower Part of the *Acetabulum*, the upper Part of the Thigh is then to be thruſt outwards with one Hand, while the lower Part is puſhed inwards by the other, and ſo the Bone may be properly reduced: but if the Head of the Thigh-bone ſhould ſtick upon the Edge of the *Acetabulum* backward, a Method contrary to the former muſt be made uſe of; *viz.* the upper Part of the Thigh muſt be thruſt inwards by one Hand, while the other Hand conducts the lower Part of the Thigh outwards.

XIV. When the Bone is replaced, it muſt be retained by the Application of a proper Bandage, called *Spica Inguinalis*, which you will find deſcribed in my Treatiſe on Bandages: and in younger Patients, where the Miſchief ariſes from a Flux of Humours, ſtrengthening Medicines muſt be given: and for the Revulſion of thoſe Humours an Iſſue ſhould be made in the Arm of the ſound Side; to which ſuch Virtues are attributed, that to this only MULLER aſcribes the Cure of theſe Luxations in young Patients. See his Diſſertation on Diſlocations of the Thigh, delivered at *Hale* in the Year 1738.

CHAP. XI.

Of a LUXATION *of the* PATELLA *and* KNEE, *or* TIBIA *and* FIBULA.

Luxation of the *Patella*.

I. THE *Patella* is uſually luxated moſtly on the internal or external Side of the Joint; but, if we may credit ſome Phyſicians, it is alſo ſometimes diſplaced both above and below the ſame: but whenever the Knee is perfectly luxated, the *Patella* can ſcarce avoid being diſplaced at the ſame Time, becauſe of its ſtrong Connection to the Thigh and *Tibia*: I muſt confeſs there are more than a few among the common Surgeons, who, from their Unſkilfullneſs in Anatomy, and particularly Oſteology, are quite doubtful and at a Loſs what to think about this Caſe; nor can they tell what is diſlocated when it happens; hence it is no Wonder if they treat this unknown Hurt of the Joint, as a Luxation made in the Knee itſelf, putting the Patient into various and painful Poſtures, and torturing him by extending and preſſing the Limb to no Purpoſe: but if one well verſed in Anatomy and the Structure of the Articulation ſhould examine the Caſe with a little more Exactneſs, there is no Room to doubt but, from comparing the diſordered and ſound Limb, he will be able to judge readily whether or no, or in what Part the *Patella* is luxated, and what Method will be proper to be taken for its Cure.

How it is to be replaced.

II. The Reduction of a luxated *Patella* is uſually no very great Difficulty, if the Patient be laid flat on his Back upon a Table or Bed, or if he be laid in that Poſture upon an even Floor, ſo as that the Leg may be pulled out ſtrait by an Aſſiſtant: for then the Surgeon may firmly graſp the *Patella* with his Fingers, and afterwards preſs it ſtrongly into its right Place; which may be alſo effected if the Patient ſtands upright: when this is done, there remains nothing but to carefully bind up the diſordered Part, and to let the Patient reſt quietly for ſome Days, ſometimes gently bending and extending his Leg to pre-

vent it from growing ſtiff: 'till the Pains are gone off, and the Limb has recovered its former Strength.

III. A Luxation of the Knee is properly ſo, when the *Tibia* recedes from under the *Femur*. The Leg is ſometimes luxated from the Baſis of the Thigh-bone, either on the out or inſide, or backwards: ſeldom or never forwards, unleſs it be forced and driven very violently that Way; becauſe forwards, the *Patella* is bound againſt the Articulation, by the very ſtrong Tendons of the Muſcles which extend the Leg: nor is it eaſy for the Bones of the Leg to be wholly diſplaced from that of the Thigh, ſo as to make a perfect Luxation; by reaſon of the great Strength of the Ligaments, and the two deep *Sinuſes* which receive the Head of the Thigh-bone: unleſs thoſe very ſtrong Ligaments ſhould happen to be broke in ſunder at the ſame Time: and this ſeems to me to be the Reaſon why People, who ſuffer a perfect Luxation of the Knee, are generally tortured with ſuch violent Pains and Convulſions, that they are wholly ſpent or waſted thereby: or, if they ſhould eſcape that, they are generally troubled with Lameneſs and Stiffneſs in the Joint: but on the contrary, the ſlighter the Luxation, or the nearer it approaches to an imperfect or Subluxation, the more eaſy it is generally to effect the Reduction and Cure: yet I myſelf reduced, ſome Years ſince, at *Helmſtadt*, a perfect Luxation of the *Tibia* backwards, and that in a robuſt and jolly Patient, without any other Inconvenience, than his perceiving a Tumor and ſome Pain in his Knee for a few Weeks: which were afterwards entirely removed by emollient Fomentations and Cataplaſms; and his Leg is perfectly ſound to this Day. Luxation of the Knee.

IV. For the reſt, as this Kind of Luxation is very eaſy to diſcover from the thin Covering of the Joint, with the Tumors and Cavities which follow; ſo, when it is diſcovered, it is as difficult to make a perfect Cure thereof, without letting the Bones join together; or leaving ſome Stiffneſs in the Knee; which firſt Accident is uſually called an *Anchyloſis:* for it is ſcarce poſſible, that this Caſe ſhould happen without greatly lacerating or contuſing the Ligaments and Glandules which belong to this Articulation, ſo that their nutritious and mucilaginous Juice being inſpiſſated in the Articulation, prevents the natural Motion of the Joint.

V. When the Knee is but ſlightly luxated, the Patient is to be ſeated on a Bed, Bench, or Table, and one Aſſiſtant holds the Thigh firm above the Knee, and the other extends the Leg; but the Surgeon in the mean Time replaces the Bones by his Hands and Slings, if neceſſary, (as in the Reduction of the Arm, *Plate* X. *N.* III.) or puſhes it by the Application of his Knee into its natural Place. If the Hands and Slings be not ſufficient for this Purpoſe, it will be neceſſary to make uſe of the Inſtruments before deſcribed in *Book* II. *Chap.* I. N. XXI. as the Girt of HILDANUS, and the *Polyſpaſton* or Pulley, *Tab.* VIII. *Fig.* 15 and 17. But we muſt be very careful here, not to make the Extenſion ſo violent in Children and young People, as to ſeparate the *Epiphyſes* from the Bones to which they are not yet firmly united: for by that Means a worſe Diſorder and Lameneſs will be brought on. After the Luxation of the Knee is rightly reduced, it is to be properly bound up, and placed in a Straw Caſe; and the reſt muſt be managed as we have before directed concerning the *Patella*, *N.* II. How it is to be replaced.

VI. Some-

Luxation of the *Fibula*.

VI. Sometimes the *Fibula* is ſeparated by ſome external Violence from the Thigh-bone, and is then diſtorted either upward or downward: and this generally happens, when the Foot has been luxated outward: therefore, whenever this is the Caſe, there is Need of Extenſion. The Bone is to be firſt reſtored to its natural Place, and then properly bound up, the reſt of the Cure being to be carried on as we directed at *N.* IV. and II. 'till it be grown firm again to the *Tibia* and Leg: laſtly, Patients ſhould be frequently cautioned not to uſe or bear any Streſs upon the diſordered Leg too ſoon; unleſs they would throw themſelves into a worſe Diſorder, an incurable Lameneſs.

CHAP. XII.

Of a LUXATION *of the* FOOT *and* ANCLE.

Luxation of the Ancle.

I. THE Ancle may be ſometimes luxated either in Jumping, Running, or Walking, and that in all four Directions, inward, outward, backward, and forward. Which of theſe Ways it happens to be luxated, may be diſcovered by the particular Poſture of the Joint: for when it is luxated internally, the Bottom of the Foot is turned outward; and on the contrary, when it is luxated outward, the Bottom of the Foot is turned inward, which latter Caſe happens much more frequently than the other: if it ſhould be diſlocated forward, the Heel becomes ſhorter, and the Foot longer than it ſhould be; if backward, the contrary Signs will appear: laſtly, the Ancle can ſcarce poſſibly be luxated outward, unleſs the *Fibula* be ſeparated from the *Tibia*, or elſe quite broke, which may happen on the external Ancle: an Example whereof may be ſeen in LE DRAN, *Obſ.* 109.

Luxation of the Ancle dangerous.

II. Nor is it uncommon for a Luxation of the Ancle to be attended with the moſt grievous Symptoms, eſpecially when occaſioned by ſome very great external Violence: for it is ſcarce poſſible for it to happen otherwiſe in this Caſe, ſince the Diſtortion of the Foot muſt overſtrain the adjacent Ligaments, Tendons, and Nerves, and thence excite moſt violent Pains: or the Veins and Arteries may be alſo lacerated; by which Means there will be a large Extravaſation of Blood about the whole Foot, which often gives Riſe to a Gangrene: of this Accident DIONIS gives an Example in his Book of *Surgery:* nay, ſometimes Death has enſued from this Kind of Luxation, as TULPIUS has remarked, in *Obſ. Med. Lib.* IV. *Obſ.* 50. who alſo quotes HIPPOCRATES on that Subject, *Lib. de Art.* XII. *Hiſt.* 48. *& Lib.* V. *Epid.*

The Ancle is ſometimes only contuſed.

III. But it ſeems to be here worth Notice, that the Ancle is not always luxated after it has been violently ſtrained by Leaping, or turning the Foot on one Side: for it ſometimes happens in thoſe Caſes, that the Ancle is not diſlocated, but retains its proper Place, only the Parts are violently contuſed and ſtrained; in which Caſe the Patient may happen to be afflicted with the moſt ſharp Pains, great livid Tumor and Stiffneſs, ſo that he can neither ſtand nor walk, but is obliged to keep his Bed for a long Time: hence, to attempt an Extenſion and Reduction in this Caſe, would be altogether uſeleſs and improper.

IV. The

IV. The Ancle is more or leſs difficult to reduce in Proportion to the Violence of the Cauſe by which it is luxated. It is a general Obſervation, that Oppoſites are uſually the Conſequences of Oppoſites: but the moſt ready Way of reducing a Luxation of the Ancle is to place the Patient upon a Bed, Seat, or Table, letting the Leg and Foot be extended in oppoſite Directions by two Aſſiſtants, while the Surgeon ſtrives to reduce the Ancle with his Hands and Fingers. When the Foot is by this Means once replaced, it is proper to bind it up carefully, after it has been well bathed with *Oxycrate* and Salt, adviſing the Patient to keep to his Bed a good while, 'till the Diſorder and its Symptoms quite leave him, and he finds his Ancle to have recovered its Strength ſo far as to ſuſtain the Weight of his Body without any Uneaſineſs or Danger. How the Ancle is to be reduced.

V. But in a Contuſion or great Strain of the Ancle, it will be not improper to plunge it ſuddenly into cold Water, and to repeat it for ſeveral Days. If any ſhould not care to undergo the Action of the cold Water, I would perſuade him to apply Compreſſes dipped in *Oxycrate* which has had Salt diſſolved in it, binding them on and renewing them often upon the diſordered Part. DIONIS runs directly into this Method of Cure: he applies what the Surgeons call a Defenſative, made of the White of an Egg and Oil of Roſes, and Allum, beat up together, which being ſpread on Linen, he binds firmly upon the Ancles, repeating it frequently: in about three Days after, he makes a Decoction of aromatic and aſtringent Medicines, as Roſes, Wormwood, Roſemary, Granate Peels, and Allom, in Wine; and with this foments the Ancle well, and applies Compreſſes dipped therein, binding them on tighter than before: this continued about a Fortnight, he then applies ſome ſtrengthening Plaſter, 'till the Pain and Weakneſs vaniſh. How a Contuſion of the Ancle is to be treated.

VI. So ſtubborn and unmoveable are violent Strains of the Ancles in ſome People, that they will give Way neither to the Skill of the Phyſician, nor Virtue of Medicines, but are only to be removed by Length of Time. Inſtances are not wanting, where the Foot has been ſo greatly diſordered, a Year after the Luxation, that the Patient could not walk in a Way the leaſt uneven, nor go up and down Stairs without great Trouble: to remedy this Diſorder, the ſame is to be carefully obſerved here, which we obſerved before at *N.* IV. The Bandages which are proper here, we ſhall deſcribe hereafter. A Contuſion of the Ancle ſometimes difficult to cure.

VII. Sometimes it happens, that only the *Os Calcis* or *Calcaneum* is luxated by ſome external Force, and that either towards the internal or external Side of the Foot. Whichever Way it happens, when there is Pain and Inequality of the Member, that is, when it has a Cavity in one Part, and a Tumor in another, there is ſtrong Reaſon to ſuppoſe a Luxation: and as ſoon as it is diſcovered, the ſame Method of Cure is required with that before recited, keeping the Limb quiet for ſome Time afterwards. Luxation of the *Calcaneum*.

VIII. Laſtly, if any other Bone in the Foot, the *Tarſus* or *Metatarſus*, ſhould happen to be luxated by ſome conſiderable external Violence, the Ligaments with the adjacent Nerves and Tendons are generally ſo affected as to excite not only moſt acute Pains, but violent Inflammations and Convulſions; and even Death itſelf has been obſerved by ſome Phyſicians to be the Conſequence, unleſs the Bones were ſpeedily replaced: it is therefore the ſafeſt Way to reduce the Luxations in theſe Bones of the Foot, by the Method we before propoſed: Luxations of the other Bones of the Foot.

propoſed for thoſe of the Hands, and that with the greateſt Expedition: ſo when any of the Bones in the Toes are diſlocated, there is nothing more required than what we propoſed before in thoſe of the Fingers: we are however, in the laſt Place, to recommend the Patient to reſt quietly in his Bed for a ſufficient Time afterwards.

An EXPLANATION *of the* TENTH TABLE.

Tab. X. *Fig.* 1. Is a Sling which may be uſed to make an Extenſion in Luxations of the Head: ſee the Chapter on Luxations of the Head.

Fig. 2. Is another Sling, to retain the Patient's Body firm in the ſame Caſe.

Fig. 3. Shews the moſt commodious Method of reducing a recent Luxation of the *Humerus*.

A, Is the *Patient*, ſeated ready to undergo the Operation.

B, Is the *Aſſiſtant*, that holds the *Patient* firm in his Seat.

C, Is the *Aſſiſtant*, that extends the diſlocated *Humerus*.

D, The *Surgeon*, reducing the diſlocated *Humerus*.

E, A Napkin, whereby the Surgeon elevates the Arm, in order to its Reduction.

Fig. 4. Is a Machine, commonly called the *Ambe* of HIPPOCRATES, uſed formerly to reduce Luxations of the *Humerus:* it conſiſts of the *Fulcrum* A A, to which is faſtened the moveable Lever B C, joined to each other by a Sort of moveable Articulation D.

Fig. 5. Shews how the former Inſtrument is to be applied to a Luxation of the *Humerus*. There is ſome Difference between the Structure of this and the former, at the Joint C D: ſome think this is preferable to the laſt.

A A, Is the *Fulcrum*; B C, the Lever, to which the luxated Arm is faſtened by the three Ligatures E E E. D, the Place where the *Fulcrum* and Lever are faſtened together by a moveable Joint. When the End of the Lever B is preſſed downwards, the luxated Arm is extended, and lifted up near its *Scapula* E.

Fig. 6. Is PETIT's *Machine* for reducing Luxations of the *Humerus*, and ſeveral others.

a a, Are two Arms or Horns, by which the Patient, and particularly his *Scapula*, is held firm, from giving Way in the Extenſion; B, the other End of it, reſting upon the Ground or Floor. C C, are the Pullies of the Machine: *d d*, the Rope, by winding up which an Extenſion is made. E, is the Handle, which being turned round, draws the Rope tight, and extends the Limb. F F, the Place where the two Horns are joined to the Body of the Machine.

Fig. 7. Is a *Retinaculum*, or Supporter, to be uſed in a Luxation of the *Humerus*. A, is an Opening or Slit in the Machine. B C, the Form of it at each End. D D, two Apertures, through which the two Legs or Horns *a a* of the Inſtrument *Fig.* 6. are to be paſſed.

Fig. 8. Is a particular Sling of Mr. PETIT's, proper for extending luxated Limbs. A A, the Part made with Leather; *b b b*, a ſilk Ligature, ſewed to the Leather in three Places at 1, 2, 3. The Part A A is faſtened round the Arm;

Fig: 9.



Fig: 6.





Fig: 5.



Fig: 4.






Fig: 8



Plate 10.

Fig: 1.





Fig: 2.










Fig: 9.



Fig: 5.




Fig: 4.




Fig: 6.





Fig: 7.



Fig: 8.



Arm; *cde*, is a ſtrong Loop faſtened to the ſilk Ligature at *ff* ſo as to be moveable.

Fig. 9. Is another Kind of Supporter to be uſed in reducing the Femur, when luxated outwards: it differs from that repreſented at *Fig.* 7. being without the Aperture A in the Middle. B and C are the two Cavities, by which the Arms are fixed to the Machine.

Fig. 10. Is an Inſtrument recommended by PETIT for the Reduction of the *Femur* when diſlocated inwards. It is to be faſtened at FF in the Machine *Fig.* 6. inſtead of the Arms *aa:* the Part A is applied to the *Os Ilium*, and the Part B to the Middle of the Thigh: but CC are fixed into the Machine *Fig.* 6. at FF.

INSTITUTIONS OF SURGERY.

PART I. BOOK IV.

Of TUMORS.

CHAP. I.

Of TUMORS *in general.*

A Tumor what.

I. ANY Part of the Body which is preternaturally enlarged, is, by Phyſicians, called a *Tumor*: but whether there be any ſuch Enlargement, in what Part it exiſts, and of what Kind it is, may be diſcovered by examining the Parts, not only by Inſpection, but more particularly by Feeling: and notwithſtanding it has been the general Cuſtom to refer *Excreſcences*, as Warts, Corns, and ſuch as grow in the Noſe and *Pudenda*, to the Claſs of *Tumors*; yet becauſe they grow not from beneath, but out of or upon the Skin itſelf, it ſeemed proper here to treat ſeparately of Tumors, properly ſo called; we ſhall take Notice of the moſt remarkable Excreſcences, when we treat of chirurgical Operations.

Kinds of Tumors.

II. There are Tumors of various Kinds, diſtinguiſhed by particular Names, according to the different Cauſes from whence they proceed, and the particular Parts wherein they are ſeated: ſome are called *hot*, others *cold* and *watery*; ſome again are termed *windy*, others *ſchirrous*; and laſtly the Phyſicians call ſome *benign*, others *malignant*: there are ſome Tumors which are contained in a proper membranous Bag, and are therefore called *cyſtic*: and if this ſhould be in an Artery, 'tis uſually termed an *Aneuriſm*; but when in a Vein, a *Varix*: in the Veins of the *Anus* or *Rectum*, the Diſorder is termed *Hæmorrhoides*: but if the Tumor be in the *Scrotum*, *Inguen*, or at the *Umbilicus*, it is generally called a *Hernia*: if any *Pus* or Matter is contained in the Tumor, it is then by the Surgeons termed an *Abceſs*: laſtly, if the Tumor is ſeated on a Bone, Phyſicians uſually call it an *Exoſtoſis*.

Kinds of Inflammation.

III. The forementioned Claſſes of Tumors are all of them generally ſubdivided into ſeveral other Kinds: thus the hot and burning Tumors, which are the ſame with Inflammations, are generally termed *Phlegmons*, when violent, and ſeated in the common Integuments; but when ſlighter, they are commonly called

called *Furuncles*. The Inflammation which is not fixed deep, but ſpreads only ſuperficially upon the Skin, is uſually diſtinguiſhed by the Name of an *Eryſipelas*; and the inflammatory Tumor that ariſes at the Fingers Ends is termed *Paronychia*. When the Inflammation fixes in the Groins or Arm-pits, the Tumor is called a *Bubo*; but when under the Ears, *Parotis*. If a great Inflammation ſeizes the Hands and Feet from extreme Cold, *Childblains* ariſe. Other Inflammations have alſo particular Names, according to the particular Part of the Body they poſſeſs: hence, in the Writings of Phyſicians, we frequently find Accounts of an Inflammation of the Breaſts, Eyes, Tonſils, Teſticles, Arms, Feet, *&c.* And this may ſuffice for a ſhort and general Account of the Kinds of Inflammations: the various other Sorts of Tumors we ſhall explain hereafter.

IV. Before we proceed farther into Conſideration and Treatment of Tumors, it will be firſt proper to take Notice that we do not intend here to handle all Sorts of Tumors to which the human Body is ſubject, but only ſuch as are external, and of the ſlighter Kind: we intend firſt to examine thoſe Tumors only which are to be cured by manual Operation, and topical Remedies, and ſo come properly under the Buſineſs of Surgery; neglecting at the ſame time ſuch Tumors whoſe Cure is to be expected chiefly from the Uſe of internal Medicines; as is uſual in ſome internal Inflammations, *Schirrus's*, Dropſies, and the like: we ſhall alſo refer thoſe Tumors which require Inſtruments and great Skill in their Treatment, to the Part of chirurgical Operations: ſuch are *Herniæ*, *Excreſcences*, *Strumæ*, *Scrophulæ*, the *Paronychia*, *cyſtic Tumors*, *Aneuriſms*, *Varices*, *Hæmorrhoides*, and others: our chief Concern here will be to treat of *Inflammations*, *Schirrus*, *Cancer*, *Œdema*, *Tumors of the Joints*, and other external Tumors. We begin with Inflammations.

CHAP. II

Of a PHLEGMON.

I. A *Phlegmon* or external Inflammation is, when any outward Part of the Body is peternaturally enlarged, and attended with a burning Heat, Pain, Redneſs, Reſiſtance, and a continual Pulſation and Pricking: upon a due Conſideration whereof, we may pretty readily perceive the Reaſon why the Diſorder came to be diſtinguiſhed by this Name. If we enquire into the proximate Cauſe of this Inflammation, we ſhall find it generally riſes from too thick or viſcid a State of the Blood, ſtagnating in the *Anaſtamoſes* of the ſmalleſt Arteries and Veins; ſo that the Blood being ſent in larger Quantities than it can paſs through thoſe Veſſels, muſt of Conſequence excite the forementioned Symptoms, and muſt occaſion great Diſorder at every Part where ſuch Stagnation is made: and though no Part of the Body, whether external or internal, nor the Bones themſelves are exempt from this Kind of Inflammation; yet it more frequently happens in the Fat and Glands than elſewhere. *What a Phlegmon is.*

II. We juſt now obſerved, that the immediate Cauſe of this Inflammation was an Obſtruction or Stagnation of the Blood in the ſmalleſt Order of Veſſels: but if we enquire into the Cauſes from whence that Inſpiſſation and Stagnation *Cauſes (1.) external.*

of the Blood in thoſe Veſſels proceeds, we ſhall, upon Examination, find them to be of two Kinds; of which, the firſt Sort may be called external, and the latter internal: amongſt the *external Cauſes*, we place in the firſt Rank, all Wounds, Fractures, Luxations, Contuſions, Punctures by Thorns and Splinters, with a too great Compreſſion of the Veſſels, whether by too ſtrict a Bandage, or otherwiſe: Each of which obſtruct the Paſſage of the Blood through its minute Veſſels, by either dividing, bruiſing, compreſſing, or diſtorting them. To the forementioned Cauſes we muſt add, Burns of all Sorts, extreme Cold, too violent Motion of the Body, the external or internal Application of ſharp and ſtimulating Subſtances, ſticking Plaſters, oily and fat Things, with Abundance of the like Nature, which ſtop up the inviſible Pores of the Skin, and impede the free Courſe of the Blood.

(2.) Internal.

III. Amongſt the *internal* Cauſes, we reckon any thing acrimonious in the Fluids, as in the Scurvy; becauſe theſe ſo irritate, corrode, prick, and contract the very ſmall Veins and Arteries, that the Blood is thence by Degrees obſtructed in them. But the ſame alſo frequently happens from the Blood abounding in too *great Quantities*, or being of *too thick a Conſiſtence*; or laſtly, when it circulates in the Body with *too violent a Motion*. For by this Means the groſſer Particles of the Blood are drove, and, as it were, wedged into much ſmaller Veſſels than they can readily find a Paſſage through; and this more eſpecially when a ſudden Cold is ſpread over a Body that is in a great Heat. In ſhort, every thing will produce an Obſtruction, which makes the Parts of the Blood too groſs and bulky, or too much contracts the Mouths of the ſmall Veſſels.

What Share an Acid and Fermentation have, as Cauſes.

IV. As this is the State of the Caſe, with regard to the Cauſes of Inflammation, I think the Opinion of ſome modern Surgeons, who ſuppoſe the chief and ſole Cauſe of the Obſtruction to be an *Acid* in the ſmall Veſſels, appears to be very evidently erroneous. For beſides their Inability to diſcover whether and where this Acid hides itſelf, it is very apparent, from what we have here delivered, that great Obſtructions may be brought on by a long Train of very different Cauſes. The ſame may be ſaid with regard to *Fermentation*, which has been formerly patronized by many as a grand Cauſe in Inflammations and Obſtructions; for there could never yet be found any ſuch Fermentation in the Blood.

The Symptoms of Inflammations.

V. We obſerved at § I. that an Inflammation was generally attended with Tumor, Heat, Redneſs, and Pain, and very often with a Reſiſtance and conſtant Pulſation: to inveſtigate the Cauſes of which Symptoms, we ſhall meet with no great Difficulty, if we ſtrictly and accurately examine the Diſorder itſelf. When the Blood is obſtructed in its Paſſage through ſome of its ſmaller Veſſels, it muſt neceſſarily move faſter through the reſt; for the ſmalleſt Arteries are never all obſtructed, but in a *Sphacelus*; the general Conſequence then muſt be a ſwifter Circulation of the Blood through all its other pervious Veſſels in the Body: hence the Arteries muſt beat quicker, ſwell larger, and thence excite great Heat. When we find a Patient in this State, we ſay he has a ſmall Fever; which is uſually accompanied, for the firſt Days, with Thirſt, Head-ach, Reſtleſſneſs, and the other common Attendants of a Fever. If we bleed the Patient in this Caſe, when his Blood is cold, it appears covered with a tough, and whitiſh Cruſt or Skin, not greatly unlike the Skin of freſh Pork: as the Diſeaſe and Heat increaſe, each of theſe Symptoms become more violent; 'till at laſt, the

the whole Maſs of Blood being deprived of its moſt fluid Parts, is converted into a tough and glutinous Body, as will be more apparent from our Obſervations at the Operation of Bleeding.

The ſeveral Ways by which an Inflammation terminates.

VI. Inflammations terminate variouſly according to their different Degrees of Violence, the Cauſes from whence they ariſe, the Parts which they affect, and the particular Conſtitution of the Patient, with ſeveral other Circumſtances, which alſo preſage to us what will be the End of the Inflammation: but the ſeveral Ways wherein an Inflammation terminates, are chiefly four: it is either (1.) ſo *diſperſed* and *reſolved* as to vaniſh without leaving any conſiderable Injury in the Part affected, which afterwards recovers its former Vigour, and is of all the beſt Courſe it can take: or elſe (2.) the Inflammation ſuppurates and degenerates into an *Abceſs*, ſo as to leave ever after ſome Damage in the Organ. Or elſe (3.) the Inflammation degenerates into a *Gangrene* or *Sphacelus*. Or (4.) laſtly, into a very hard Tumor, commonly called a *Schirrus*, which grows more compact in the Part affected as the Inflammation remits or goes off.

How it may be (1.) diſperſed, or (2.) ſuppurated.

VII. As to the Reſolution or *Diſperſion* of an Inflammatiou, that is uſually practicable when it is only of a milder Kind, in a ſound Habit of Body, when the Blood is not yet too viſcid nor vehement in its Motion: but *Suppuration* follows, when the Inflammation is more violent, the Circulation more rapid, but yet the Maſs of Blood ſomewhat temperate, and without much Acrimony; that is, when the Blood becomes ſomewhat more inſpiſſated, and its larger Particles ſticking in the minute Veſſels can find no Paſſage; by which means the very ſmall Veſſels are diſtended and burſt by the Preſſure and Impulſe of the obſtructed Blood, ſo that their Contents are extravaſated in the Fat, Fleſh, and adjacent Parts: upon this Extravaſation the more ſubtle Parts of the Fluids putrify by the great Heat, they become fœtid, acrimonious, and corrode all around them: the Fluids thus changed and corrupted, are then by the Surgeons called *Pus* or Matter; of which there are ſeveral Kinds, according to its different Colour and Conſiſtence; being either white, yellow, greeniſh, reddiſh, or party-coloured.

Or end in (3.) *Gangrene*, or (4.) *Schirrus*.

VIII. The Inflammation generally terminates in a *Gangrene* (which CELSUS and the *Latins* term *Cancrum*), when the forementioned Symptoms are much more violent, and when the Blood is at the ſame time more acrimonious and rapid than it ought to be; for, in that Caſe the ſmall Arteries and Veins are corroded, and burſt or broke: hence all the adjacent Parts are diſſolved and corrupted by the acrimonious and extravaſated Humours; and particularly the Skin is very ſubject to be filled with Puſtules, when its *Cuticle* has been ſeparated, as in Burns. The *Sanies* contained in theſe Puſtules and elſewhere, is uſually termed *Ichor*; which is generally of a pale reddiſh Caſt, being ſometimes fleſh-coloured, and ſometimes brown or livid, which is the worſt of all: for unleſs the Patient in theſe Circumſtances be timely aſſiſted, the forementioned Symptoms of Inflammation go off, the Tumor, Reſiſtance, Heat, Redneſs, Pain, and Pulſation gradually diſappear, and the afflicted Member grows flaccid and cold; it afterwards turns pale, becomes dead and inſenſible, and the Inflammation creeps to ſome other Part. If this Caſe ſhould chance to be treated with Medicines too hot, aſtringent, cooling, fat, acrimonious, or narcotic; or if the Parts ſhould be bound up too tight, the Fleſh then quite dies, its Paleneſs turns to a livid or leaden Colour, ſometimes reſembling the Rind of Bacon: In the mean Time the incloſed *Sanies* finds no Vent, becomes more acrimonious, and

ſo

ſo greatly corrodes the adjacent fleſhy Parts, as wholly to deſtroy all Senſe and Motion throughout the Limb; whereupon follow a *Sphacelus*, or entire Corruption of the Member: but if the inflamed Part be full of Glands, and the Blood very thick, glutinous, and inſpiſſated; the ſmall Blood-veſſels are then ſo ſtrongly ſtuffed up with the glutinous Blood, that are compacted together, the Parts loſe their Senſation, and are converted into a *hard* Tumor, which is thence called a *Schirrus*: this may be ſufficient concerning the four ſeveral Ways wherein a *Phlegmon* may terminate: but it remains that we ſhew the particular Method of Treatment and Cure proper in each of thoſe Stages.

Of the Reſolution or Diſperſion of INFLAMMATIONS.

In what a Diſperſion thereof conſiſts.

IX. Though the Methods uſed to cure Inflammations may be various, according to the ſeveral Cauſes and ſupervening Symptoms, with other various Circumſtances; yet, as the Inflammation conſtantly ariſes from an Inſpiſſation of the Blood in its ſmalleſt Veſſels, the grand Intention of each of thoſe Methods ſhould be, to open ſuch ſmall Veſſels as are obſtructed, and to reſtore the Blood to its natural Conſiſtence and free Circulation: this has been commonly termed Reſolution or Diſperſion: therefore, whenever the inflammatory Signs, mentioned § V. are gentle, it is much the beſt Way ſpeedily to conclude about diſperſing it: the right Method of performing which, we are now going to lay down.

Removal of the external Cauſes.

X. If the Cauſe of the Inflammation is found to be external and obvious to the Senſes, as Thorns, Splinters, the End of a Sword, Bullets, or any other foreign Body ſtuck in the Part; nothing can be more ſerviceable than to ſpeedily and carefully remove whatever is lodged there, if it can be done with Safety: ſo alſo when the Inflammation proceeds from a too ſtrict Bandage in Wounds, *&c.* or from a Luxation or Fracture; the firſt and principal Buſineſs is, to ſpeedily relax the Bandage, or elſe to ſet the Fracture, or reduce the Luxation.

Treatment of a ſlight Inflammation.

XI. When the external Cauſes are once removed, the next Thing to be conſidered is, whether the Inflammation be mild, or violent. If mild, there is no Occaſion for Bleeding, or ſtrong Purges; the End is generally anſwered, by applying Compreſſes, dipped in Spirit of Wine, with a little *Sal Ammoniac*, or Lime-water, and ſome Spirit of Wine camphorated, to the inflamed Part, and repeating them warm at proper Diſtances: likewiſe Oxycrate, or Cabbage-pickle, or a Decoction of Wormwood, Southernwood, *&c.* in Wine or Sea-water, applied in the ſame Manner, are of wonderful Efficacy in removing ſlight Inflammations.

Bleeding and Purging to be uſed.

XII. But where the Inflammation is violent, it is very uſeful to open a Vein, either in the Arm or Foot, and to draw off a large Quantity of Blood, proportionable to the Strength and Habit of the Patient; giving afterwards a briſk Purge, not one that heats the Body, but judiciouſly accommodated to the Age and Conſtitution of the Patient: both theſe are very neceſſary here, and if the Symptoms do not remit and grow milder, they muſt be repeated at Diſcretion: but I would adviſe the Surgeon in this Caſe, where he can, to call in the Advice of ſome prudent Phyſician; becauſe it may be otherwiſe carried on to Exceſs, as many do among the *French*, or elſe not made ſufficient to anſwer the Intention. The moſt proper Purges for theſe Caſes, we have mentioned before (at

Book I. *Chap.* XV. § XIV. *ſeqq.*) in ſpeaking of Inflammations ariſing from Contuſions: but in very mild Inflammations, or where the Patient is of a weak Habit, or has loſt much Blood by a Wound, or any other Cauſe, Phlebotomy and even Purging itſelf ſeems to be quite improper: on the contrary, when the Inflammation is great, and the Patient ſtrong, it is almoſt incredible of what great Service a prudent Adminiſtration of laxative and diſcutient Medicines may prove.

Internal Medicines proper.

XIII. To reſolve and attenuate the inſpiſſated Blood in the ſmall Veſſels, exceeding great Benefit will be found by giving internal Medicines, which are watery, diluent, cooling, and attenuating; becauſe Bleeding alone, which the *French* rely too much upon, is frequently inſufficient, unleſs it be joined with a proper Regimen and Diet; by which means any Acrimony in the Blood may be mollified and taken off: but all Aliments which are of a difficult Digeſtion, ſuch as are pickled or ſalted, with all Spices and fermented Liquors, or any thing elſe that may heat the Blood, are to be ſtrenuouſly and altogether avoided: ſuch internal Medicines are moſt proper here to cool and qualify the Blood, as are commonly given with Succeſs in continual ardent Fevers, or internal Inflammations, as the Pleuriſy, Meaſles, *&c.* ſuch are the abſorbent Powders of *Lap. Cancro. Conch. pp.* neutral and nitrous Salts, cooling and diaphoretic Mixtures and Juleps, made of diſtilled Waters, ſubacid Juices and Syrups, alſo thin Emulſions, made of the four cold Seeds: but the bezoardic and ſpirituous Tinctures preſcribed and recommended by ſome in this Caſe, are ſo far from being ſerviceable, that they encreaſe the Inflammation in the Blood, and raiſe a new Fire.

A regular Diet to be obſerved.

XIV. With regard to the particular Regimen and Diet, the moſt proper *Aliment* ſeems to be Broths and Drinks, made with Barley, Oats, or Flower, alſo Viper's Graſs, Succory, Chervil, Sorrel, Lettice, Endive, Apples, and Vegetables of the like Nature; in the Decoction of which may be mixed the Juice of Citrons or Vinegar, to communicate a grateful Sharpneſs, and temperate the inflammatory Heat: hence roaſted Apples, or Cherries, and Plumbs boiled, are very wholſome for inflammatory Caſes, where they ſit eaſy upon the Stomach: the moſt proper *Drinks* are ſuch as are thin, watery, and cooling, made of a Ptiſan or Decoction of Barley, Oats, or Bread, and to give it a pleaſant Taſte, Apples may be uſed, or ſome acid Syrup: but when the Inflammation is violent, it will be proper to add a ſmall Portion of Nitre[a]: of theſe may be drank plentifully in Proportion to the Thirſt and Heat: but Care ſhould be taken not to let the Patient over-drink himſelf: Ale and ſtrong Wine ſhould be wholly abſtained from: but if they are of the ſmalleſt Sort, and the Patient has a ſtrong Deſire for them, he may be gratified without any great Danger; eſpecially if a Slice or two of a Citron be infuſed therein: beſides the foregoing, it may be not amiſs, for Variety, to uſe Coffee and Tea, *&c.* If the Patient ſhould happen to be of a cool and phlegmatic Habit, it may be not improper to add ſome of the milder Sort of Spices to his Drink, as Cinnamon, Saſſafras, Mace, Anniſeeds, and the like: or the Patient may be ordered to infuſe ſome proper medicinal Herbs in the Manner of Tea, or a very weak Decoction of Saſſafras; the

[a] HARRIS recommends the following Drink, as of great Service in Inflammations: ℞. *Nitri* ℥j. *Pulv. Coccion. gr.* xii. *Aquæ ſimp.* ℔iij. *Sacch.* ℥ j. *m. f. Potio*; *Cujus bibat æger* ℥v. *vel* vj. *ſtiq. vel 4ta quaqua hora.*

drinking of which will promote a gentle *Diophoreſis* or Perſpiration: for by this Means whatever is glutinous in the Blood, will be readily attenuated and reſolved, and the Blood will recover its free Circulation.

External Medicines.

XV. Nor is there leſs Care required in the Application of *external Medicines:* for though ſome Phyſicians uſe nothing but heating Remedies, and others only cooling Medicines, to appeaſe the Inflammation; yet both of them, when applied indiſcriminately, may prove both uſeleſs and pernicious: for one Medicine is not to be applied to every Patient, but particular Remedies are to be ſuited to the Strength and Conſtitutions of particular Patients; or elſe Injury might follow upon the Application of hot Medicines to hot Conſtitutions, and the contrary: I therefore look upon it to be Matter of Conſequence to obſerve diligently, that cooling Medicines be applied to ſuch as are of a hot Temperature. Among the Coolers, the principal are *Acetum Lithargyriſatum*, applied warm by linen Rags folded together; or *Acetum calidum Minio Bolove permixtum*; or *Oxycratum ex æquis Aceti et Aquæ Portionibus confectum*: of each of thoſe Liquors may be taken, for Example, ℥vi. *Salis communis* ℥j. *Nitri vel Salis Ammoniaci* ℥ij: let them be mixed, and applied to the affected Limb with linen Cloths: among the vulgar, common, or domeſtic Medicines, the *Stercus bubulum recens atque calidum Aceto calidiori admixtum*, is an Application very eaſy to be had, and of no ſmall Efficacy: pickled Cabbage-leaves, Broth, Brine, *&c.* are alſo ſometimes uſed with Succeſs to the inflamed Limb: ſome prefer cooling Plaſters, as the *Emp. ad Ambuſta, de Minio, de Lithargyro, Diapompholygos, Saturninum, &c.* theſe Plaſters may do pretty well in the ſlighter Inflammations, for ſuch Patients as have a good Opinion of Plaſters; particularly they will do very well in the Night-time, when the Preparation and Application of Fomentations are difficult and troubleſome.

Remedies proper in cold or phlegmatic Habits.

XVI. In cold and phlegmatic Patients, *Sp. Vin. rectificat. Sp. Vin. Camph. vel pauxillo Theriac. permiſt.* are very ſucceſsful in diſperſing Inflammations, being often applied by means of hot Cloths; ſo is alſo the *Aqu. Calcis, vel Mera, vel cum Sp. Vin. Camph. Aq. Reg. Hungar. Bolo, Ceruſſa, Lap. Calaminari, Sale Ammoniaco, aut Lythargyro permixta:* a Mixture of *Sp. Vin.* ℔j. and *Sapon Venet. vel Hiſpan.* ℥ij. being applied warm, gives Place to hardly any Medicine for diſperſing an Inflammation. Laſtly, there are many Herbs proper for this Purpoſe, as *Scordium, Abſinthium, Mentha, Sabina, Abrotanum, Matricaria Flor. Tanaceti, &c.* which may be diſcretionally made into a Decoction with *Aq. Salſa, Marina, vel Calcis:* with this Decoction may be mixed *Sp. Vin. Rect. vel Camph. & Sap. Venet.* by which means its Virtue will be increaſed: the forementioned Herbs may be alſo commodiouſly boiled and made into a Cataplaſm, and applied in the ſame Manner with the reſt of the Medicines, *i. e.* by linen Cloths folded together, and bound round the diſordered Member.

External Applications how to be made.

XVII. There remains one Thing to be eſpecially obſerved, with regard to the Application of external Medicines in this Caſe: namely, that each of them muſt be always applied hot, and never be permitted to grow firſt cold: the Inflammation alſo diſperſes generally much more ſpeedily, when the diſordered Limb is firſt rubbed well with a Cloth dipped in ſome warm diſcutient Fomentation, before any freſh Cataplaſm be laid on: and this Method is to be continued, till the Inflammation be either wholly diſperſed, or elſe brought to an End by Suppuration or a Gangrene.

XVIII. In

XVIII. In the mean Time the Surgeon ſhould carefully obſerve, that the Apartment where the Patient lies be neither too hot nor too cold, but be kept as near as poſſible to the Degree of temperate Air, alſo to procure Reſt and Sleep to the Patient, and to let him not be kept awake too long: laſtly, to let the Patient keep his Mind free from pernicious Paſſions, as Anger, Fear, Care, great Thought, *&c.* What Sort of Life the Patient ſhould lead.

CHAP. III.

Of SUPPURATION *and* ABCESS.

I. WE obſerved before, that the ſecond Way in which an Inflammation went off, was, by Suppuration; that is, a Converſion of the inſpiſſated Blood, and the ſoft adjacent Parts (as the ſmall Veſſels and Fat) into *Pus* or Matter; which Diſorder, when it has not yet found an Opening, is generally called by Surgeons an *Abceſs*. Suppurating what.

II. An Inflammation may be known to tend to Suppuration from the Signs before-mentioned at *Chap.* II, § VII, which generally happens, when the Inflammation has been of long ſtanding when the Surgeon is called in, or when it cannot be diſperſed by the Uſe of the forecited Remedies. Signs of Suppuration.

III. As ſoon as we find it tend to Suppuration, we muſt wholly lay aſide the Uſe of reſolving Medicines; and we muſt ſtrive, (1.) to forward the Inflammation to Maturity, *i. e.* to convert the ſtagnating Blood into laudable Matter; then (2.) to procure a Diſcharge or Vent for this ſuppurated Matter: (3.) to let the diſordered Part be well cleanſed from all that is corrupted: and laſtly, (4.) to incarn, agglutinate, and heal the wounded Part. What is to be done in Caſe of Suppuration.

IV. As to forwarding the Inflammation to *Suppuration*, that is to be promoted by particular maturating Remedies: among which, the beſt ſeem to be ſuch of the Emollients as obſtruct the Pores of the Skin, as Fats, Oils, glutinous and ſlippery Medicines; as alſo the Application of ſharp, pungent, and ſomewhat cauſtic Medicines, made up and uſed in the Form of a Cataplaſm: or Plaſters of the like Kind may be applied to the diſordered Part. Maturation, how promoted.

V. Among the emollient Medicines for this Purpoſe, there are ſeveral Kinds of Herbs, Fruits, Seeds, and Meals that may be here enumerated; as the *Althæa, Malva, Lilia, Parietar. Verbaſc. Branca Urſina, Solanum, Hyoſcyamus, Ficus, Semen Lini, Fœnu-græci, ejuſdemque Seminis Farina; Farina item triticea aut ſiliginea, Panis primarii & ſecundarii Micæ, Vitelli Ovorum, Butyrum, Mel, variorumque Animalium Pinguedines vel Adipes, Oleum Lini, Olivarum, Liliorum alborum, Chamæmel.* and many others of the like Kind: as to the other Claſs of Maturaters, which are ſharp, pungent, and ſtimulating, but alſo emollient at the ſame Time, there may be reckoned *Chamæmelon, Melilotum, Cepæ ſub Cinere toſtæ, Allium, Crocus, Terebinthina; variaque Gummata, Galbanum inprimis, Ammoniacum, Bdellium, Opopanax, Sagapenum, in Vitellis Ovorum reſolutum; & denique Fermentum Panis.* Simple maturative Remedies.

VI. From a proper Mixture of the now-recited Simples, may be made various and uſeful Cataplaſms and Plaſters for this Purpoſe: it may be not unacceptable here to inſtance a few of the moſt proper and efficacious of theſe compound Maturatives. Compound Maturatives.

1. ℞ *Herb. Malv. Alth. Parietar. Chamæmel. aa* M j. *Farin. Sem. Lini vel Fœnugræci* ʒ ij. *Coq. leni igne Aqua vel Lacte, ad Conſiſt. Cataplaſmatis, poſtea add. Fermenti Panis* ʒ ij, *Gum. Galban. in Vitell. Ovi reſoluti* ʒ j. *Dein Linimentorum convolutorum Adminiculo, calida, & quam ſæpiſſime quidem ſupra læſum Membrum dilegantur: vel,*
2. ℞ *Fol. Malv. Branc. Urſin. aa* M ij. *Caricar. Pinguium contuſar.* N° vj. *His eadem, ut modo retulimus, ratione decoctis adde Butyri recentis, nec non Ceparum ſub Cineribus toſtarum. aa* ʒ ij. *& denique Farinæ Sem. Lini, quantum quidem ad Cataplaſma conficiendum ſufficit, admiſcendum: vel,*
3. ℞ *Rad. Lilior. alb.* ʒ ij. *Herb. Parietar. Mercurial. Melilot. aa* M j. *Ficuum recent. contuſ.* N° vj: *hæc in Aqua penitus coccoquantur, admixtiſque Gumm. Ammoniac. & Sagapen. in Vitellis Ovorum ſolutor. ut & Aceti boni aa* ʒ j ß. *in Cataplaſma quoddam convertantur: vel,*
4. ℞ *Farinæ Siligineæ, aut Triticeæ* M ij. *vel* iij. *coq. in ſ. q. Lactis admiſc. Gumm. Bdellii & Opopanacis cum Vitellis Ovor. ſubactor. aa* ʒ j. *ut & Croc.* ʒ j. *in Cataplaſma tranſmutentur: vel,*
5. ℞ *Fermenti Panis* ʒ iij. *Mellis* ʒ j. *Saponis Veneti comminuti* ʒ ß. *Olei Lilior. alb. q. ſ. F. ſuper leni igne Cataplaſma: vel,*
6. ℞ *Mellis* ʒ iv. *ad lentum ignem ex Aqua decoquantur; his poſtea Olei Lini aut Chamæmel. pauxillum, ut & Farinæ Siligineæ aut Sem. Lin. quantum ad Malagma parandum ſatis eſt, admiſceatur.*

Theſe Cataplaſms, or others of the like Nature, are to be often applied hot to the Part affected, till the Matter within appears to be ſufficiently digeſted or maturated by the Softneſs and Whiteneſs of the Tumor: but when the Abceſs is of the ſmaller Kind, it is every Way more commodious to apply ſome maturative Plaſter, as *Empl. Diachyl. cum Gumm. vel & Emplaſtrum ex Melle & Farina compoſitum:* theſe may be applied to the Part affected, till Suppuration enſues.

Internal Medicines proper to be uſed.

VII. In the mean Time, when the Patient's Condition requires it, we muſt be careful to temperate the Motion of his Blood, not by external Applications only, but alſo by internal Medicines and a proper Regimen: when the Blood moves too ſlowly, as may be known by the Pulſe, the Patient ſhould moderately uſe Meat, Drink, and Medicines, which are warm and ſtimulating: by which Means the inſpiſſated Blood contained in the ſmall Veſſels may be the more eaſily converted into Matter, by the increaſed Motion of the Blood: ſtrong Broths, Wines, and Ale are alſo very effectual for the ſame Purpoſe: but where theſe are inſufficient, and the Pulſe indicates that the Motion of the Blood is ſtill ſlower, it will be proper to order the *Theriaca, Diaſcordium,* or *Alkermes* to be taken a Bit upon the Point of a Knife ſeveral Times in a Day, or diſſolved in Wine, Cinnamon Water, or ſome other cordial Liquor: in the mean Time we muſt not neglect the *Tinctura Bezoard. Eſſent. Alexipharm. Diaphoret. Eſſent. Cinnam.* with other warm cardiac and comfortable Eſſences, Spirits, and medicated Teas, by infuſing a few Saſſafras Chips, red Sanders, Cinnamon, *&c.* but on the contrary, when the Motion of the Blood appears by the Pulſe and great Heat to be too violent, then cooling Medicines muſt be directly ordered, to aſſwage and temperate the Heat and Motion, and to prevent a Gangrene:

to

to this Head belong all Sorts of thin and watery Drinks, with ſubacid Medicines and abſorbent Powders with Nitre, as we mentioned in *Chap.* II. § XI: it is alſo ſometimes proper in this Caſe to open a Vein, and bleed a little. Laſtly, when the Strength of the Conſtitution is not impaired, but remains firm, and the Motion of the Blood and Pulſe appear to be neither too ſwift nor too ſlow; unleſs there be ſome urgent Symptom, the Uſe of internal Medicines to promote the Suppuration, ſeems to be wholly unneceſſary, if the Patient keeps up to a proper Regimen.

When the Abceſs ought to be opened.

VIII. With regard to opening the Abceſs and diſcharging its Matter, it is a Caution very neceſſary to be obſerved, that the Opening be not made too ſoon, before the Matter has arrived at a perfect Maturity: for elſe, the Diſcharge of the Matter will be not only impeded, but the Part will, in all Probability, be flung into a greater Inflammation. This has indeed been the common and conſtant Practice hitherto; but ſome of the Moderns (among whom is GOUEUS, a *Frenchman*, p. 259, of his Surgery) will have it proper to open the Tumor directly, without waiting a Suppuration, if it cannot be quickly diſperſed: which I alſo find to have been the Advice of CELSUS (*p.* 408.) formerly. But to return, an Abceſs is known to be ſufficiently ripe, when the Tumor, which before reſiſted, feels ſoft and pliant; when it turns pale or of a yellow Colour; when, upon applying the Fingers, you perceive a Fluid to be lodged within; when the Pain, Redneſs, Heat, and Pulſation go off, wholly or in Part, and the Senſation of a Heavineſs or Weight ſeizes the diſordered Part in the Room of the former: yet ſometimes there are Abceſſes, as I myſelf have more than once experienced, when the Skin does not change its Colour; but on preſſing the Tumor you perceive a Softneſs, and a Fluctuation within. In theſe Caſes the other Surgeons and Phyſicians were either doubtful of the Abceſs, or flatly denied it: nevertheleſs, when I opened the Tumor, there flowed out immediately a large Quantity of Matter. When the above-mentioned Signs appear, the Abceſs muſt be opened in the moſt prominent and depending Part without more Delay; for Delay generally proves of a worſe Conſequence than opening it too ſoon, tho' both of them are bad: for when the Matter is retained longer than it ſhould be, in a large Suppuration and nervous Part, there is Danger leſt the corrupted Matter ſhould corrode the adjacent Parts, and produce *Fiſtulæ* or a *Caries* of the Bones; or by inſinuating itſelf into the ſmall Veſſels, and corrupting ſuch Parts of the Blood as it mixes with, it may excite ill-conditioned Fevers; or laſtly, by diſturbing the Functions of the Brain, Lungs, Liver, and Kidneys, it may bring on Inflammations and Suppurations, and at length Death itſelf. Sometimes the moſt ſubtile Part of the Matter perſpires, and only the groſſer Parts are retained behind, which gives Riſe to hard Tumors, eſpecially in glandular Parts; ſince theſe muſt therefore be the Conſequences, if the Abceſs be not timely opened, the Surgeon's great Care muſt be to uſe the proper Opportunity, and to make an Opening by the uſual Methods where the Skin appears to be the thinneſt: the Methods for making this Opening are principally two, either by *Inciſion* with the *Scalpel*, or by making an *Eſchar* with a *Cauſtic*.

How the Inciſion is to be made.

IX. The Parts which are now ſuppurated are to be inciſed in the following Manner: the Surgeon is to graſp the Baſis of the Tumor with one Hand, preſſing the Matter outward towards the Skin, to avoid hurting any Veſſels or

Nerves in the ſubjacent Parts; he is then to make the Inciſion by the ſharp Scalpel (*Tab.* I. A or B) in his right Hand; making the Opening in the ſofteſt and moſt depending Part of the Abceſs, that the Matter may have the freer Exit. When the Abceſs is large, the Scalpel is not to be taken out as ſoon as the Opening is made, but the Inciſion in the Skin is to be further enlarged with it; but with ſo much Caution as to avoid the larger Veſſels and Nerves, with the Muſcles of the ſubjacent Parts. The Abceſs thus opened, the putrid Matter is to be let out; and when it is glutinous and thick, it may be gently preſſed forth with the Hands: but if the Quantity of Matter contained be very large, and the Patient not bold enough to bear the Knife, but faints away, which is often the Caſe; then the beſt Way ſeems to be, to diſcharge the Matter in Part, and fill up the Cavity with Lint: and after the Patient has been recovered by *Aq. Reg. Hungar.* or ſome other Cordial, to complete the Dreſſing with a Plaſter, Compreſs, and Bandage, leaving the perfect Diſcharge and Cleanſing thereof to the next Dreſſing: but if no *Deliquium* happens, the Matter may be all diſcharged at one Time. The remaining Treatment of this Ulcer is to be the ſame as we have directed before in Wounds: in the firſt Place, the Abceſs is to be cleanſed with Digeſtives: afterwards ſarcotic or balſamic Medicines are to be applied, till the Wound is filled up internally with new Fleſh, and externally cloſed or cicatrized: Tents, particularly of the harder Kind, muſt be here cautiouſly avoided, as they generally produce Ulcers which are very difficult to cure: it is much ſafer to fill up the Cavity with Doſſils of Lint, and to remove them once or twice a Day as there is more or leſs Matter.

How the Matter may be diſcharged by a Cauſtic.

X. The other Method of opening an Abceſs is, by Means of a Cauſtic or corroſive Medicine; and is generally uſed for Children and ſuch as are of a tender Conſtitution, who are very much affrighted at the Approach of the Knife or Scalpel for Inciſion. Among theſe cauſtic or corroſive Medicines, the moſt commendable and proper are, the *Lap. Cauſt. ex Cineribus clavellatis & Calce viva vel ex Lixivio Saponariorum paratus:* alſo the *Lap. Infernalis, Butyrum Antimonii*, and ſuch like, of which there are ſuch Abundance, that almoſt every Apothecary and Surgeon has now his proper Cauſtic, made after his own particular Method, which is ſuppoſed to excel the reſt. The *Lapis cauſticus* is to be applied to the Abceſs either whole in the Lump, or elſe beat ſmall, as may beſt ſuit the Occaſion: but then a defenſative Plaſter muſt be firſt applied to the Abceſs, perforated with an oblong narrow Aperture, much as we have delineated in *Tab.* II. *Fig.* 11: for thus a proper Proviſion is made againſt the ſpreading of the Cauſtic beyond its due Bounds, making its Way through the Skin only in a ſmall or narrow Compaſs: over the Cauſtic is to be applied a Compreſs of Lint or Linen, over the Compreſs a large Plaſter, and over the Plaſter a ſtill larger Compreſs of Linen: and to keep all on firm, a proper Bandage muſt be applied. Things being thus managed, the Patient is to compoſe himſelf to reſt for a while, and the Dreſſing ſhould not be taken off from the Abceſs for the Space of ſeveral whole Hours: three Hours is the leaſt, but ſometimes it requires four, five, or ſix Hours to make an Outlet to the Matter by Cauſtics, in Proportion to the Thickneſs of the Skin and Strength of the Medicine. When the Cauſtic is judged to have remained long enough upon the Abceſs, the Dreſſing muſt be then taken off, that the noxious Matter may be diſcharged: but

if

if the Cauſtic has not ſufficiently penetrated, the Opening may be forwarded and enlarged by gently applying the Scalpel, Probe, or Spatula, that all the Matter may have a free Paſſage: but as ſoon as the Cauſtic has made an Eſchar or Cruſt, it muſt be mollified by applying *Butyrum recens*, *Ung. digeſt. vel Baſilic.* to be retained by a Plaſter and Bandage: when the Eſchar is found looſe or ſeparated, the reſt of the Treatment muſt be the ſame with that we mentioned before, in opening the Abceſs by Inciſion: but to ſay Truth, without Diſſimulation, I muſt acknowledge in my Opinion and Advice, that the Knife is greatly preferable to the Cauſtic, as being more neat, expeditious, and ſafe, and the Aperture heals with a ſmaller and neater Cicatrix: ſo that moſt prudent Surgeons do, with Reaſon, always propoſe the Knife before a Cauſtic, uſing the latter only in Caſes of great Timidity, and where the firſt cannot be conveniently admitted.

XI. That our Reader might not be at a Loſs for the Compoſition of the *Lapis cauſticus*, we thought it would not be amiſs here to lay down a ſhort and approved Method of making the ſame: ℞. *Ciner. clavellat. & Calc. viv. fortiſſ. a a* ℥vj. *vel. Ciner. clavellat.* ℔j. *Calcis vivæ* ℥vj. theſe being pulveriſed ſeparated[a], and afterwards mixed together in a large Glaſs, or earthen Veſſel, are there to be diſſolved in a good deal of Water, letting them ſtand an Hour or two to melt perfectly: then the Liquor, with what it has diſſolved, is to be filtrated through a linen Cloth from its groſs Sediment, evaporating it afterwards in an Iron Pan over the Fire: the conſiſtent Maſs, left after Evaporation, is to be put in a Crucible, and melted with a ſtrong Fire, ſo that it may flow like Oil: it may then be caſt into a Mortar or broad Pan, and either cut or beat into ſmall Pieces before it is quite cold, which are to be put into a Glaſs very cloſely ſtopped, and preſerved in a dry Place for Uſe. When an Abceſs is to be opened, a ſufficient Quantity of this is to be taken and applied, either whole or in Powder, and bound upon the Skin, as we obſerved before: if the Cauſtic be wetted, it generally acts a great deal ſooner, ſo as to corrode the ſubjacent Parts, and make an Eſchar in an Hour or two: but when it grows old, by long keeping, it commonly loſes its Force, ſo that at length it cannot corrode at all. Another and no contemptible Method of preparing this Cauſtic may be ſeen in the Chemiſtry of LEMERY, in the *Leyden* Chemical Collections, and the Surgery of DIONIS, *Edit.* 2. *p.* 709.

The Cauſtic how prepared.

CHAP. IV.

Of TUMOR *and* INFLAMMATION *in the* BREASTS.

1. WE have been hitherto treating of Suppuration: our next Buſineſs was to have proceeded to a Gangrene: but as there are ſeveral Kinds of Inflammation and Suppuration which do not commonly terminate in a Gangrene, it was proper firſt to treat of theſe ſeparately, before we came to the Conſideration of a Gangrene. We begin with thoſe Inflammations which

What happens in an Inflammation of the Breaſts.

[a] BOERHAAVE, in his *Materia Medica* Sect. 412. takes *Ciner. Clavell.* ℥iv. *Calcis. vivæ* ℥vj. and uſes another Method of Preparation, which did not ſucceed with me: you will find a more compendious Preparation of it in the *London* Diſpenſatory: *Ex Calcis vivæ* ℔ *& Cinerum Clavellat.* ℔j.

uſually

uſually afflict the Breaſts, being a Diſorder moſt incident to Child-bearing Women, and almoſt conſtantly happens in a few Days after their Delivery: if the Milk ſhould be impelled into the Breaſt too plentifully and forceably, which at ſuch Times frequently happens, and if the Mother ſhould then be ſeized with great Cold, Fear, or Anger, the ſanguiferous and lactiferous Veſſels being thence obſtructed, the Breaſts muſt then become inevitably tumified, and at the ſame time they will be afflicted with great Heat, Redneſs, Reſiſtance, and violent Pain: the ſame Accident ſometimes happens to Women that give ſuck, even a long Time after their Lying-in: which proceeds from the ſame Cauſes which we juſt now mentioned; and is alſo ſometimes the Caſe of thoſe who have no Milk. I have even obſerved the ſame Caſe in a Man of a weak Habit, which aroſe from a great Fright: one Breaſt was vaſtly tumified, and turned to an Abceſs, from which, upon the firſt Opening, I extracted above two Pounds of Matter, to the great Surprize of the Patient and the Byſtanders. This Kind of Inflammation is uſually attended with a Fever or great Heat all over the Body, followed with a quick Pulſe, Thirſt, Head-ach, and difficult Reſpiration: and this in ſuch a Manner, that a Shivering generally proceeds in its Invaſion.

The Cauſes of an Inflammation in the Breaſts.

II. The general Cauſes of Inflammation in the Breaſts of Child-bearing Women, are uſually, as we hinted before, a ſudden Cold taken when the Body is very hot or in a Sweat, cold Drink, Anger, Fear, Grief, and any other violent Perturbation of the Mind, from whence the Blood and Milk may become inſpiſſated and obſtructed in the ſmall Veſſels of the Breaſt: and tho' the Inflammation of the Breaſts happens moſt frequently in Women, eſpecially ſuch as have lately lain in, and either will not ſuckle the Infant or cannot; or when the *Fœtus* died in the Womb, or ſoon after the Birth, in which Caſes it proceeds from the Stagnation of the Milk brought on by Fear and Grief: yet it may frequently happen from the ſame Cauſes in ſuch as have left off giving ſuck for a conſiderable Time, as alſo from a Blow, Contuſion, or ſome other external Injury of the like Nature.

The Difference of theſe Inflammations.

III. This Imflammation does not always happen to be equally intenſe and violent: for ſometimes it ſeizes the whole Breaſt, ſometimes only one Side, and greatly tumifies it with violent Pain; but then again, at other Times it occupies only a ſmall Part of the Breaſt: in one Patient the Inflammation lies very near the Skin; in another ſpreads very deep: at one Time the Inflammation has very urgent Symptoms, as violent Pain, Heat, Redneſs, and Tenſion; but at other Times it ſits very eaſy upon the Part.

Prognoſis.

IV. He that is deſirous to be an able Preſager in the Events of this Kind of Inflammation, ſhould firſt carefully conſider the ſeveral Symptoms of the diſordered Part now mentioned: for as the Tumor is leſs, and the Inflammation and Fever ſlighter, the more gentle and happy is like to be the Conſequences, and the leſs is the Danger: for in that Caſe there is room to hope it may be diſperſed, without coming to Suppuration: but on the contrary, the more violent the Symptoms, the greater is like to be the Suppuration: ſometimes it turns to a *Scirrhus*, and a *Scirrhus* commonly ends in a Cancer of the Breaſts.

Inflammations of the Breaſts, how to be treated.

V. This Diſorder may be very readily prevented in Women of Condition, and ſuch as cannot, or will not, ſuckle their Children, if ſome of the *Emplaſt. de*

Spermate Ceti ſpread on Linen be applied warm all round upon the Breaſt ſoon after Parturition, being perforated in its Middle to tranſmit the *Papilla* or Nipple; the Acceſſion of the Milk being alſo repelled by a pretty ſtrict Bandage: it may be alſo not improper in this Caſe to hang the *Galactites*, or ſome *Argentum vivum* incloſed in a Nutſhell, about the Patient's Neck, down the Back; and to apply *inter Scapulas Emp. ex Spermate Ranarum, Saccharo Saturni, Oleoque Hyoſcyami permixtum.* Among the internal Medicines, the moſt proper are ſuch as bring down the *Lochia Puerperarum*, when they do no not flow in ſufficient Plenty of themſelves: the principal for this Purpoſe are *Eſſent. Myrrhæ, Succin. Eſſen. Croc. Elix. proprietat. &c.* taken now and then in a proper Doſe: laſtly, with reſpect to the proper Diet, it muſt be carefully obſerved to diminiſh the Quantity of Milk by the Smallneſs and Poverty of the Meat and Drink: upon which Account the Patient ſhould be recommended to drink nothing but ſmall Broth, Tea, or the like watery Liquors, for many Days together, till the Afflux of Milk to the Breaſts is found to be ſufficiently weakened: but if the lying-in Mother be deſirous of ſuckling the new-born Infant herſelf, there can be no better Preſervative for her againſt Inflammations of the Breaſts, than to keep free from Colds, and to cautiouſly avoid all violent Affections of the Mind, letting the Child ſuck frequently at proper Seaſons, to prevent the Milk from Stagnation: beſides this, Care muſt be taken to uſe Plenty of ſmall Broth and thin Fluids for the firſt Week or two; by which Means the Milk will not be ſo abundant, nor apt to be inſpiſſated in the lactiferous Ducts of the Breaſts.

VI. But when Inflammation and Tumor have already fixed themſelves in the Breaſts, the Surgeon's principal Buſineſs is, to uſe all Endeavours to diſcuſs whatever ſtagnates in the ſmall Ducts and Veſſels with the utmoſt Expedition, both by internal as well as external Medicines; in order to prevent the Tumor from running into Suppuration or *Scirrhus:* for when it ſuppurates, there generally remains an ugly Cicatrix, which is very diſagreeable to moſt Women, but eſpecially the more noble and elegant: as to the internal Medicines proper to be given to Child-bed Women, to diſperſe Tumors in the Breaſts, which are generally accompanied with a Fever, I would adviſe the Surgeon and Patient to conſult ſome prudent and ſkilful Phyſician on that Head; leſt the lacteal Fever (as it is generally called) carry off the lying-in Patient under an injudicious Treatment. Cure, (1) by Diſperſion.

VII. As to the external Remedies, in which the Surgeon ought to be particularly ſkilled, the ſtrongeſt Diſcutient that I have frequently found to excel others for theſe Tumors, is, the *Emplaſtrum ex Sperm. Ceti præparat:* in the mean Time it may be of ſome Service to lay over the Plaſter a diſcutient Bag, made warm and ſtuffed *ex Furfure ac Sale, vel Flore Samb. Chamæmel. Melilot. Lavend. vel ex Sem. Fœmin. Cumin. Aniſ.* &c. There are ſome who put Lambs Skin over the Plaſter inſtead of diſcutient Bags, which not only defends the Breaſts from external Cold, but is alſo no improper Diſcutient for what ſtagnates in them: but there is ſtill a uſual and very effectual diſcutient Application for theſe Tumors, which is a Calf's Bladder filled with a warm Decoction of *Flor. Samb. & Chamæm.* in Milk, which is to be often applied to the Breaſt, its Warmth being renewed as it is impaired: of nearly the ſame Virtue is the *Emp. Diachyl. ſimp.* either alone or mixed with *Emp. de Spermate Ceti:* the *Rob. Sambuci* External Reſolvents.

Sambuci or *Theriaca* mixed *cum Sale Abſinthii*, being ſpread upon Linen, and applied in the Way of Liniment, prove of great Efficacy in diſperſing theſe Tumors, eſpecially if they are applied warm, and covered with warm diſcutient Bags; but they are hard to be put up with among the rich and very nice Women, becauſe they uſually dawb the Skin, Cloaths, and Bedding: to theſe we may add the Uſe *Acet. Lithargyr. Acet. cum Semine Carui Aq. Calcis*; which are of very eaſy and conſiderable Uſe; being applied to the Breaſts by Means of linen Compreſſes dipped in the Liquors while hot, and often repeated: a great many eſteem it a ready and effectual Remedy to expreſs the Milk upon burning Coals; nor do I think it proper to raſhly reject this Method as wholly uſeleſs: for though this Sort of Cure ſeems to be ſympathetical and ſuperſtitious; yet as it may excite a ſtrong Imagination of drying up the Milk in the ſuperſtitious Woman, and that Imagination may have a conſiderable Influence, we ſee no ſufficient Reaſon intirely to condemn it: but if the Breaſts are internally very much diſtended with Milk, it will be proper to diſcharge it by the ſucking either of an Infant, an old Woman, or a Puppy, or elſe by the Application of a Glaſs Inſtrument which we ſhall hereafter deſcribe: the Milk ſhould be thus diſcharged till the Tumor ſubſides and the Pain vaniſhes.

(2) by Suppuration.

VIII. But when the Inflammation is greater than can be diſperſed in the Space of four or five Days; or when, as it frequently happens, the Surgeon is conſulted too late; the beſt Way is to forward it to Suppuration as faſt as poſſible, rather than hazard its turning to a *Scirrhus* or Cancer by Delay: if therefore the noxious Matter be not arrived at a State of Maturity by the Uſe of the diſcutient Medicines, in order to accelerate the Suppuration there ought to be a ſpeedy Application of an *Emp. Diachyl. cum Gumm.* or *Emp. de Hyoſcyamo:* but more effectual Cataplaſms are to be alſo made Uſe of to digeſt the Matter, ſome of which we mentioned in the preceding Chapter, § 5 and 6, and others we ſhall alſo propoſe here: as,

1. ℞ *Farinæ Siligin.* ℥ ß *vel* ℥ j. *Melliſque quantum ad Cataplaſma conficiendum ſufficit: tum Lactis & Croci pauxillum admiſceatur, calefactumque in Patella quadam linteamentis obducatur, Mammiſque ſuperimponatur, ac ſæpius poſtea renovatur.*
2. ℞ *Farinæ Siligin.* ℥ iv. *Gummi Galbani Vitello Ovi reſoluti* ℥ j. *Aceti* ℥ iij. *his Aquæ tanta Portio admiſceatur, quanta Cataplaſmati coquendi ſufficit: vel,*
3. ℞ *Fermenti Panis* ℥ ij. *Mellis* ℥ ß. *Saponis Venet. comminuti, & Oleo Chamæm. ana* ℥ ij. *quæ ſibi invicem commixta in Patellam conjiciantur, atque Igni admota in Pulticulam ſive Malagma convertantur.*

How the Abceſs is to be opened.

IX. Theſe Cataplaſms are to be applied hot, and very often, to the Breaſts; keeping them on by linen Compreſſes or Bolſters, the better to retain the Heat, till the Tumor breaks of itſelf, which it often does in this Part, from the Thinneſs of the Skin: or elſe, when ripened, it may be conveniently opened by the Scalpel: but the Inciſion ought always to be made in the lower Part of the Breaſt, unleſs Neceſſity obliges it to be otherwiſe, leſt there ſhould be left a viſible Cicatrix after the Cure: though there are not wanting ſome Surgeons who uſe the Cauſtic for opening Suppurations of the Breaſt, yet, as they uſually occaſion diſfiguring *Cicatrices*, we think the Knife is greatly preferable to ſuch Medicines.

X. After

X. After the noxious Matter has been diſcharged from the Breaſts, the reſt of the Treatment is to be the ſame with what we propoſed in the Cure of other Wounds and Abceſſes: the Ulcer is to be firſt cleanſed with ſome digeſtive Ointment, and afterwards healed with ſome Balſam, as the *Peruvian* for Example; with Oil of Eggs and Wax: but when the Suppuration has run very deep, the beſt Way is to inject the Wound with a cleanſing Decoction of the *Sanicula* or *Alchimilla* mixed with a little *Mel Roſarum*; and to prevent the Lips of the Wound from cloſing, before the Bottom is filled up with new Fleſh, it will be proper to introduce a ſoft Tent, or ſome ſcraped Lint: as the new Fleſh grows up from the Bottom, the Tent may be gradually leſſened or made ſhorter, and, at laſt, wholly removed when there is little or no Occaſion for it. Treatment after the Diſcharge of Matter.

XI. But it ſometimes happens that Tumors in the Breaſts of Child-bed and ſuckling Women will neither yield to Diſperſion nor Suppuration, but will retain their ill Condition for the Space of ſeveral Months or Years: if this happens in young and healthy People, it occaſions little or no Diſturbance to the Œconomy: nor is there great Danger of the Tumor's turning to a *Scirrhus* or Cancer, which the poor female Patient is often vaſtly afraid of: the Surgeon's Buſineſs here is to take Care to keep the afflicted Patient in good Heart by his Perſuaſions: and to the Tumor itſelf is to be applied *Emp. de Spermate Ceti cum Pauxillo Camphoræ, vel Diaſapon cum Camphorâ*, to be conſtantly kept on, and the Breaſt muſt be carefully defended from the external Cold; by which Means Tumors of long ſtanding have grown gradually leſs, and at laſt vaniſhed: but the Caſe is uſually otherwiſe in Women, who are advanced in Years, and of a melancholy or ſorrowful Diſpoſition: for in ſuch there is great Danger of the inveterate Tumor turning to a *Scirrhus* or Cancer. What is to be done when the Tumor can be neither diſperſed or ſuppurated.

CHAP. V.

Of INFLAMMATION *in the* TESTICLES.

I. SOMETIMES an Inflammation and Tumor happens in one or both of the Teſticles: which, if it be any thing violent, generally tortures the miſerable Patient with moſt ſharp Pains. Inflammation does ſometimes happen in the Teſticles.

II. This Diſorder may ariſe from two Cauſes: either from ſome great *external Violence*, as by a Fall, Blow, or Contuſion; to which ſome are liable from mounting a Horſe with too much Haſte and little Thought: or from a *venereal* Cauſe; chiefly when ſome of the venereal *Virus* inſpiſſates the *Semen*, and obſtructs its Courſe through the ſmall *Tubuli* of this Gland. The Cauſes of Inflammation in the Teſticles.

III. An Inflammation of the Teſticle is diſtinguiſhable from any other Diſorder in theſe Parts, and particularly from a *Hernia Scroti*, when the Patient has previouſly ſuffered any of the Cauſes § II. and complains of a great Swelling, Heat, Redneſs and Pain in his Teſticle, the ſame being confirmed to the Surgeon by Inſpection: his Feeling will alſo acquaint him with the Nature of the Diſorder: for, upon applying the Hand, one or both of the Teſticles are found to be ſwelled conſiderably larger than they ought to be, exceeding ſometimes the Size of one's Fiſt. *Diagnoſis.*

Prognoſis. IV. This Diſorder is not of ſo ſlight a Conſequence as is generally thought; for it very frequently turns out ſo as to deprive the Man either of his Life or Virility, by degenerating into an Abceſs or *Sphacelus:* or elſe it turns to a *Schirrhus* or Cancer, which have alſo Death for their uſual Conſequence: or, laſtly, it is followed by a *Sarcocele* or *Hydrocele*, which are little leſs troubleſome and fatiguing to the Patient.

Cure by Diſperſion. V. The ſame external Medicines will ſerve to reſolve the Inſpiſſations which happen in an Inflammation of the Teſticle, which we before oppoſed to Inflammations in the Breaſts: and above others we prefer *Acet. Lithargyriſat. Aq. Calcis cum Sp. Vin. Camph. Ceruſſa, Tutia, Lap. Calamin. permixt. vel & Farin. Fabarum cum Acet. Decoct. in Cataplaſma.* But in the Night-time, when the Application of Fomentations is not ſo convenient, it will be proper to apply *Emp. de Ranis cum duplici Mercurio, vel Emp. Diachylum.* Nor are internal Medicines to be here neglected: for, if the Tumor aroſe from ſome external Violence or an Inſpiſſation of the Blood, he ſhould often take of the *Pulv. ex Lap. Cancror. præp. Teſt. Oſtreor. Mat. Perlar. Cinnab. Arcan. duplicat. &c.* together with thin Drinks, as Tea, Decoctions of the Roots, Woods, and diſcutient Herbs: plentiful Feeding, Things which heat the Blood, and Aliment of difficult Digeſtion, are to be carefully avoided: and if the Inflammation ſhould be of the more violent Kind, it will not be amiſs to mix a little Nitre with the forementioned Powder; and to drop ſome *Sp. Vitriol. Sulphur, &c.* into his Drinks: not neglecting to open a Vein in plethoric Habits.

Treatment when from a Venereal Cauſe. VI. If the Diſorder take its Riſe from ſome venereal Taint, it ſeems neceſſary to adminiſter good briſk Cathartics, always adding a Quantity of *Merc. dulc.* to them: at the ſame Time ſuch other Medicines ſhould be uſed as are calculated particularly againſt the venereal Diſeaſe itſelf: warm Drinks made of Tea, or a Ptiſan of Barley, Liquorice and Aniſe boiled in Water muſt not be here neglected: by taking theſe, the Blood uſually becomes temperate and attenuated, and the Tumor frequently diſperſed.

How a Suppuration is to be managed. VII. Laſtly, if the Surgeon be called in too late, or if the Inflammation prove ſo violent as not to give Way to the preceding Remedies for Diſperſion, a Suppuration or Gangrene is generally the Conſequence: therefore the Application of the ſame maturating Remedies will be here proper, which we propoſed in the preceding Chapter for an Inflammation of the Breaſts: and when the Matter is ſufficiently digeſted, and the Abceſs does not ſoon break of itſelf, it will be proper to open it carefully by Inciſion: the Matter being diſcharged, the Wound is to be firſt well cleanſed by ſome digeſtive Ointment, injecting ſome ſtrong ſpirituous Fomentation which reſiſts Putrefaction, and at laſt healing it with ſome vulnerary Balſam: but firſt, to digeſt the Matter, and mitigate the Pains, it is found extremely ſerviceable to apply *Emp. de Hyoſcyamo, vel Diachyl. cum Gummis*; which are alſo ſtrongly recommended by LUDOVICUS in his *Chirurgical Works, pag.* 718: while theſe Applications are properly uſed, we muſt ſtrive to extirpate the venereal Diſeaſe itſelf: and notwithſtanding, in many of theſe Caſes, the *Scrotum* happens to be conſumed ſo as to leave the Teſticle quite bare; yet the Loſs of Subſtance in the *Scrotum* may be generally reſtored again, by a proper Treatment with digeſtive and balſamic Remedies, as I myſelf have frequently ſeen.

CHAP. VI.
Of an ERYSIPELAS.

I. A *Eryſipelas* is an Inflammation ſeated in the exterior Part of the Skin and *Membrana Adipoſa* beneath it, which wanders and ſpreads ſometimes to a very great Extent, being accompanied with great Redneſs, Heat, and often Pain: upon preſſing the Part afflicted with the Finger, it looks white; but upon removing the Finger it turns red again: this Inflammation has been obſerved to fix itſelf oftenest upon the Arms or Legs; but ſometimes it ſeizes the Head, Neck, Shoulders, and Face[a]; often the Noſe, and ſome other Parts: it generally ſeizes the Patient with a *Horror* or Shivering, after which a great Heat ariſes, equal to what is uſually felt in burning Fevers; and hence it has been diſtinguiſhed, as well by the Antients[b] as Moderns, by the Name of *Ignis Sacer*, or *St. Anthony's Fire*. An *Eryſipelas*, what it is.

II. Any Cauſe that can produce other Inflammations may alſo occaſion an *Eryſipelas*: more eſpecially expoſing the Body to ſudden Cold, when it is in a great Heat or Sweat; an obſtructed Perſpiration, the drinking too much fermented and ſpirituous Liquors; a Surfeit, or Over-feeding; and laſtly, a hot and ſharp State of the Blood: from all which, either aſunder or together, the Blood may be eaſily inſpiſſated, the ſmall Veſſels contracted, and an Obſtruction, with its conſequent Inflammation, be brought on. Cauſes of an *Eryſipelas*.

III. With regard to the Event of this Diſorder, it is obſerved that there is no great Danger, when the Inflammation is but ſmall and properly treated. On the contrary, when the Inflammation is violent, the Habit of the Body ill and infirm, the Diet and Way of Life irregular, or the Part affected expoſed to Cold, neglected, or improperly treated; it is no Wonder if the Inflammation turns to an ardent Fever, an ill-conditioned Exulceration, Gangrene, or *Sphacelus*. But an *Eryſipelas* is more particularly dangerous, when treated with external Applications which are cooling, fat, or oily; and when internal Medicines are taken which heat the Blood, whether Wine, Cordials, Spices, or the like. *Prognoſis.*

IV. In order to cure an *Eryſipelas*, the grand Intention is, to dilute the inſpiſſated Blood, and divide it where it ſtagnates and obſtructs: to effect which, there ſeems to be no better Way than that of giving Plenty of thin watery and warm Drinks, by which, a gentle and laſting Sweat may be excited: for by this Means all Viſcidities in the Blood will be diluted, any Acrimony will be temperated, and what heſitates or obſtructs will be reſolved: and laſtly, the uſeleſs and corrupted Part of the Blood will be ejected by the inviſible Pores of the Skin; by which natural Tranſpiration, the *Eryſipelas* will be happily carried off, as by an inſtant Remedy: heating Medicines of all Kinds, eſpecially the *Tinct. Bezoardica Aq. Epidem.* and other ſuch ſtrong, heating, and ſpirituous Medicines, are, in my Opinion, wholly foreign and improper for this Cure; becauſe the Inflammation is generally more increaſed than abated by the Internal Treatment.

[a] An Example of an enormous *Eryſipelas* in the Face and Eyelids, which laſted two Months, may be ſeen deſcribed by VERDUC on Bandages, *Chap.* III. and another exulcerated *Eryſipelas* in both the Thighs is obſerved by SCULTETUS, *Obſ.* 92.

[b] CELSUS, *variis in locis.*

Uſe of them. On the contrary, Medicines which are temperating and moderately cooling, are here much more ſafe and uſeful: particularly Preparations from Elder, as *Rob. Sambuc.* ℥ ß. *vel Cochlear.* j. diluted in *Aq. Flor. Samb.* In the mean Time may be uſed Tea, Coffee, or a Diet Drink of phyſical Herbs. The Patient's Body is to be carefully defended from the external Cold, and to be kept in a gentle and conſtant Sweat: when the Patient is troubled with great Thirſt, he may drink thin Barley Gruel, and for Variety, a little warm ſmall Beer; for the Main of the Cure generally depends upon moderate Warmth and ſmall Drinks: but if the *Rob. Sambuci* ſhould not be liked by the Patient, ſome diaphoretic Powder may be given in its Room, or together with it, made of the *Teſtacea, Antimon. diaphoret. cum Nitri portiuncula,* in order to excite a gentle Sweat: but then the warm thin Drinks ſhould not be neglected in the mean Time. Laſtly, the Regulation of the Non-naturals proper here, we ſuppoſe to be ſufficiently evident from what we have already ſaid of Inflammations in general, *Ch.* II. *N.* XIII, *&c.*

External Treatment.

V. If the Inflammation in an *Eryſipelas* ſhould be but ſlight, it may then be often cured only by external Warmth: but when violent, external Warmth will not be of itſelf ſufficient, without the Application of Medicines: the diſordered Part is therefore to be covered with *Rob. Sambuci,* ſpread on blue Paper or Linen, over which are to be laid warm Cloths, or diſcutient Bags, as we propoſed before in Inflammations: but the Uſe of the *Rob.* as well as the *Theriaca cum Sale Abſinthii* is ſeldom complied with, becauſe of their Uncleanlineſs, though very effectual in mitigating Inflammations, as we obſerved under Inflammations of the Breaſts: upon which Account, the Uſe of diſcutient Powders is much more frequent: among which, the following ſeems to have the Preference, compoſed *ex Flor. Samb. Glycyrrhiza contrita, Creta præparata, Ceruſſa item ac Myrrha, aa admixtis cum pauxillo Camphoræ:* this is to be applied to the Part between ſoft blue Papers or linen Cloths, over which are to be put little warm Bags: to this we may add the *Pulv. contra Eryſipelas Mynſichti,* which is very efficacious, though not much uſed amongſt the Apothecaries. Laſtly, we need not ſay much here of the green internal or middle Bark of Elder, whoſe eminent diſcutient Virtue in Inflammations is almoſt known by every body, and has been this long Time confirmed by conſtant Experience.

A Caution concerning an *Erpſipelas.*

VI. Notwithſtanding there are ſome who judge liquid Medicines wholly improper for the Cure of an *Eryſipelas*; it muſt yet be allowed, that *Sp. Vin. Camph.* uſed warm, either alone, or mixed *cum Croco vel Theriaca* applied warm with coarſe Paper or linen Rags, are of very great Service here: nor can I paſs by a Mixture, which I have frequently experienced in this Caſe, *ex Aq. Calc. viv. cum Sp. Vin. Camph.* SCULTETUS (*Obſ.* 94.) greatly extols the following liquid Remedy againſt an œdematous *Eryſipelas*; he aſſerts, that he never found any Thing anſwer like it:

℞ *Lixiv. mediocr. ex cinerib. vitis* ℔ j. *Nitri* ℥ j. ß. *Salis commun.* ℥ j. *Aceti vini opt.* ℥ j. *M.*

Univerſals being premiſed, this Mixture may be applied to the Part affected by Means of double Compreſſes warmed and retained on with Bandage; by which Means it has ſurprizingly diſperſed, in three or four Days Time, ſuch large

Tumors of this Kind, as have threatened a Gangrene: in the mean time, other liquid Medicines which are over acid, and almoſt all Obſtruents and Aſtringents, together with fat and oily Things, ſhould be cautiouſly avoided: for it can ſcarce be imagined how vaſtly theſe ſtop up the Pores, and, by hindering the Blood from throwing of its Feculencies by Tranſpiration, fling the Patient into imminent Danger.

VII. Bleeding and Purging ſeem not to be ſo neceſſary in an *Eryſipelas* as in a *Phlegmon:* for whatever is corrupted of the Juices in an *Eryſipelas*, as it lies near the Skin, ſeems to be much more eaſily diſchargeable by Sweat: but when the Heat is too great, the Pulſe too high, and the Blood too abundant, Bleeding in that Caſe cannot but be judged proper: but to keep the Bowels open, Glyſters ſeem preferable to ſtrong Purges. Bleeding and Purging when to be uſed.

VIII. It here frequently happens that an *Eryſipelas* comes to Suppuration; from whence uſually ariſe the very worſt of untractable and ſpreading Ulcers: when this is the Caſe, the Ulcer is always to be carefully cleanſed, and dreſſed with *Ung. Saturnin. vel de Lithargyro vel de Ceruſſa, una cum Emplaſtro Saturnino,* to temperate the Acrimony of the *Serum:* but it is alſo at the ſame time proper to take ſuch internal Medicines as will temperate and ſweeten the Blood, uſing ſometimes ſuch as diſcharge ſharp Humours by Stool: and laſtly, a ſtrict Regimen of Diet muſt be obſerved, till the Ulcers are healed again, which is even then a very difficult Matter to effect: eſpecially when ſeated in the Legs of old cachectical or valetudinary People: ſee SCULTETUS on this Head, *Obſ.* 90. An *Eryſipelas* ſometimes comes to Suppuration.

CHAP. VII.

Of a FURUNCLE, *or* BOIL.

I. A BOIL or Furuncle is a ſmall reſiſting Tumor, with Inflammation, Redneſs, and great Pain, ariſing in the *Membrana adipoſa* under the Skin: as there is no Part of the Body free from being the Subject hereof, ſo the whole is ſometimes ſo miſerably infeſted with them, that the Patient can hardly tell how to ſtir himſelf, or on what Part to lie: not only Adults, but alſo the younger, even new-born Infants are obnoxious to this dreadful Diſorder, which occaſions in them moſt fatiguing Clamour and Reſtleſſneſs. A Boil, what.

II. The *Signs* proper to a Furuncle we ſuppoſe to have been ſufficiently evident in what we but now propoſed concerning its Nature: and although it be apparent from what has been ſaid, that there is no great Danger in this Diſeaſe, when it happens to Adults; yet it ſometimes happens, when they are very numerous in tender Infants, thut they excite not only violent Pains, Reſtleſſneſs, and Toſſings, with Weakneſs, Convulſions, and Epilepſies, but at length even Death itſelf follows: indeed moſt of the bad Symptons that attend Boils, as in all other Inflammations, are owing to a thick and viſcid Blood: the more glutinous therefore the Patient's Blood is, the more Boils will infeſt him, and thoſe of worſe Conſequence. Signs and Cauſes.

III With

Cure.

III. With regard to the Cure, it ſeems to conſiſt chiefly in reſtoring the inſpiſſated and ſtagnated Blood to its former Circulation and free Motion, and that as ſoon as poſſible, by proper Remedies: if the Boils are few in Number, they are ſeldom treated with internal Medicines, the Means generally uſed being only external Remedies: but when they are very numerous, or return again, it is neceſſary to uſe internal purging Medicines, and ſuch as attenuate and cleanſe the Blood: ſo that in adult Patients it ſeems proper to bleed both by the Lancet and Scarification, with Cupping: at the ſame time a ſtrict Regimen of Diet ſhould be uſed, drinking frequently and plentifully of a Decoction of the Woods, and ſuch like Attenuaters of the Blood: the Patient ſhould alſo entirely abſtain from drinking fermented and ſpirituous Liquors, particularly Wine and its Spirit, and from the too frequent Uſe of Tobacco.

External diſcutient Remedies.

IV. When the Diſorder is recent, external Medicines only will frequently ſuffice for the whole Cure, if the Patient obſerve a ſtrict Regimen: for this Purpoſe the following Mixture is of great Service, made of Honey acidulated with Spirit of Vitriol, till the Mixture has acquired a conſiderable Sharpneſs, which is then to anoint the Furuncles: of no leſs Virtue is the frequent touching them with mere *Spirit. Vitriol. aut Sulphuris:* and laſtly, diſcutient Plaſters are often found very ſerviceable here, as *Emp. Diachylum ſimplex, de Meliloto, de Spermate Ceti, vel Diaſaponis.*

How they are to be brought to Suppuration.

V. But if the Remedies hitherto propoſed prove inſufficient to diſperſe the Tumor, either through ſome Neglect, or any other Cauſe; the only Means then left is, to bring it to Suppuration: and indeed the Maturation of the peccant Matter is found a very difficult Taſk in ſome Caſes; inſomuch that the Tumor ſometimes remains wonderfully hard and troubleſome, even after ſeveral Weeks Treatment: ſometimes the ſtagnating Matter becomes ſo acrimonious, from its great Inſpiſſation and long Stay, that the Inflammation degenerates into Ulcers, which grow gradually worſe and worſe, till they end in incurable *Fiſtulæ:* in the mean time, to promote and quicken the Suppuration, it is generally found of great Service to apply *Emplaſtrum ex Melle & Farina confectum, necnon Empl. Diachylum cum Gummis:* and where theſe are inſufficient, to make Uſe of the maturating Cataplaſms, which we before recommended in a *Phlegmon, Book* IV. *Chap.* II. § XVI. and in Inflammations of the Breaſts, *Book* IV. *Chap.* IV. § VIII. though we muſt obſerve here, that Plaſters are much more commodious for Uſe in Infants, than Cataplaſms: laſtly, when the Furuncle is ſufficiently maturated, which we may learn from its Softneſs and yellow Head, we muſt have Recourſe directly to the Scalpel, and having made an Opening, we muſt diſcharge whatever corrupted Matter is therein contained: after this, is to be applied *Emplaſt. Diachyl.* and the Ulcer is to be daily cleanſed from its Matter, till being freed from all Malignity, it is to be healed with Balſamics.

Furuncles in Infants how to be treated.

VI. When ſucking Infants are afflicted with Furuncles, it is proper to give the Mother, or Nurſe, ſome purging Medicine, and to order a ſtrict Regimen and Diet: at the ſame time the Infant ſhould take ſome gentle laxative Medicine, with abſorbent Powders, *ex Lap. Cancror. conch. Mat. Perlar. Pulv. Aniſi & Antimon. &c,* to allay the Acrimony of its Juices: laſtly, thoſe Puſtules and Pimples, which ariſe in the Skin of the Face of ſome People, are no leſs than

ſmall

ſmall Furuncles, and therefore ought to be treated like them: the drinking of Whey and the mineral Waters is extremely uſeful for People who are troubled with theſe.

CHAP VIII.

Of the BUBO *and* PAROTIS.

I. THERE are ſome Kinds of Tumors which ariſe with Inflammation, only in certain or particular Parts, to which they are proper, as in the Armpits, in the Groins, and under the Ears; and theſe are called *Parotids*, when under the Ears; in the other Parts, *Bubos*. The *Bubo* and *Parotis*, what they are.

II. The Diviſion or Diſtinction of theſe Tumors, the *Parotis* and *Bubo*, is generally twofold; into ſuch as are *benign*, or ſuch as are *malignant*: which Diſtinction, as it regards the different Method of Cure, we ſhall explain a little more at large: they are ſaid to be *benign*, (1.) when they ariſe ſpontaneouſly, without any preceding contagious and peſtilential Diſeaſe, as they frequently do in Infants: (2.) thoſe are alſo of this Kind which come after benign Fevers, being a critical Diſcharge of the Diſeaſe: but the *malignant* are ſuch as happen in the Peſtilence or venereal Diſeaſe, and are therefore commonly termed *peſtilential* or *venereal Bubos*. The Kinds of theſe Tumors.

III. With regard to the *Cauſes* of benign *Bubos*, we muſt obſerve, that they ariſe from the ſame internal Cauſes with all the reſt of the Inflammations; that is, from an Inſpiſſation and Obſtruction of the Blood: ſo that they differ from other Inflammations only in the particular Part where they are ſeated, as in the Groins, under the Arms and Ears, where there are many ſmall Glands and much Fat. Cauſes of benign *Bubos*.

IV. Nor is the *Diagnoſis* of theſe Tumors difficult, if we do but conſider whether there has preceded any peſtilential or venereal Cauſe, to occaſion that Tumor and Inflammation in thoſe Parts: but great Care muſt be taken to diſtinguiſh a *Rupture* from a *Bubo* in the Groin; for by an imprudent Inciſion in ſuch a Caſe, the Life of the Patient may be endangered. *Diagnoſis.*

V. When theſe Tumors are benign, their Conſequences are uſually milder and leſs dangerous: becauſe they may be generally either diſperſed or ſuppurated: but a ſpeedy Diſperſion or Suppuration of theſe Tumors is found to be more difficult and of pernicious Conſequences in Patients of an ill Habit: inſomuch, that a Suppuration of them ſometimes produces *Fiſtulæ*, which are very difficult to cure: laſtly, the *Parotides* are the moſt difficult to cure, the inguinal *Bubos* not ſo difficult, and the axillary *Bubos* are the eaſieſt of all, as they generally tend to Suppuration. *Prognoſis.*

VI. In *Bubos*, which are unaccompanied with any other Diſeaſe, eſpecially thoſe of Infants, the frequent taking of ſome cathartic Medicine, with an Addition of *Merc. dulc.* is found to be of great Service; as it draws off the glutinous and inſpiſſated Blood from the Part affected, and at the ſame time thins the whole: other Medicines, which attenuate the Blood, ſhould be alſo uſed, ſuch as we before propoſed for Furuncles: but if there ſhould be any thing of a Fever, the Internal Treatment.

the Advice of ſome prudent Phyſician ought to be called in, who will take Care of the Fever, and treat it with proper Medicines.

External Reſolution or Diſperſion.

VII. When the Inflammation is ſo gentle as to gives Hopes of Diſperſion, it may be proper to apply diſcutient Plaſters externally: as *Emp. Diachyl. ſimplex, de Spermate Ceti, de Galbano, Diaſaponis, vel de Ranis cum Mercurio, &c.* ſince by theſe Means both *Parotides* and *Bubos* have been frequently diſperſed.

Suppuration how promoted.

VIII. But when the Inflammation proves more violent, the Pains more intenſe, and the diſcutient Plaſters avail nothing, we muſt then ſtrive to bring it to Suppuration, by the Application of *Empl. Diachylon cum Gummis*, which is effectual here: if violent Pains alſo afflict the Patient, the frequent Application of digeſting Cataplaſms warm to the Part, will generally not only mitigate the Pain, but alſo greatly promote a Diſperſion, or elſe a Digeſtion and Maturation: Cataplaſms of this Kind may be made of the Crum of Bread and Milk, boiled to a proper Conſiſtence, mixing afterwards a little Butter and Saffron therewith: or Meal with Honey and freſh Butter, reduced to the Conſiſtence of a Cataplaſm over the Fire, may be frequently applied warm, and a little Quantity of *Theriaca* may be added to it with Advantage.

The Treatment after Suppurations.

IX. Cataplaſms like the former, or ſuch as we recommended in a *Phlegmon*, and Inflammation of the Breaſts, ſhould be thus frequently applied warm to the Tumor, till the ſtagnating Matter appears to be ſuppurated: as ſoon as we find this, we muſt directly make an Opening, either with the Scalpel or Cauſtic: ſee before, *Chap.* III. § 10. but great Care muſt be taken in the Opening, not to wound any of the large Veins and Arteries which are near the Abceſs, as the Jugulars and Carotides in the Neck, the Axillaries under the Arm, and the Crurals in the Groin: for a fatal Hæmorrhage might by that means be brought on: as ſoon as the Abceſs is opened, the Remainder of the Treatment is to be the ſame with what we have ſo frequently adviſed in other Abceſſes: more eſpecially it is of Service here to apply *Empl. Diachyl.* as it readily diſperſes or ſoftens any remaining Hardneſs that may adhere to the Mouth of the Ulcer.

CHAP. IX.

Of PESTILENTIAL BUBOS, *as alſo of* CARBUNCLES.

The Kinds of Peſtilential Tumors.

I. PESTILENTIAL Tumors are eaſily diſtinguiſhed by Phyſicians into *Bubos* and *Carbuncles*: and here, by the Name of *Bubo* they comprehend all Tumors, not only ſuch as ariſe under the Ears, Arms, and in the Groins, but alſo in the Neck, Breaſt, Arms, Legs, and other fleſhy Parts of the Body, which ſwell and inflame in peſtilential Fevers; whilſt Nature endeavours to drive out the peſtiferous Matter, which lay concealed in the Body.

Diagnoſis.

II. Peſtilential *Bubos* are diſtinguiſhable from other Tumors, by their happening at a Time, and in Conjunction with the Plague, and from their being accompanied in the Patient with the Symptoms proper to that Diſtemper: for it muſt be here obſerved, agreeable to the Teſtimonies of the beſt modern Writers, who have lived in Time of the Plague [a], that People who are ſeized and in-

[a] As by GOD's Providence I never ſaw the Plague, I cannot write any thing of it on my own Experience; yet I was unwilling to be ſilent on ſo conſiderable a Diſorder, and not mention what has been obſerved and confirmed by the beſt modern Phyſicians: I therefore carefully peruſed ſuch

fected

fected by the Distemper, if they do not die quickly, are shortly to expect these Tumors in several Parts of their Bodies. They appear sometimes sooner, at other times later. In some the Tumors appear before they are taken sick by, or ever perceive, the pestilential Venom: in others, the Tumors are two, three, and four Days, after the Appearance of the Distemper, before they come out; but they are seldom observed to come out later. These Tumors or *Bubos* are sometimes joined with Carbuncles: but though the *Bubos* frequently arise without the Carbuncles, yet the Carbuncles seldom arise without Tumors.

III. It has been this long time observed, particularly in the later Plagues, that such Patients as had Tumors come out, without any very bad Symptoms, had them maturate speedily, and were the soonest free from the Distemper: hence it is not without Reason affirmed, by some of the more learned and modern Physicians, that *almost the whole Business of curing the Plague* consisted in carefully promoting the Eruption of *Bubos* and Tumors; nor that any one could be preserved but by means of those Tumors: while those who rightly cure these *Bubos* do also at the same time rightly cure the Pestilence: the Case being thus, resolving, discutient, and repelling Medicines, together with Bleeding and Purging, are so far from proper in the Cure of the Plague, that by throwing the Venom again into the Blood, they destroy the poor Patient: therefore the chief Business of the Physician or Surgeon here, is, carefully to assist Nature in her Endeavours to throw out the Tumors as soon as possible, and to bring them speedily to Suppuration and Maturity. *Prognosis.*

IV. That this may be effected the more readily, it seems to be much the best Way to order the Patient to keep House upon the first Appearance of the Tumors, or rather to keep in a warm Bed, to be more secure from the Air; for by this Means the Patient rests more securely from the external contagious Air, and by the Use of proper external and internal Medicines, the *Bubos* may be more regularly expelled, and brought to Suppuration. General Treatment.

V. Externally it is very serviceable to rub the tumified Part pretty strongly with the Hands or Cloths, and, what is still preferable, to apply external maturative and emollient Medicines, whereby they will come out the sooner. And we shall also here find great Benefit from the Use of a Cataplasm made *ex Fermento Panis calido, vel solo, vel et cum Sale atque Sinapi contrito.* By means of this, the tense Parts are relaxed and stimulated; whereby the pestilential Matter may be received and cast off from the Blood, and come afterwards to Suppuration. Of the like Virtue are not only the Cataplasms, which we before recommended for suppurating other Tumors, in *Chap.* II. § 16. and *Chap.* IV. § 8. but more particularly those which are made *ex Cepis sub Cineribus tostis, atque cum Theriaca & Butyro subactis, vel etiam ex Pane Triticeo sive Semilagineo interiori, cum Lacte atque Croco probe concocto.* But there are some Surgeons who prefer emollient Plasters to Cataplasms; because the frequent Renewal of the Cataplasms requires the Body to be often uncovered, whereby the Perspiration is impeded and disturbed. The emollient Plasters used instead of the Cataplasms, are the *Empl. Diachylum simplex vel compositum*, or such as follow. The excellent External Treatment.

as had observed the last of this Distemper in *Austria, Bavaria, Silesia, Prussia, Poland, Holsatia, Denmark*, and *Marseilles*, endeavouring to reduce what they had observed with regard to the Symptoms, &c. to a Sort of *Compendium*, that my Reader might rely on them afterwards.

BARBET, in his Treatiſe *De Peſte*, particularly recommends the following Plaſter, which ſeems very efficacious:

℞. *Empl. Diachyl. c. Gummis, de Mucilaginibus ana* ℔ ß. *Semines Sinapi pulveriſati* ʒ iij. *Unguenti Baſilici* ʒ iv. *m. f. Empl.*

A Plaſter of this is to be applied to the tumified Part, after it has been firſt well rubbed, and to be renewed every or every other Day: the celebrated Dr. HODGES, in his Deſcription of the great Plague in *London*, *An.* 1665, greatly recommends the following:

℞. *Empl. Oxycroc.* ʒ iij. *Gum. Galban. colat. Carannæ ana* ʒ j. *Picis Naval.* ʒ ij. *cum Ol. Chamæmel. liquato f. Empl.*

This may be uſed like the former: nor is the Uſe of that Plaſter to be deſpiſed here, which is made of Honey, Meal, and the Yolks of Eggs: but the Bliſtering with *Cantharides* and dry Cupping, uſed by the Antients to forward Suppuration, are wholly rejected by the moſt expert of the modern Phyſicians in the Cure of the Plague [a].

A particular Obſervation of BEINTEM's.

VI. But what the celebrated *German* Phyſician, BEINTEM, obſerves, is not a little ſurpriſing, and worthy of our Conſideration: he aſſerts, in the laſt Book of his *latin* Treatiſe on the Plague, that peſtilential *Bubos* were frequently diſperſed and cured without any Danger, merely by the Application of warm Aſhes: though there is ſcarce any body beſides him, that adviſes to diſcuſs or cure peſtilential *Bubos*, without bringing them to Suppuration, or that ever found ſuch a Method ſafe and ſucceſsful: but, in the Judgement of BEINTEM, the peſtilential Venom was not drove into the Blood again in the Diſcuſſion, but was rather attracted and carried off by the Aſhes.

Internal Method of Cure.

VII. To theſe external Applications it will be proper to join internal Medicines; by the Help of which, the Venom lurking in the Body may be expelled by a gentle Sweat: but ſuch ſudorific Medicines, as are very ſtrong and heating, have been always found dangerous and pernicious by the modern Phyſicians. Warm and watery Drinks have generally been found more ſafe and uſeful in this Caſe, as being particularly adapted to temperate the Blood, and excite a gentle Sweat: among theſe Drinks, we may reckon common Tea, with the Addition of a little Saffron; or Infuſions of other alexipharmic Herbs, as *Salv. Scordium, Ruta, Millefol. Betonica, &c.* or elſe the plentiful drinking of ſome warm Ptiſan, made with or without *Rad. Scorzoner.* taken till it excite a conſtant but very gentle Sweat: and as the more vehement Sort of Sudorifics are improper, ſo the drinking of cold Liquors is generally found equally pernicious; for they not only wonderfully ſuppreſs the gentle Sweat, but alſo ſtrike in the *Bubos*, in whoſe Eruption a happy Cure chiefly conſiſts. The Air of the Patient's Chamber ſhould be temperate, neither too hot nor too cold; his Bed ſhould alſo be the ſame, and made as convenient as poſſible. If the Patient ſhould find himſelf very weak, but without any great Heat, it will not be improper to give a few Drops of *Elix. Proprietatis vel Mixtur. ſimplicis, Tinctur. Bezoart. Eſſent. Myrrhæ, Eſſ. Scordii, &c.* about thirty or forty Drops for a Doſe two or three Times a Day, in ſome warm Liquor: or, it may be

[a] Yet SCHREIBER, a very modern Writer on the Plague, declares, that in *Ruſſia* Bliſters were applied to peſtilential *Bubos* with great Succeſs, laying on afterwards the emollient Cataplaſms juſt now recommended. See his *Obſ. on the Plague*, *p.* 23.

requiſite

requiſite to give ſome proper bezoartic Powder[a]: on the other Hand, in warm Conſtitutions, where the Heat is too violent, it will be proper to give *Nitrum depuratum cum Lapidibus Cancrorum Conchiſque præparatis:* alſo temperate Acids, as *Succ. Malor. Citreor. Ribeſior. Granator. &c. vel Syr. ejuſd. cum Aqua Borag. Bugloſſ.* or any thing that is temperately cooling, to which the Patient has a Fancy; and if the Heat be ſtill more vehement, it may be neceſſary to drop in *Spiritus Vitrioli dulcis aliquot Guttulas.*

How the Abceſs is to be opened.

VIII. The Medicines hitherto propoſed are all of them allowed to be the moſt proper to be often taken, and ſufficiently powerful to drive out any peſtilential Venom that may lurk in the Blood, agreeable to the Writing and Practice of the moſt expert Phyſicians, who have lately wrote in *Poland, Pruſſia, Denmark, Auſtria, Hungary, Ratiſbon,* &c. the Uſe of theſe ſhould therefore be continued 'till the Tumors are either diſperſed (which they allow to ſometimes happen) or ſuppurated and brought to Maturation, which is the common and conſtant Practice. In ſome Caſes the Tumor turns ſuddenly to Suppuration: and in others it remains for ſome Weeks without being any thing ſofter: when this is the Caſe, it is neceſſary to continue the Uſe of the foremention'd Remedies, 'till the Tumor either breaks of itſelf, or is fit to be opened like other Abceſſes by Inciſion with the Scalpel, that the peſtilential Matter may be diſcharged, and prevented from returning into the Blood.

Treatment after Aperation.

IX. When the Abceſs is thus opened, we muſt proceed directly to the cleanſing of it; and after the cleanſing, the Wound is to be healed with ſome vulnerary Balſam, as we before propoſed: to deterge and cleanſe, the beſt that can be uſed here, is *Ung. digeſt. cum Theriac. Balſ. Sulph. Terebinth. Portiuncula permixtum.* At each Dreſſing the Matter is to be gently diſcharged from the Ulcer, and, when cleanſed, it is to be treated with the foremention'd Ointment; but without Tents, unleſs its Opening ſhould be very narrow; then applying ſome proper Plaſter, it may be bound up again as before: the beſt Plaſters for this Purpoſe are, the *Emp. Diachyl.* or that made *ex Melle & Farina*; the Uſe of which may be continued till it is perfectly healed up.

The Inciſion is not to be made too ſoon.

X. With regard to the Time of opening the Abceſs by Inciſion, Phyſicians are not agreed upon it; for there are many, eſpecially of the modern Authors, who have wrote on the Plague, that forbid the Opening of peſtilential *Bubos* till they are perfectly ripe and ſoft: beſides, theſe *Bubos*, agreeable to the Obſervation of many, do generally ſuppurate and break of themſelves; inſomuch, that, in the Opinion of theſe Gentlemen, an Opening made by Inciſion too ſoon, may greatly endanger the bringing on ill-conditioned *Fiſtulæ*, a Stiffneſs in the Limb, and even a Gangrene[b]: others, on the contrary, will have it, that an Opening made by Inciſion in the very Beginning of the *Bubo*, is not only without Danger, but even directly ſuited to preſerve the Patient, and recovering him the ſooner from his dreadful Diſeaſe: *vid. Ephem. Nat. Curioſ. Cent.* VII. *Obſ.* 69. *pag.* 170.

[a] Here SCHREIBER recommends the following Powder to be given every three Hours, having in the firſt Place vomited the Patient with *Ipecac.* R. *Antimon. Diaphor. Nitrat. Gr.* xv. *Merc. dulc. Gr.* i. *Camphoræ Gr.* ij. *M. F. P. in Sero Lactis vel Hordei decoct. tepid. ſumendus, ſuperbibendo ejuſdem, oz.* v.

[b] The Author of a *French* Treatiſe, entitled, *Obſervations ſur la Saignée de Pied*, obſerves, that in the Plague of *Marſeilles*, too early an Inciſion of *Bubos* was generally unſucceſsful.

XI. Notwithſtanding ſeveral of the antient Phyſicians have contended for a ſpeedy and entire Extirpation of peſtilential *Bubos* by the Knife, in order to diſcharge the contagious Venom; yet the Moderns do not without Reaſon diſſent from their Opinion: for ſuch a Method of Cure is not only found to be too harſh, but alſo of very dangerous Conſequence in many Parts of the Body: in like Manner, all Emetics, Cathartics, Bleeding, and hot cordial Medicines are, by the unanimous Conſent of the Moderns, condemned as Things very pernicious in the Peſtilence; notwithſtanding they were held in ſo great Eſteem by the Antients, ſuch were the bezoardic Tinctures, hot Eſſential Oils, and volatile antipeſtilential Spirits, together with the *Theriaca* and *Mithridate*.

CHAP. X.

Of the PRESERVATIVES *particularly neceſſary to defend and preſerve the Phyſician or Surgeon from* PESTILENTIAL CONTAGION.

Preſervatives hitherto uncertain.

I. HITHERTO we have been treating of *Peſtilential Bubos*: but, before we proceed to Carbuncles and *Anthraces*, it will be proper to ſay ſomething of the Means that may be uſed by the Surgeon to defend himſelf from the peſtilential Contagion, that he eſcape free in viſiting the infected: but before we take upon us this Taſk, it will be firſt proper to inform our Reader that we believe there has not ever been yet found a certain Preſervative for this Purpoſe; ſo far from it, that many of the Remedies purpoſely contrived and recommended, are wholly uſeleſs and improper; even ſome of them are very dangerous when lodged in imprudent Hands, and are therefore to be cautiouſly avoided.

Which of theſe Methods are to be avoided.

II. There are many, who aſſert frequent Purging to be wonderfully adapted to carry the peſtilential Contagion off the Body, and prevent it from getting into the Blood; there are others, who lay great Streſs upon ſudorific Medicines, Scarifications, and frequent Bleeding, as of great Service to defend the Body from the peſtilential *Virus*: whereas all of them, unleſs the Body is habituated to them, are great Deſtroyers of the Strength; and by that means, rather than defend, they make the Body more obnoxious to, and ſuſceptible of, the contagious Venom: others again believe nothing more effectual as a Preſervative againſt the Contagion, than the frequent and plentiful drinking of certain hot Spirits or Waters, dignified commonly with the Title of epidemic or antipeſtilential: but we ſhall be ready to judge the Uſe of theſe alſo to be equally foreign, and altogether improper, if we do but conſider what violent Heats the plentiful Uſe of ſuch ſpirituous Liquors will excite in the Blood, beyond what it ſhould naturally ſuffer, and by that means it may be rendered more liable to fall into a peſtilential Fever; unleſs the Perſon has been accuſtomed to the Uſe of ſuch Liquors before, or elſe uſed them with great Moderation: the ſame Judgment we muſt always paſs upon the common Spirit of Wine, *Aqua Vitæ*, and the alexipharmic Electuaries and Oils, with all other heating Medicines, ſince their Nature and Effects are directly the ſame: laſtly, there are ſtill others who confide in things hung about the Neck, as Arſenic, Mercury, Sand, Camphire, and

and *Rad. Colchici*; or elſe the keeping open large Iſſues, from all which they expect a ſecure Defence from the Plague: when at the ſame time there can be found little or no Virtue in either, or all of them, to reſiſt the peſtilential *Virus*. It has been obſerved by antient Writers, that Perſons afflicted with Ulcers of any ſtanding, eſcaped the Infection of the Plague; which Obſervation is confirmed by SCHREIBER amongſt the Moderns: in that Caſe thoſe Ulcers ſhould by no means be healed; on which Account ſome Phyſicians have adviſed Iſſues as Preſervatives; but to no Purpoſe.

III. The beſt and readieſt Defence againſt the Plague ſeems in general to conſiſt in this, that ſuch as are able ſhould remove out of the peſtilential or infected Air into ſome healthy Part of the Country; or, wherever they are, they ſhould keep from the Company of ſuch as are already infected, and not meddle with their Cloaths, Bedding, Meat, Drink, or Veſſels; and above all, if poſſible, not to make themſelves over afraid of the Diſeaſe; but let them always keep a chearful and confident Mind, with a proper Diet; but for the Phyſician and Surgeon, whoſe Buſineſs is to relieve the Sick, and for that Purpoſe muſt enter dangerous Places, it is beſt for them to keep up a couragious Mind, and not be anxiouſly afraid of Diſeaſes, nor even the Plague; for it is to be hoped that thoſe, who riſque themſelves with theſe Precautions to ſuccour peſtilential Patients, will be preſerved in Safety by a Divine Providence: SCHREIBER recommends, as a ſafe Preſervative, the taking every Evening, *Merc. dulcis & Camphor.* of each 1 *Gr.* The beſt Defence againſt the Plague:

IV. But beſides, there are ſeveral human Cautions and Obſervations neceſſary to be regarded by the Phyſician and Surgeon: the chief of theſe are, that *they ſhould never go faſting to viſit a Patient ſick of any contagious Diſeaſe, and much more of the Plague:* but they ſhould always eat ſomething and drink ſome ſtrong Liquor before-hand, in order to defend themſelves from the peſtilential Contagion and infected Air: ſome Phyſicians therefore always *eat Bread and Butter, and drink a Draught of* Spaniſh *or Wormwood Wine*, or ſome other ſtrong Wine, before they offer to ſet a Foot in the Patient's Houſe: by this Method, the celebrated Dr. HODGES writes, that he preſerved himſelf from Infection in the violent Plague at *London*, chiefly by drinking *Spaniſh* Wine: DIEMERBROER tells us, that by the Benefit of *Rheniſh* Wine he eſcaped the Infection in the Plague at *Neuberg:* others prepare themſelves in a Morning, *by eating a Slice of Bread ſoaked in good Vinegar*, either ſimple, or wherein Rue has been infuſed: SYLVIUS has contrived an acidulated Medicine purpoſely for this Uſe, which the Apothecaries call *Aqua Prophylactica Sylvii*: and is to be drunk to the Quantity of one or two Spoonfuls in a Morning, either alone or with a Slice of Bread, by ſuch Surgeons as are going to viſit peſtilential Patients: others again aſſert it to be confirmed by Experience, that ſome good Broth or Suppings, eſpecially of Chocolate, are of great Service in keeping off the peſtilential Venom: CARDILUC mentions his eating a Citron every Morning, *Lib. de Peſte, p. m.* 70.---Upon the whole it is my Opinion, that in hot Conſtitutions, Acids, and Coolers, are the beſt Preſervatives; in cold, the reverſe, at leaſt moderate Cordials. (1). before we viſit the Patient.

V. Being come to the Patient's Apartment, great Care muſt be always taken *that we neither eat nor drink there, nor even ſwallow our Spittle*; for there is no ſmall Danger in that Caſe, of ſwallowing the volatile peſtilential Exhalation (2). while we are with the infected Patient.

or

or Effluvia, by which means our internal *Viſcera* and Blood would be infected: for which Reaſon we cannot approve of the Cuſtom of ſome who are continually chewing and ſwallowing Myrrh, Cinnamon, Angelica, Zedoary, or the like, all the Time they are in an infected Place; for as ſuch Things excite a plentiful Diſcharge of *Saliva* into the Mouth, it is hardly poſſible but ſome of the infectious Effluvia will be intangled therein, and ſo go down into the Stomach, and get into the Blood: but the chewing of ſuch Aromatics may be very proper at home, as they are in their own Nature wholſome; the Uſe of them in the former Caſe being improper only as to Time and Place: we ought alſo to be particularly careful *not to ſtay longer in the infected Place or Apartment of the Patient than our Buſineſs really requires:* for there is great Danger that the Strength of our Conſtitution, however conſiderable, may be overcome by the too great Quantity and Force of the peſtilential *Virus*; whereas we might have eaſily reſiſted and ſuſtained a ſmall Quantity of the ſame infectious Effluvia.

(3.) when we are returned home from the Patient.

VI. After we are returned home from the Patient, it is much the ſafeſt Way to waſh our Hands and Mouth well with Vinegar mixed with Water: for if there be any thing prevailing againſt the peſtilential Venom, Vinegar ſeems to be the chief: the Cloaths are to be changed for others, and expoſed to the free Air, and to be afterwards perfumed: then Suppings of Coffee, or Tea of *Scordium*, Sage, and the other alexipharmic Herbs, ſhould be plentifully uſed; for theſe excite a gentle Sweat, and ſo drive out ſuch contagious Particles as might happen to be mixed with the Blood, keeping it free and temperate.

A proper Diet is to be ſtrictly obſerved.

VII. As an accurate Regimen of the Diet is always healthful in other Caſes, ſo alſo in Places where the Peſtilence rages it is found to be altogether neceſſary: therefore ſo much Aliment, ſolid and fluid, is always to be taken at one Time, as is requiſite to keep up the Strength of the Body, and may be conveniently and perfectly digeſted; but Care muſt be taken not to burthen Nature therewith: for it can ſcarce be ſaid how vaſtly Intemperance weakens the Stomach and Body, and renders it liable to contagious Diſtempers; from the Crudities and undigeſted or corrupt Matter, which is by that means lodged in the Blood. Modern Phyſicians obſerve, that there is no Occaſion for chooſing a particular *Diet:* ordinary or common Food may be taken as uſual, if it be not againſt Cuſtom and Temperance: in Broths and Suppings ſhould be always mixed, whenever it can be done conveniently, ſome Vinegar, or the expreſſed Juice of Lemons or Citrons, a few Capers, or ſome other ſubacid Thing of the like Kind: for the Uſe of every Thing gently acid is uſually very ſafe and beneficial in the Peſtilence; ſo that a moderate Plenty of all Sorts of Pickles are in this Caſe found very ſalutary[a]. There is no need of any great Change in the common and daily Drinks: but this I muſt obſerve, that thoſe conſult beſt for their Stomach and the Strength of their Conſtitution, who drink *Spaniſh*, *Rheniſh*, or any other good Wines at their Meals: if any one be accuſtomed to *Tobacco*, I would adviſe him to keep up the Habit: but I would not perſuade ſuch as diſlike it, or are of a hot Conſtitution, to take Tobacco againſt their natural

[a] The Phyſicians at *Marſeilles* were of Opinion, that an Acid was the Cauſe of peſtilential Diſorders; becauſe DEIDIER, in the Diſſection of Subjects who had died of the Plague, had found the Hearts exceeding large, and the Blood in them coagulated: but this to me is no Proof at all.

Appetite

Appetite as a Preſervative from the Peſtilence: for I think it has been this long Time obſerved that Lovers of Tobacco have been equally as often and eaſily ſeized by the Plague as others who do not uſe it: laſtly, where Perſons have been before accuſtomed to the Uſe of Stomachics, Sudorifics, Vomiting, Purging, Scarification, Bleeding, and the like, at certain Times or Seaſons, they muſt be cautious not to break off too ſuddenly from ſuch Habits, but rather to continue them at their ſtated Times[a]: but for Coition, as it greatly weakens and even ruins the Conſtitution at ſuch an unfavourable Time, eſpecially if the Habit of the Body be naturally infirm, that ſhould be equally avoided with the Peſtilence itſelf.

VIII. In the laſt Place, in order to keep off or correct the peſtilential Effluvia, it will not be improper frequently to hold a Sponge to the Noſe which has been firſt wetted with ſimple Vinegar, or that wherein Rue or Lavender has been infuſed: the Chamber ſhould alſo be fumigated with Juniper Chips, Gunpowder and Brimſtone, or with Vinegar, ſprinkled upon a red hot Tile or Iron, in order to expel and correct the peſtilential Air. External Helps againſt the Plague.

CHAP. XI.

Of CARBUNCLES, *or* ANTHRACES.

I. A CARBUNCLE is ſaid to be an Inflammation which ariſes in Time of the Plague with a Veſicle or Bliſter, almoſt like thoſe produced by burning, or the Application of Cantharides: but this Sort of Inflammation generally terminates in a *Sphacelus*, and putrifies the ſubjacent Parts down to the Bone, they becoming as black as a Coal, inſenſible and dead: and this ſeems to be the Reaſon why they are by the *Latins* termed *Carbunculi*, and by the *Greeks Anthraces*[b]. A Carbuncle what.

II. A Carbuncle always breaks out very ſpeedily, even in the Space of an Hour or two, attended with Heat and Pain: upon opening it, there is diſcharged a darkiſh and ſometimes limpid or watery *Sanies:* within, the Fleſh is of a black Colour, a *Sphacelus* having then ſeized the Parts, which ſpreads more and more by Degrees: but the putrid Fleſh in thoſe who recover, ſuppurates and parts from the ſound: the Size of theſe peſtilential Bliſters is various, more or leſs, as is alſo their Number in the ſame Patient; for there is no Part of the Body which they do not infeſt: and they generally appear in Company with *Bubos*; indeed they are ſeldom or never to be obſerved without *Bubos*. The Nature of Carbuncles.

III. The immediate and uſual Cauſe of Carbuncles is doubtleſs a violent Inflammation, excited in the Blood by the peſtilential Venom: the Inflammation is ſpeedily and ſuddenly followed by a Corruption and Sphacelation of the Parts; but the Parts and Juices do not ſuppurate into Matter, as is uſual in other Tumors; but whatever is internally corrupted ſeparates and intirely falls off: for the inflamed Parts ſuppurate at the Margin or Extremity of the Inflammation; ſo that if the Patient does not die ſuddenly, the ſphacelated Parts which have Cauſes.

a This CELSUS very judiciouſly recommends, *Lib* I. *Cap.* 10. *de Peſtilentia.*

b As to the *Term Carbuncle*, Vide CELSUS, *Lib.* V. *Cap.* 28. *de Carbunculo. Lib.* VI. *Cap.* 6. § 10. & *Lib.* VI, *Cap.* 18. § 5.

the

the Carbuncle are by that means ſeparated from the ſound and living Parts, and are by Degrees wholly caſt off.

Prognoſis. IV. Experience witneſſes that the Events of a Carbuncle are very doubtful, and much worſe than thoſe of *Bubos*; eſpecially if the Eruptions turn directly either livid or black: but when the Puſtules are red at firſt, and then gradually turn to a citron Colour, the Danger is much leſs: thoſe Carbuncles which ariſe in the Face, Neck, Breaſt, or in the Arm-pits, are obſerved to be of the worſt Kind; for they generally kill the Patient.

Internal Treatment. V. As for the internal Treatment of Carbuncles, whether by Diet or Medicines, the very ſame is to be obſerved in this Caſe, with what we recommended in *Chap.* IX. § 7. of peſtilential *Bubos*; for the chief of the Cure conſiſts in keeping the Patient in a gentle and conſtant breathing Sweat: but SCHREIBER, in his *Obſerv. on Peſtilential Carbuncles*, conſidering them as a Species of *Sphacelus*, ſtrongly recommends the *Peruvian* Bark, as given in Fevers.

External Treatment. VI. The chief Deſign of the external Treatment is, to quicken, as much as poſſible, the Separation of the Parts ſphacelated with the Carbuncle from the ſound: therefore ſome of the modern Phyſicians uſe only Scarification in this Caſe, with very good Succeſs: for by cutting away frequently the corrupted Parts quite to the ſound, they let out the acrimonious and peſtilential Matter with the corrupted Blood[a]: others only open the Eruptions with a Pair of Sciſſars, and having diſcharged the Matter, they often waſh the Carbuncle with *Sp. Vin. Camph.* or *Sp. Vin.* wherein has been digeſted a little *Theriaca*: they afterwards apply a maturating Cataplaſm, like the following.

℞. *Mellis cochlearia* iv. *Fermenti panis cochlearia* iij. *Vitell. Ovor. N°* ij *Sapon.* ʒ ß. *Quæ probe commiſceantur, calidaque ſuperimponantur: Vel,*

℞. *Farinæ Siligin. vel Tritic.* ʒ iij. *Aceti* ʒ ß. *quæ ex Aqua vel Lacte ebutyrato decocta atque in Cataplaſma converſa cum Mellis* ʒ j. *Crocique contriti* ʒ j. *miſceantur, calidaque ſæpiſſime ſupradentur.*

Whether the Carbuncle ſhould be cut out. VII. The Application of the forementioned Cataplaſms is to be continued till the Carbuncle ſeparates or caſts off from the ſound Parts: for it is better to diſſolve the Carbuncle gradually from the adjacent ſound Parts, than to cut it out all at once: nor are Inſtances wanting where the Patient has been killed by an unſeaſonable and entire Extirpation of the Fleſh and Carbuncle; for we learn by Obſervation, that moſt ſharp Pains and other dangerous Symptoms uſually follow ſuch an over-powerful Remedy; but where the greateſt Part of the Carbuncle is already ſeparated from the live Fleſh, the Remainder may be ſafely divided by the Scalpel.

VIII. But if an ill-conditioned luxuriant Fleſh grow internally either of itſelf, or from the Extirpation being made too ſoon, it is upon all Accounts neceſſary to entirely conſume it by the Application of *Ung. Ægyptiacum vel fuſcum Wurtzii*, or elſe by the Ointment following:

℞. *Mellis cochlearia* ij. *Vitel Ovor, N°.* ij. *Alum. uſti pulv. Gentianæ Ariſtolochiæ ana* ʒj. *m. f. Unguentum.*

[a] SCHREIBER adviſes ſcarifying, as very advantageous, *Lib.* IX. *p.* 23.

IX. If the Inflammation inclines the adjacent Parts to a Gangrene, which is not unuſual, it will be proper to uſe the following Ointment: How to remove a ſupervening Gangrene.

℞. ☉ *Abſinth.* ℥ ß. *Herb. Scord. Flor. Sambuc. Chamæmel. ana M* j. *Aq. ſimpl.* ℔ ij. ß.

When theſe have been well boiled and ſtrained, mix of the beſt *Sp. Vin. Camph.* ℥ vj. *Theriac.* ℥ ij. then let it be applied very often and hot to the Parts, by means of linen Rags folded together, or Compreſſes, till the Violence of the Inflammation abates.

X. But when theſe very bad Symptoms are abſent, after a Separation of the Carbuncle from the live Parts, it will be proper to cleanſe the Ulcer with *Ung. Fuſ.* WURTZII, or the digeſtive Ointment before deſcribed in *Chap.* IX. § V. *Of peſtilential Bubos.* And this ſhould be done perfectly, leaſt any of the peſtilential Venom ſhould remain behind, and excite the former Symptoms again. Therefore the Deterſion of the Ulcer ought to be continued till there remains nothing of theſe peſtilential Symptoms; and when that is effected, the Wound may be healed like other Abceſſes: more eſpecially, it ſhould be dreſſed with Lint dipped in *Eſſent. Myrrhæ & Aloes*, applying over an *Emplaſt. de Lithargyro*, or the like, till the Ulcer is perfectly cured. What is to be done after a Separation of the Carbuncle.

XI. There are many of the more celebrated Phyſicians, who, with CELSUS, allow nothing to be more effectual in extirpating and curing Carbuncles, than the actual Cautery, or a red-hot Iron: with this they order the dead Parts to be burnt till the Fleſh becomes in every Part ſenſible of the Pain; by which means there ſeems to be no Reliques left of the Carbuncle. This Method was obſerved by Dr. HODGES to be the readieſt Way of Cure for Carbuncles in the great Plague at *London*: but there are abundance of Circumſtances which prohibit the forementioned Method of Cure by the Cautery from being uſed in many Caſes; as the Dread of the Patient, the Tenderneſs and Conſequence of the Parts, *&c.* that rather perſuade ſuch Methods of Treatment as we have before propoſed; which are therefore to be made uſe of here. SCULTETUS ſays, that a dying Carbuncle will ſtart out afreſh (which is of dangerous Conſequence) on holding a red-hot Iron at a Diſtance. Whether Cauteries are to be applied.

XII. The celebrated SYLVIUS thought *Butyrum Antimonii* an efficacious Remedy to extirpate Carbuncles, if the circumjacent Parts were anointed with it. For, in the Opinion of SYLVIUS, it not only prevents the Diſorder from ſpreading, but it alſo readily makes an Eſchar that divides the ſound Parts from thoſe which are corrupted, and at length wholly ſeparates them. But ſuch of the modern Phyſicians as have wrote profeſſedly on the Plague at *Vienna* and *Ratiſbon*, do by no means agree with him: for if we may believe theſe, the *Butyrum Antimonii* is ſo far from being ſerviceable in Carbuncles, that it rather excites the worſt of Symptoms, and often brings ſudden Death. In the mean time, we find BOTTICHERUS aſſenting to the Opinion of SYLVIUS, in his *Loimographia Hafnienſis:* where he frequently praiſes and recommends the *Butyrum Antimonii* as an excellent Remedy for this Purpoſe: and SCHREIBER tells us, that many Surgeons applied the *Lapis Infernalis* to the Lips of the Carbuncle; and that, in Conſequence thereof, it ſeparated with great Eaſe, by means of a digeſtive Ointment, and an emollient Cataplaſm: but whichever be the Caſe, the Method by uſing *Butyrum Antimonii* is, in my Opinion, more ſafe, and preferable to the Way Whether it be proper to apply *Butyrum Antimonii.*

of Cure by the Cautery. Laſtly, whichever of theſe Methods of Cure is practiſed, the Buſineſs afterwards will be always firſt to perfectly cleanſe the Wound, and then to heal it up.

CHAP. XII.

Of VENEREAL BUBOS.

What a Venereal *Bubo* is.

I. A VENEREAL *Bubo* is a Tumor with Pain and Inflammation ariſing in the Groins or Arm-pits, after Contact with an impure Woman. *Bubos* of this Kind are diſtinguiſhed into two Sorts. (1.) Such as ariſe without any other Symptoms of the venereal Diſeaſe: or, (2.) thoſe which are accompanied with the other uſual Attendants of the Diſeaſe, as a *Gonorrhœa*, and venereal Ulcers, uſually termed Chancres.

Symptoms.

II. *Bubos* of this Kind uſually ariſe, as we before obſerved, after Contact with an impure Woman, who is afflicted with the venereal Diſeaſe: after which, they ariſe ſometimes ſooner, and ſometimes later: that is, within a few Days after Infection. The Tumor then ariſes in the Patient with Hardneſs, Redneſs, and Pain, either in one or both the Groins, and ſometimes in the Arm-pits: ſo that if we regard the Colour of venereal *Bubos* there is little or no Difference between them and the benign Sort. See *Chap.* VIII. foregoing. Care muſt therefore be always taken, that we do not miſtake one for the other: for ſuch as take benign *Bubos* for venereal ones, generally treat the Patient with an unjuſt Suſpicion, Contempt, and a harſh Method of Cure. On the other hand, when venereal *Bubos* are miſtaken for benign ones, there is Danger leſt the Patient, being treated in the mild Method ſuited to the benign *Bubos*, ſhould be unhappily brought into a confirmed *Lues*.

Diagnoſis.

III. The moſt certain Signs that theſe *Bubos* are venereal, are the Patient's having had to do with unclean Women, and from their being, or having been, accompanied with a *Gonorrhœa*, Chancres, or other Symptoms of the venereal Diſeaſe: when any of theſe are preſent, they give ſtrong Reaſon to ſuppoſe the *Bubo* to be venereal: but when they are abſent, they take off, or at leaſt greatly diminiſh, the Probability of the *Bubos* being virulent. As ſoon as it appears from the Patient's Confeſſion or other Circumſtances, that the *Bubos* are venereal, we muſt proceed accordingly with Expedition to a proper Method of Cure. Though this Diſorder generally admits a pretty eaſy Cure at the Beginning, yet, when it has gained Ground, either from Delay, improper Treatment, or an irregular Courſe of Life, a Cure becomes then extremely difficult, and it frequently turns into the *Lues* itſelf.

Whether a Diſperſion be ſafe.

IV. With regard to the Cure, there are many Phyſicians who hold a Diſperſion of venereal *Bubos* equally improper with the peſtilential; becauſe, by that Method, the venereal Venom returns, contrary to the Deſign of Nature, into the ſmall Veſſels, and, by infecting the Blood, brings on a Pox: they therefore judge it neceſſary to abſtain entirely from Bleeding and Purging, and to forward the Tumor to Suppuration as faſt as poſſible: but with Submiſſion to theſe Authors, I cannot be of their Opinion: for the Cure by Suppuration is not only

only ſlow and tedious, but alſo attended with many Inconveniencies; whereas I have frequently experienced, with the greateſt Safety, much better Effects from the taking of cathartic and mercurial Medicines, together with a Decoction of the Woods, and other ſuch Purifiers of the Blood: for by this means the Virulency may be diſcharged from the Body much ſooner than by Suppuration; and the Tumors may be ſafely diſperſed without Danger of a *Lues*, or other bad Symptoms.

How the Diſperſion is to be effected.

V. Whether the Patient have a *Gonorrhœa* or not, the beſt way is to purge him with frequent and large Doſes of *Merc. Dulc.* as is uſual in carrying off *Gonorrhœas:* for in curing a *Gonorrhœa*, you alſo cure *Bubos* generally at the ſame Time, and by the ſame Means: nor can *Bubos*, be happily cured till the Body is firſt quite freed from the venereal Venom. When there is a conſiderable Inflammation, eſpecially in young plethoric Habits of Body, it ſeems to be altogether neceſſary to bleed, and give mercurial Purges afterwards, with a Decoction of the Woods, and Eſſences which purify the Blood. Externally to the Tumor ſhould be applied ſome diſcutient Plaſter; as *Emp. de Meliloto, de Ranis cum Mercurio, Diachylum*, or the like: at the ſame Time the Patient ſhould keep ſtrictly to a regular Diet and Courſe of Life; taking ſcarce any thing but Ptiſans made with Barley, Oats, or the like: in the room of ordinary Drink may be taken a Ptiſan, made of Barley, Liquorice, and Aniſe or Fennel: for a Change may be drank a Decoction of the Woods; and for a greater Variety, a little clear and very ſmall Beer. Wine and all other ſtrong fermented Liquors ſhould be carefully avoided, as they generally increaſe the Inflammation: if the Patient be kept up carefully to theſe Reſtrictions, venereal *Bubos*, which are not yet inveterate, may be diſperſed very commodiouſly, and without Danger.

Suppuration how to be promoted.

VI. But if Advice ſhould be called in too late or the *Bubo* prove ſo obſtinate as not to give way to Diſperſion; or if upon any other Account the Surgeon is deſirous to effect a Cure in the Way of Suppuration, in order to diſcharge the *Virus* and prevent a *Lues*, he is to diligently promote and quicken the Maturation as faſt as poſſible. But the moſt powerful Medicines to promote Suppuration have been mentioned at *Chap.* III. § IV. and *Chap.* IV. § VIII. Though it is beſides not improper here to rub the *Bubo* with linen Rags, or the Fingers greaſed with Butter or Oil, till they grow red with Pain; adding afterwards a maturating Plaſter; for by this means a Suppuration is greatly promoted and accelerated: the Plaſter to be afterwards applied may be of *Diachylum cum Gummis, vel Emplaſtrum de Galbano*, particularly when the Patient can as yet walk pretty well: the Plaſter may be taken off, and the *Bubo* rubbed well, three or four times a Day, more or leſs, agreeable to the ſeveral Circumſtances. Violent Dancing, Boxing, Fencing, and other ſuch Exerciſes, are alſo here very ſerviceable for promoting the Suppuration: but if the Patient cannot walk any longer from his Pains, which is frequently the Caſe, it may be proper to apply a maturating Cataplaſm inſtead of a Plaſter, ſuch as we have deſcribed in the Chapters juſt now mentioned; which are uſually much more effectual than Plaſters. The beſt of theſe Cataplaſms for this Caſe, are thoſe *ex Cepis ſub Cinere toſtis, vel Farina et Melle, vel ex Fermento, vel denique ex Mica Panis Siliginei cum Lacte atque Croco decocta*; which are to be now and then applied warm to the Parts, after they have been firſt well rubbed.

Internal Treatment.

VII. While the former are carrying on, internal Medicines muſt be alſo called in to Aſſiſtance. The Patient ſhould take a warm Draught of a Decoction

of the Woods two or three Times in a Day, about eight, ten, or twelve Ounces at a Time, with thirty or forty Drops of *Eſſent. Lignor. Pimpinellæ albæ. Fumariæ, val Scordii, vel his ſimilium, & Mercurii dulcis aliquot Granis quotidie.* For as theſe greatly attenuate the Blood, drive it towards the Skin, and correct the venereal Venom, they alſo greatly promote either a Diſperſion or a Suppuration.

The Manner in which *Bubos* are to be opened.

VIII. Theſe Methods are to be followed till the *Bubo* comes either to a Diſperſion or Maturation. When the Tumor appears to be perfectly ſuppurated, the Scalpel is to be taken in hand, in order to make an Inciſion upon the *Bubo*; but then it muſt be done with Caution, to avoid hurting any of the large Blood-veſſels in either the *Inguen* or *Axilla*; from whence might enſue a very dangerous Hæmorrhage. The better to avoid injuring theſe Veſſels, the protuberant Part of the *Bubo* ſhould be preſſed outwards by the Fingers: but with regard to the Time in which it is proper to make the Inciſion, it muſt be always carefully obſerved not to let it be too ſoon nor too late; becauſe both are dangerous; for when they are opened too ſoon, it occaſions Pains, violent Inflammation, and other bad Symptoms; as when they are delayed too late, it generally occaſions (as HILDANUS witneſſes) the corrupt Matter to return into the Blood, and by infecting the whole Maſs, brings on a confirmed *Lues*: if the Patient dreads the Knife, the *Bubo* may then be opened by a Cauſtic. Here the Reader ſhould turn to what we have ſaid before on Abceſſes, *Chap.* III. § X, *ſeq.* When the Matter is once diſcharged, it will be proper to cleanſe the Ulcer with ſome digeſtive Ointment, mixed with ſome *Theriaca* and a little *Merc. præcip. rub.* Afterwards may be applied a Plaſter of *Diachylum cum Gummis*; by which means the Lips of the *Bubo* will be ſufficiently ſoftened and cleanſed; and then it may be healed with ſome vulnerary Balſam, applied on ſcraped Lint.

Whether and when the actual Cautery ſhould be uſed.

IX. Sometimes the ulcerated *Bubo* becomes ſo ſtubborn, that it will neither incarn nor cicatrize, by the Help of any Medicines; but always affords a copious Diſcharge of Matter. When this is the Caſe, and the forementioned Medicines have been uſed to no Purpoſe, *viz. Præcip. rub. & Alum. uſt.* prove alſo to be of no Service, there then remains no other probable Method, in my Opinion, than to cauterize the corrupted Parts to the quick by the actual Cautery: for by that means the Communication of the infected Lymphatics may be cut off. From what we have hitherto propoſed, it ſeems to be ſufficiently apparent, that it is always ſafer and more convenient to bring venereal *Bubos* to a ſpeedy Diſperſion or Reſolution, when a Cure may be that way effected, than to bring them ſlowly to a Suppuration: but when the Blood is found too much infected, and already corrupted by the venereal Venom, ſo that a confirmed *Lues* begins to ſhew itſelf, the Cure by Suppuration may be then both proper and requiſite.

CHAP. XIII.

Of CHILBLAINS.

Chilblains what they are.

I. WE generally give the Name of *Chilblains* to thoſe Tumors which happen in the Hands and Feet from violent Cold; they being at the ſame time accompanied with Inflammation, Heat, Redneſs, pricking Pain and Im-

Immobility in the Limb. Sometimes they are of a livid or leaden Colour, and ſometimes they break out with Scabs or elſe with Chaps or Slits, which afterwards penetrate deeper, and become ulcerous: the Humour which they diſcharge is ſometimes a little fœtid, and pretty much reſembles *Pus* or *Sanies*. The Inflammation alſo frequently turns to a *Spachelus:* ſo that I think we may readily conclude hence, that Chilblains wholly belong and ought to be referred to the Tribe of Inflammations; the more becauſe they excite the ſame Senſe of Heat or Burning with other Inflammations[a], and do, like them, terminate in either Diſperſion, Suppuration, Gangrene or *Sphacelus*.

Diagnoſis.

II. Chilblains may be known and diſcovered by ſeveral Means; for (1.) we may obſerve the common Signs of Inflammation which we have but juſt now mentioned: (2.) we muſt enquire whether the Patient afflicted with them has been ever previouſly affected in thoſe Limbs with vehement Cold or Froſts, to which Travellers and Soldiers, who are engaged in Winter Expeditions and Sieges, are often greatly expoſed. Laſtly, (3.) it is alſo a Sign that they are Chilblains, when the Patient feels Pricklings or Shooting in the Part, with Heat and violent Itching; and when the Part affected is found inflexible and almoſt inſenſible.

The Degrees of this Diſorder.

III. While the Chilblains are yet tumified and red, and the Part retains its Senſe and Motion without any great Heat and Pain remaining, the Diſorder is then of the mildeſt Kind: on the contrary, when they turn livid, occaſion the Limb to become ſtiff and inſenſible, or excite pricking Pains therein; there is then Danger of a worſe Conſequence, leſt it ſhould degenerate into a Gangrene, or, at leaſt, a deep Exulceration. When the Skin riſes into Puſtules or Bliſters, like what frequently happens in Burns and violent Scalds, it is a Sign that there is an incipient Gangrene upon the Part. Laſtly, when the Member loſes its Senſibility, turns livid, ſoft, and flaccid, there is great Reaſon to ſuſpect that it is then dead, and ſphacelated.

Cauſe.

IV. We have no Room to doubt but that the real Cauſe of Chilblains is the Cold: for by violent Cold, the Mouths of the ſmall Blood Veſſels are not only greatly contracted, but the Blood is alſo by the ſame Means rendered too thick; which are the two great Cauſes of all Inflammation: nor is there any Symptom that attends this Diſorder, but what may be readily explained as a Conſequence of theſe Cauſes.

The Nature of Cold.

V. Though Naturaliſts are not yet well agreed among themſelves concerning the true Nature of Cold, yet I cannot conſent to the Opinion of thoſe who look upon Cold to be only the Effect of a Privation or Abſence of Heat: but I rather judge it to conſiſt in[b] certain hard, ſharp, rigid, and ſaline Particles, which float in the Air; which are by the Preſence of Heat, rendered very minute, ſoft, flexible, and volatile; but upon the Approach of Cold, they coaleſce and become rigid. Now when theſe Particles inſinuate themſelves into the ſmall Pores of the Body, they conſtringe the ſmall Veſſels, and by wounding them, either inſpiſſate or ſtop the Blood. Hence (in my Judgment) we may perceive the Reaſon why the Cold ſlits or cleaves the Skin of the Face, Lips, and

a So TACITUS very elegantly, *Annal.* xiii. *cap.* 15. Many of the Men, upon the Watch, had their Limbs quite *burnt* with the Extremity of the Cold.

b *V.* HAMBERGERI *Diſſ. de Frigore.*

other

other external Parts, and afterwards afflicts them with continual Prickings and Shootings: for the leſs Motion and Heat the Blood has in any Part, it is the more apt to be inſpiſſated generally: ſo that it is no Wonder if the Hands, Feet, Heels, Fingers, Toes, Noſe, Ears, *&c.* are more frequently afflicted with Chilblains than any other Parts of the Body; being ſometimes ſlight, but often very violent. Sometimes the Cold is ſo great as to quite ſtop the Courſe of the Blood throughout the whole Body; which then quickly kills the Patient; and we ſay commonly, that he was frozen to Death, or periſhed with Cold.

Prognoſis. VI. Though all Chilblains are in the general ſomewhat dangerous, yet they are more or leſs ſo in Proportion to the Extremity and Violence of the Cold which occaſions them: in Conſequence whereof, more or leſs grievous Symptoms ariſe. When the whole Hand or Foot is ſeized by the Cold, the Danger is generally greater than when it affects only a Finger or Toe: but nothing can be more fatiguing, than that thoſe who have once been afflicted with Chilblains ſhould afterwards become liable, almoſt every Year, to Inflammations, Pains, Ulceration, and even Gangrene, upon the Approach of any great Froſt. Laſtly, when Chilblains are ill treated, by ſuddenly expoſing the Part from the Cold to a Fire, or any thing hot, or by wrapping it up in hot Things, there is great Danger of the Part's becoming black, ſoft, and putrid; and at length, loſing all its Senſation, it may contract a *Sphacelus*.

External Treatment. VII. Having found this to be the State of the Caſe, it readily follows, that the Cure of all Chilblains muſt conſiſt chiefly in reſtoring the Blood to its former Fluidity and free Circulation as ſoon as poſſible: but the inſpiſſated Blood requires to be reſolved in this Caſe by Methods very different from thoſe generally uſed in other Inflammations: for the warm Medicines, which are very beneficial and even abſolutely neceſſary in other Inflammations, are found to be extremely pernicious for Chilblains: nor can it ever be ſafe for thoſe who have ſuffered extreme Cold to expoſe themſelves preſently to Heat or a Fire; for Death has been often the Conſequence of ſuddenly expoſing the Body to the Viciſſitudes of Heat and Cold: it is therefore much more ſafe and convenient to expoſe the Patient firſt to an Air that is either cool or temperate, and to order him to continually exerciſe his Limbs as much as he poſſibly can; and laſtly, to advance him gradually to a ſtill greater Warmth or Heat. When the Patient is too weak to exerciſe himſelf, it will firſt be proper to bathe the Parts affected with Snow, or cold Water, which will ſeem to be hot to the Patient; by which means the ſharp ſaline *Spicula*, which ſtick in the Pores of the Skin, will be drawn out, and the Blood reſtored to its natural Circulation. Afterwards, when the Limb is become ſenſible, we may, by Degrees, apply comforting Medicines; ſuch as *Sp. Vini, meri vel cum Theriaca, Oleum item Petræ, Balſ. Sulph. &c.* When the Parts affected have been well rubbed and bathed with theſe, the Patient may then be advanced towards the Fire, or put to bed; endeavouring afterwards to excite a gentle Sweat.

Internal Treatment. VIII. To anſwer this Intention, great Service will be had from a few Glaſſes of hot Wine, wherein has been boiled ſome Cinnamon and Sugar: for by drinking, or rather gradually ſupping of this, the Patient generally revives and grows warm, and the Blood recovers its Circulation: though it may not be improper to give alternately with this, a ſmall Quantity of a ſudorific Mixture: as,

℞. *Aq.*

℞. *Aq. Galeg. Rutæ, Scord. ana* ℥ij. *Theriacal. Vit. Matthiol. ana* ℥ vj. *Prophylact. Sylv.* ℥ ß. *Mixtur. Simpl. vel Tinct. Bezoard.* ℈ ij. *Syrup. Cinamon. Caryophillor. ana* ℥ ß. *Misc.*

A little Draught of this, about three Spoons full, ſhould be given to the Patient every Quarter of an Hour, and the hot Wine as often, till we find the Appearance of a Sweat: if Wine be not at hand, good Ale boiled with Cinnamon, Cloves, and Sugar, may well enough ſupply its Place. Such Suppings as theſe ſhould be continued ſo as to keep up a Sweat for an Hour, or leſs, according to the ſeveral Circumſtances: for it can ſcarce be imagined how certain and expeditious this Method of Cure is for the moſt grievous Chilblains, which even threaten a Gangrene: but if the Diſorders which proceed from Cold are much ſlighter, this Method is then not ſo directly neceſſary, but may be laid aſide, though it is much preferable to any other.

How a Suppuration or Gangrene is to be treated.

IX. When Chilblains tend to Suppuration, it is proper to treat them like other recent Abceſſes. Firſt, to cleanſe the Wound with ſome digeſtive Ointment, as *Ægyptiacum*, *&c.* then to dreſs it with *Ol. Ovor. Ceræ*, *&c. vel Balſ. Peruvian. Eſſent. Aliès*, *Myrrhæ*, *&c.* and laſtly, to apply *Emplaſt. Saturnin. vel de Lithargyro.* Sometimes we ſhall find Benefit from *Oleum Myrrhæ per Deliquium*; as alſo from *Mures aduſti*, if we may believe the *Ephemerides Naturæ curioſorum.* Laſtly a Mixture of *Aq. Calcis cum Sp. Vin. Camph.* will be frequently found of great Service here; or *Rapæ coct.* which many Writers recommend; particularly CELSUS, *L.* V. *C.* 38. And PLINY, *L.* XX. *C.* 3. if a Compreſs dipped therein be bound upon the Part, either alone, or after the Application of the forementioned Medicines: but if a Gangrene or *Sphacelus* appear, the Parts affected are then to be treated in the Method we ſhall propoſe in the following Chapter.

To prevent Chilblains.

X. If a Patient has before been troubled with Chilblains, which are uſed to return every Year, in the Winter; to prevent the Diſorder from returning again, he may arm himſelf by proper Medicines. The beſt Preſervation for this Purpoſe, is, to anoint the Parts affected with *Petroleum* or Oil of Turpentine, before and while the Severity of the Winter comes on; but when the Diſorder has begun to ſhew itſelf again by Tumor, Inflammation, and Pain, the diſordered Heel or Finger may be wrapped up in a Swine's Bladder, dipped in the forementioned Oils. But the Cold itſelf ſhould be always carefully avoided, by defending himſelf well with proper Cloths or Coverings. The Reader may conſult at his Pleaſure M. A. SEVERINI *Diſſert. de Pernionibus in Lib. de Abceſſibus.*

CHAP. XIV.

Of a GANGRENE *and* SPHACELUS.

What a Gangrene and *Sphacelus* are.

I. HITHERTO I think we have ſufficiently conſidered the *Exit* of an Inflammation by the Way of Diſperſion or Suppuration. It follows, that we now examine the third and laſt Method whereby an Inflammation terminates, *viz.* a Gangrene and *Sphacelus*, to which Diſorders the antient Phyſicians

Phyſicians gave the Name of *Cancrum* a. By a *Gangrene* we underſtand that moſt great and dangerous Degree of Inflammation wherein the Parts affected begin to corrupt and put on a State of Putrefaction. But by a *Sphacelus* we underſtand not an incipient, but an abſolute and perfect Corruption, or Death of the Parts.

The Signs. II. A Gangrene may be diſcovered generally from the following Signs: namely, the Inflammation, with its Symptoms, which have all along been very violent, do generally undergo a ſudden Change, as if they were going off. The Parts which were before ſwelled and tenſe, do now become ſoft and flaccid; and upon preſſing with the Finger upon the Skin and Fat, its Impreſſion remains behind, as in an *Œdema*; at length the *Cuticula* ſeparates from the *Cutis*, often riſing up in Bliſters like thoſe in Burns, filled with a reddiſh, yellowiſh, and ſometimes black Humour; and the Senſe of the Limb is, in ſome Degree, diminiſhed. The chief Mark whereby we diſcover a *Sphacelus* is, when after a previous Gangrene the Parts turn livid, and entirely loſe their Senſation, in ſuch a Manner, that the Fleſh may be pricked and cut without giving any Pain; and if the Gangrene penetrates deep, ſo as to affect the Nerves and Muſcles, the Limb alſo loſes its Power of Motion: afterwards the Colour of the Part turns black by Degrees, and the Skin feels cold and flaccid; and at length it adheres ſo looſely to the Fleſh, that it may be eaſily pulled up and off from it. Sometimes the Skin becomes hard and dry, like the Rind of Bacon. Laſtly, it yields a moſt intolerable cadaverous Stench, and the *Sphacelus* ſpreads by degrees through the adjacent ſound Parts: unleſs there ſhould happen to be a Separation of the dead Parts from the ſound; though it frequently ſtops of itſelf, and by forming a circular Suppuration, the mortified Parts are caſt off from the ſound. I have ſeen ſome Caſes, where from vitiated Humours a Bladder has ariſen, without any previous Symptoms, ſometimes of a larger, ſometimes a leſs ſize, chiefly in the Feet, more generally in the Toes; which Bladder was full of Water or *Serum*, and ſpread itſelf gradually, blackening and mortifying the Fleſh underneath, as in peſtilential Carbuncles. There have been other Inſtances, where the Toes of a ſudden have turned firſt livid, then black, with an utter Privation of Senſe and Motion; nor could any Inciſion draw Blood from the Part.

Cauſes. III. The *Cauſes* of a *Gangrene* and *Sphacelus* are either external or internal. Among the *internal Cauſes* we reckon an *Eryſipelas*, and all other Inflammations which ariſe ſpontaneouſly, and can by no means be diſperſed nor brought to Suppuration. Inflammations of this Kind uſually proceed from the Blood's being too acrimonious or corrupted by the Bile, or in a *Scorbutus*; or when the Circulation of the Blood is too quick or too ſlow, by reaſon of old Age or any other Weakneſs; or laſtly, when the Patient indulges himſelf in a bad Courſe of Life, with reſpect to his Diet, or is ſubject to violent Paſſions, (eſpecially Anger, Grief, and Fear,) during the Time of the Inflammation. By *external Cauſes* we mean Injuries from the Air, cold Water, and the Application of topical Remedies externally to the inflamed Parts, which are either cooling, aſtringent, fat, oily, or the like; together with all great external Hurts or Accidents

a Vid. CELSUS, *Lib.* V. *Cap.* XXVI. *N.* 31. 34.

which

which frequently happen to the Body through Falls, Blows, *&c.* as in Wounds, Fractures, Luxations, *&c.*

IV. A Gangrene is, for the generality, never without Danger; becauſe it eaſily changes into a *Sphacelus* or entire Mortification, which never admits of a Cure but by taking off the dead Parts. But a Gangrene which is ſlight, incipient, and not ſpread far, but only affects the Skin and Fat, is not very difficult to cure; eſpecially when it happens in a young and ſtout Patient, in a mild and temperate Seaſon, and does little or no Injury to the Muſcles and Nerves[a]. But the larger, more violent and confirmed is the Gangrene, and the faſter it ſpreads, the more difficult is it generally to effect a Cure; eſpecially in an old or weak Patient[b], or in an ill Habit of Body from a Dropſy, *Phthiſis*, or *Scorbutus*: the Weather alſo being too hot or very cold, or the Parts affected being near the *Thorax* or *Abdomen*, may make the Caſe more dangerous: nor can this Caſe be neglected without the utmoſt Danger of Life; for the putrid Matter being abſorbed by the ſmall Veins, and mixed with their Blood, is conveyed to the Heart and Brain, and corrupts the whole Maſs; from whence all the vital Actions are diſturbed, the Appetite goes off, and Phrenzy with Death follow: ſo alſo in large inveterate Ulcers, in the Extremities and Feet of old People, when they become dry and livid, it is almoſt a conſtant Sign that a *Sphacelus* and Death are at hand. Death is alſo preſaged in great Inflammations attended with Spaſms, continual Hiccoughs and Belchings, cold Sweats, Faintings, a *Delirium*, and continual Reſtleſſneſs or Drowſineſs, eſpecially if they happen in a Patient who is then afflicted with a Gangrene or *Sphacelus*: and laſtly, if the Gangrene be not directly treated with proper Medicines, it commonly turns ſuddenly into a *Sphacelus*; and if the ſphacelated Parts are not timely removed or amputated, the Diſorder ſpreads through the adjacent Parts, and brings on a ſpeedy Death: but in a *Sphacelus* from an internal Cauſe, and eſpecially at the Verge of Life, the Amputation of a Limb is generally ineffectual; for either the Patient thro' extreme Weakneſs dies ſhortly after; or from a bad Habit of Body the Mortification ſeizes ſome other Part, and ſoon terminates in Death. *Prognoſis.*

V. We muſt therefore always endeavour to treat the Gangrene ſo as that it may not determine in a *Sphacelus*. Firſt of all therefore, in plethoric and ſtrong Habits, we are to bleed largely, and to repeat the Operation at Diſcretion; but in weak Habits, it ſhould be omitted: the Remainder of the Treatment will conſiſt chiefly in obſerving the three following Directions: The Cure conſiſts in three Things.

(1.) *To be careful in the Beginning to remove all violent external Cauſes of the Inflammation*; as too ſtrict a Bandage in Wounds and Fractures, all foreign Bodies which are ſtuck in the Parts, as Thorns, Splinters, Needles, *&c.* improper Medicines externally applied, as Ointments, Oils, and Plaſters, with cooling and aſtringing Things, as we before obſerved; all which ſhould be removed as ſoon as poſſible. (1.) A Removal of the external Cauſes.

VI. (2.) The other Obſervation reſpects chiefly the *keeping up of the Patient's Strength*, eſpecially in weak and old People: this may be beſt effected by ordering a Diet which not only affords good Juices, but is alſo well accommo- (2.) A proper Diet and internal Medicines.

a See CELSUS, *De Medic. Lib.* VI. *Cap.* 26. § 34.

b New Inſtances may be ſeen of Death from a Gangrene in old People in LE DRAN's *Obſ.* 100 & 101. I have alſo been Eye-witneſs to many of the like Caſes.

dated to the Age, Conſtitution, and other Circumſtances of the Patient: if the Patient is weak and in Years, is naturally of a cold Habit, has loſt much Blood, or abounds with Acidities, the moſt ſuitable Diet will be Soops, and ſtrengthening Broths, ſuch as are made of Chicken or Capons, Beef, or ſome other good Fleſh, boiled with Mace, Ginger, or other Spices; as alſo Suppings of Ale boiled with the Yolks of Eggs, Cinnamon, and Sugar; Eggs themſelves poached ſoft, ſo as to be potable; ſtrong Gellies of Calves Feet, Hartshorn, and Ivory Shavings; old and rich Wines, as *Rheniſh*, *Hungarian*, *Spaniſh*, *Canaries*, &c. and laſtly, fine Ale may do very well, eſpecially for the Poor. With reſpect to Medicines, the moſt proper are the *Corroborantia*, uſually termed *Cordial*, as the Spirits, Eſſences, Powders, and Electaries of that Tribe, eſpecially made up or mixed with *Confect. Alkermes*: at Intervals may be drank hot, Tea of Sage, *Scordium Veronica*, and Herbs of the like Nature, with the Addition of a little Cinnamon, or a few Shavings of *Lign. Saſſafræ Santal. Citrin.* &c. for by theſe Means the ſtagnating Blood will be wonderfully reſolved and attenuated, its ſound and healthy Parts will be retained in a due Circulation, and its noxious Parts will be diſcharged and diſſipated: it is alſo not improper, in this Caſe, frequently to apply a Sponge to the Noſe or Carpal Arteries, which has been dipped in *Aq. Reginæ Hungar.* alſo to bind it upon the Temples. In like Manner we ſhall find almoſt equal Benefit from the Crumb of Rye Bread mixed up with powdered Cloves; if it be firſt macerated in very ſtrong Vinegar, or *Hungary* Water, then made into a globular Form, wrapped up in a Piece of linen Cloth, and frequently applied to the Noſe. For Patients who are of a more warm, ſanguine, or bilious Habit, Soops and Ptiſans mixed with the acid Juice of Citrons or Lemons will be very proper Strengtheners; alſo Barley Gruel mixed with *Syr. Mali Citrei vel Mori, vel Rubi Idæi, vel Ribeſiorum aut Ceraſorum acidor.* to be taken daily as a common Drink. When the Heat is ſmall, the Patient weak, or before accuſtomed to Wine, it may be allowed to mix a little Wine with the Gruel, eſpecially *Rheniſh*, and ſometimes a Glaſs of rich Wine may be taken unmixed at proper Intervals; at the ſame time not neglecting the other Medicines which are proper to be uſed in Fevers, ſuch as are mild, temperating, cooling, and cordial: but the *Cortex Peruvianus* [a] is by many celebrated in this Diſorder beyond any other internal Medicine; they look upon it as the only Medicine in this Caſe, and adminiſter it in the ſame Manner as in intermitting Fevers. I have myſelf ſeen the good Effects of this Medicine; though indeed it has failed in ſome Caſes, which proceeded from an internal Cauſe, and where the Patient was advanced in Years. See *Act. Acad. Nat. Curioſ.* Vol. II.

(3.) External Remedies.

VII. 3. The third and laſt Obſervation concerning the Treatment of a Gangrene is chiefly *to diſcharge the ſtagnating and corrupted Blood from the Parts affected, as ſoon as poſſible*, and to prevent the neighbouring Parts from being affected thereby. The principal Means to effect this are (1.) to make uſe of proper internal *Corroborantia*, or ſtrengthening Medicines; (2.) to make *Scarifications* (*pro re nata*) by the Scalpel upon the Parts affected, making the Inci-

[a] Conſult WERLHEFFII *Obſ. de Febrib. p.* 332. taken from the Obſervations of RUSHWORTH, AMIAND and DOUGLAS. See alſo a particular Treatiſe publiſhed by DOUGLAS on Mortifications,

ſions

ſions very numerous lengthways upon the Parts, and of a ſufficient Depth, in order to diſcharge the ſtagnating and corrupted Blood, and to make way for the Ingreſs of the Virtues of the diſcutient Medicines which are applied externally; by which means they can the better penetrate through the ſmall Wounds to the internal Parts. Laſtly, (3.) diſcutient, ſtimulating, and balſamic *Fomentations* and *Cataplaſms* which reſiſt Putrefaction, are to be carefully applied to the diſordered Parts; of which Kind is the following Fomentation:

℞ *Aquæ Calc. viv.* ℔ j. *Sp. Vin. Camph.* ʒiij. *Sal. Ammoniac.* ʒß. *M.*

This may be applied hot with Compreſſes, it being what I have very frequently experienced and ſtill continue to uſe with very good Succeſs in theſe Caſes, and in other Inflammations. A very extraordinary and uſeful Mixture is alſo made *ex Aq. Calc.* ℔ j. *cum Mercur. dulc.* ʒ j. to be applied like the other: in the Hoſpital at *Amſterdam* the following excellent Fomentation was uſed with Succeſs in Gangrenes, within my Remembrance [a]:

℞. *Spirit. Vini* ʒiij. *Pulv. Aloës, Myrrh. ana* ʒ ß. *Ung. Ægyptiac.* ʒij. *M.*

Or, *Sp. Vin. cum Aloï, Myrrha, & Croco leniter coctus; vel Sp. Vin. Camph. cum Theriaca mixtus; vel Sp. Theriacalis aut Matricalis cum ſexta quaſi parte Elix. Proprietat. roboratus*; or, what GARENGEOT greatly extols, *Vinum calidum, Sp. Vin. ſimplic. vel Camphora roboratum, vel Sp. Vin. Camph. Sale Ammoniaco acuat.* which he extols as an excellent Remedy to revivify Parts which ſeem to be dying [b]. Or,

℞. *Fol. Scord. Abrotan. Abſinth. Rut. recent. ana M.* ij. *Flor. Chamæmel. M* j. *coq. in ſ. q. Aq. ſimpl. colatur.*

℞. *Hujus* ℔ ij. *adde Spirit. Vin. Theriacal.* ʒiv. *Sapon. Venet.* ʒij. *Salis Gemmæ* ʒ ſs. *M. f. Fomentum.*

This Fomentation is to be applied hot ſeveral Times in a Day to the Parts affected, by means of linen or woollen Cloths; and, to give a laſting Warmth, we may apply a hot Tile wrapped up in a thick Cloth, or a Bag of hot Sand.

VIII. For the Poor in this Caſe, there is a cheap and domeſtic Remedy, but of great Efficacy, recommended by SIMON PAULUS and others, *viz.* the *Pickle of Cabbages*. VALESIUS DE TARANTA has long before taught us, that Horſe or Cow-dung, boiled in Vinegar or Wine, makes an excellent Fomentation for this Purpoſe: but a long Time after him, we are told that SYLVIUS and BARBET held the ſame Remedy as a Secret in this Diſeaſe; but the Filthineſs of the Medicine makes it unworthy a Phyſician, it being fitter for the Poor and Vulgar than People of Faſhion: but there is a neat, as well as a very efficacious Fomentation for a Gangrene, to be made of *Scordium*, Wormwood, and Southernwood, either ſeparated or mixed, to be boiled in Sea-water, or, where that is not to be had, Salt-water or Vinegar, to be applied hot like other Fomentations ſeveral Times in a Day, giving a laſting Warmth by hot Bricks or Tiles, till the Diſorder diſperſes or diminiſhes: thus there will be no Occaſion ſo frequently to unbind the Part, and expoſe it to the Air, to apply more of the warm

A domeſtic Remedy for the Poor.

[a] Vid. KOENERDING *in Libello de Gangræna & Sphacelo, Belgico Sermone edito, Amſt.* 1698, 8vo. *Chirurgical Operations*, in the Chapter of a *Gangrene*.

Fomentation; but it is ſufficient, nay even preferable, to ſoak the Compreſſes well in the Fomentation, and to keep them hot upon the Parts by the forementioned Contrivance [a].

An obſtinate Gangrene, how to be treated.

IX. But the more obſtinate and nearer we find a Gangrene is to a *Sphacelus*, the more potent Remedies are we obliged to make uſe of. Such principally are the very numerous, long, and deep Inciſions and Scarifications of the Parts affected down to thoſe which are ſound. The Inciſions are alſo made not only longitudinally, but alſo tranſverſly, where they may be ſo with Safety, as in the Arm, Leg, and Thigh; by which means the Humours which lodge in the membranous Coverings of the Muſcles may be the better diſcharged, and the Tenſion of the Membranes taken off, and ſuch as ſtop the Motion of the Fluids by their Stricture be relaxed: afterwards the injured Parts are to be well rubbed and ſoaked with the ſtimulating, diſcutient, and balſamic Medicines, at § VII; to which may be added the Oil of Cloves, or Spirit of Turpentine, both powerful Remedies in this Caſe: then is to be applied a penetrating and diſcutient Cataplaſm, that the Blood in the vitiated Parts may be reſtored as much as poſſible to its free Motion. The following may ſerve for a Cataplaſm of this Kind:

℞. *Herb. Scord. Malv. Abſinth. Matricar. ana M* ij. *Menth. Abrotan. ana M* j. *Coquantur in ſ. q. Oxycrati, Vaſe clauſo, ad conſiſtentiam Cataplaſmatis ſive Pultis, eique poſtea admiſce Salis Ammoniaci* ʒ ß. *Farin. Lin.* ʒij. *Ol. infuſ. Rut. vel Chamæmel.* ʒ ß. *M. f. Cataplaſma.*

Always before the Cataplaſm is applied to the Part, it ſhould be mixed with ſome *Sp. Vin. Camph. aut Theriac.* to increaſe its Virtue: or, inſtead of this Cataplaſm, we may uſe the following, recommended by the forecited KOENERDINGIUS [b].

℞. *Mic. Pan. alb.* ℔j. *pulv. Abſinth. Scord. Rutæ ana M* j. *Vini q. ſ. ad Conſiſt. Cataplaſmatis, poſt levem Ebullitionem adde Sp. Vini* ʒ iv.

This is to be applied warm. In the mean Time, it is a neceſſary Caution to be obſerved in the Application of Fomentations and Cataplaſms, *viz.* that they ſhould not be renewed too often, but only two or three Times in a Day; for Experience has taught us, that the Humours may by that means be diſperſed and attenuated ſooner and with more Eaſe than by uncovering the affected Parts every Hour, as is cuſtomary [c]: but we muſt alſo carefully obſerve, that Cataplaſms and Fomentations ſhould not only be as warm as poſſible when they are firſt applied, but are alſo to be kept warm all the while upon the Parts, by covering them with hot Cloths, Tiles, or a Bag of Sand; by which means they

[a] HARRIS *(in Diſſ. Chirurg.* II.) adviſes, that, where an Inflammation is juſt turning to a Gangrene, the Part affected ſhould be dipped, if poſſible, in red Wine made hot, and fomented with the ſame.

[b] *In Libello de Gangræna, ſupra citato.*

[c] GARENGEOT will have the Dreſſing not to be opened above once in the Space of four and twenty Hours, in this Diſeaſe, (in his *Operations*, Chapter of a *Gangrene.*) But becauſe the Parts affected may ſuffer great Alterations in that Time, and as the Virtue of Medicines will ſcarce laſt ſo long, I think it more adviſeable for the Surgeon to inſpect the Parts two or three Times in a Day, that he may renew the Medicines, know how it goes forward, and what is to be further done, and that he may prevent any bad Accident.

will

will penetrate, ſtimulate, move, and attenuate much better; for if they become cold, they prove not only uſeleſs, but very pernicious: all things well conſidered, we can hardly affirm, that we have any thing that will cure a Gangrene, or prevent a *Sphacelus*; but if the *Cortex Peruvianus* has the Effects attributed to it in this Diſeaſe, we need not be troubled with ſuch a Train of ineffectual Remedies, nor charge our Heads with ſo many irkſome Cautions and Obſervations thereon [a].

How a Sphacelus is to be treated.

X. But if the Parts are already become quite dead, ſo as to be entirely without Senſe, and ſoft, ſo as to retain the Impreſſions of one's Fingers Ends, and appear to be fœtid and corrupted; in that Caſe, all the Medicines in the World will be inſufficient to reſtore the Parts to Life again: but there remains one, though a miſerable Remedy, to preſerve the reſt of the Body, by amputating the dead Parts, that the Diſeaſe may not ſpread through the reſt which are ſound: but a different Courſe muſt be taken in this Amputation, according to the Degree of Corruption, and the particular Nature of the Parts ſo affected: for if only ſome Extremity of the Foot, *Tarſus*, *Metatarſus*, Ancle, or Inſtep, or only the bare Skin and Fat, are ſphacelated, the whole Member or Foot ought not in that Caſe to be amputated; but, preſerving the Limb entire, we are to remove only that Part which we find vitiated, and that, in my Opinion, by means of Suppuration, as we taught in *Chap.* XI. § VI. *ſeq. Of Peſtilential Carbuncles*, or elſe to be taken off by cauſtic Medicines. Thoſe who undertake the Cure of a *Sphacelus* by *Suppuration*, are to take three Things chiefly into Conſideration; (1.) to effect the Suppuration as ſoon as poſſible; then (2.) to remove the dead Cruſt or Eſchar of the Ulcer, and ſeparate it from what is ſound; and (3.) to cleanſe the Ulcer, to preſerve the ſound Part, *&c.* and heal the Wound.

A Suppuration, how to be promoted.

XI. To expedite and quicken the Suppuration, nothing equals the making long and deep Scarifications or Inciſions, eſpecially near the ſound Parts: for by making innumerable Inciſions ſo deep, till we find that we every way touch the ſenſible Parts, ſo as to excite Pain, the noxious Matter lodged under the Eſchars may thereby be more eaſily diſcharged, proper Medicines will more readily penetrate the Parts, and the dead Parts will, by that means, be more ſpeedily ſuppurated, and the ſooner ſeparate from the ſound: but the moſt efficacious Medicines to promote this Separation of the vitiated Parts from the ſound, are Emollients and Balſamics which reſiſt Putrefaction, uſed in the following Method; *viz.* the inciſed Parts are to be firſt well anointed with *Unguent. digeſtivum*, and then to be carefully treated with the balſamic Cataplaſms and Fomentations: to this Place belongs the following Fomentation, beſides thoſe mentioned, § VII, VIII, IX.

℞. *Decocti Hordei vel Scordii* ℔ j. *Acet. Rutac.* ℥ vj. *Spir. Vin. Theriacal.* ℥ v. *Sal. Marin. aut Vulgar.* ℥ j. *vel* ij. *Miſc.*

a I made Trial of the *Cortex* lately upon a corpulent female Patient of near ſixty, who was afflicted with a Gangrene from an internal Cauſe, about the lower Part of the *Tibia*, *Tarſus*, and *Metatarſus*, wherein the common Integuments of the Body were already ſphacelated and corrupted: but ſhe always threw up the Remedy by Vomit, ſoon after every time ſhe took it, as ſhe had likewiſe done other Medicines for ſome time before; ſo that I was obliged to lay it aſide; but after many other Things tried in vain, I at length reſtrained her Vomiting by the *Pyrmont Waters*, drank cold (for ſhe threw them up when warm) and performed the reſt of the Cure by the Medicines hereafter recommended for the Cure of a *Sphacelus*; whence it appears, that all Gangrenes and *Sphaceli* from internal Cauſes are not incurable, as ſome Authors have aſſerted.

this is to be applied hot with Compreſſes to the inciſed Parts, and frequently repeated, till the Diſorder appears to ſpread no further. We know the *Sphacelus* ceaſes to ſpread, when the Tumor of the vitiated Parts ſubſides and the Lips of the adjacent ſound Parts become tumid all round; and on the ſecond or third Day after, a Suppuration is gradually formed, and the ſound Parts ſeparate from the vitiated: but to ſoften and promote a ſpeedy Separation of the Eſchar afterwards, the following Cataplaſm will be found very ſerviceable:

℞. *Folior. Scordii M.* ij. *Malvæ Hyoſciam. Alth. ana M.* j. *Flor. Lavendul. M.* ß. *coquantur cum Aceto vel Oxycrato ad Conſiſtentiam Cataplaſmatis, cui tandem admiſce Farin. Lin.* ʒiij. *Ol. Lin.* ʒj. *Sal. Ammoniac.* ʒij. *F. Cataplaſma.*

This is to be applied warm over the whole, and it is to be retained in that Condition of Heat as long as requiſite, by the Means before mentioned at § VII, IX. This will wonderfully promote the Separation of the vitiated Parts from the ſound; eſpecially if the Bark be taken inwardly at the ſame time. We read in the *Acta Edinburg.* that a *Sphacelus* is often ſtopped, and the Ulcer healed by the Uſe of the Bark inwardly, and the external Application of the Spirit of Turpentine only. This Spirit has long ſince been much extolled by HARRIS, and I myſelf have found it extremely ſerviceable.

The Separation and Cleanſing of the Parts how to be effected.

XII. After theſe Medicines have been uſed, and when the whole ſurrounding Skin is gently tumified, with Redneſs, a Cruſt or Eſchar is formed by Degrees, and the ſound Fleſh begins to ſeparate from the reſt; this is then a Sign that the Diſorder has done ſpreading, and that an entire Separation of the vitiated Parts will ſhortly follow: therefore, whenever this Separation ſhews itſelf, it ſhould be promoted as much as poſſible, by the Uſe of ſome ſuppurating Ointment, ſuch as is commonly termed digeſtive; which may be applied either alone or mixed with ſome *Theriaca*; to be retained on between the ſound and dead Parts (which may be ſometimes a little divided by the Lancet), after which the preceding Cataplaſm ſhould be applied: but in all future Dreſſings, whatever of the dead Parts is found looſe, or ſeparated, ſhould be removed every Day; or, if any of the vitiated Parts ſhould in ſome Meaſure adhere to the ſound, they may be ſeparated by the Sciſſars or Scalpel, without any great Pain or Danger. After this, it will be proper to remove the Cataplaſm, and apply ſome digeſtive Ointment, or *Empl. Diachyl. vel Saturnin.* in the room thereof, till the corrupted Parts are entirely caſt off, and the Ulcer appears to be well cleanſed. The Separation of the corrupted Parts from the ſound, may be wonderfully promoted by keeping the diſordered Limb in a conſtant Warmth, by Cataplaſms covered with hot Bricks or Tiles, to retain the Heat and avoid the frequent uncovering of the Parts to apply freſh Cataplaſms. When the ſound Parts are ſufficiently deterged or cleanſed, we muſt then proceed to their Agglutination or Cure: in order to which we ſhall find great Benefit from *Ung. Digeſtiv. vel Baſilicum, vel Balſ. Arcæi*, together with the forementioned Plaſters.

Cauſtics, which and how to be applied.

XIII. But there are many Surgeons, who, to avoid the Length of Time which is uſually taken up in forming a Suppuration, and for ſome other Reaſons, have Recourſe directly to *cauſtic Medicines* in this State of their Diſorder. Their Method of Treatment is this: they anoint the Lips only, or elſe the whole, of the corrupted Parts every Day with *Butyr. Antimon.* or *Lap. Cauſt. liquefact.* till the

the living Parts are ſurrounded by a Sort of Eſchar: and always afterwards they apply the forementioned (§ IX and XI.) Fomentations and Cataplaſms; in order to prevent the Diſorder from ſpreading, and to make the corrupted Fleſh ſeparate from the ſound: to this Place belongs the *Aqua Phagedænica* and the *Lixivium rodens* BOERHAAVII *in Mater. Med.* § 462:

℞. *Calc. viv. fortiſſ.* ℥ iij. *Ciner. Clavellator.* ℥ ix.

Theſe are to be firſt ground ſeparately, and to be afterwards mixed together, adding a little Water; then let them be put in a Glaſs, and ſtand in a moiſt Cellar to diſſolve: as ſoon as they are become fluid, filtrate them through coarſe and ſpongy Paper, and then let the Liquor be preſerved for Uſe. When there is Occaſion to uſe it, leta Bruſh or Feather be dipped into it, and afterwards rubbed over the Part, once or twice in a Day, as you ſhall ſee Occaſion: or you may wet fine linen Rags with this Liquor, and lay them upon the Part, not neglecting the Uſe of the emollient Cataplaſms at N°. VIII. or IX. at the ſame time: This Method of Dreſſing ſhould be continued till the Parts ſhall ſuppurate or fall off in Cruſt or Scales. If this Application has ſo far anſwered your Intention, you may proceed to cleanſe the Wound with Digeſtives, and afterwards heal with a vulnerary Balſam, as we juſt now directed above at N°. XII. But if any Miſchief ſhould remain underneath after you have healed, you muſt again have Recourſe to corroſive Medicines; and, as to the reſt, proceed as we have directed above: the beſt Form of a corroſive Application that I have ſeen, is deſcribed by BELLOSTE, in his *Hoſpital Surgeon:* He is not ſhort in commending it himſelf; he ſays, when you are furniſhed with this, you may ſpare yourſelf the Trouble of ſearching for a better Remedy. The following is the Deſcription of it:

℞. *Spirit. Nitri vel Aquæ Fort. P.* ij. *Argenti Vivi P.* j. *m. f. lento calore Mercurii ſolutio.*

The mortified Part is to be wetted with this corroſive Liquor, which will occaſion a ſpeedy Separation of it from the ſound Parts. But I myſelf have ſeen, where the vitiated Parts wete not ſeparated from the ſound by this Corroſive; and, what is worſe, even the ſound Parts were deſtroyed by the Application of it.

Of the actual Cautery in this Caſe.

XIV. Several Phyſicians and Surgeons, particularly the famous[a] BOERHAAVE, adviſe cauteriſing or dividing with the Knife down to the Bottom where it is ſound, and this Method they prefer to all others: but as this Kind of Treatment carries great Cruelty with it, and cannot be performed without giving the Patient violent Pain, and is frequently attended with Danger, I cannot help preferring the Uſe of *Suppurants* or *mild* Corroſives, as a gentler and ſafer Method of Cure: and indeed the Surgeons of the preſent Age, in general, are not ſo fond of calling for the actual Cautery as their Fathers were, eſpecially where they can find Remedies of equal Efficacy.

Amputation when to be performed.

XV. Laſtly, when the *Sphacelus* is ſo deeply fixed in any Part of the upper or lower Extremity, that it has penetrated through the Muſcles, as far as to the Bone, and has reſiſted all the Force of Medicines, or the proper Time for applying them has been neglected; in this Caſe, for the Preſervation of Life in the

a. BOERHAAVE's Aph. *De cognoſc. & curand. Morbis*, No. 462.

Parts

Parts that remain untouched, the injured Part muſt be ſeparated from the Body, with proper Inſtruments. We ſhall fully deſcribe the Method of doing this in each particular Part of the Body, when we come to treat of *Chirurgical Operations*. In the mean time, I cannot give the Surgeon a more ſeaſonable Piece of Advice than this: that when ever he thinks the Amputation of a Part neceſſary, he cannot more effectually conſult his own Reputation and his Patient's Safety, than by calling in a prudent *Phyſician* or two, that may confirm his Opinion of the Neceſſity of the Operation; and may give him their Aſſiſtance if any bad Accident ſhould happen, ſuch as Hæmorrhage, Faintings, Fever, and the like, which are very common Conſequences of theſe great Operations: he ſhould, above all Things, conſider the Strength of the Patient; whether he is able to undergo the Operation: and where the *Sphacelus* ariſes from an internal Cauſe, and the beſt Remedies have proved ineffectual, the Amputation ſhould not be haſtily undertaken: for in this Caſe, it is generally unſucceſsful; as GARENGEOT and many others have teſtified: the Surgeon ſhould alſo be very careful in keeping up the Strength of the Patient as much as poſſible, leſt he ſhould ſink under the Diſcharge of Matter.

CHAP. XV.

Of BURNS *and* SCALDS.

A Burn, what.

I. I BELIEVE no one will be offended at our treating of *Burns* as a Species of Inflammation, ſince the Appearance, as well as Conſequences, of both are exactly the ſame. Injuries that are received in any Part of the Body, either by Fire itſelf, or by the Inſtruments heated with Fire, we call a Burn or Scald. Therefore we do not reckon Fire alone as the Cauſe of Burns and Scalds, but any other Bodies, whether ſolid and hot, as live Coals, Iron, or other Metal, redhot or melted; Gunpowder, or boiling Liquors, as Water, Beer, Wine, Oil, *&c.* are all to be reckoned under this Head.

The Nature of a Burn or Scald.

II. When any thing of this Kind is applied to the Body, Fibres and ſmall Veſſels of the Parts that are touched by it will inſtantly corrugate and burſt, whilſt the Blood and other contained Fluids will be extravaſated, ſtagnate, and corrupt. The Burns that we receive from ſolid Bodies are always attended with more grievous Conſequences than thoſe which are occaſioned by boiling Liquors (which we call *Scalding*); therefore there are different Degrees of this Injury, as there are of Inflammation.

Four Degrees of Burns.

III. We may very fairly therefore divide Burns or Scalds into four Degrees. The *firſt*, and the ſlighteſt, is that which occaſions Heat, Pain, and a ſmall Veſication on the injured Part, in a ſhort Time. The *ſecond* Degree is, when the Part is inſtantly affected with great Pain and Veſication. The *third* is, when the common Integuments and ſubjacent Fleſh are ſo burnt, that they form a Cruſt. The *fourth* and laſt is, where every thing is deſtroyed quite down to the Bone. The third Species is nearly allied to the Gangrene, and the fourth to a *Sphacelus*. This illuſtrates the near Relation between Burns and Inflammations.

Prognoſis.

IV. By conſidering the Degree of the Burn, and the Uſe and Conſequence of the Part burnt, you may prognoſticate in what Manner the Injury will terminate. A Veſication raiſed in the Hand by the Fire, is leſs to be dreaded, than

than a ſlighter Burn upon the Eye; for that very tender and uſeful Part of the Body can ſcarce receive any Injury by Fire, without endangering the Loſs of Sight: we ſhould alſo conſider the Extent of the Burn, what Length of Time it has been upon the Part, before we can form a true Judgment of the Conſequences that will attend it: for the Danger will be greatly increaſed by the length of Time that the Part has been injured, and in Proportion to the Degree to which the Injury has ſpread itſelf: for where the whole Surface of the Body is burnt with Gunpowder, or ſcalded with any boiling Liquor, though the Injury, conſidered in any particular Part, ſhall be looked upon as a very ſlight one, yet by being ſpread to ſo great an Extent, it is a Diſorder of the laſt Conſequence: in this Caſe, it is impoſſible for the Patient to lay down, or change his Poſture, without horrid Pain and Torture, which will prevent his Sleep, increaſe his Fever, and by Degrees bring on a *Sphacelus* and Death itſelf: and this is the Caſe more particularly in Infants, ſince they have leſs Strength and Patience than Adults, and want Reaſon to diſcover which would be the moſt convenient Situation for them. The Danger of the Burn will likewiſe be increaſed, in Proportion to the Depth to which it has penetrated: Burns of the Face are not only to be dreaded for the Deformity which they occaſion, but chiefly for the Inconveniences that they may produce by cauſing the Eyelids to grow together. Deep Burns of the Neck, if not timely remedied, occaſion a Wryneſs of that Part: you will eaſily be able to foretel what Danger or Inconvenience will ariſe from Burns of any other Part, if you diligently conſider what we have here ſaid, and are well ſkilled in the natural Uſe of the injured Parts.

V. As we obſerved above that Burns nearly reſembled inflammatory Diſordets in their Degrees, ſo do they in the Method of Cure. In the ſlighteſt or firſt degree of a Burn, the Intenſion is to diſperſe it by the Remedies which we adviſed for a *Phlegmon*, (*Chap.* II. § IX) Of theſe there are two Sorts, *Aſtringents* and *Emollients:* the beſt ſlight Aſtringent is, *Spiritus Vini*[a] *vel vulgaris bonæ notæ, vel rectificatus, vel & camphoratus:* this may be applied to the Part with linen Rags: with the ſame Intention alſo you may order *Acetum Lithargyriſatum, Muria Braſſicæ conditæ, vel & Oxycratum cum Sale decoctum calidumque:* theſe may be applied in the ſame Manner with the foregoing, and ſhould be repeated as you ſhall ſee Occaſion. *Oleum Terebinthinæ* has very good Effects is this Caſe, if you apply it in Time, and repeat it frequently: the vulgar Method of applying the burnt Part to a Candle or the Fire, and keeping it in that Poſition as long as you can bear it, repeating this Proceſs till all Senſe of Heat and Pain is entirely removed, is frequently attempted with Succeſs, where the Injury is in one of the Fingers or on the Hand: for the ſtagnating Fluids are by the Force of the Fire, driven back into their proper Channels, and by this Means the Veſication and other troubleſome Symptoms which uſually ſucceed, are happily prevented: from hence it appears, that the firſt Degree of Burns is eaſily remedied.

Cure of the firſt Degree by diſperſing Remedies.

VI. There is another Method of Cure, which is equally efficacious with the former, though it is founded upon a contrary Intention. This is by *emollient Remedies,* which remove the Tenſion of the Fibres and Veſſels, and reſtore the

By Emollients.

[a] This is highly and deſervedly recommended by the great SYDENHAM, in his *Practical Works.*

Blood to its natural Courſe, before any bad Symptoms come on. The injured Part may be fomented with Water, as hot as the Patient can bear it, till the Pain and Heat entirely diſappear: SYDENHAM highly recommends this Practice, and, in my Opinion, with great Juſtice: but this Fomentation will be improved if you boil ſome emollient Ingredients in the Water, as *Althæa*, *Malvæ*, *Verboſcum*, *Sem. Lini*, *Fœnu Græc. Mali Cydonii Semina*, or others of this Intention. But emollient Cataplaſms are of the higheſt Service in this Caſe, made of any of the abovementioned Ingredients for a Decoction, and frequently laid on upon the affected Part as warm as they can be endured: emollient Oils alſo have their Uſe in forwarding this Intention, as *Oleum Lini*, *Amygdalarum dulcium*, *Olivarum*, *Liliorum alborum*, *Hyoſciani*, and the like. Theſe Oils are to be uſed either by dipping Rags into them, and applying them to the burnt Parts; or they may be laid on frequently with Feathers as faſt as they begin to dry away. We muſt not omit in this Place to mention a famous Liniment of MYNSICHTUS, which he calls his *Unguentum ad Ambuſtiones*. This is compoſed *ex Oleo Lini vel Olivarum cum Albumine Ovi mixto*, and applied as the Oils above: *Mali Cydonii Mucilago* is properly enough preſcribed in this Caſe: the Remedies which we have here recommended never give effectual Relief, unleſs frequently repeated: therefore, when the Face is burnt they ſhould be ſpread upon a linen Maſk, which you muſt keep continually moiſt by freſh Applications of the Remedy. (See *Plate* XXXVII.) Where the Neck is burnt, to prevent it from contracting, you muſt have Recourſe to a particular Kind of Bandage, which you will find deſcribed below, when I come to treat of Bandages.

Cure of the ſecond Degree.

VII. When the Burn is of the ſecond Degree, which I have deſcribed above, attended with Veſication or Puſtules, I would by no means adviſe opening the Veſications, or ſcarifying the lacerated *Cutis*, becauſe this Practice brings on very ſharp Pains: you will always find it more adviſeable to apply one of the Remedies preſcribed above; the neareſt at hand, ſuppoſe warm Water, burnt Wine, or Spirits of Wine; and renew the Application of it frequently: by this Means you will find the Heat and Pain quickly go off, and the Cuticle will ſeparate from the *Cutis*, without leaving any Deformity: but if, notwithſtanding the repeated Applications of theſe Remedies, ſome Pain ſhall ſtill remain, dreſs the Part with Emollients. The moſt eligible of theſe are *Oleum Lini*, *Unguentum ad Ambuſtionem* MYNSICHTI, *vel Nutritum*, *de Lithargyrio*, *vel Diapompholygos*: theſe ſhould be either rubbed into the Part frequently, or ſpread upon a linen Rag and applied to it. After the Heat and Pain are removed by theſe Applications, lay on the *Empl. ad Ambuſta*, *vel de Minio*, which will keep the Skin ſmooth, and forward the Renovation of the Cuticle. If the Injury is very conſiderable as to its Extent, and great Part of the Body is ſcalded or burnt, it will be neceſſary to open a Vein and bleed plentifully, even *ad Animi Deliquium*, and afterwards you ſhould preſcribe a briſk Purge, of the ſame Kind which we directed for Contuſion. (*Book* I. *Chap.* XV. § XIII.) This Method may poſſibly prevent ill Conſequences which uſually attend Burns of large Extent, ſuch as foul Ulcers, large Cicatrices, and Gangrene itſelf: the ſame external Dreſſings are to be applied in this Caſe which we adviſed above. When *Infants* are the Subjects of this Diſorder, their tender Age prevents us from Bleeding plentifully: therefore the Revulſion muſt be made by repeated Purging. That ſtrict Regularity in Diet which we enjoined above in treating of

Wounds and Inflammations, is never more requiſite to be obſerved than in this Caſe; all Intemperance is of the laſt Conſequence, as it increaſes the Fever and Pain. According to the Opinion of the famous DIGBY, nothing takes off the Heat ſooner than *Spiritus Salis* given from *Guttæ* x to xv. in any Liquor, and repeated at Diſcretion: theſe Methods, being timely and diligently proſecuted, heal and reſtore the burnt Parts of the Body in a moſt wonderful Manner.

Cure of the third Degree.

VIII. In the third Degree of Burns, where the injured Part is covered with a Cruſt or Eſchar, the Cure cannot be performed without Suppuration: when this happens in the Face, we ſhould uſe all our Attention to prevent Deformity, which may be occaſioned by a large Cicatrix: therefore, in this Caſe, the Uſe of all Plaſters and Ointments whatſoever is to be avoided, even though they ſhould be eſteemed as valuable Secrets, and highly commended for their Virtue in curing Burns and Scalds: for the Miſchief of theſe Kinds of Remedies is, that they dry up the Wound too faſt, and at the ſame time contract the Fibres and the Skin, and by that Means leave a very unequal Cicatrix: for the ſame Reaſon you cannot be too ſolicitous in forwarding the caſting off of the Eſchar, and the Evacuation of the Matter that is concealed under: but to diſcover the happieſt Means of performing theſe Intenſions, *hoc Opus, hic Labor eſt:* they who attempt this by tearing away the Eſchar with their Hands, or endeavour to ſeparate it with the Knife, by no means conſult the Good of their Patients: the eaſieſt and moſt ſucceſsful Method, in my Opinion, is, by the Uſe of Emollients: any of the Emollients we mentioned above may be applied warm, and repeated till the hard Cruſt ſeparates from the live Fleſh: the Part ſhould be dreſſed two or three Times in a Day, and at each Dreſſing, if you ſhould obſerve any Portion of the Cruſt tending to the Separation from the reſt, you ſhould remove it with your *Forceps*,. and anoint the remaining Cruſt with Butter, at the ſame time being never neglectful of the Uſe of Fomentations: this Method ſometimes takes up two, ſometimes three, ſometimes four Days before it performs its Office: the Cruſt being now entirely caſt off, our next Intention is, to cleanſe and heal the Wound: the firſt of theſe Offices may be very well executed by any mild digeſtive Ointment, mixed up with *Mel Roſarum*; the Medicines uſed for healing the Wound, are principally *Unguentum Diapompholygos, vel de Lithargyrio, necnon Emplaſtrum ad Ambuſta*; but if any Portion of the Eſchar is left under theſe Ointments and Plaſters, Experience ſufficiently teſtifies the Danger that will follow, of making a deformed Cicatrix, from the Conſtruction of the neighbouring Parts, and from the Acrimony of the confined *Sanies*. Whoever proſecutes this Method of Cure, ſhould always obſerve, that if the Eſchar does not ſeparate in two or three Days, it will be neceſſary for him to make a deep Inciſion into it, that the *Sanies* may have Room to diſcharge itſelf, as we adviſe in the Caſe of Gangrenes, (*Chap.* XIV. § VII.) and then the Fomentations above mentioned are to be diligently applied, the Evacuations by Bleeding and Purging being always premiſed: proper Regulations with regard to Diet are never more neceſſary to be complied with, than in this Caſe: the beſt Method of encouraging the Renovation of the Skin, is, by frequently holding the burnt Part over the Steam that ariſes from boiling Water: where the Part ſkins over very ſlowly, it may be proper to dreſs the Part with a Cerate made *ex Cerâ & Ovorum Oleo.*

Cure of the fourth Degree.

IX. But what is to be done in the fourth Degree, which we have deſcribed, which is always attended with extreme Danger? for when the Burn has penetrated ſo deep as to deſtroy all the Parts, quite down to the Bone, Medicine can take no Place; therefore there remains but one Remedy, and that a dreadful one, to wit, to amputate the injured Limb, that the ſound Parts may be ſaved, as we adviſed above in treating of a *Sphacelus* (*Chap.* XIV.)

CHAP. XVI.

Of a SCHIRRHUS.

A *Schirrhus* what.

I. WE have already taught, that the fourth Manner in which an Inflammation terminates is a *Schirrhus*: we uſually call a hard Tumor of any Part of the Body, that is void of Pain, a *Schirrhus*; this almoſt always ariſes from the Inſpiſſation and Induration of the Fluids contained in a Gland, though it may appear in other Parts, particularly in the Fat [a].

Seat of a *Schirrhus*.

II. The Seat of a *Schirrhus* is very various; for this Diſorder is not confined to the internal Parts alone, to wit, to the Liver, Spleen, Lungs, Meſentery, Pancreas, and in Females to the *Uterus*: but it frequently happens alſo to the external Parts, as to the Lips, Tongue, Tonſils, *Fauces*, Palate, Gums, Neck, *Mammæ*, *Axillæ*, *Inguina*, *Penis*, and Teſticles; and that generally after a previous Inflammation of any of thoſe Parts. A *Schirrhus* ſometimes appears without any previous Inflammation; eſpecially in Subjects of a heavy, phlegmatic, melancholic Habit of Body: ſometimes it is occaſioned by an external Injury, as by a Fall or Blow, *&c.* it is no difficult Matter to determine the principal Cauſe of the Diſorder.

Effects of a *Schirrhus*.

III. As ſoon as a *Schirrhus* is formed, it is an immediate Conſequence that not only the indurated Part becomes unfit to perform the Function allotted it by Nature, but the neighbouring Parts alſo will ſuffer Preſſure, and be impeded in the Performance of their Offices: therefore it ought to appear no Wonder that the neighbouring Parts ſhould be ſubject to Inflammations, Exulcerations, Cancer, Gangrene, *Tabes*, Stiffneſs, Immobility, or the like, according to the Nature of the injured Part.

Signs.

IV. You will be at no great Difficulty in determining the Caſe to be a *Schirrhus*, when you diſcover a hard Tumor, on the external Parts (more particularly in thoſe Parts where the Glands are moſt frequent) and the Tumor is entirely free from Heat, Redneſs, and Pain: as I am ſpeaking to *Surgeons*, I only treat of external *Schirrhi*; for thoſe which are ſituated in the internal Parts, fall very juſtly under the Province of the *Phyſician*.

Prognoſtic.

V. In order to form a proper Prognoſtic of this Diſorder, ſeveral Things are to be obſerved: (1.) the more inveterate the *Schirrhus* is, ſo much the more dangerous will it be, and more difficult of Cure: (2.) a *Schirrhus* happening to young Perſons, and to thoſe of a firm Habit of Body, is much more ſafe and tractable, than when it falls upon Perſons advanced in Years: particularly

[a] See a learned Diſſert. of GRASHUSIUS, an eminent Phyſician at *Amſterdam*, on the *Schirrhus* and *Cancer*.—*Amſt.* 1741.

where

where Children have indurated Glands in the Neck, but are in all other Reſpects in perfect Health, they are ſeldom attended with any Miſchief; and you uſually find they outgrow it; but in Valetudinarians, or where you have Reaſon to ſuſpect the Pox to be at the Bottom, the Caſe is far otherwiſe: (3.) A *Schirrhus* is of more or leſs Conſequence in Proportion to the Conſequence of the Part it falls upon, in performing the neceſſary or noble Offices of Life: for this Reaſon, internal *Schirrhi* are always more dangerous than thoſe which happen upon the external Parts[a]. Laſtly, (4.) the greater Miſchiefs the *Schirrhus* brings on, by ſo much the more grievous will it be: for as long as it lays quiet, and produces no Pain, ſo long will it remain without Danger; but as ſoon as it becomes painful, or is ulcerated, it generally threatens an approaching *Cancer*: it may be proper to inform you in general, that the Cure of *Schirrhi* by Medicine, is uſually attended with the greateſt Difficulty; therefore you ſhould never flatter your Patients with the Promiſe of certain Relief: but ſometimes they do admit of a Cure with the Knife or with Corroſives, eſpecially in younger Subjects that are otherwiſe of a good Habit of Body.

Method of Cure.

VI. When the *Schirrhus* is of long ſtanding, and the Patient infirm, it is far better to abſtain entirely from any Attempt to cure it, than to pretend to bring it to Digeſtion; for in this Caſe it is much to be feared, eſpecially in the Breaſts of Women, that whilſt you are proſecuting your Intention, the diſeaſed Part may ſhew its bad Diſpoſition, and become apparently cancerous: On the other hand, where the *Schirrhus* is but newly formed, and you have no Signs of vehement Pain or Hardneſs, where your Patient is otherwiſe of a ſound Habit of Body, I ſee no Reaſon why you ſhould not uſe both external and internal Remedies, to ſet the confined Fluids at Liberty. The internal Remedies, which are found principally ſerviceable in anſwering this Intention, are the Decoctions of the Woods, digeſtive Tinctures or Eſſences, and mild Mercurials, giving between whiles relaxing Medicines, to reſolve the thick inſpiſſated Humours[b]: it is very dangerous to truſt to the Uſe of external Remedies alone; therefore a prudent Phyſician ſhould always be conſulted in this Caſe, who may not only preſcribe proper internal Remedies, but direct the Patient alſo what Sort of Regimen will be moſt uſeful for him to obſerve, with regard to his Diet.

Diſperſing by (1.) Plaſters.

VII. With regard to external Reſolvents, *Plaſters* claim the firſt Place; ſuch I mean as are made of the warm Gums, as *Gum. Ammoniac. Galban. Opopon. Sagapen. Bdell. &c.* which may be applied alone or mixed together; ſometimes with the Addition of *Radix Bryoniæ atque Ariſtolochiæ rotundæ* finely powdered: of the ſame Intention are *Empl. de Galbano, de Gum. Ammoniaco, de Cicutâ, de Ramis* VIGONII, *vel Diachylon cum Mercurio*; or the following:

℞. *Gumm. Galban. Opoponac. ana* ℥ j. *Ammoniac. Bdell. ana* ℥ ij. *Ol. Olivar.* ℔ ij. *Ceræ Citrin.* ℔ *ſs. Pulv. Ariſtoloch. long. ver. & rotund. ver. Lapid. Calaminar. Myrrh. Thur. ana* ℥ j. *Terebinthin. Venet.* ℥ iv. *m. f. ſ. a. Emplaſtrum.*

[a] Where a *Schirrus* falls on the internal Parts, particularly the Meſentery or the Lungs, it is generally attended with ſuch Tumors, as deſtroy the Patient.

[b] For the internal Method of Cure, ſee my *practical Compendium*, Cap. vii, Sect. 29 & 30.

(2.) By Cataplaſms.

VIII. The next Place to Plaſters is held by *Cataplaſms:* amongſt the principal of which may be reckoned the following:

℞. *Rad. Bryon. alb.* ℥ iv. *Ariſtoloch. rotund. Angelic. ana* ℥ j. *Herb. Sabin. Rut. Scord. Abſinth. Flor. Chamæmel. ana M.* j. *Melilot. Sambuc. Altheæ, Centaur. minor. ana M. ſs. coq. cum q. ſ. Aquæ ſimplic. ad conſiſtentiam Cataplaſmat. Vaſe clauſo, ſub Finem addendo Galban.* (*Vitell. Ov. q. ſ. ſolut.*) ℥ iij. *Farin. Lini* ℥ ij. *Ol. Lini q. ſ. f. Cataplaſma.*

This Cataplaſm, or, if you rather chooſe it, a Fomentation made of the ſame Herbs boiled in Vinegar, is to be applied warm, and repeated as you ſhall ſee Occaſion, not neglecting at the ſame time the Uſe of internal Remedies.

(3.) By acid Vapours.

IX. Some highly recommend acid Vapours in this Caſe: ſometimes it has been found ſerviceable to receive the Steam of boiling Vinegar upon the diſeaſed Part, either of common Vinegar, or of that made with Lavender, Alder, Rue, or *Theriaca*: ſome ſprinkle the Vinegar upon a hot Stone, and receive the Steam through a Funnel: others ſet *Sulphur* on Fire, and hold the Part over the Fume: others again are fond of Fumigations of *Cinnabar*: great Care muſt be taken in this Caſe not to raiſe too large a Fume, nor to repeat it too frequently, and the Patient muſt be cautioned not to admit it at the Noſe or Mouth; for it can ſcarce be ſaid how injurious theſe Steams are to the Lungs, and the Quantity of Mercury contained in Cinnabar makes it very apt to raiſe a Salivation.

(4.) By Mercurials.

X. Mercurial Medicines perform Wonders in this Caſe, either adminiſtered in the Beginning, or after other Remedies have failed. Beſides giving Mercurials internally, you may make an excellent Ointment, *ex Hydragyro cum Adipe Suillâ, necnon modico Terebinthinæ, quantum ad eum ſubigendum ſufficit, admixtis in Mortario vitreo vel lapideo:* the *Schirrhus* ſhould be anointed twice or thrice a Day with this, covering it with the *Emplaſtrum Vigonis cum Mercurio:* but to prevent this Method from raiſing a Salivation, it will be neceſſary to preſcribe an opening Medicine every fourth or fifth Day, ſuch as *Rad. Jalap. præp.* or *Extract. Rud.* in ſmall Doſes: whilſt the Patient is in this Courſe, his Jaws ſhould be very diligently inſpected, and if you find the Glands enlarge and grow painful, you muſt entirely omit the Uſe of Mercurials, and repeat your purging Medicines, till all theſe Symptoms of an approaching Ptyaliſm entirely diſappear: by obſerving theſe Cautions, you may have very good Reaſon to promiſe yourſelf Succeſs, where you are called in Time, before the Caſe is become deſperate.

(1.) By the Knife.

XI. If all the abovementioned Remedies prove unſucceſsful; if the *Schirrus* is free and moveable, and its Situation threatens no great Danger from the Neighbourhood of conſiderable Veſſels; if you ſhall judge the Strength of the Patient to be ſufficient to undergo the Operation, you may very fairly call the Knife in Aid, to prevent the Caſe becoming cancerous (which too often happens;) when you have taken out the *Schirrus*, dreſs the Wound with a digeſtive Ointment, and afterwards with the *Linimentum Arcæi*, or any other vulnerary Medicine, and heal as we have directed in other Wounds: this Method DEIDIER particularly recommends, *Lib. de Tumor.* p. 129.

When the Caſe is entirely to be let alone.

XII. Where the *Schirrus* is fixed, knotty, uneven, and deeply rooted; where the Patient is of a bad Habit of Body, is ſubject to form *Schirrhi* from ſome hereditary

hereditary Taint, or perhaps has formed ſeveral already; or laſtly, where the Situation of the Diſorder is ſuch, that, from the Vicinity of conſiderable Veins and Arteries, you are in apparent Danger of bringing on an Hæmorrhage which may prove fatal; then all Attempts to cure, whether by the Knife, or by digeſtive or corroſive Applications, are to be neglected[a]: for this Kind of *Schirrhus* is almoſt conſtantly attended with very ſharp Pains, and often degenerates into a *Cancer*: in this Caſe therefore the Pains are to be aſſuaged, if poſſible, and the Cancer to be prevented.

Internal Medicines to prevent a *Cancer*.

XIII. When you are under Apprehenſions of an approaching Cancer, your Buſineſs is not only to attempt to correct the Acrimony of the Blood, by the Uſe of both internal, as well as external Remedies, but a ſtrict Regimen with regard to Diet muſt alſo be moſt religiouſly enjoined. Conſtitutions of this Sort are much mended by the Uſe of Broths and Soops of various Kinds, made from the Fleſh of younger Animals, with the Addition ſometimes of a few Pot-herbs, *ſc. Hordeum, Avena, Oryza, Milium, Spinachia, Aſparagus, Scorzonera*, &c. The moſt wholeſome common Drink in this Caſe, is either fair Water, or a Ptiſan made *ex Decocto Radicis Chinæ, Sarſaparill. Gramin. Polypod. Veronicæ, Linguæ Cervinæ, Agrimon. Solidagine Saraſenica, Herb. Parietar. Capillor. Veneris*, and others of this Kind: when the *Schirrhus* is attended with violent Pains, you may add to the Materials of your Decoction, *Sem. Papaver. albi*, and if the Patient has no Objection to it, you may ſweeten it *cum Sirupo Papaver. albi*. It will be very proper alſo at this Time to correct the Acrimony of the Blood by giving two or three Times every Day, a Doſe, *e Pulv. Lap. Cancr. Sale Abſinthii, Cinnabari Nativâ, Antimonio crudo, Antimonio diaphoretico*, adding to each Doſe, as you ſhall ſee Occaſion, *Laudani Opiati gr.* ß. to aſſuage the Vehemence of the Pain: Wonders are alſo effected in this Caſe by the *Pulvis Succuſve recens ex Millepedibus*, with *Spermr Ceti ad* ʒj. to be given with any of the foregoing Powders; by Purges even of the mercurial Kind, and by bleeding and cupping frequently in Spring and Autumn.

External Applications.

XIV. A thin Plate of Lead, well impregnated with Quickſilver, may very conveniently be faſtened on the Part, and worn there conſtantly with ſome Benefit: for this Method frequently leſſens the Senſe of Heat and Pain, not to ſay that it frequently prevents the Cancer: but if the Application of a Plate of Lead ſhall ſeem to be unequal to the Intention for which it was deſigned, then you may apply Plaſters and Ointments compoſed of ſuch Ingredients as are moſt likely to aſſuage the Pains. Of this Kind are the following:

℞. *Unguenti Diapompholygos* ʒij. *Opii puri* ∋ ß. *m. f. Ung. quocum pars affecta ſæpius inungatur. Vel,*

℞. *Amalgam. Mercur. & Plumbi* ʒj. *Unguenti Roſati q. ſ. m. f. Unguentum cum Linteo inſtar Emplaſtri applicandum. Vel.*

℞. *Aceti Lithargyriſat.* ʒj. *Olei expreſſ. Sem. Hyoſcyam. Papav. alb. Olei infuſ. Roſar. ana* ʒij. *m. f. ſ. a. Nutritum, cui ſub Finem add. Opii puri gr.* vj. *ad* x. *quod Linteolis illitum aliquoties quotidie ſuper Schirrhum applicetur.*

[a] But if the Surgeon is expeditious in ſtopping the Blood and dreſſing the Wound, he may ſometimes attempt the Cure of the moſt inveterate Schirrhi, and not without Succeſs. I have myſelf often undertaken very extraordinary Schirrhi in the parotid and ſubmaxillary Glands, where I was obliged to cut the large Ramifications of the external Carotid Artery; nor did any one of thoſe Patients miſcarry.

If

If your Patient diſlikes the Application of theſe Ointments, and prefers a neater Application, you may ſubſtitute refrigerant Plaſters in their room; ſuch as *Emplaſtrum Saturninum* MYNSICHTI. *De Minio*, *Diapompholyges*; or laſtly, that excellent Plaſter for alleviating Pain, which is preſcribed in the following Manner:

℞. *Succ. recent. expreſſ. & purificat. Fol. Hyoſcyam. Papav. Hortenſ. Phellandr: ana* ℥ iv. *coquendo leni igne inſpiſſa*; *ſub Finem add. Ceræ alb.* ℥ viij. *Ol. infuſ. Roſar.* ℥ j. *m. f. Emplaſtrum. Vel,*

℞. *Sacch. Saturn. Cerüſſ. præparat. Amalgam Mercurii & Saturni, Ol. expreſſ. Hyoſciam. infuſ. Roſar. ana* ℥ij. *m. f. Emplaſtrum.*

If the Pains are very violent, you may add a diſcretional Quantity of *Opium* to either of theſe Plaſters, and apply it to the Part.

What is to be concluded concerning Suppurants, Corroſives, and the actual Cautery.

XV. Notwithſtanding many Phyſicians and Surgeons of Eminence at this Time recommended the Uſe of *Suppurants*, *Corroſives*, and even the *actual Cautery*, for the Cure of ſchirrhous Tumors, yet I cannot help being of Opinion, that the Danger of a Cancer enſuing from the Uſe of Suppurants or Corroſives [a], and the natural Dread that moſt People are ſtruck with at the Sight of a red hot Iron, beſides innumerable other Inconveniencies, ought to diſſuade us from attempting ſuch ſlow, hazardous, and cruel Methods of Cure. For this Reaſon it will appear, that the ſafeſt and readieſt Method of deſtroying a large or painful *Schirrhus*, is, to cut the indurated Part entirely out, whether it be ſituated on the Lips, Salivary Glands, *Mammæ*, or Teſticles, provided you run no Riſque of a mortal Hæmorrhage, (§ XI, XII.) If you leave any Part of it behind, there is great Danger you may lay a Foundation for a Cancer; nay, what is hardeſt of all, though the *Schirrhus* be entirely rooted out, it frequently happens that another ſprings up without any Fault to be laid to the Surgeon. I can by no means approve the Practice of ſome Phyſicians, who order the Bottom of the Wound to be cauterized, to prevent any Return of the *Schirrhus*, and to take off the Hæmorrhage. In this they are doing nothing [b], ſince it is of very little Conſequence in preventing the Return of the Diſorder, and there are many milder and ſafer Remedies at hand to ſtop the Hæmorrhage: therefore, when you have finiſhed your Operation, dreſs as in other Wounds.

CHAP. XVII.

Of a CANCER.

A Cancer, what.

I. WHEN a *Schirrhus* can neither be diſperſed, ſoftened, (See *Chap.* XVI. *N.* XII and XIV.) or taken out with the Knife, whether it be occaſioned by the Vehemence of the Diſeaſe, or the Ignorance and Maltreatment of the Surgeon, the Patient will complain of pricking Pains in the Part, and the Tumor will ſpread itſelf unequally. This malignant and worſt State of a *Schirrhus*

[a] There are ſome Inſtances, where Schirrhi and Cancers have been cured by Cauſtics; but in general they are unſucceſsful.

[b] This was obſerved in the moſt antient Times. See CELSUS *Book* V. *Chap.* XXVII. § 2

was

was called formerly *Carcinoma*, by us a *Cancer:* for the Veins about the Part are diſtended, and form Incurvations, which ſome imagine bear a Reſemblance to a Crab's Claws. As long as the Tumor is entirely covered with Skin, it is called an *occult Cancer*; but when the Skin breaks and is ulcerated, it is termed by the Phyſicians an *ulcerated Cancer*.

Beginning and Increaſe of the Diſeaſe.

II. The Beginning and Increaſe of the Diſeaſe afford pretty near the following Appearances. At firſt there appears a very ſmall Tumor, which ſometimes maintains the ſame Size for a conſiderable Time, without any apparent Increaſe; on a ſudden it enlarges beyond all Conception: at firſt it is attended with little or no Pain, and that only by Intervals; upon the Increaſe of the Tumor the Pain becomes intolerable, ſometimes ſo violent as not to be born without Fainting. If you apply repelling or aſtringent Remedies to the Part, the Diſorder increaſes wonderfully; inſomuch, that one Month will produce more Increaſe of Pain and Tumor, than a Year without any medicinal Applications: the Uſe of Medicine will ſo far irritate this Diſorder, that the Skin will preſently break, and form a foul ſtinking Ulcer.

Seat of a Cancer.

III. A Cancer, as well as a *Schirrhus*, will ariſe in almoſt any Part of the Body; but moſt frequently in the upper Parts; as the Noſe, Ears, Lips, and the Breaſts of Women, nay ſometimes of Men; a very memorable Inſtance of which you will find recorded by BIDLO. *(Excercit. Anat. Chirurg.)* But beſides theſe Parts, the Gums, *Fauces*, and Tongue, and even the Parts of Generation are ſometimes the Seat of a Cancer.

Cauſes.

IV. The Cauſes of a *Schirrhus* and *Cancer* are common to both; only theſe ſeem to have acquired ſome additional Acrimony: the malignant *Stimuli* of a Cancer are not only produced by the Application of digeſtive, acrimonious, or cauſtic Medicines, but they are alſo occaſioned by ſundry other Cauſes. That Sort of Diet is moſt miſchievous which is moſt apt to produce Acrimony in the Blood: therefore all Perſons that are by Habit of Body liable to Diſorders of this Kind, ſhould religiouſly abſtain from Lard and Pork Meats: Grief and Trouble of Mind are very apt to create a cancerous Diſpoſition of Body: it is obſerveable, that old Maids and even married Women that do not breed, are very ſubject to Cancers in the Breaſt. This generally happens to them when they are turned of forty Years of Age, at the Time when the menſtrual or hæmorrhoidal Diſcharge begins to decreaſe or diſappear; though I have frequently known this Caſe happen to Perſons not ſo far advanced in Years, even between twenty and thirty.

Diagneſis.

V. The Signs of an *occult Cancer* are as follow: the Patient perceives an Itching, Heat, or pricking Pain, in or about the *Schirrhus*, the neighbouring Parts grow livid: the Tumor has an unequal Surface, increaſes in Size, and grows conſiderably harder than before: the Veins enlarge and become livid, though this Circumſtance does not always happen: if the Caſe is an *ulcerated Cancer*, you will diſcover it not only by the Ulceration of the Part, by an occult Cancer having preceded it, but the following Symptoms will make it evident to Demonſtration.

Symptoms.

VI. A thin *Sanies* flows from the Ulcer in great Quantities; ſometimes ſo corroſive and acrimonious, that the Dreſſings ſeem as rotten as if they had been eaten by *Aqua Fortis:* the Stench is intolerable, eſpecially to thoſe who are not accuſtomed to it, and fills the whole Chamber. The Diſorder continues to

ſpread itſelf wider, the Lips of the Ulcer enlarge, are wonderfully diſtorted and turned in; are ſometimes pale, ſometimes red, purple, green, livid, black, or variegated: Pains attended with a Senſation of burning, pricking, gnawing, come on at Times with ſuch Vehemence, that through Anguiſh and want of Sleep, the Patients are driven to almoſt Diſtraction and Deſpair, which greatly waſtes their Strength: their Appetite and Senſe of Smelling entirely fail them, till at laſt Death delivers them from a miſerable ſtinking Carcaſe. The Urgency of the Symptoms which we have recounted, depends upon the Patient's Habit of Body, and upon the Situation of the Part affected.

Prognoſis. VII. An occult Cancer, which is not attended wirh any conſiderable Degree of Pain, may be endured for a conſiderable Length of Time, without any great Inconvenience, by a Perſon endued with Strength and Temperance: but theſe very ſame Perſons, by an Irregularity of Diet, or medical Application, will be ſubject to the ſame grievous Symptoms which we have juſt enumerated: notwithſtanding what has been ſaid, many have imprudently boaſted that they have been poſſeſſed of infallible Secrets for the Cure of Cancers; though at the ſame time it muſt be confeſſed with [a] HILDANUS, and other capital Phyſicians, who confirm the Opinion of [b] HIPPOCRATES, and [c] CELSUS, that no Phyſician has yet been happy enough to diſcover a Medicine from which he could promiſe any Certainty of Cure in this Caſe. We have a very memorable [d] Example of this in ANNE of *Auſtria*, Mother to LOUIS XIV, late King of *France*, who laboured under a cancerous Breaſt, and was not only attended by the Court Phyſicians, but by almoſt every one in that Kingdom who had any Pretentions to the Practice either of Phyſic or Surgery; particularly by thoſe that boaſted of their ſecret Art in curing Cancers: but notwithſtanding all the Attempts of Art, which the Deſire of gaining a royal Reward could excite, no Help could be obtained for her; from which we may very fairly conclude, that there is no Help to be expected from any thing but the Knife. The Hopes we may entertain from Extirpation, depend upon the Degree of the Diſorder, the Urgency of the Symptoms, and the Strength and Habit of the Patient. When you ſhall be of Opinion that the Cancer is ſo deeply rooted, that it will be impoſſible to extirpate it entirely, it is far better to lay aſide the Operation, than to torment miſerable Patients without any Hopes of relieving them: for inſtance, when this Caſe falls upon the *Uterus*, *Fauces*, *Uvula*, *Tonſils*, *Axillæ*, and *Inguina*, it is ſcarcely ever curable. But Cancers of the Lips, *Palpebræ*, and *Mammæ* are extirpated with ſafety, and ſometimes admit of Cure; but there is great Danger of their returning: ſome believe a Cancer to be contagious, but I could never obſerve any Foundation for this Opinion, though I have been pretty converſant in theſe Caſes.

KORTHOLTUS's Remedy for a Cancer. VIII. In *Ephemerid. Breſlavienſ. Phyſico-Medicis*, which were ſometime ſince publiſhed in *High Dutch*, in ſeveral Volumes, and alſo *in Praxi Medicâ* NENTERI, you will find great Recommendations of a *Noſtrum* of KORTHOLTUS's which is corroſive and emetic. But I muſt tell you at the ſame time, that I

[a] *In Lib. de Gangræna. cap.* VII. [b] *Aphoriſm.* xxxviii. § 6. *quibus occulti Cancri ſiunt, eos non curare (ſive attingere) melius eſt. Curati enim cito pereunt, non curati vero longius tempus perdurant.* [c] *Lib.* V. *Cap.* XXVII. § 2. [d] See *Memoirs de Madame* DE MOTEVILLE, *Tom.* V.

am

am informed by Men of the greateſt Credit, that it is of no Efficacy in the true Cancer.

When a Cancer is incurable.

IX. When a Cancer yields to no Medicine; when it happens in old Age or to a bad Habit of Body; when it is ſituated under the *Axillæ*, or near large Blood-veſſels, or has ſpread itſelf to a great Extent, and is of long Standing; or where the Patient is afflicted with a Cancer in more Parts than one; in either of theſe Caſes the Knife is foreign to our Purpoſe: for as the diſeaſed Parts can never be entirely extirpated, the Surgeon, by attempting the Operation, will only make Matters deſperate, and haſten the Death of the Patient: therefore the beſt Method of treating an incurable Cancer, which is not yet broke or ulcerated, is, (1.) to endeavour, without uſing any violent Means, to prevent it from degenerating into an Ulcer: (2.) to relieve and aſſuage the moſt threatening Symptoms: in this Manner we may prolong the Proſpect of Death, and many other Miſchiefs, by a palliative Method.

Palliative Method.

X. If any one is deſirous of palliating this dreadful Caſe, he muſt look for Aſſiſtance, not from Medicine alone, but principally from a diligent Obſervation of Rules with regard to Diet; which we have already explained at large in diſcourſing of a *Schirrhus*. (*Chap.* XVI. § VI, &c.) The Patient ſhould loſe Blood in Spring and Autumn, but if of a plethoric Habit, oftner; and the Bowels ſhould be conſtantly kept open. It would not be amiſs alſo to adviſe the Uſe of *Goats Milk*, unleſs the Patient has a particular Averſion to it; you may give it either alone or boiled with vulnerary Herbs or Crayfiſh: by this Method you may very ſucceſsfully prevent very dangerous Symptoms: but if notwithſtanding this, violent Pains ſucceed, it will be proper to give him a Doſe of *Opium* now and then, or boil *Sem. Papav.* in his Drink; or you may make an Emulſion of them: theſe Medicines, by giving Sleep, are excellent Remedies againſt Pain and Weakneſs: the ſame Method is to be obſerved with regard to external Treatment, which we preſcribed in the above cited Place.

How an ulcerated Cancer is to be treated.

XI. Almoſt the ſame Method is to be obſerved in treating a Cancer that is broke or ulcerated; only in this Caſe the Part is to be kept clean, the *Sanies* frequently wiped off, and the Ulcer to be filled with ſoft dry Lint; or, in order to leſſen the Pain, the Part may be anointed before it is dreſſed, with ſuch Medicines as obtain moſt Credit for anſwering this Intention. The principal of theſe are, *Ol. Myrrhæ per deliquium, vel ejus Eſſentia cum Eſſentia Succini, vel Aqua Calcis ſola, aut pauxillo Sacchari Saturni admixto. Vel,*

℞. *Aceti Lithargyriſati* ℥j ß. *Olei Roſacei aut Solani* ℥j. *m. f. in mortario plumbeo aut vitreo Unguentum, quod Nutritum appellatur, Vel,*

℞. *Aq. Roſar. Flor. Sambuc. Papav. erratic. ana* ℥ij. *Sacch. Saturni, Eſſent. Opii ana* ℥j. *Spirit. Vini Theriacal.* ℥ij. *M. Vel,*

℞. *Aqua Sperm. Ranar. Solan. ana* ℥iij. *Plumb. uſt.* ℥j. *Sacchari Saturni* ℥ß. *M.*

In the Place of theſe you may ſubſtitute a vulnerary Decoction *ex Herb. Marrab. Agrimoniæ, Veronicæ*, &c. or *Succ. Solan. & Plantægin.* The Ulcer may be very eaſily waſhed with any of theſe at every Dreſſing, and the Lint may be wetted with them: but if the Pain ſhould be very violent, you may then in-

creaſe your Doſe of *Opium* or Eſſence of *Opium*, or you may moiſten the Pledgets with Eſſence of *Opium* at every Dreſſing; ſince it will be impoſſible to aſſuage the Pains with a leſs powerful Medicine: the Eſſence of *Opium* to be uſed in Caſe, is not to be prepared *cum Spiritu Vini*, but rather *ex Aquis deſtillatis, ex Solano, Floribus Papav. Erratic.* DIONYSIUS adviſes a raw Piece of Veal to be laid on the Part: dry Powder ſhould never be ſprinkled upon a Cancer, as it is cuſtomary on other Ulcers: the Dreſſing with *Plumbum uſtum cum Sem. Lini aut Pſyllii Mucilagine miſt.* mitigates the Pain in a ſurpriſing Manner: varying the Application in this Caſe is very uſeful, but we ſhould ſtick moſt to thoſe Remedies which ſeems to agree beſt with the Patient. Laſtly, the *Aqua Vulneraria ſive Sclopetaria*, commonly known by the Names of *l'Eau d'Arquebuſade* prepared with *Aqua Solani*, rather than with Wine or Spirits, being laid on warm and frequently repeated, is of eminent Service.

How a Cancer is to be extirpated.

XII. When the Cancer is ſo circumſtanced that you may venture upon Extirpation, without Danger of any conſiderable Miſchief; you are firſt to adminiſter mild cathartic Medicines to cool and correct the Acrimony of the Blood; (§ XI.) but more particularly to prepare the Patient by an exact Regimen with regard to Diet, before you attempt the Operation. The Inſtruments which are uſed in taking off Cancers of the Lips, Eyes, *Mammæ*, and Parts of Generation in the Male, you will find deſcribed below in their proper Place, when I come to treat profeſſedly of chirurgical Operations: the Wound is to be dreſſed in the ſame Manner which we have directed for treating other Wounds; with a digeſtive Ointment, and vulnerary Balſam: the Dreſſings ſhould be laid on lightly, and but ſeldom repeated, which will greatly conduce to the Cure: when the Wound is healed, the Patient ſhould obſerve a very ſtrict Regimen with regard to Diet through the remaining Part of his Life; he ſhould entirely abſtain from all acrimonious, ſalt, acid, or ſpiced Meats; he ſhould frequently take gentle cooling Purges, the beſt of which are the purging medicated Waters; not omitting to loſe Blood by Cupping or the Lancet whenever he perceives any Fulneſs, particularly at Spring and Fall: for if theſe Rules are neglected, the *Schirrhus* and Cancer eaſily return.

CHAP. XVIII.

Of an OEDEMA.

An *Oedema*, what.

I. HITHERTO we have been treating of Tumors that ariſe from Inflammation, and of the ill Conſequences that attend them: we proceed now to deſcribe that Sort of Tumor which is attended with Paleneſs, Cold, and yields little Reſiſtance, retaining the Print of your Finger when preſſed with it, and accompanied with little or no Pain. The Name proper to this Tumor is *Œdema*, or a *phlegmatic Tumor*: it obtains no certain Situation in any particular Part of the Body, ſince the Head, Eyelids, Hands, ſometimes Part of the Body, ſometimes the whole Body is afflicted with it. When the laſt mentioned is the Caſe, the Patient is ſaid to be troubled with a Cachexy, *Leucophlegmatia*, or Dropſy; but if any Part of the Body is more ſubject to this Diſorder than another, it is certainly the Feet; which are at that Time called

ſwelled or œdematous Feet: we ſhall treat diſtinctly of them in this Place, that it may appear what is the true Nature and rational Treatment of phlegmatic Tumors, in whatever Part of the Body they ſhall be found.

II. The proximate Cauſe of an *Œdema* is doubtleſs to be found in the too great Seroſity or Viſcidity of the Blood, which ſtagnates in the very minuteſt Veſicles of the Fat, or *Tunica Celluloſa*, and by this means ſtretches out the Skin with which it is immediately covered. This vitiated State of the Blood chiefly ariſes in Men, (1.) who are either of a cold and phlegmatic Habit of Body, or are advanced in Years: it chiefly falls upon them in cold Weather, or in the Winter, when the Inclemency of the Seaſon heightens the Diſorder of Nature: it is no wonder therefore that Perſons whoſe Legs ſwell greatly in the Day, frequently find themſelves much lighter and ſlenderer in thoſe Parts every Morning, which certainly proceeds from the Warmth they receive in Bed: (2.) another Cauſe of this Diſorder is, *an Irregularity in Diet*, by over eating or drinking, and by the conſtant Uſe of crude, cold, and hard Meats: (3.) *Intermitting Fevers or Agues* conduce very much to this Diſorder; eſpecially if the Patient indulges himſelf in an intemperate Uſe of cooling Liquors whilſt the hot Fit is upon him, and his Thirſt very urgent: (4.) This Diſeaſe frequently owes its Riſe to *too plentiful a Diſcharge of Blood* from a Wound, from the Noſe, or Lungs, by vomiting, or from the hæmorrhoidal Veſſels or *Uterus*: or, (5.) ſometimes to *Obſtructions of the menſtrual Diſcharge* in Women: or, (6.) to *a Compreſſion of the Vena Cava*, by the Weight of the *Fœtus* in Women far gone with Child, or by any ſchirrhous Body in the *Abdomen*; which greatly hinders the Return of the Blood from the lower Limbs: or, (7.) to *too ſedentary a Way of Life*, or to too great an Indulgence in lying in Bed or ſleeping: or laſtly, (8.) to a *Phthiſis and Difficulty of Breathing*; or to any other Diſorder or Fatigue of Body, which diſturbs or deſtroys the natural Force of the Heart in maintaining the Circulation with due Vigor. *Cauſes.*

III. From what has been delivered, it plainly appears by what Signs an *Œdema* manifeſts itſelf: therefore this Obſervation alone remains to be added; that the harder the Tumor is, and the longer the Pitting which is made by the Finger remains viſible, the ſtagnating Fluid is in ſuch Proportion thicker and more tenacious. *Diagnoſis.*

IV. Œdematous Tumors that come with other Diſeaſes, as a Dropſy, Conſumption, Aſthma, intermitting Fever, or with an Increaſe, or at the going off of the menſtrual Diſcharge, can ſeldom be cured but by curing the Diſtemper from whence they ariſe: œdematous Tumors of the Legs are of very little Conſequence in Women with Child; eſpecially if they are naturally of a good Habit of Body: for the Preſſure being taken off the *Vena Cava* by the Delivery of the Woman, the Tumor quickly diſappears in Conſequence: but weakly Women do not come off ſo well in this Caſe, particularly if the Tumors remain long after Delivery; for they are, in this Caſe, frequently the Forerunners of Dropſy, Aſthma, and Death: the more inveterate theſe phlegmatic Tumors are, by ſo much the more dangerous and doubtful are they to be eſteemed: on the other hand, thoſe that are recent and attended with no other Diſeaſes are very eaſily cured: thoſe that are Attendants on an intermitting Fever, are cured with much greater Eaſe, than thoſe which are the Conſequence of a large Profuſion of Blood, or of any other Weakneſs: Thoſe which ariſe from an *Prognoſis.*

Obſtruction

Obſtruction of any natural Diſcharge, are cured by the Return of that natural and cuſtomary Diſcharge of Blood: young Perſons are more readily cured of theſe Tumors in the lower Limbs than old; for indeed Perſons advanced in Years are generally incurable in this Caſe; when Tumors of the Legs and Feet are treated with improper Remedies, eſpecially externally, Aſthma and Death will, by Degrees, be the neceſſary Conſequences.

Method of Cure.

V. The Method of treating œdematous Tumors is ſurprizingly different, according to the different Cauſes to which they owe their Riſe: therefore we are firſt to make diligent Search after the genuine Cauſe of the Diſorder before we attempt its Cure: and as from the Nature of the Diſtemper, the internal Parts are to be ſet right, we muſt by no means put our whole Truſt in external Remedies; but are chiefly to expect Help from internal Medicines preſcribed by a *prudent Phyſician*[a]: the external Method of treating theſe Tumors in the Legs and Feet, is uſually, (1.) to have Recourſe to frequent *Frictions* with warm Cloths, to be repeated Morning and Evening till the Parts grow red and hot; (2.) then the Limbs are to be diligently preſerved from the Injuries of the cold Air; for which End he may wear Stockings made of ſome warm Furr; and at Night he ſhould keep hot Bricks about his Legs and Feet, to attenuate the Blood: (3.) after this you may apply a proper Bandage, which is to aſcend gradually from the Feet up to the Knees; this ſtrengthens the Limb, and prevents a Collection and a Stagnation of the Blood in any Part of it: (4.) after the Uſe of proper internal Medicines, and the external Methods which we have juſt mentioned, it will be very proper to uſe ſtrengthening Remedies externally: to this End you may place the Limb over burning rectified Spirits of Wine, wrapping it up in Cloths, in ſuch a Manner that it may receive the Steam; this will incline the ſtagnating Fluids to eſcape through the Skin, or render them fit to return into the Circulation, and at the ſame time reſtore the natural Tone to the Limb; (5.) many, eſpecially amongſt the common People, apply, as a family Medicine, the *Chelidonium majus*, firſt bruiſing it, and then laying it on as a Cataplaſm: others apply in the ſame Manner the *Perticaria acris*, either alone, or mixed with the forementioned Remedy, and from this Method they frequently find great Relief; for they are very active Medicines, and powerful Reſolvents: there are ſtill others again who uſe *Raphanum Ruſticanum Raſum*, or *Lepidium*, which they boil in Wine, and apply hot for the ſame End: but the moſt excellent Remedy to execute this Intention ſeems to be the Cataplaſm which is prepared *ex Columbarum Fimo, Sale atque Aceto inter ſe invicem commixtis, calide ſæpius impoſitum:* of the ſame Virtues are Fomentations made *ex Cineris Querni Lixivio parat. cum Aq. Fabri Ferrar. addendo Spirit. Vini Uncias aliquot, Aluminiſque Portiunculam*: this may be applied with Stuphs, or the Legs may be bathed in the Liquor as warm as it can be well born, twice every Day: *Aqua Calcis* is ſaid to be of equal Service, uſed in the ſame Manner either alone, or mixed *cum Spiritu Vini & Alumine*: the following Mixture alſo anſwers the ſame Intention.

℞. *Spirit. Vini, Aceti Vin. ana* ℔ j. *Alum. Crud.* ʒ j. ß. *Vitrol.* ʒ j. *M.*

This is to be applied as we directed above: but you muſt carefully take Notice, that after rubbing and fomenting, the Legs are to be well covered with Bandages

[a] See my *Practical Compendium, Chap.* II. *N.* 14.

and

and Stockings: the Patient ſhould drink ſparingly, uſe moderate Exerciſe frequently, and be very diligent in the Uſe of proper internal Remedies. Sometimes the medicinal Waters, particularly thoſe of the ſulphureous Kind, are found very ſerviceable in this Caſe, but not always: GARENGEOT adviſes ſcarifying the Feet, firſt in the inward and middle Part of the *Tibia*; and when thoſe Wounds are healed, afterwards to repeat it on the oppoſite Side; and to apply to them the *Emplaſtrum Norimbergenſe*, as a moſt excellent Medicine; cap. *De Paracenteſi. Abdominis.* HARRIS, a celebrated *Engliſh* Phyſician, in *Diſſert. Chirurg.* IX. relates, that he has cured the moſt dangerous of theſe Caſes, *cum Croco Martis aperitivo, Cortice Peruviano miſt.* Others affirm, they have done it with the *Cortex* alone; others again are confident, that this is a hurtful and dangerous Method: the beſt Way is, to conſult ſome prudent Phyſician, who beſt knows how to adviſe you what Methods to purſue, and what to avoid.

CHAP. XIX.

Of [a] FUNGOUS TUMORS, *and* DROPSY *of the* JOINTS.

I. *OEDEMATA* are nearly reſembled by fungous Tumors of the Joints: theſe are Diſorders of very bad Conſequence, and therefore deſerve a particular Diſquiſition: that they have been entirely omitted or ſlightly paſſed over by many chirurgical Writers, ſeems to proceed from their Ignorance of the true Cauſe from which they ariſe; for whether they owe their Origin to a Collection of Blood or ſerous Fluids, corrupted Matter, *Pus*, *Flatus*, or to any other Cauſe; they could not pretend to diſtinguiſh. When we ſpeak of a *Fungous Tumor of the Joints*, we mean that Tumor of the Limb which ariſes at the Joint, looks pale, is void of Heat and Pain, eaſily yeilds to the Preſſure of the Fingers, but riſes again inſtantly, like a *Fungus*, upon removing the Finger, leaving no Pit behind. Though no Joint either of the upper or lower Limbs can be ſaid to be ſecure from this Diſorder, yet the *Knees* are moſt ſubject to it, as they are particularly ſubject to Injuries from Falls, or other Accidents, and becauſe they abound in a large Quantity of fat and glandular Bodies, which are concealed amongſt the Ligaments and Tendons. There are ſeveral Species of this Tumor; for ſome are ſmaller, ſome larger, ſome ſofter, ſome harder, ſome more, ſome leſs glutinous with regard to the State of the inſpiſſated Fluid; again, ſome are attended with Pain, ſome entirely free from Senſation: [b] in ſome the noxious Humours are ſituated without the Joint, which Kind of Tumors are properly the fungous Bodies we are now treating of; but in others they are collected and retained in the Joint itſelf, as the *Serum* is contained in the Teſticle in an *Hydrocele*, many of which I have ſeen and cured: this laſt-mentioned Diſorder may not improperly be called a *Dropſy of the Joint*, and may properly be diſtinguiſhed from the fungous Tumor of the Joint, by the Inlargement that ap-

A fungous Tumor, what.

[a] In *England* they are known to us by the Name of White Swellings, or ſcrophulous Tumors of the Joints.

[b] PURMANNUS, in his *Chirurgia Curioſa*, has given us a Deſcription of a very large *Fungus*.

pears all round the Joint; whereas the fungous Tumor is ſituated more on one or the other Side of it: from what has been already ſaid of theſe two Caſes, I think it plainly appears, that it is no difficult Matter to diſtinguiſh one from the other.

Cauſes. II. The proximate Cauſe of fungous Tumors is, without Doubt, the viſcid glutinous *Serum*, which is found about the Ligaments of the Joints, and is apt to ſtagnate after the Ligaments have received any conſiderable Violence from a Fall or Blow: ſometimes the Tumor riſes in the external Parts, ſometimes in the Articulation itſelf, by which the Ligaments being weakened, the Part loſes its natural Motion; but when the Nerves or Blood-veſſels are greatly preſſed upon by the Tumor, the Parts below are uſually deprived of Nouriſhment, and the Joints by Degrees being greatly inlarged, the neighbouring Parts diminiſh and waſte.

Prognoſis. III. We have already obſerved, that in fungous Tumors of the Joints, the Ligaments are too much lengthened and relaxed, and the natural Strength and Motion of the Limb are leſſened in Proportion to the Degree of the Diſorder; and as the loſt Vigor of the Part is very difficult to be reſtored, and the Tumor will not readily yield either to Suppurants or diſperſing Remedies, any one will be ſenſible that the Surgeon has no eaſy Taſk upon his Hands, when he undertakes the Cure of a fungous Tumor upon the Joint; the Suppuration of the Part is not only difficult to bring about, but it is generally a very dangerous Attempt; for by this Means *Caries* and incurable *Fiſtulæ* are ſometimes produced, which require Amputation of the Part. When the Tumor is recent, and not very large or hard, it ſometimes admits of Cure by the Application of digeſtive and corroborating Remedies; whereas they will be greatly irritated by emollient Applications: but where the Tumor is large and inveterate, no Succeſs is to be expected from any thing but the Knife; and even that is ſometimes unequal to the Cure, or improper; if the noxious Fluids are contained in the Joint, they may be let out by Inciſion; but, upon healing the Wound, the Tumor will generally return.

Cure of recent fungous Tumors by Diſperſion. IV. In order to hinder the Cure of recent and mild fungous Tumors the eaſier by diſperſing Remedies, it will be beſt to rub the diſordered Part well every Day with warm Cloths, fomenting it afterwards with *Spirit. Vini tartariſat.* this Method is to be conſtantly obſerved, till the natural Strength and Form of the Limb are reſtored: PURMANNUS's Fomentation is excellently calculated for this Purpoſe:

℞. *Muriæ Halecum* ℔ ij. *Acet. Vini fortiſſim.* ℔ j. *Fol. Salv. M.* ij. *Vitriol. Rom.* ℥ j. ß. *Alumin. Crud.* ℥ vi. *M.*

Theſe Ingredients are to boil together for half an Hour, and to be uſed in the Manner we have above deſcribed. When the Tumors begin to diſperſe, and the Parts to recover their Strength, it will be very beneficial in perfecting the Cure, to foment the Limb well ſeveral Times every Day *cum Spiritu Vini Tartariſati, vel cum Oleo Tartari fœtido*; laying on the Bandages immediately afterwards, to keep the Part warm, and defend it from the Injuries of the cold Air, of which it is very ſuſceptible: laſtly, I cannot help adding a Form under this Head, by the Aſſiſtance of which I have frequently made very happy Cures of fungous Tumors.

℞ *Lithar-*

℞. *Lithargyr.* lib. ß. *Boli Armen.* ℥j. *Maſtichis, Myrrhæ ana* ℥ ß. *Aceti Vini* lib. j. *m. & coque hæc omnia per horæ quadrantem, tinctiſque in iſto decocto linamentis craſſis calida ſemper & matutino & veſpertino tempore in lectulo applicentur.*

At the ſame time proper purging Medicines, Attenuants, and Sudorifics ſhould be diligently attended to.

Cure of inveterate Fungus's (1.) by the Knife.

V. If the fungous Tumor is of long ſtanding, and will not give way to the diſperſing Remedies which have been preſcribed, almoſt the only Hope left is, to make an Inciſion into the dependent and moſt convenient Part of the diſeaſed Joint, taking great Care to avoid wounding the Ligaments or Tendons: you are well juſtified in following this Method, by the Examples of thoſe two celebrated Surgeons[a] WURTZIUS and PURMANNUS: by this means the ſtagnated *Serum* is inſtantly evacuated, if it is contained in one Cavity; but if it is contained in different Cells, it will all eſcape in a few Days. Tents daubed with ſome digeſtive Ointment, and ſprinkled with Allum, are ſerviceable in this Caſe. Before you make your Inciſion, you ſhould pull the Tumor down as low as you can with your Fingers, and make a tight Bandage above to retain it in this Situation; by this means the moſt convenient Part for the Inciſion to be made in, will lie fair; and when the Opening is made, the *Serum* will readily burſt out like Blood at the opening a Vein, or Lymph in tapping for the *Hydrocele* or *Aſcites*. When this is done, if any Tumor ſtill remains, dreſs the Part with *Emplaſtrum Diachylum vel Oxycroceum, vel* WURTZII *Rubrum, vel Aqua Calcis, vel Spirit. Vini*; by continuing any of theſe Applications, what remains inſpiſſated in the Tumor will entirely diſperſe. When the Limb is reſtored to its natural Shape, heal the Wound with vulnerary Balſams, diligently avoiding the Uſe of fatty or oily Medicines, as being very hurtful to the Tendons and Ligaments, with which thoſe Parts abound: if the *Serum* contained in the Tumor is ſo glutinous, that it cannot diſcharge itſelf for Want of Fluidity, you muſt throw up attenuating Injections at every Dreſſing: the beſt calculated for this Purpoſe are thoſe which are prepared *ex Decocto Agrimoniæ, Ariſtolochiæ, aut Alchymillæ cum Roſarum aut Chelidonii Melle miſto*: Injections of this Kind will quickly diſſolve the ſtagnating *Serum*, and diſperſe the Tumor.

(2.) By Corroſives.

VI. Though thoſe fungous Tumors, which are opened with the Knife, are more readily diſcharged and healed, yet ſome Surgeons prefer the Application of cauſtic Medecines to the Knife, diſcharging the collected *Serum* upon the falling off of the Eſchar; after which they proceed in the ſame Manner which we adviſed above: whilſt the Part is healing, in either Caſe, I think, it would be very proper to warm and invigorate the Ligaments and Tendons, eſpecially when the Injury falls upon the Knee, by the Uſe of ſome nervous Ointment, or aromatic Spirit.

Remedies to prevent the Return of the Tumor.

VII. It very frequently happens, that after you have evacuated the inſpiſſated *Serum*, and cicatriſed the Wound, you ſhall have a freſh Collection of a vitiated and corrupt Fluid, which I am an experienced Witneſs of; to prevent this Accident, the following Method will be ſerviceable: let the Patient continue in a ſtrict Courſe of proper purging, ſudorific and attenuating Medicines, and

[a] *Chirurg.* p. 268. [b] *Chirurg. P.* III. p. 46. *it. Chirurg. Curieſ.* p. 622.

keep the Wound open with Tents for a conſiderable Time, cleanſing it every Day by throwing up an Injection prepared in the Manner we directed in the preceding Section: PURMANNUS highly commends this Manner of keeping the Wound clean, and atteſts that, after the ſixth Time of Injecting, he has not only ſeen the Wound clean, but filled up with new Fleſh: it will be proper alſo to inject *Aqua Calcis vivæ*, or *vulneraria Gallorum* ſometimes; and to cover the external Part with a warm Plaſter, or to foment it with ſome Liquor of the ſame Intention, eſpecially in the Knees: this Method is recommended by that experienced Surgeon FELIX WURTZIUS, as the moſt likely Means of preventing the Return of the Diſorder.

When an Opening is not to be made.

VIII. Before I leave this Head I muſt inform you, that *it is not every fungous Tumor of the Joints which is ſo ſituated, that it can be opened with Safety*; for if the Tumor is of very long ſtanding, hard, of a great Size, or the Patient is of an infirm weakly Habit of Body, you muſt entirely lay aſide the Knife; for this Method of Treatment would produce more Miſchief than Good, by laying a Foundation for new Diſorders; to wit, *Caries*, *Fiſtulæ*, and *Gangrene:* as to the other Species of lymphatic or phlegmatic Tumors, which require the Knife, ſuch as *Dropſy*, *Hydrocele*, *Hydrocephalus*, and *Ranula*, I ſhall treat more fully of them in their proper Place, when I come to deſcribe Chirurgical Operations; in the mean time, with regard to Swellings in the Knee, conſult SCULTETUS's *Obſervat.* 79. WURTZIUS's *Surgery*, p. 268. and MEEKREN and ROONHUYSEN, *Obſerv. Chirurg.*

INSTI-

INSTITUTIONS OF SURGERY.

PART I. BOOK V.

Of ULCERS.

CHAP. I.

Of ULCERS.

I. AN Ulcer is a Diſorder ſo well known to every one, that, when I have mentioned the Name, it would be impertinent to illuſtrate it with a Deſcription; a Definition in this Caſe would only ſerve to make the Matter more obſcure: you have a very juſt and clear Notion of an Ulcer, when you are told, that it is a *Solution of the ſoft Parts of the Body and the Skin*, ariſing from an internal Cauſe, *ſc.* an Inflammation, Abceſs, or ſharp Humours: Wounds alſo and Contuſions, by Length of Time, degenerate into Ulcers, and properly aſſume that Name. An Ulcer what.

II. The proper Seat therefore of an Ulcer is, any ſoft Part of the Body, *ſc.* the Skin, Fat, Glands, muſcular Parts, and even the *Viſcera:* if any of the hard Parts of the Body, that is, of the Bones, are ulcerated or corroded, the Diſorder is rather called a *Caries* or *Spina Ventoſa*, than an Ulcer: but, from the Similitude there is between both Caſes, I think they may very properly be treated of under the ſame Head, and I have therefore joined them together. Its Situation.

III. If you deſire to know how Ulcers differ from *Abceſſes*, *Contuſions*, and *Wounds*, a diligent Examination into the Nature of each will give you full Satisfaction upon this Head: though *Wounds* and *Contuſions*, as well as *Ulcers*, conſiſt in a *Solution of Continuity of the ſoft Parts of the Body*, yet they widely differ in this Circumſtance; to wit, Wounds and Contuſions always ariſe from an external Cauſe, and are produced in a Moment; whereas Ulcers owe their Riſe chiefly to internal Cauſes, § I. and come on by ſlow Degrees: *Abceſſes* are as it were the firſt Beginnings of Ulcers, or rather are immature Ulcers; which is the Caſe when Inflammations come to Suppuration, the Skin ſtill remaining whole: but, as ſoon as an Opening is formed in the Skin, and the maturated *Pus* diſcharges itſelf, from Abceſſes they become Ulcers, whether the Skin is eroded by the *Pus*, or the Opening made by the Surgeon's Inſtrument. Difference from other Diſorders.

Various Kinds of Ulcers.

IV. Ulcers cannot be confined to one Species, for they differ, (1.) in the Part of the Body which they infeſt; for ſometimes they are found in the Skin, Fat, and Glands, ſometimes in the muſcular Parts: (2.) in their Size; for ſome are ſpread wide, others occupy but a ſmall Space; ſome are deep, others ſhallow; thoſe which penetrate deep, and are narrow, particularly if they are very ſmall at their Opening, are called *Sinuſes* or *Fiſtulæ:* (3.) in their Standing; whence they are called *recent* or *inveterate:* (4.) in Number and Degree of Symptoms, or accidental Diſorders that attend them; ſome are very mild, and are thence called *benign:* others are *malignant*, that is, either *attended with very acute Pains*, or *fetid, putrid, fatty, diſcharging great Quantities of Ichor, ſpreading wide, cancerous, callous, fiſtulous*, or *attended with Worms:* (5.) in their Cauſes; hence Ulcers are called *ſcorbutical, venereal, cancerous, peſtilential*, or are ſaid to be occaſioned by *Faſcination:* laſtly, (6.) Ulcers differ in their Situation, and are called *Ulcers of the Noſe, Fauces, Breaſt, Anus*, and *Fiſtulæ lacrymales*, as they attack this or that Part.

Cauſes.

V. I think thoſe Phyſicians amongſt the Moderns draw too haſty a Concluſion, who aſſert, that the *principal Foundation of Ulcers* is owing to a *foreign acid Humour*, which corrodes and deſtroys the Parts of the Body which it falls upon, in the ſame Manner that *Aqua fortis* would; ſince there is no acrimonious Humour, whether it is of a ſalt, lixivious, alcaline, or acid Nature, but would corrode the Body, and raiſe an Ulcer of ſome Kind: and, to ſay Truth, the ſtagnating Blood generally degenerates into an Acrimony of the alcaline Kind, and is by no means, according to the Opinion of ſome, converted into an Acid; this you may collect from the fetid Smell of Ulcers. The Phyſicians have pronounced an *Alcali* to be any Saltneſs or Acrimony, which is adverſe to all Kinds of Acids, as Salt of Tartar is to Vinegar, Oil of Tartar *per deliquium* to Spirit of Vitriol: as there are many Kinds of Poiſons, ſo of acrimonious Things, and therefore of Ulcers: the more Virulence the Acrimony is poſſeſſed of, by which the Body is corroded, ſo much the worſe will be the Conſequence of ſuch Corroſion; the Ulcers will be the more fetid, the more dangerous, and perhaps incurable, which is the Caſe in Cancers: but Ulcers do not ariſe from Acrimony alone; but from any other Cauſe by which the Blood may be made to ſtagnate and corrupt: upon this Principle you frequently ſee *Tumors, Inflammations, Wounds, Contuſions, Fractures, Luxations, Schirrhus, Cancer*, and *Caries* degenerate into Ulcers; which though they begin with very ſlight Symptoms, yet, either from a bad Habit of Body, Irregularity in Diet, or Ignorance in the Surgeon, they very often become extremely dangerous.

Diagnoſis of inveterate Ulcers.

VII. Although moſt Ulcers may be diſcovered by the Sight; yet, in order to have a thorough Knowledge into the Depth and Tendency of the *Sinus*, and whether it is accompanied with a *Caries* of the ſubjacent Bone, you muſt have Recourſe to the Uſe of the Probe: you will learn beſt from the Patient whether it be recent or of long ſtanding: from him alſo you may collect the Cauſe of the Inveteracy of the Diſorder; whether it is owing to a ſubjacent *Caries*, to an irregular Courſe of Life, or to the unſkilful Treatment of the Surgeon. The Ulcer is ſaid to be *benign*, if it is recent, and attended with no violent Symptoms; if the *Pus* is of a moderate Conſiſtence, whitiſh, without Acrimony, and of no bad Smell; and laſtly, if the Patient is otherwiſe healthy, and of a good Conſtitution: on the contrary, it is called *malign*, if the Patient is of a weakly

ſcorbutical, or hydropical Habit of Body; if the *Pus* is too fluid, acrimonious, fetid, yellow, brown, green, or blackiſh, or of the Conſiſtence of Lard: the Diſorder is equally dangerous, where the Patient ſuffers very intenſe Pain, or where the Ulcer is ſo formed that it cannot admit of being treated like Wounds and recent Abceſſes, with Digeſtives and vulnerary Balſams.

Nature of putrid, running, corroding, fiſtulous and callous Ulcers.

VII. *Ulcers* are ſaid to be *unclean and putrid*, in which the Fleſh appears corrupted, ſoft, white, yellow, or livid; where the Matter is thin and glutinous, and at the ſame time green or variegated: they are called *running* or *rheumatic Ulcers*, when there is a very plentiful Diſcharge of a thin *Sanies*: we term them *corroding and ſpreading Ulcers*, if the Matter is corroſive enough to deſtroy the adjacent Parts, ſometimes ſlower, ſometimes faſter, in Proportion to the Degree of Acrimony of which it is poſſeſſed. *Fiſtulous Ulcers* are thoſe which penetrate deep, under the Skin, or between the Muſcles, eſpecially if the *Sinus* is wide, and the Opening very narrow. In *callous Ulcers* the internal Parts are lined with a hard and almoſt cartilaginous Subſtance.

Nature of venereal, cancerous, carious Ulcers, and of thoſe which are occaſioned by Charms.

VIII. *Ulcers* are termed *venereal*, when they are the Conſequence of Familiarity with an infected Perſon, and either accompany or ſucceed other venereal Diſorders: they are confined to no particular Part; but more frequently ariſe in thoſe Parts which are the Seats of venereal Buboes, or in the Noſe or Throat, ſometimes alſo upon the *Penis*; Ulcers of this laſt-mentioned Part are called by the *French*, *Chancres:* in the other Sex the *Labia Pudendi*, or Neck of the Womb, are chiefly obnoxious to this Symptom of the Pox. *Cancerous Ulcers* are either Cancers themſelves burſt out, the Sign of which we have given you above, in *Book* IV. *Chap.* XVII. § V. VI. or very nearly approach the Nature of *Carcinomata*, if you regard the Degree of Pain with which they are affected, or the Quickneſs of their Increaſe. *Ulcers* are called *carious*, when any neighbouring Bone is deprived of its *Perioſtæum*, or affected with a *Caries*; but we ſhall treat more fully of this Caſe below: *Ulcers* are, by the Vulgar, believed to ariſe from *Faſcination*, when Needles, Hairs, Threads, Rags, Egg-ſhells, Coals, or any preternatural Body of this Kind is found in an Abceſs or Wound: but, in good Truth, it is my Opinion, that not only the Sign by which the common People pretend to diſcover Charms, but even Faſcination itſelf, is an Impoſition which can be ſwallowed by none but Perſons loaded with Superſtition; for many Ulcers have been ſaid to be owing to Faſcination and Witchcraft, which have evidently proceeded from natural Cauſes.

Prognoſis of malignant inveterate Ulcers.

IX. Recent and benign Ulcers, like recent Abceſſes, are generally attended with no great Difficulty in the Cure; eſpecially if they happen to young and robuſt Subjects: the Difficulty of the Cure will ariſe in Proportion to the Malignity of the Symptoms and the Inveteracy of the Diſorder; therefore *putrid*, *running*, *fiſtulous*, *callous*, *carious*, and *cancerous Ulcers*, require great Skill and Addreſs in the Cure: thoſe quackiſh Perſons who boaſt of a ſecret Plaſter, or Ointment, for the Cure of Ulcers of ever ſo great Inveteracy, or attended with the worſt of Symptoms, egregiouſly impoſe upon themſelves and their credulous Patients. If the Patient is weak and infirm, advanced in Years, has great Acrimony in his Blood; if the Ulcer has a very offenſive Smell; if the *Pus* is of a bad Colour and full of Acrimony; any of theſe Circumſtances will render the Cure of the Ulcer very difficult: if there are many Ulcers, or if an Ulcer ſpreads very wide, the Diſcharge will be very plentiful, and reduce the Patient wonderfully. It is never good Practice to heal old Ulcers of the Legs, eſpecially

eſpecially in weak Habits, or in Perſons advanced in Years; for Experience teaches us, that they are always in the beſt State of Health whilſt this Drain is kept open in their Legs: but if you heal the Ulcer, and ſtop up the Diſcharge, the worſt of Diſorders follow, to wit, Pains in the Head, Vertigines, Apoplexy, Epilepſy, Difficulty of Breathing or Aſthma, Diarrhœa, Dyſentery, and Inflammations on the internal Parts, and many other Diſorders of this Kind, till Death brings up the Rear: that excellent Phyſician CRATO deſerves to be conſulted upon this Head, in his *Epiſtolæ Medicæ*, where he treats this Point very judiciouſly. Where inveterate Ulcers dry up upon old Subjects, and the Lips grow hot and livid, there is immediate Danger of *Sphacelus* and Death itſelf: the Cure of inveterate Ulcers is much eaſier in young and robuſt Subjects; but you will always find it a uſeful and indeed neceſſary Obſervation to you, that in Ulcers of this Kind, you are not only to remove the immediate Cauſe of the Diſorder, but you are alſo to reſtore the Blood to its priſtine Purity; and in doing this you will uſually meet with great Difficulty: therefore, if the Diſorder is very inveterate, and the Patients are tired with the continual Uſe of Medicines, and with the ſtrict Regimen to which they are enjoined, it is no Wonder if theſe Ulcers often fail of a Cure, even in robuſt Perſons.

Prognoſis of venereal, fiſtulous, callous, carious, and cancerous Ulcers.

X. *Venereal Ulcers* cannot be cured till you have thrown the venereal Poiſon out of the Body by proper Remedies; till you have done this, external Remedies are to no purpoſe. *Fiſtulous*, *callous*, and *carious Ulcers*, are never cured without the Knife; for if you heal the Ulcer and bring on a Cicatrix, it will burſt out again, and afflict the Patient with greater Vehemence. A Caries, eſpecially if it is large, and ſituated in the Joint, will bring on ſo large a Diſcharge of Matter, that if the Limb is not taken off in Time, the Patient will be entirely run down; this will appear very plain to you when you read what will follow in its Place, on the *Caries* and *Spina Ventoſa:* the ſame may be ſaid of cancerous Ulcers; for if the Part affected is not taken off, there remain no hopes of Cure, as we declared above, treating of a *Cancer*; but even after taking off the Part, Cancers frequently return, and entirely deſtroy the Patient. When Ulcers fall upon the *Viſcera*, they are generally deemed incurable, becauſe out of Reach, both of the Hand and of immediate medical Applications.

Cure of recent Ulcers.

XI. The Method of treating Ulcers differs greatly according to the different Nature of the Diſorder: when the Ulcer is quite recent, it may be treated as a recent Abceſs or Wound: firſt, therefore, it is to be cleanſed, then to be filled with new Fleſh, and laſtly to be covered with an even Cicatrix.

(1.) How the Ulcer is to be cleanſed.

XII. The Ulcer, if recent, is to be cleanſed in the following Manner: firſt, the Matter is to be diſcharged; if it does not flow freely enough of itſelf, you muſt preſs it gently with your Fingers: if there is a deep *Sinus*, you may cleanſe it with an Injection; or, if it lays fair enough, with Lint: any ſmall Membrane, or fatty Body, that remains corrupting at the Bottom of the Ulcer, will readily enough caſt off afterwards, at the Removal of every Dreſſing, which ſhould be a digeſtive Ointment ſpread upon Lint, and ſecured upon the Part with *Diachylon*, *Diapalma*, or any other Plaſter of that Kind, covering the whole with proper Compreſſes and Bandages: this Method is to be continued till the Ulcer appears to be entirely cleanſed, that is, till the Fundus of it becomes florid, and it appears to be filled with new Fleſh.

XIII. Having proceeded thus far, the next Intention is to fill the Ulcer with new Fleſh; this Intention is ſatisfied with thoſe Medicines which are vulgarly called *Sarcotics:* the beſt of this ſort appears to me to be the common *Unguentum Digeſtivum*; for where there is no remarkable Impediment, I have never found it neceſſary to uſe any other ſarcotic Remedy than this: it is not eaſy to ſay what ſhould induce almoſt all Phyſicians to cry up certain balſamic Remedies, as having a peculiar Virtue in generating new Fleſh: beſides, our Digeſtive is endued with a true balſamic Power: but, to ſay the Truth, the Generation of new Fleſh is not ſo much owing to the Uſe of any particular Medicines, as to the Benefit of Nature; the whole Buſineſs of the Surgeon in this Caſe is, only to remove any thing that may impede the Cure: if any ſhall think that the *Unguentum Digeſtivum* is not equal to this Intention, they may have my free Conſent to ſubſtitute in its room *Balſamum Arcæi*, *Balſamum Peruvianum*, *Balſamum de Mecchâ*, *Balſamum Sulphuris*, *Eſſentia Myrrhæ & Aloë*, *Oleum Myrrhæ per Deliquium*, *Oleum Ovorum*, or any vulnerary Balſam of this Kind, till the Wound is entirely healed. (2) How the Ulcer is to be filled up.

XIV. If the Ulcer penetrates very deep, ſo that you can neither reach the Bottom of it with your Eye, nor apply your Medicine to it, it will then be proper, at every Dreſſing, as ſoon as you have preſſed the Matter out of it, to inject ſome cleanſing healing Liquor to it; ſuch as *Decoctum Agrimoniæ vel Ariſtolochiæ cum Melle Roſarum, vel & Myrrhæ atque Aloes Eſſentia miſtum*; or that which Belloste cries up, in his *Hoſpital Surgeon*, *Decoctum ex Nucum Foliis cum admixto Saccharo:* this Method of injecting is to be continued till the Bottom is entirely healed: afterwards you may proceed to fill up the Ulcer in the Manner we adviſed above. What is to be done in deep Ulcers.

XV. The Ulcer being filled up with new Fleſh, it remains that we bring on a proper Cicatrix; this is beſt done by dreſſing the Part daily with dry Lint, till the Cicatrix is formed: but if, notwithſtanding this Method of dreſſing, the Fleſh becomes luxuriant, and the Ulcer is moiſt, it muſt be ſprinkled with drying Powders, *ex Maſtiche*, *Thure*, *Sarcocollâ*, *Colophonia*, *Lapide Calaminari ac Tutiâ*, covering it with dry Lint and ſecuring all with ſome Plaſter, till it is entirely healed: but, if the luxuriant Fleſh has grown above the Skin, the beſt Way is, to eat it down with *Vitriolum cæruleum*; or, if that is not ſtrong enough, you may uſe the *Pulvis ex Præcipitato rubro atque Alumine uſto* till it becomes quite even: then you may proceed as directed above. (3) How the Cicatrix is to be formed.

XVI. Laſtly, it is ſcarce poſſible to ſay what great Relief the Patient will receive from obſerving a proper Regimen with regard to his Diet: Practitioners in Surgery have in all Times obſerved, that Ulcers of the moſt malign Kind have been ſometimes cured by this means, almoſt without the Aſſiſtance of any other Remedy; whilſt, on the other hand, the moſt benign Ulcers have ſo far degenerated, as to become altogether incurable by an irregular way of living: in this Caſe, therefore, the Patient ſhould moſt diligently avoid all acrimonious, ſalt, acid, fatty, or heating Meats, or thoſe that are hard of Digeſtion; according to the Directions which we gave you above when we were treating of *Wounds* (*Book* I. *Chap.* I. § XLV, and the following): if your Patient already labours under a bad Habit of Body, which obſtructs the Cure, it is your Duty to call for the Aſſiſtance of ſome ſkilful Phyſician, that may take off the conſtitutional Complaint, by preſcribing proper internal Medicines. A proper Regimen to be obſerved.

CHAP.

CHAP. II.

Of the Method of treating [a] FISTULOUS CASES.

The Cure of *Fiſtulæ* performed, (1.) by cleanſing.

I. WHEN you diſcover, either by your Eye or the Probe, that Ulcers are attended with *Fiſtulæ* [b] not yet become callous, your readieſt Way of curing them is, to lay them open with the Knife to the Bottom, if you can do it with Safety, and afterwards cleanſe and heal them: but, ſince Patients are very unwilling to conſent to the Uſe of the Knife, you may cleanſe them with a proper Injection, or dreſs them with *Ung. Digeſtivum* upon Lint, as we adviſed in the foregoing Chapter. Many Surgeons are for conveying their Medicine to the Bottom, by the Aſſiſtance of *Tents*; but, as they are very apt to do Miſchief by their Hardneſs, or too great Length, bringing on a *Callus*, Inflammation, or too great Flux of Humours upon the Part; therefore I think it moſt adviſeable either to throw them entirely aſide, or at leaſt to guard as ſtrongly as poſſible againſt any of theſe Inconveniences, by making them very ſoft, and as ſhort as the Caſe will admit of. BELLOSTE, and MAGATUS before him, both Men of great Name in Surgery, have been ſo offended at the miſchievous Abuſe of Tents, that they have abſolutely forbid the Uſe of them; and I am ſo far from diſagreeing with theſe Authors, that I readily join with them in Opinion. I think the Uſe of Tents is never to be juſtified, but where the Opening of your *Fiſtula* is ſo ſmall, that you are in conſtant Fear of its healing; and even in this Caſe your Tents can ſcarcely be too ſhort, and ſhould be made of the ſofteſt Materials.

(2.) By Preſſure.

II. The next Thing to be obſerved in treating *Fiſtulæ* is, to preſs the *Fundus* as near to the Opening as poſſible. When the Ulcer is cleanſed, and the proper Dreſſings applied, you muſt clap a ſmall Compreſs, or a Slip of Plaſter doubled up in the Form of a ſmall Compreſs, upon the Part where you judge the *Fundus* of the *Fiſtula* to be ſeated; ſecuring all with a Bolſter, Plaſter, and Bandage, as uſual: in rolling up the beſt Method will be to place the Beginning of the Roller upon the *Fundus* of the *Fiſtula*; or at leaſt to make your Faſtening tight upon that Part: this will direct the contained Matter towards the Opening, and the Bottom will heal before the Reſt of the *Sinus:* this happens beſt in *Fiſtulæ* of the upper or lower Extremities; eſpecially if the *Fundus* is in the upper Part of the Limb, and the Opening in the lower Part.

III. When *Fiſtulæ* penetrate ſo deep that you cannot come at the Bottom of them with your Dreſſings, you muſt inject ſuch ſort of Liquors as we adviſed in the foregoing Chapter: you may alſo very properly add the following:

℞. *Ung. Digeſtiv. ex Terebinth. & Vitell. Ovi parat:* ℥j ß. *Mell. vulgar. vel Roſar. vel Chelidon.* ℥j. *Spirit. Vini vulgar.* ℥ix. *M. Vel,*

℞. *Decoct. Scord. vel Abrotan. vel Agrimon.* ℥viiij. *Spirit. Vini Simpl.* ℥iiij. *Elixir Proprietat. vel Eſſent. Myrrh. & Aloës* ℥j. *Mell. Roſar.* ℥ij. *M.*

Theſe are to be injected at every Dreſſing, and the Opening of the *Fiſtula* ſhould be kept cloſe, that the Medicine may be retained as long as poſſible, which will

[a] See FABRIC. AB AQUAPENDENTE, MARCHETTI, and a Treatiſe by ASTRUC, who treat fully and judiciouſly on *Fiſtulæ* of the *Anus*.

[b] In *England* we call this Caſe a *Sinus*; never a *Fiſtula*, till it becomes callous.

haſten

haſten the Agglutination of the Part; afterwards you are to proceed as we directed above, treating of *Ulcers*, *Chap.* I. § XIII. and the following.

IV. If the Method of Cure, which we have hitherto deſcribed, is unequal to the Intention of cleanſing and healing, you will find greater Aſſiſtance from the Knife than from any other Remedy; and that chiefly where the *Fiſtula* tends downwards, or takes a very irregular Courſe, ſo that the *Fundus* of it cannot be preſſed towards the Opening; in this Caſe, you muſt lay open to the Bottom. Seldom cured without the Knife.

V. You ſhould gently paſs a grooved Probe or Director down the *Fiſtula*, and directing your Knife down the Groove, lay open the Fleſh and common Integuments as far as you think ſafe and neceſſary: all the Sinuſes of the *Fiſtula* being laid open, a free Paſſage is made for a Diſcharge of the corrupted Matter, and you can come at the diſeaſed Parts with your Remedies: this Operation may be performed without the Uſe of a Director, if your Knife has a Button at the Point. (See *Plate* V. *Fig.* 4 and 5.) Some divide the Fleſh with a *ſtrong Pair of crooked Sciſſars*. (*Plate* I. *Fig* D.) But this Method of cutting is attended with far greater Pain and Inconveniency than the other, except the Skin and Fleſh are exceeding thin. Inciſion how to be made.

VI. If the Operation is ſucceeded by a large Diſcharge of Blood, which frequently happens; at the firſt Dreſſing you muſt fill up the Wound with dry Lint; afterwards you may dreſs with *Unguentum digeſtivum cum Ægyptiaco*, *vel Præcipitato Mercurii rubro*, till the Wound is cleanſed: every thing elſe is to be done as we adviſed above, treating of recent Ulcers. The Method of treating *Callus*, *Caries*, and thoſe ſort of Diſorders which attend *Fiſtulæ*, ſhall be delivered ſeparately below. CELSUS *Book* VII. *Chap.* IV. upon *Fiſtulæ* in general, and particularly on the *Fiſtulæ Coſtarum*, *Ventris*, *& Ani*, deſerves a diligent Peruſal. Conſult likewiſe SCULTETUS's *Armament. Chirurg.* and MUYS's *Podalir. Rediviv.* p. 41, 42. What to be done after the Operation.

CHAP. III.

Of the Method of treating STUBBORN ULCERS.

I. HITHERTO we have treated of mild and well-conditioned Ulcers. It follows that we now deſcribe Ulcers of a more malignant Nature, which will not admit of a Cure by any of the Methods we have hitherto laid down; from the Stubbornneſs of their Diſpoſition they are called in the medical Schools, *Ulcera Dyſepulotica*, *Chironia*, *Cacoëthica*, *Rebellia*, *Contumacia*. No Man in his Senſes will deny that they have all their proper Cauſes to which they owe this bad Diſpoſition. Theſe malign Ulcers uſually appear in Subjects of a bad, ſcorbutical, cachectical, and hydropical Habit of Body; or where you have the Pox, a *Caries*, or *Callus*, where there is great Acrimony of Blood, or a Cancer at the Bottom of the Caſe: whoever expects to be attended with Succeſs in treating theſe Caſes, ſhould diligently ſearch out and extirpate, if poſſible, theſe Cauſes of the Diſorder: but this in moſt Caſes is ſo difficult, that it will foil the moſt experienced Surgeon or Phyſician; nor will quackiſh Various Cauſes of Stubborn Ulcers.

Empirics get any Credit here, let them boaſt ever ſo long of the ſecret Virtues of their famous Plaſters and Unguents.

(1.) When there is a bad Habit of Body.

II. When you can diſcover nothing of a *Fiſtula*, *Callus*, *Caries*, putrid Fleſh, or of Worms concealed in the Ulcer, it owes it Obſtinacy to the bad Habit of the Patient, either from a Redundancy of a glutinous, acid, acrimonious, or bilious Quality in the Blood; or from ſome venereal Taint; or from Irregularity in Diet; or, in Women, from an Obſtruction of the menſtruous Diſcharge; in Men, of the Hæmorrhoids: in order to correct this vitiated Habit of Body, you muſt not only have Recourſe to internal Remedies, but muſt alſo inſiſt ſtrongly on a ſtrict Regularity in Diet: this is of ſo great Conſequence, that I have ſeen the worſt of Ulcers yield to a [a] proper Regimen of Diet, without the Uſe of one internal Remedy; only cleanſing them daily with proper Medicines, or dreſſing them with any common Ointment, Oil, or Balſam, covering the Dreſſing with any Plaſters in Uſe, as the *Emplaſtrum Saturninum vel Diapompholygos*; or indeed with the ſimple Application, of a Plantain Leaf, or Birthwort; with regard to eating and drinking. thoſe Things which ſit lighteſt upon the Stomach ſhould be preferred, and ſhould be given in very ſmall Quantities at a Time; for every thing that is too ſalt, acrimonious, acid, hard, or crude; all ſorts of Fat, Lard, or Swines Fleſh, every thing even of the lighteſt Kind taken intemperately, muſt be looked upon as Poiſon in theſe Circumſtances. Perſons of a ſanguine Habit ſhould avoid warm Things; thoſe of a phlegmatic cold Habit, cooling Things: a proper Regimen or Abſtinence is very much aſſiſting in the Performance of the Cure, by attending diligently to the Application of proper external Remedies: therefore the Ulcer ſhould be kept very clean; that the corrupt Matter, by lying long upon the Part, may not get an additional Acrimony, and ſo occaſion the ſpreading of the Diſorder. After it is well cleanſed, it is to be dreſſed with *Unguentum digeſtivum*, to which may be added, *Myrrha Maſticha, aut Colophonia*; or a Decoction *ex Juglandi Foliis cum injecto pauco Saccharo*; or *Decoctum Viridis æris cum Vino*. In ſome Caſes, *Spiritus Vini ſimplex, vel Aqua Calcis cum Linimentis immiſſa, vel Lapide Medicamentoſo* CROLLII *roborato*, has great Power in healing and drying up Ulcers: if you diſcover any *Sinuſes* or *Fiſtulæ*, they are to be laid open, and to be cleanſed afterwards in the Manner we taught above, and to be healed with *Balſamum Peruvianum, Copaibæ, Sulphuris Terebinthinatum*, or with any other agglutinating Medicine; laſtly, if internal Remedies are not neglected, there is no Doubt but that the very worſt of Ulcers may generally be cured.

(2.) Running Ulcers.

III. When theſe ſtubborn Ulcers are accompanied with a large Diſcharge, there is Reaſon to apprehend that the Blood abounds with too large a Quantity of thin acrimonious *Serum*; and they are from thence called *Rheumatic Ulcers*. This cannot be drawn off more properly than by cathartic Medicines. Where the Strength will admit of it, your Intention may be executed by preſcribing Cathartics and Diuretics, to be repeated frequently; at the ſame time cautioning your Patient againſt drinking too freely. *Millepedes præparati, Eſſentia Succini, Myrrhæ, Balſami Peruviani, Tinctura Tartari, Tinctura Antimonii tartariſata*, or any other Tinctures or balſamic Eſſences, of known Virtues for promoting the Secretion of Urine, are very properly preſcribed in this Caſe.

[a] Amongſt many others, conſult DOLÆUS in his *Encyclopædia* upon this Subject.

Large

Large and frequent Draughts of ſmall Liquors, which are frequently the Cauſes of theſe Diſorders, are diligently to be avoided: on the contrary, ſtrong Ale, or old Wine is tobe uſed, but ſparingly, for common Drink at Meals; but the Patient ſhould drink nothing between Meals: with regard to a proper Choice for Diet, thoſe Meats are beſt which have the feweſt Juices in them, and are moſt roaſted; Flummery, Calves Feet, and Calves Foot Jelly are very proper Diet: the external Medicines, after proper Evacuations, ſhould be thoſe that obtain the greateſt Reputation as Dryers: the principal of theſe are, *Aqua Calcis, Lapis Calaminaris, Tutia præparata, Creta, Maſtiche, Thus, Colophonium, & Cinnabaris nativa:* when you have ſprinkled any of theſe finely powdered upon the Ulcer, you are to lay over it the *Emplaſtrum Diapompholygos, Saturninum, vel de Lapide Calaminari.*

IV. Ulcers which ſpread and corrode the neighbouring Parts, are in the medical Schools called *Phagedænic Ulcers*, and betray a great Degree of Acrimony in the Blood; which is to be temperated as much as poſſible by the Phyſician, by the Uſe of lenient Medicines: the principal among theſe are, *Decocta ex Rad. Chin. Sarſaparill. Sympht. polypod. Lignit. Scorzon. Lapath. acuti, Herb. Malv. Alth. Hyperic. Sanicul. Agrimon. Marrub. alb.* and the like. With regard to Diet, you may obſerve the Directions we gave above at § III. All ſeaſoned Meats are bad in this Caſe: the Patient will receive great Relief by taking a purging Medicine ſometimes with the Addition of ſome *Mercurius dulcis*; this will not only leſſen the foul Diſcharge of the Ulcer, but will alſo deſtroy the Acrimony of the Blood, and forward the Cure: the ſame external Remedies are to be uſed here as we recommended at § II, III. eſpecially with the Addition of Mercury: the Uſe of them is to be diligently obſerved, till the Cure is perfected. (3) Phagedænic Ulcers.

V. *Cutaneous Ulcers* that attack the Skin of Adults as well as Infants, particularly about the Face, approach very near to the Nature of Phagedænic Ulcers: for they not only ariſe from an Acrimony in the Blood, but are apt alſo to ſpread abroad: therefore, in both theſe Caſes, thoſe Medicines will prove moſt effectual which keep open the Bowels, and ſoften the Acrimony of the Blood; (§ III and IV.) Adults in particular ſhould be adviſed to drink freely of what we call the Decoction of the Woods, or *Decoctum Radicis Lapathi acuti, aut Herbæ Fumaræ:* either of theſe Decoctions ſhould be drank by the Patient to the Quantity of ℥ viij. or ℥ x. three or four Times in a Day, as hot as he can bear it: the firſt Draught ſhould be taken in Bed, and a Sweating ſhould be endeavoured to be raiſed: to theſe you may very properly add *Eſſent. Fumariæ, Lignorum, Succini, vel Tinctura Antimonii tartariſata ad guttas* xxx *vel* xl. You may alſo preſcribe abſorbent Powders to be taken with theſe Decoctions, *ex Antimonio & Flor. Sulphur. parat.* A proper Regimen of Diet ſhould be ſtrictly obſerved in this Caſe: *in Infants Caſes who are yet at the Breaſt*, you ſhould preſcribe Medicines that will conſtantly keep the Body open and alleviate the Acrimony of the Blood; and at the ſame Time the Mother or Nurſe ſhould obſerve the Courſe we have preſcribed above, and be very exact in her Diet: with regard to external Application, you will receive great Benefit from *Oleum Tartari per Deliquium*, if you dip a Pencil or Feather into it, and dawb the Part three or four Times every Day, either with this alone, or with the Addition of *Oleum Ovorum atque Ceræ:* over this you muſt lay a Plaſter, as the *Empl. Saturninum*, (4) Cutaneous Ulcers.

turninum, vel de Minio, vel de Spermate Ceti cum Camphora, to correct the Acrimony, and to prevent Injuries from the external Air: if the whole Face ſhould be affected, which is frequently the Caſe in Infants, a Plaſter will be very improper; but you may make a linen Maſk, ſuch as we deſcribed above, treating of Burns: you will find the Uſe of the following Medicines in this Caſe, by no means to be deſpiſed; *Ol. Philoſophorum cum Oleo Ovorum, necnon Aqua Calcis, vel & Aqua ex Edulcoratione Antimonii diaphoretici*; the Ulcer ſhould be daily waſhed and cleanſed with one of theſe: if you pleaſe, in the room of theſe you may anoint the ulcerated Parts with *Unguentum de Lithargyrio vel Diapompholyg. vel de Enula*, with which in very ſtubborn Caſes may be mixed *Argenti vivi vel Mercurii Præcipitati rubri portiuncula:* If theſe Ulcers are attended with a large and foul Diſcharge, it will be proper to ſprinkle them with ſome abſorbent or drying Powders, as *Pulv. Tutiæ, Lapid. Calaminar. Ceruſſa, Creta*, &c. *cum Cinnabari nativa, aut Præcipitato rubro miſt.* or you may work any of theſe up into an Ointment *cum Cremore Lactis*, and uſe it as ſuch.

(5.) Cancerous Ulcers.

VI. *Cancerous Ulcers* are the moſt grievous of all the corroſive Kind: in theſe Caſes the ſame internal and external Remedies are to be uſed which we directed for the ulcerated Cancer; (*Book* IV. *Chap.* XVII. § XII.) Nevertheleſs, according to the Opinion of that great Phyſician and Surgeon M. A. SEVERINUS, there is more to be expected from manual Operation than Medicine in this Caſe; for many have been cured by the Knife or actual Cautery, where Medicine has availed nothing: but whenever you ſhall think it adviſeable to uſe the Knife or Cautery, remember that you go to the Bottom, and leave no Part of the diſeaſed Matter behind you; if you ſhould, all your Work would be in vain: ſome preſcribe here an *Aqua Phagedænica* made in the following Manner: ℞. *Aq. Calc. Viv.* ℔ j. *Mercurii Sublimati* ʒ j ß. *M. aut hujus loco Mercurii Præcipitati albi* ʒ j. *vel* ʒ j ß. which they apply upon Lint: ſome make this ſtronger of the Sublimate; others add *Spirit. Vini* ʒ j. *vel* ʒ ij: in the room of the Sublimate I have frequently ſubſtituted with Succeſs, *Mercurius dulcis*, mixed with *Aqua Calcis*, which is a much ſafer Method: digeſtive and balſamic Ointments are to be avoided in Cancerous Ulcers, as not only foreign to the Purpoſe, but extremely miſchievous.

(6.) Putrid and fetid Ulcers.

VII. When Ulcers are *putrid* or *fetid*, this Circumſtance ariſes either from the Patient's very bad Habit of Body, or from the Negligence or Unſkilfulneſs of the Surgeon: therefore it is the Buſineſs of the Phyſician to correct the Habit, by the Adminiſtration of proper internal Remedies, and of the Surgeon to clean the Ulcer frequently, eſpecially if it is attended with intenſe Heat: for where Wounds are dreſſed and cleanſed but ſeldom, which muſt frequently happen in the Army after ſmart Engagements, where great Numbers are wounded, it can ſcarce happen but that the injured Parts will be annoyed with Heat, Putrefaction, or Worms: you cannot more readily prevent theſe Inconveniences, than by carefully dreſſing the Parts with *Unguentum digeſtivum cum Ægyptiaco ſeu Fuſco* WURTZII *permixtum*; *aut Aqua Phagedænica*; *aut Mercurius præcipitatus ruber, vel ſolus, vel cum Alumine uſto miſtus, vel cum Unguento digeſtivo ſubactus:* theſe Dreſſings are to be continued till the putrid Fleſh ſeparates and leaves the *Fundus* of the Ulcer with its own natural roſy Colour: whilſt this is doing, it will be proper to cover the Part with Lint diped in Spirits of Wine, which is a very powerful Remedy againſt Putrefaction:

When

When the putrid Parts are caſt off, you muſt proceed in Healing as you do in other Caſes: but Myrrh ſhould be particularly added to the healing Medicines, as a moſt excellent balſamic. The Surgeon ought always to take Care in this Caſe to call in a ſkilful Phyſician, who, by proper Remedies, may keep up his Patient, and preſerve him from ſinking before the Cure is perfected. Ulcers attended with Worms are to be treated in the ſame Manner; for whatever prevents Putrefaction, will deſtroy Worms.

VIII. Some Ulcers are ſo very malign and obſtinate, and notwithſtanding they have no Alliance with any venereal Taint, yet they will not yield to any of the foregoing Remedies: when this happens, the only Method of Cure is, by adminiſtering mercurial Medicines, or raiſing a gentle Ptyaliſm, as I have frequently experienced: for ſome Men's Blood is ſo foul, that their Ulcers will not even be palliated, much leſs cured, without the Aſſiſtance of Mercury: but if they ſhould be attended with any venereal Diſorders at the ſame Time, the Uſe of Mercurials will then be abſolutely neceſſary, as we ſhall ſhew in the next Chapter. Some Ulcers require Salivation.

CHAP. IV.

Of the Method of treating VENEREAL ULCERS[a].

I. VENEREAL Ulcers, as we have already declared, are almoſt always ſituated in the *Inguina*, after the Suppuration of venereal *Bubos*; or, in the *Prepuce*, *Frænum*, or *Glans Penis*, which is uſually termed a *Chancre*: in Females they are frequently ſituated upon the *Vagina*, or *Labia Pudendi*: ſometimes the Noſe, Palate, Lips, Fauces, Tongue, and *Uvula*; ſometimes the *Os Frontis* and other Bones of the Head, and elſewhere, are ſubject to them: if they are neglected or ill-treated, one Ulcer of this Kind will produce an univerſal Pox: therefore the principal Intention to be obſerved in this Caſe is, to expel the venereal Poiſon by proper Remedies, both internal and external. Seat of Venereal Ulcers.

II. The Cure by internal Medicine, is to be performed by the Adminiſtration of purging Powders or Pills mixed with *Mercurius dulcis*: you may alſo adviſe your Patient at intermediate Times to drink Decoctions of the Woods, or to take *Eſſent. Lignorum*, *Pimpinell. Alb. Succin. Tinctura Antimonii*, &c. in a proper Vehicle: theſe Medicines have great Efficacy, if you take them before you riſe in the Morning and encourage a moderate Sweat. A ſtrict Regimen to be obſerved in Diet, is very neceſſary: Wine, and vinous or ſpirituous Liquors, Aromatics, Spices, Salt, acrimonious or acid Things, are Poiſon in theſe Circumſtances: if the Diſorder has acquired ſo great a Degree of Inveteracy, that theſe Medicines are not equal to the Cure; you muſt have Recourſe to the ſtrongeſt Sudorifics, eſpecially to ſtrong Decoctions of the Woods: or you may give Mercury in ſuch Quantities as to raiſe a Salivation, by which you will cure both the Ulcers and the Pox which was the Cauſe of them. Internal Treatment.

III. Whenever the Ulcers are ſituated in the Mouth, *Uvula*, Fauces, Tonſils, or Tongue, external Remedies become neceſſary as well as internal: the Patient ſhould frequently uſe a Gargle, made *ex Decocto Lignorum, vel ſimplicis, vel Melle* External Treatment.

[a] On this Subject read ASTRUC, *De Morbis Venereis*.

Roſarum

Roſarum temperato. The vitiated Part ſhould often be touched *vel Aqua viridi* HARTMANNI, *vel Roſarum Melle cui ad lenem uſque acorem, Spiritus Vitriolis pauxillum inſtillatum eſt:* after this it is to be healed, *per Eſſent. Succini et Myrrhæ, vel per Oleum Myrrhæ per Deliquium.* If the Ulcers appear on external Parts, it will be proper to deſtroy them with *Unguentum digeſtivum aut Baſilicon Mercurio vel vivo vel albo aut rubro præcipitato permixtum*; theſe Dreſſings are to be covered with the *Emplaſtrum de Ranis* VIGONIS, or with the *Diachylon cum Mercurio.* When the Ulcer is cleanſed, you may dreſs with the Eſſences we adviſed above, or ſprinkle it with the abſorbent Powders we have ſo often recommended, (ſee *Chap.* I. *N.* XV.) but you muſt add a ſmall Portion of red Præcipitate. An equal Power with the foregoing, in cleanſing and healing theſe Ulcers, is held by the *Aqua Phagedænica, vel Aqua Calcis Mercurio dulci imprægnata:* either of theſe may be applied frequently every Day, touching the Part ſometimes with the *Lapis Infernalis*, or *red Præcipitate:* when the Ulcer is thoroughly cleanſed, you may heal either after the Method recommended by HARRIS *in Diſſertat. Chirurg.* that is, with a ſimple Ointment compoſed *ex Mercurio vivo cum Tereb. q. ſ. ſubactum:* or you may uſe the following *Formula:*

℞. *Ung. Mundificativ. vel Diapompholyg. Mercur. crud. pauca Terebinthin. extinct. ana* ℥j. *vel* ℥ß. *M. in Mortario Vitreo.*

℞. *Amalgam. Mercur. et Stanni* ℥j. *Bol. Armen.* ℥ij. *Ung. Roſat. q. ſ. M. f. Ung.*

If at the ſame Time you have a *Caries* of the Bone, you are to treat it with the Remedies which we ſhall deſcribe below at *Chap.* VIII. particularly *cum Euphorbio vel Oleo Caryophyllorum, vel Aqua Phagedænica, vel Spiritu Nitri, in quo Mercurius ſolutus fuerit*; or laſtly, if you can do it with Safety, apply the actual Cautery. Sometimes, when theſe Ulcers fall upon the ſoft Parts of the Body, particularly on the *Inguina* they ſpue out ſuch large Quantities of Lymph, that all the Medicines you can invent, for cleanſing or drying them up, will avail nothing: this is occaſioned by the Rupture or Eroſion of ſome lymphatic Veſſels: In this Caſe, we ſhould try what we can do by the Application of proper[a] Compreſſes and a tight Bandage: but if theſe afford you no Aſſiſtance, you muſt call the actual Cautery in Aid, and apply it frequently, with Caution, to the vitiated Parts.

Venereal Ulcers attended with great Danger.

IV. If venereal Ulcers of the *Penis*, or its *Glans*, are negligently treated, an univerſal Pox will frequently be the Conſequence; the *Urethra* will often be perforated in various Places, and the Urine be diſcharged as through a Sieve: ſometimes the whole *Glans* and *Penis* will be eaten off, or ſo miſerably afflicted with *Scirrhus* and [b] *Cancer*, that you will be forced to extirpate them with the Knife: when the Noſe is affected with theſe Ulcers, it is frequently demoliſhed by them; the Diſorder in this Part is called *Ozæna*, of which we ſhall treat more fully when we come to deſcribe *Chirurgical Operations:* ſometimes the Palate with its Bones are ſo eroded and perforated, that an open Communication is made between the Mouth and Noſtrils; that the fluid Part of our Aliment makes its Way out at the Noſe: theſe Paſſages can ſcarcely ever be cloſed again, eſpecially if they are large; but when the Extremities of them are healed, they may

[a] In this I have followed RUYSCH, *Obſ.* 41. with Succeſs.

[b] Conſult RUYSCH *Obſ.* 30. *et* DOEBEL. *Hiſt. Penis Cancroſi.*

be

be closed with a small a Plate of Silver or Gold: the Tonsils, the external Coat of the *Uvula*, and the whole *Uvula*, are very frequently destroyed by the Virulence of these Ulcers: Decoctions of the Woods and Mercurials are the principal Antidotes to this Poison; lastly, the *Cranium* itself, particularly on the frontal Bone, is frequently, as I have often seen, so eroded and perforated by a *Caries*, that the Brain lays bare, and you may plainly see the Pulsation of the Arteries; from whence arise grievous Symptoms, and frequently Death, unless timely prevented by a proper Method of Cure.

CHAP. V.

Of Callous Ulcers.

I. THE Cure of a *callous* Ulcer is attended with great Difficulty: to say the Truth, it will admit of no Cure till the *Callus* is extirpated. A *Callus* may be extirpated three ways: the mildest Method which is to be used to a recent *Callus*, that is not yet become very hard, is performed by *corrosive Medicines*, and those of the mildest Kind: amongst many others you may use *Alumen ustum*, *Præcipitat. rub.* either separately, or mixt in equal Proportions, or made up with *Unguentum digestivum*, or *Basilicon*; The *Unguentum Ægyptiacum seu Fuscum* Wurtzii will answer this Intenrion, especially if you add a little *Præcipitatum rubrum* to it. If the *Callus* does not yield to these Applications, you may destroy it with *Lapis Infernalis* or *Butyrum Antimonii*; the same End is also well answered by the Medicine which is made by a Solution of *Argentum vivum in Spiritu Nitri vel Aqua forti*. A recent Callus how to be treated.

II. Le Dran has taught us a still milder Method of destroying Callosities, *in Observat. Chirurg. N.* CXV. *Tom.* II. which is as follows; for four or five Days he applies a Plaster, made *ex Emplastro Diachyl. cum Gummis, et Vigonis cum quadruplici Mercurio, ana*; and this he renews Morning and Night, in order to soften the callous Lips in some Measure: after this he makes frequent Incisions that pass so deep as to penetrate through the whole Thickness of the *Callus*, and stops the Blood that succeeds these Incisions with dry Lint: then he applies the same Plaster again to the Ulcer, so that it may touch the naked incised Lips: after about four Days he repeated the *Sacrifications*, and this to a third or fourth Time, if it is necessary; that is, if they are not destroyed before: by this Method he affirms, that Callosities by Degrees give way, and a Cicatrix will succeed, without the Use of any other Remedy; I have often tried this Method, and with good Success. Le Dran's Method.

III. If *callous* Ulcers are accompanied with *Fistulæ*, then the Sinus must be laid open, before we attempt to destroy the *Callus*, as we advised above when we treated professedly on *Fistulæ*; after this the *Callus* is to be consumed in the same Manner as we directed above: but if we shall think the Use of the Knife unsafe, or if the Patient will not admit it, it will be proper to form Tents, and daub them with *Unguentum Ægyptiacum vel Fuscum* Wurtzii, and thrust them up the *Sinus*; by this Method a *Callus*, that is not of long standing, may be When Fistulæ are accompanied with *Callus*.

[a] See Part II. *Operat. on the Palate.*

destroyed;

deſtroyed; eſpecially if you dip the End of the Tent, in *Præcipitat. rub. Lapid. Infernal. vel Butyr. Antimonii*, before you paſs it up the *Sinus*; and continue this Method till the *Callus* is deſtroyed: but when you cannot reach the *Callus* with the corroſive End of the Tent, you may uſe the following Method; you may inject *Aqua Phagedænica*, or a Solution of *Ung. Ægyptiacum aut Fuſcum* WURTZII *in Spiritu Vini*, up the *Sinus*, and cloſing the Aperture, confine it as long as you can conveniently; repeating it as you ſhall ſee Occaſion: when you have removed the *Callus*, the Ulcer may be cured in the Manner we have directed above *Chap.* II.

When the *Callus* is hard.

IV. Sometimes you will be obliged to uſe the Knife; as in *callus Ulcers* or *Fiſtulæ*, that are of long ſtanding, and have formed Variety of *Sinuſes*, where you can do nothing with corroſive Medicines; or, where they affect and corrode the Nerves, or Veins and Arteries, and bring on violent Convulſions, or Hæmorrhages, before they affect the *Callus*; in this Caſe, the ſafeſt Way is to lay open the *Fiſtula* in the Manner we deſcribed above, (*Chap.* II. *N.* V.) taking great Care not to wound Nerves, Tendons, or Arteries: when you have laid open the *Sinuſes* of the *Fiſtula*, you may preſently deſtroy all the *callous* Bodies, either by the Uſe of Corroſives, or by LE DRAN'S Method, recommended at *N.* II; healing the Ulcer afterwards in the Manner we have already adviſed.

How a very bad *Callous* Ulcer is to be treated.

V. Laſtly, if even this Method of Treatment ſhall not anſwer the deſired End; if the Patient is well ſtocked with Strength and Courage; if the Situation of the Nerves and Arteries is favourable; you may cut out all the *callous* Parts with your Knife, or deſtroy them with the actual Cautery; this Operation, though attended with great Pain, will bring the moſt obſtinate *callous* Ulcer to the State of a recent Wound; and unleſs a Caries, bad Habit of Body, Pox, Scurvy, Dropſy, or ſome other conſtitutional Complaint is in the Way, it may be cured by the moſt common Remedies: therefore there is no Reaſon why we ſhould fall into Admiration at, or doubt the Veracity of, M. A. SEVERINUS, when he affirms, that he has happily ſucceeded in the Cure of the moſt deſperate Ulcers, by this Method.

CHAP. VI.

Of ULCERS *ſuppoſed to be produced by* MAGIC *or* WITCHCRAFT.

Medicine foreign to this Caſe.

I. THE Remedies that PARACELSUS, HELMONT, AGRICOLA, and many others have, with great Induſtry, invented to cure Ulcers, which are the Effect of Magic, and always contain ſomething unnatural in them, as Thread, Nails, Needles, are entirely uſeleſs, and therefore ridiculous and abſurd: but if any are to be preferred to the reſt, we ſhould give the firſt Place to the following Remedies; *Folia Quernea, aut Salignea, Adianthum, Hypericum, vel Fuga Dæmonum, Mercurius vivus, Aſſa fætida:* theſe are hung round the Neck, or applied in ſome idle Manner, ſo that they can do no Miſchief: ſome preſcribe the Aſhes of a Witch that has been burnt; others burn *Stercus Humanum*, and ſprinkle the Ulcer with the Aſhes; HEERIUS and HORSTIUS are high in the Commendations of *Unguentum de Viſco Corylino* CARICHTERI; MYNSICTH preſcribes his *Emplaſtrum fætidum*: others, different Remedies of equal Efficacy.

II. Thoſe Phyſicians who conſult their own Reputation, and the Health of their miſerable Patients, ſhall I ſay, or infatuated Patients, will preſcribe natural Remedies, ſuch as are beſt ſuited to the Nature of the Ulcer, and the Patient's Habit of Body, as we have taught in the foregoing Chapters: for although we ſhould make ever ſo large Conceſſions, concerning the Power which Devils and Sorcerers are by ſome ſuppoſed to have over Men, yet we ſhould never be juſtified in aſſerting, that Diſorders, thus produced, were not to be treated by natural Remedies, but that we ought to have Recourſe to ſuperſtitious, naſty, and ridiculous Methods of Cure: to ſay the Truth, thoſe Ulcers are uſually affirmed to be the Effect of Magic by unſkilful and ſuperſtitious Barbers and Medicaſters, which evade their Art, though at the ſame time they are eaſily to be cured by an experienced Surgeon, who can thoroughly inveſtigate the true Cauſe and Nature of the Diſorder. There have been, even amongſt the Surgeons, ill-minded Men, who have falſely affirmed Ulcers to be the Effect of Magic, in order to inhance the Price of the Cure. What is to be done.

CHAP. VII.

The Method of treating old ULCERS, *eſpecially thoſe that affect the* LEGS.

ALTHOUGH there is ſcarce any Part of the Body free from inveterate and obſtinate Ulcers; yet the Legs are found to be much more ſubject to them than the reſt. As we before (*Chap.* III.) treated of malignant and inveterate Ulcers in general, we ſhall here only conſider thoſe which are ſeated in the Legs, or lower Extremities: but the general Cauſes of obſtinate Ulcers in the Legs are almoſt always the ſame with thoſe of malignant Ulcers in general; for theſe, like the former, uſually ariſe either from a bad Habit of Body, too great Thinneſs or Acrimony in the Juices; or from being attended with Calloſity, and Caries of a Bone; or, laſtly, from the Obſtruction of ſome uſual Evacuation, as of the Menſes in Women, or from other Cauſes of the like Nature: in order therefore to remedy theſe Ulcers, the Surgeon ſhould give a particular Attention to their Cauſes, that he may be thereby led to a rational Treatment of them. In what the Cure chiefly conſiſts.

II. Before we enter into an Inquiry, what are the moſt likely Means to be uſed to cure theſe Ulcers, it will not be amiſs to examine, whether they can be healed without Danger to the Patient: for we are furniſhed with frequent Examples, in the Writings of Phyſicians of the greateſt Experience, where the worſt of Diſorders, and even Death itſelf, has been the Conſequence of healing theſe Ulcers. The Anſwer to this Queſtion, if I am not miſtaken, is very clear, from what I have delivered above, in *Chap.* I. N°. IX. to wit, in Perſons advanced in Years, or labouring under an infirm Habit of Body, it is moſt adviſable not to attempt to heal them; ſince they are in this Caſe to be looked upon rather as a Relief of Nature, than a Diſorder, as they ſerve to drain off all noxious Humours from the Body: but I would not have this Rule extended to young robuſt Subjects, without ſome very material Reaſon; for, in theſe, the firſt Cauſe of ſtubborn Ulcers may be removed by Abſtinence, or a regular Way Whether Ulcers in the Legs may ſafely be healed.

of living, by opening Fontanells, or by proper internal Remedies, without any Danger: and the Cauſe being removed, the Ulcer may be healed with great Safety.

What is to be obſerved in general in old Ulcers.

III. Although we have declared above, that it is improper to heal inveterate Ulcers in old Subjects; yet I am very far from affirming, that no Care at all ſhould be taken of them: on the contrary, I think it abſolutely neceſſary that they ſhould be attended to. The Surgeon is to obſerve two Things in this Caſe: firſt, to relieve the Pain, and other violent Symptoms; next, to prevent the Ulcer from ſpreading, and new Symptoms from coming on.

Internal Remedies.

IV. In the firſt Place, Abſtinence and a ſtrict Regimen in Diet is to be obſerved: they ſhould abſtain from Pork, from all ſalt or ſeaſoned Meats, or of hard Digeſtion, and eat but ſparingly of the moſt innocent Food: gentle Purges are to be frequently repeated, to carry off the redundant Humours by Stools: proper internal Medicines are alſo to be preſcribed, ſuch as are moſt likely to remove the Cauſe of the Ulcer. In Perſons advanced in Years, balſamic and bitter Medicines are requiſite, to temper the violent Acrimony of the Blood: as *Elixir Proprietatis*, *Eſſent. Myrrh.* *Eſſent. Succini*, *Eſſent. Balſami Peruviani*, and others.

External Remedies.

V. With regard to the external Treatment of the Ulcer, Care muſt be taken that it be cleanſed from its Sanies, once or twice every Day: you may then dreſs it, either with dry Lint, or with Lint dipt in *Decoct. Fol. Nucis Juglandis vel Ariſtolochiæ.* Over this you may lay the *Emplaſtrum ad Ulcera antiqua* BAUHINI, *Diaſulphuris* RULANDI, *Diapompholygos*, *Saturninum*, *de Lapide Calaminari*, or any other of this Kind: theſe Rules being nicely complied with, if the Patient is well guarded from external Cold, and particularly from a moiſt or damp Air, there is no room to doubt but theſe Ulcers may become very mild, and convenient for the lengthening his Life. The Phyſicians amongſt the Antients, obſerving the ſalutary Effects of Ulcers upon old Perſons, thought Nature to be the beſt Guide, and therefore opened Fontanells in many Caſes, which anſwer the End of Ulcers, in draining off the noxious and redundant Humours.

How Inflammation and Pain is to be treated.

VI. Whenever Inflammations and violent Pains come on, as they frequently do, either from a Blow, or Cold, or putting the Leg into cold Water; or from Paſſions of the Mind, or Irregularity in Diet; it will be proper in this Caſe, firſt, to take away ſome Blood, in plethoric Conſtitutions: then to apply a linen Compreſs, dipt in *Aqua Reginæ Hungariæ vel Spiritu Vini Theriacali*, *aut Camphorato*, *vel & Aqua Calcis & Spiritu Vini Camphorato calidis.* The Patient ſhould keep his Bed, and defend the injured Limb as much as poſſible from Cold: and in the Morning he ſhould be ordered to drink plentifully of ſmall green Tea, white Wine Whey, or any other ſmall Liquors that may be likely to promote a Sweat. By theſe Means the Inflammation and Pain will quickly go off: but there is great Danger, when the Patient is of a bad Habit of Body, if the Inflammation runs to a great Height, and begins to degenerate into a Gangrene: in this Caſe the ſame Remedies are to be uſed, both internal and external, which we preſcribed above when we were treating of a Gangrene (*Book* VI. *Chap.* XIV. N°. V. *and the following.*) But above all, you are here to be very careful to keep up the Spirits of weak and aged Perſons with proper Remedies (particularly the Bark), and to provoke gentle breathing Sweats. If theſe

Rules

Rules are neglected, there is very imminent Danger that *Sphacelus* and Death will by Degrees ſteal upon you.

How to treat Ulcers that dry up ſpontaneouſly.

VII. When theſe Ulcers dry up ſpontaneouſly, in old and infirm Perſons, a Horror, Nauſea, and a great Weakneſs uſually ſucceed, which declare Death to be at Hand: (*Chap.* I. N° IX.) the firſt Intention is, to ſupport the remaining Strength of Nature as much as poſſible by proper Diet and Medicines: there ſhould inſtantly be applied to the Ulcer *Radix Gentianæ, vel Iridis Florentinæ contrita*; or if theſe ſhall be thought of too little Force, *Radix Hellebori nigri in Pulverem aut Globulos redacta*; or laſtly, *Pulvis Cantharidum, aut Globulus ex Emplaſtro Veſicatorio Officinarum:* theſe Applications will produce ſo great a Stimulus, that the Ulcer will frequently run again, to the great Relief of the Patient: when this happens, you muſt treat it as before: but when it reſiſts all Remedy, and ſtill continues dry, you have no Hopes of Life remaining.

CHAP. VIII.

Of CARIES *of the* BONES.

I. THE *Caries* or Corruption of the Bone may very juſtly be eſteemed one of the principal Cauſes of the Depravity and Inveteracy of Ulcers: for you will find it ſcarce practicable to heal an Ulcer, or if you do bring it to heal, it will not remain long in that State, where you have a carious Bone concealed at the Bottom.

A Caries what.

II. We call that Diſorder of the Bone a Caries, where the Bone, from whatſoever Cauſe it ſhall proceed, is deprived of its Covering, or *Perioſtæum*, and having loſt its natural Heat and Colour, becomes fatty, yellow, brown, and at length black: this is the firſt and lighteſt Degreee of this Diſorder, and is called by the Antients, according to CELSUS, *Lib.* VIII. *Cap.* II. *Os Vitiatum* and *Nigrities:* but the greater Degree of this Diſorder is, when the Bone is eroded and eaten, and becomes uneven like a Pumice Stone, from the Number of ſmall Holes, of which it is full; when it diſcharges a filthy Sanies, whoſe Acrimony ſoftens, relaxes, and deſtroys the fleſhy Parts that grow round it: this is a true *Caries*, or *Ulcer of the Bone*, and every Bone in the Body is ſubject to this Diſorder: and although this Ulcer may ſometimes appear to be very happily healed, yet, after the Cicatrix has been brought on for ſome Time, you have an Abceſs formed: the Diſorder will return afreſh, and the acrimonious corrupted Matter, which continually ſpues out from the carious Bone, being collected within, will produce various grievous Symptoms, ſuch as Shivering, Vomiting, and Fever, and deſtroy the neighbouring Fleſh again.

Different Denominations.

III. There are many Names and Species reckoned of this Diſorder, and of others that bear a near Relation to it: for it is called a *Caries*, [a] *Spina Ventoſa* or *Spinæ Ventoſitas*, a *Gangrene* and *Cancer* of the Bone by [b] CELSUS, ſometimes by the *Greek* Term [c] *Teredo*, and ſometimes *Pædarthrocaces* [d]; though ſome Au-

[a] We have a Treatiſe on the *Spina Ventoſa*, by PANDOLPHUS, an *Italian*, republiſhed with the learned Notes of MERCKLINUS, *Norimberg*, 1674, 12*mo*.

[b] See the Book laſt cited, *p.* 258. [c] Ibid. *p.* 64, 104, 143, 264. and the following.

[d] M. A. SEVERINUS treats on this Subject in his Book *De Abſceſſibus*, and there are ſeveral *Academical Theſes* on this Head, by different Authors.

thors conſtitute as many diſtinct Species of a *Caries* as we have reckoned up Names, yet I think there is not ſo material a Difference between them, that we ſhould multiply them into ſo many ſeparate Species: therefore I think it beſt to diſtinguiſh them into two Sorts: the firſt where the Diſorder begins in the internal Part of the Bone; the other, on the Outſide, or from an external Cauſe. I would call this a *Caries*, and that a *Spina Ventoſa:* or when it happens in Children, I would comply with SEVERINUS, and call it *Pædarthrocaces:* but of theſe we ſhall preſently treat more fully, in a particular Chapter for that Purpoſe, and explain their Differences more accurately.

Cauſes. IV. We find two Cauſes of the [a] *Caries* of the Bone: for, 1. a *Caries* ariſes, when the Bone is deprived of its *Perioſtæum*, by a Wound, Fracture, Bruiſe, or any other Accident, and either is expoſed to the Injuries of the external Air, or is corrupted by greaſy Dreſſings, or the common vulnerary Oils which are uſually applied to ſimple Wounds, ſuch as *Oleum Hyperici*, *Lilior. albor. Balſamum Samaritanum*, &c. or, 2. a *Caries* ariſes, when the Fluids are interrupted in their Circulation, by any external Violence, or internal Cauſe whatſoever, from whence Inflammation and Suppuration ſucceed; by which the *Perioſtæum* and Bone ſuffer to ſuch a Degree, that the Veſſels which are ſent to theſe Parts for the Nouriſhment and Support of the Bone and *Perioſtæum*, being inflamed and corrupted, the Bone is brought into Conſent, and ſoon becomes carious; this Diſorder, if not quickly remedied, ſpreads and communicates itſelf to the neighbouring Parts of the Bone, making the ſame Progreſs with Ulcers in the ſoft Parts.

Degrees. V. From whence it evidently appears, that there are ſeveral Degrees of Eroſion or *Caries* of the Bone: the firſt and mildeſt Degree is, when the Bone is laid bare, looks greaſy, and turns yellowiſh: but as ſoon as it becomes truly yellow, brown, or black, the incipient *Caries* degenerates into a worſe State: the third Degree is, when the Bone becomes uneven, rough, and rotten: the greater Eroſion the Bones have ſuffered, the more rough and uneven will they appear. When the *Cranium* is perforated through both Tables, or the *Tibia* or *Femur* are eaten through to the *Medulla*, this is a *Caries* of a very bad Kind: but the worſt Kind of *Caries*, where indeed the Caſe may almoſt be pronounced deſperate, is, that which falls upon the Joints, or any Parts of the Bones that lie deep: becauſe you can have no Acceſs to it with your Hands, to clean the Bone, and the Caſe admits of no Remedy but Amputation of the Limb.

Diagnoſis. VI. A *Caries* may be diſcovered two Ways; as it is concealed, or as the diſeaſed Bones are expoſed to View: 1. When the Bones lie open to the Sight, the *Caries* diſcovers itſelf by the following Signs: the Bone looks greaſy, and degenerates from its natural Colour, to yellow, brown, or black; it is bare, and the *Perioſtæum* deſtroyed: if you apply your Finger or Probe to the Bone, it will diſcover itſelf to be rough, uneven, and ſpongy: 2. but where the Bone is covered with Fleſh, it will then diſcover itſelf by the following Signs: the Matter that flows from it will appear greaſy, brown, or blackiſh, and ſtink like rank Lard: when you take off the Dreſſings, they will be tinged with a blackiſh Hue, from the Colour of the Diſcharge: when you have room to paſs your Probe to the Bone (which is not always the Caſe) you will find it to be rough

[a] HAYNE, in his Book *De Oſſium Morbis*, treats ingeniouſly on the Formation and Cauſes of a *Caries*.

and

and uneven: the neighbouring Fleſh will appear flaccid, ſoft, looſe, ſpongy, and ſtink like rank Lard: laſtly, in Caſes where you can neither ſee the Bone, nor get at it with your Probe, you may very reaſonably ſuſpect it a foul Bone, or when the Ulcer frequently breaks out afreſh, after it has been healed, without any other manifeſt Cauſe; and eſpecially if any Fragments of Bone are diſcharged from it.

VII. From what has been laid down, it plainly appears, what Dangers the *Caries* is attended with, and what Event we may expect from each different Degree of it. Ulcers of this Kind give great Trouble in healing: they are very apt to ſpread, eſpecially where we cannot conveniently come at the *Caries* to deſtroy it: and when they are healed, they frequently break out again, as was juſt obſerved. Where the Diſorder increaſes, and extends itſelf to the Joints, particularly to the Knee, there is ſcarce any Remedy, but Amputation of the Limb: where the Circumſtances are ſuch that it ſhall not be thought adviſeable to take off the Limb, the Patient is followed with great Weakneſs and a feveriſh Diſorder, and by Degrees, with Death. Caries in the *Femur*, *Coccyx*, *Os Sacrum*, *Carpus*, *Tarſus*, and *Oſſa Palati*, meet with extreme Difficulty in the Cure. When the *Cranium* is affected with this Diſorder, it is frequently eat through even to the *Dura Mater*: from whence proceed acute Pains of the Head, great Watchfulneſs, Vertigo, a diſturbed Imagination, and many other Diſorders of that Kind, with great Danger of Death. *Prognoſis.*

VIII. With regard to the Cure of a *Caries*, many Methods have been attempted: [a] the firſt and mildeſt Method is applied to the ſlighteſt Degree of a *Caries*, and is performed by the Application of ſpirituous Remedies; ſuch as *Spiritus Vini*, or *Aqua Reginæ Hungariæ*: with which Applications alone I have cured ſlight *Caries*: or by the Balſamics, ſuch as *Pulv. Ariſtolochiæ, atque Iridis Florentinæ vel Pulv. Myrrhæ atque Aloës*: One of theſe Powders is to be ſprinkled upon the Part, after you have diligently wiped away the Sanies with dry Lint: this Method is to be continued till the diſeaſed Part of the Bone is caſt off, and new ſound Fleſh ſprings up in its ſtead. In a Caries that penetrates ſomewhat deeper, ſtronger Remedies take Place; ſuch as [b] *Pulv. Euphorbii, vel Eſſentia Euphorbii, cum Spiritu Vini optimo parato, vel Oleum Caryophyllorum Cinnamomi aut Ligni Guaiaci*: theſe may be applied with a Pencil, or ſpread upon Lint, and laid on the Part affected: others apply corroſive Medicines, as the *Aqua Phagedænica, aut Spiritus Vitrioli aut Sulphuris*, and with the ſame Succeſs: In the room of all theſe, you may very well ſubſtitute, *Soluti Mercurii in Aqua forti vel Spiritu Nitri.* We have enumerated theſe as the principal, from a great many other Remedies of the like Nature, that have been preſcribed for the ſame End: we purpoſely paſs by ſuch as are either too weak for the Intention, or too vehement to be admitted with Safety: ſuch as *Arſenicum vel Mercurius ſublimatus* in Subſtance: when you have procured an Exfoliation of the diſeaſed Part of the Bone, your Buſineſs is, to complete the Cure with Balſamics: therefore the next Dreſſings to take Place are, *Aqua Reginæ Hungar. Eſſentia Maſtich. Myrrhæ, Succini, Aloës, Ariſtolochiæ, Balſamum Peruvianum vel Capivi*, or any Balſam of this Kind: cover theſe with a Plaſter, and proceed after- Cure 1. By Medicine.

[a] The Antients uſed the Cautery or Raſp in the ſlighteſt Caſes, as you may ſee in CELSUS, *Lib.* VIII. *Cap.* II. but at preſent we never uſe theſe violent Methods but in deſperate Caſes.

[b] This is highly extolled by many. See MERCKLINI *Lib. De Spinæ Ventoſitate*, p. 473.

wards as you are directed above in the Cure of Ulcers in general (*Chap.* I. N° II. *and the following.*) Le DRAN has given us Obſervations on *Caries* of the Bones very well worth our remarking: particularly on a *Caries* of the Cubit, *Obſ.* 51, 52, 53: in the Loins, *Obſ.* 69: after the Small Pox, *Obſ.* 70: in the *Os Illium*, *Obſ.* 95: in the *Trochanter major*, *Obſ.* 97: in the Knee, *Obſ.* 102, 103: and in the *Tibia*, *Obſ.* 104.

2. By the Trepan.

IX. A Second Method of Cure for a greater Degree of *Caries*, conſiſts in [a] perforating the Bone after it is laid bare, with the Trepan or Inſtrument deſcribed in *Plate* VII. *Fig.* 2. *or Fig.* 7. A. *or Plate* XV. *Fig.* 8. in the ſame Manner as we adviſed in another Place to be done with the *Cranium*, after it had been laid bare by a Wound. *Book* I. *Chap.* XIV. N° 17. After this is done, the Part is to be dreſſed either with Lint, or with the balſamic Medicines which we have recommended above: by theſe Means the Exfoliation of the foul Bone is forwarded, and new Veſſels puſh through the Foraminula that you have made, which, joining with the neighbouring Fleſh, make a new Covering for the Bone.

3. By the Raſpatory or Chiſſel.

X. The third Method of Cure is performed by ſcraping away the diſcoloured or vitiated Part of the Bone, with a *Raſpatory* or Chiſſel (*Plate* VII. *Fig.* 3, 4, 5.) till, all the corrupted Parts being deſtroyed, the Bone appears white or ruddy and ſound. CELSUS adviſes this Operation of raſping the Bone, to be done boldly and expeditiouſly: SCULTETUS is of Opinion [b], that you ſhould never begin to ſcrape, till the Bone lies fairly expoſed, or rather not till it begins to ſeparate from the ſound Parts; and that you ſhould dreſs the Part with nothing but dry Lint, till this happens: but this Rule is not conſtantly to be obſerved. Others in particular Caſes uſe a Chiſſel and Mallet, (*Plate* VII. *Fig.* 10, 11.) by the Aſſiſtance of which, they ſtrike off the corrupted Parts from the ſound: but both theſe Methods, N° IX, and X. have been pretty much neglected by the modern Surgeons: though PETIT affirms, in his Book *de Morbis Oſſium*, when he is treating of a *Caries*, that where you have fungous Fleſh continually ſprouting up, the beſt Method is, to raſp the Bone, and afterwards to uſe the Cautery. In certain Tumors of the Bone, which are called by us *Spinæ ventoſæ*, which refuſe to yield to any medical Application, he adviſes not only to make frequent Perforations, but to take off the Tumors with the Chiſſel and Mallet: but we ſhall treat of this Caſe in the the following Chapter.

4. By the Cautery.

XI. The fourth, which is the moſt antient, ready, and certain Method of Cure, eſpecially in the greater Degrees of this Diſorder, is performed by burning down the vitiated Part of the Bone with the actual Cautery: ſee different Sizes and Figures of Cauteries in *Plate* III: great Care muſt be taken in performing this Operation, that you do not injure the Fleſh or other ſoft Parts that lie near: to prevent Miſchief of this Kind, your Aſſiſtant ſhould keep back the Lips of the Ulcer with his Hands: if the Opening is too narrow, it ſhould be enlarged with a ſponge Tent, or widened by the Knife, till the Bone lies fair: the Bone itſelf ſhould be well cleanſed with dry Lint: and if there is any fungous Fleſh, it ſhould be removed before you go to work with your Cautery. One Application of the Cautery will ſeldom be ſufficient for your Purpoſe, where the *Caries* is conſiderable: it muſt be frequently repeated, at longer or ſhorter Intervals as you ſhall think proper, if the *Caries* has ſpread itſelf

[a] See the Method of perforating, by CELSUS, *Lib.* VIII. *Cap.* ii, and iii.
[b] *In Armament. Chirurg.* pag. 42.

ſo

ſo wide, that you cannot deſtroy it with one Cautery, the firſt Iron ſhould be applied to the Middle of it, proceeding afterwards to its Lips: this Operation is not attended with great Pain, if you take Care not to hurt the ſoft Parts; for the Bones have no Senſe of Pain. [a] When the Bones of the *Cranium* are become carious, a cautious Surgeon will never riſque his Reputation on this Operation, from the apparent Danger there is of injuring the Membranes of the Brain, or the Brain itſelf: the ſame Caution may be obſerved in ſome other ſoft and ſpongy Bones, as in the *Sternum*, or a *carious Rib*, where, for the like Reaſons, the Cautery is to be avoided. The *Carpus* and *Tarſus* will not well admit of cauteriſing, and other ſpongy Bones of this Kind: and that more particularly from the Neighbourhood of the Tendons and Ligaments, which will neceſſarily be in great Danger of ſuffering.

What is to be done after cauteriſing.

XII. When you have cauteriſed the Parts in the Manner I have deſcribed, you ſhould dreſs at firſt with dry Lint: but if the Patient complains of a Senſe of Heat in the Part, you ſhould moiſten your Lint *cum Spiritu Vini*. You may afterward dreſs with Balſamics, ſuch as we deſcribed above at *N.* VIII. till the Exfoliation ſucceeds: and the Vacuity will ſhortly be filled up with new ſound Fleſh, which will be a Teſtimony of the Recovery of the Part: but where it happens otherwiſe, and the Bone is left bare, uncovered with Fleſh; or if the Fleſh with which it is covered, is ſoft and ſpongy, and does not adhere ſufficiently to the ſubjacent Bone, or where the Bone remains diſcoloured; in either of theſe Caſes, your original Diſorder is not extirpated. In theſe Circumſtances your Work is to be done over again, the ſpongy Fleſh muſt be removed, either with the Knife or Cathæretics, ſuch as the *Alumen uſtum et Mercurius præcipitatus ruber*, or ſtronger, if they ſhall be found neceſſary: and the actual Cautery muſt be again called for, or you cannot expect your Cure to ſtand.

When a Caries penetrates to the *Medulla*, the Trepan is ſometimes neceſſary.

XIII. When the *Caries* penetrates even to the *Medulla* in the larger Bones, [b] PETIT adviſes us, after the Example of [c] MEEKRENIUS, to make a Perforation, or two, or more, in the Bone with the Trepan; and furniſhes us with an Inſtance where he made three Perforations in this Manner, in the *Tibia*, after he had tried the Cautery, and was juſtified by Succeſs: but this Method can ſcarcely be put in Practice upon any other great Bone than the *Tibia*, becauſe you will be obſtructed by the great Quantity of muſcular Parts which you will meet with. He further informs us, that the *Os Pectoris* or *Sternum* may be perforated in this Manner, to make a Paſſage for the Diſcharge of Matter, which is ſometimes confined under it; and to make Way alſo for the immediate Application of Medicines to the diſordered Parts: but the Performance of this Operation on the *Sternum* requires the greateſt Caution and Deliberation; becauſe Reſpiration may be injured by it, or other grievous Diſorders may be produced. It is to be obſerved in this Place, that the Caries of the Bone which penetrates to the *Medulla*, or begins in the *Medulla* (which we term the *Spina Ventoſa*) does not always ariſe from an internal Cauſe, but frequently from an external Violence; by which the Veſſels which are diſtributed on the internal Part of the Bone are burſt, and Blood extravaſated: thus by its Stagnation in the Cavity quickly forms *Pus*, erodes the Bones, and produces a Caries, which extends itſelf from the *Medulla* to the external Parts.

[a] CELSUS has given the ſame Caution, *Lib.* VIII. *Cap.* ii. [b] *Lib. de Morb. Oſſium, cap. de Carie.* [c] *Obſ. Med. Chirurg.* 72. *edit. Latinæ, & 69 Belgicæ.*

XIV.

Admonitions of CELSUS.

XIV. When the Blackneſs or Caries extends to the other Side of the Bone, ſo that the whole Bone ſeems to be corrupted, CELSUS adviſes to take it entirely out, *Lib.* VIII. *Cap.* 2, 3. If the lower Part remains ſound, you muſt remove only as much as is corrupted: if a Bone of the *Cranium*, or the *Os Pectoris*, or one of the *Coſtæ*, is carious, the Cautery is not to be uſed, but it muſt be cut out: and in this no Delay is to be ſuffered, but you are to take it out the Inſtant you have laid it bare, before any inflammatory Symptoms come on, by which means you will do it with greater Safety. When a Cartilage is become carious, you muſt pare off the carious Parts with your Knife, according to CELSUS; to whom I am obliged for this Section, not having met with any modern Surgeon, who has treated ſo well on this Subject.

In what the Cure of a Caries conſiſts.

XV. Upon a diligent Attention to what has been delivered, we may very reaſonably conclude, that the principal Buſineſs in curing a Caries of the Bone, conſiſts in a ſpeedy Extirpation of the carious Parts of the Bone: this is done in very ſlight Caſes by the Application of *Spiritus Vini*, or *Aqua Reginæ Hungariæ*; in Caſes of more Conſequence, by a Solution of *Argentum Vivum in Aqua forti*: but in Caſes of the laſt Conſequence, by the Cautery or Knife. The reſt of the Cure is performed in the ſame Manner as other Ulcers are treated, by the balſamic Remedies which we have ſo often recommended.

How Bones that are very rotten are to be treated.

XVI. Where the Bone is exceeding rotten, or where the Diſorder has communicated itſelf to the Joint, for Inſtance, to the Knee, or to any Joint of the Arm or Leg, ſo that the diſeaſed Part cannot be extirpated, and the reſt of the Limb preſerved; you have only one Remedy left, and that a melancholy one, which is the Amputation of the diſeaſed Limb: otherwiſe your Patient will drag on a miſerable Life; yet at laſt perhaps, worn down with Pain and Weakneſs, attended with a long Train of grievous Symptoms, he muſt yield to Death [a]. In the large Bones, where the whole Bone is not carious, but only Part of it, as the external Part of the *Maxilla*, *Os Humeri*, *Tibia*, or *Clavicle*; or any Part of the *Rib*, *Ulna*, *Radius*, or *Fibula*, &c. you muſt not immediately proceed to the Amputation of the Limb, but only remove, in the moſt convenient Manner you can, either by medicinal Applications, or by the actual Cautery, the diſeaſed Part of the Bone, dreſſing afterwards as we taught above at *Sect.* XII. till the Bone is covered with ſound Fleſh, and the Ulcer healed. Sometimes Part of the diſeaſed Bone ſeparates ſpontaneouſly from the reſt of the Bone: if you can lay hold of it, and the Ulcer is wide enough, you ſhould remove it with your Fingers or the Forceps: if the Ulcer is not wide enough to admit of this, you muſt enlarge it with your Knife. You will meet with a remarkable Caſe of this Kind in MEEKREN. *Obſerv. chirurgic.* 66. *Edition. Belgic. et Obſerv.* 69. *edit. latina*, where a large Portion of foul Bone ſeparated and caſt off from the Arm: and another in RUYSCH. *Obſervat.* p. 94. *ac Theſaur. Anatomic.* VIII. *Tab.* III. where the ſame Caſe happened in the *Tibia*.

[a] LE DRAN gives us ſeveral Inſtances of Amputations in a *Caries*; particularly *Obſ.* 101, 102, 103, 104. But many of theſe Patients died, for the Reaſons juſt mentioned.

CHAP.

CHAP. IX.

Of the SPINA VENTOSA, PÆDARTHROCACES, *and* EXOSTOSIS, *which may be called* TUMORS OF THE BONES.

Spina Ventoſa, what.

I. THAT Species of Corruption of the Bones, which takes its Riſe in their internal Parts, and by Degrees enlarges the Bone, and raiſes it into a Tumor, is at this Time called, by Phyſicians and Surgeons, a *Spina Ventoſa*, by ſome [a] *Spinæ Ventoſitas:* Though the Antients were entire Strangers to theſe Terms, and diſtinguiſhed them by the Names of *Sideratio*, *Gangræna*, or *Cancer Oſſis*, or ſometimes by the Word *Teredo*. Some amongſt the *French* call it an *Exoſtoſis* [b]. Though this Term more properly belongs to certain [c] Eminencies or preternatural acuminated Excreſcences in the Bones, which happen after a Fracture or other Accident, and are ſometimes accompanied with a *Caries*, yet I have frequently ſeen this Caſe of the Bones, and have now Bones of this Kind in my Collection, where there is not the leaſt Appearance of *Caries*. This Diſorder ſeems to have borrowed the Term *Spina* from the Reſemblance which the Eminences of the Bone in this Caſe bear to Thorns, continually pricking the Fleſh, and producing grievous Pains: And the Epithet *Ventoſa* is added, becauſe the Tumor appears upon touching to be filled with Wind or Air, though in Fact it is never, or [d] very ſeldom diſtended with Air. Afterwards ſeveral Writers, and particularly PANDOLPHINUS, barbarouſly diſtorted the Word into *Spinæ Ventoſitas*.

Pædarthrocaces, what.

II. When this Diſorder happens to Children, many, with M. A. SEVERINUS, call it [e] *Pædarthrocaces*, from the *Greek* Words παις a Child, αρθρον, a Joint, and κακον an Evil: To ſignify that this Diſorder is moſt frequently found in the Joints of Children: For as the Bones of Children are ſofter and more ſpongy than the Bones of Adults and old Perſons, they are therefore ſo much the eaſier diſtended by Humours, and more frequently form [f] Tumors. SEVERINUS made another Diſtinction between the *Spina Ventoſa* and *Pædarthrocaces*. For ſome of theſe Tumors which we call *Spinæ Ventoſæ*, are very painful, frequently look red, and have all the Appearances of Inflammation: Others are free from Pain (at leaſt in any conſiderable Degree) in the Beginning, particularly in rickety Children, and theſe he called *Pædarthrocaces*. But at preſent theſe Names are pretty much confounded, and are deſervedly, as [g] MERCKLINUS has taught us, uſed for one and the ſame Diſorder; only with this Difference, that this Diſorder in Children begins with little or no Pain, but is almoſt always attended with Pain in its Progreſs.

III. There

[a] By the *Arabians*, witneſs JOS. PANDOLPHINUS, *Lib. de Ventoſitate Spinæ*.

[b] See MERCKLINUS, Annotations on PANDOLPHINUS, and what we ſaid above at *Sect.* III.

[c] See GORÆUS in *Definit.* JO. A. VIGO *in Chirurg.* and PETIT *Lib. de Morb. Oſſium, cap. de Exoſtoſi & Carie.* [d] MERCKLINUS relates a Caſe of this Kind, where, upon opening a Tumor, nothing was diſcharged but a Flatus, and the Patient died. [e] See M. A. SEVERINUS's Book *de Pædarthrocace*, contained in his excellent Work *De recondità abſceſſuum naturâ*; alſo the academical Theſes of AMMANNUS, TANSIUS, MÆBIUS, CHUNIUS, and others. [f] Caſes of this Sort may be ſeen in M. A. SEVERINUS *de Abſceſſ.* p. 144. and p. 467. RUYSH. *Epiſtol. Anatomic.* XIV. BIDLOO *Exercitat. Exoſtoſi.* [g] *Lib. de Spinæ Ventoſitate*, p. 53, 54, 248, et ſeq.

Symptoms. III. There are other Names of a *Caries*, which we have recited above *Sect.* I. and in the foregoing Chapter *Sect.* III. which agree much better with this Diſeaſe of the *Spina Ventoſa*, than with that Diſorder, which is vulgarly and ſtrictly ſpeaking called a *Caries*; as *Cancer Oſſis*, *Gangræna*, *Sphacelus Oſſis*, which Terms are frequently uſed by the Tranſlators of HIPPOCRATES; and the *Greek* Word [a] Τερηδων, which they tranſlate *Teredo*, from the Similitude of thoſe Worms which are called *Teredines*, which eat into and deſtroy Wood. It is very probable, that theſe are all ſynonymous Words for the *Spina Ventoſa*, different perhaps only in Degree: But I ſhall ſpend no Time in Defence of this Opinion, becauſe MERCKLINUS, in my Judgment, has ſufficiently demonſtrated not only this, but that the Diſeaſe itſelf was well known by the [b] Antients, contrary to the [c] Opinion of ſome. Whoever deſires farther Satisfaction upon this Head, may turn to this Writer's Notes on PANDOLPHINUS's Book, which we have ſo often quoted. Laſtly, we muſt obſerve in this Place, that PETIT, in his Book *De Oſſium Morbis*, *Chap.* XVI. ranks all theſe Names and Diſeaſes under the Name of *Exoſtoſis*, and at the ſame Time entirely neglects to mention the other Names, which are more vulgarly known, and in conſtant Uſe amongſt medical Writers. Whether he has judged well in this Caſe, I leave others to determine. For my own Part, I ſhall chiefly uſe the Term *Spina Ventoſa*, as the moſt received Name amongſt us at this Time.

Differences. IV. But theſe Diſorders, particularly their *Differences* and *Degrees*, are, in my Opinion, not deſcribed with ſufficient Accuracy by moſt Writers. I intend to deſcribe them as clearly as I can; for great Numbers of theſe Caſes have fallen under my Care: And nothing can tend more to an Improvement in the Method of treating theſe Diſorders, than an accurate Knowledge of their Differences. A *Spina Ventoſa* is by us underſtood to be a Corruption and Eroſion, or Caries of the Bone, occaſioned by a Depravity of the contained Fluids, and ariſing generally ſpontaneouſly, without any external Cauſe, beginning, not upon the external Face of the Bone, but between its *Lamellæ* or *Cells*, or in its internal Cavity; and extending itſelf by degrees, to the external Parts, at length affects either the [d] whole Bone, or a [e] greater or ſmaller Part of it, expanding it to a greater Width, or raiſing it into a Tumor. (*See Plate* XII. *Fig.* 16. A. B.) It is frequently hard, and ſometimes without Pain; at other Times it appears as if it was filled with Wind, and is attended with a greater or leſs Degree of Pain, pricking, ſhooting, at laſt it grows red, and is attended with other bad Symptoms; till the diſordered Bone being, by degrees, corroded, the common Integuments, and other ſoft Parts that lay over it, remaining at firſt entire, but at laſt partaking of the Diſorder, foul Ulcers of the moſt ſtubborn Sort break out. When Tumors of the Bone are hard, and the ſoft Parts not inflated, and are free from Redneſs, Inflammation, and Pain, as is frequently the Caſe in rickety Subjects, in this Caſe they are not attended with ſuch bad Symptoms as we have deſcribed above. SEVERINUS has given the Name of *Pædarthrocaces* to theſe Tumors, as we have already obſerved, becauſe this Caſe chiefly happens to Children,

[a] See GORÆI *definitiones ſub hoc vocabulo* Τερηδων. [b] Pag. 52, 63, 257, et ſeq.

[c] HEYNE was of this Opinion, *Lib. citat.* p. 62. He affirms, that this Diſeaſe was not known till the Appearance of the *Lues Gallica*.

[d] This happens to the ſmall Bones, ſuch as the Bones of the Fingers, *Carpus*, or *Tarſus*.

[e] This to the larger Bones, ſuch as the *Oſſa Cranii*, *Tibiæ*, *Femoris*, *aut Brachii*.

Children, and in order alſo to diſtinguiſh it from the *Spina Ventoſa* of the *Arabians*. But the painful, red, inflated Tumors that happen equally to Children and Adults, are called *Spina Ventoſa* [a], *Cancer vel Gangræna Oſſis*, *aut Teredines*. By an *Exoſtoſis* I mean a preternatural Eminence of the Bone, which is ſomewhat acute, or if you pleaſe, an Excreſcence of the Bone, whether it is attended with Eroſion or not. A *Spina Ventoſa* differs from a *Caries*, by being accompanied with Tumor; and is to be diſtinguiſhed from the *Rickets*, becauſe rickety Subjects are attended with various deformed Tumors on the *Epiphyſes* of the Bones, without Pain or Eroſion.

The Part affected.

V. Each of theſe Diſorders generally begin about the Heads or *Epiphyſes* of the larger Bones, where they are moſt tender and ſpongy, and where the noxious Matter may not only have ſufficient Room to lodge in the cellular Subſtance, but where it will alſo meet with the leaſt Reſiſtance in ſoftening and expanding the Parts [b]. Nevertheleſs, I have ſometimes ſeen this Diſorder ariſe in the Middle of theſe Bones, between their *Lamellæ*, eſpecially in the *Tibia*. *Tophs*, and *Venereal Gummata*, as they are called, which ariſe in the *Os Frontis*, and on other Parts of the *Cranium*, and frequently on other Bones, particularly on the *Tibia*, may all be ranked under this Claſs, as they owe their Origin to an internal Cauſe, and are only diſtinguiſhed from the others by being particularly painful in the Night. Yet we ſometimes ſee this Kind of Tumor in very chaſte Perſons, and where there is nothing venereal in the Caſe. Thus you ſee the *Spina Ventoſa* is not confined to the Bones of the Extremities, but ſeizes even upon the Bones of the Head, Face, Neck, and Breaſt: Though the Bones of the Arms, Legs, Fingers, *Carpus* and *Metacarpus*, *Tarſus* and *Metatarſus*, are more frequently the Subjects of this Diſorder. You may ſee various Caſes of this Kind in MERCKLINUS's Notes on PANDOLPHINUS, *p.* 227, *et ſeq.*

Generally from internal Cauſes.

VI. They ariſe generally, as we have declared above, from internal Cauſes, from acrimonious, ſcorbutical, rickety, or variolous Humours [c]. But principally from a venereal Taint; for they were not ſo frequently [d] obſerved in *Europe* before the Appearance of the venereal Diſeaſe. In the mean time it is reaſonable to ſuppoſe, nor is it contradicted by Obſervations, that this Diſorder may ſometimes owe its Riſe to external Cauſes, eſpecially in Perſons conſtitutionally diſpoſed to theſe Diſorders: When, for Inſtance, the Veſſels between the *Lamellæ* of the Bone, or in the *Medulla* itſelf, are, by a Blow, Fall, or any other external Violence, injured or torn, and the Fluids extravaſated. By degrees they putrify, corrupt and deſtroy the *Medulla*, and ſoften and corrode the Subſtance of the Bone: Whence proceed Pains, Tumors, Ulcers, and *Fiſtulæ* of Bones and the adjacent Parts, and all the ſame Miſchief which is uſual to ariſe from internal Cauſes.

VII. The

[a] Vid. CELSUS *L.* viii. *C.* 10. TULPIUS *Obſ. Med. L.* iv. *C.* 12.

[b] I have diſſected ſeveral, who died of this Diſorder; and in ſome I found the Bones ſofter than a Cartilage. [c] HILDANUS gives you Inſtances of this Kind, *Cent.* 4. *Obſ.* 95, 96.

[d] Some are of Opinion, that this Diſorder was abſolutely unknown till the Appearance of the Venereal Diſeaſe; as HEYNE, *in Lib. de Morb. Oſſ.* p. 62. but MERCKLINUS, in his Notes on PANDOLPHINUS, *Cap.* I. has plainly evinced the contrary, and ſhewn that it was known to HIPPOCRATES, GALEN, CELSUS, and others, who have deſcribed it under the Names of *Siduratio*, *Gangræna*, *Cancer Oſſis*, *&c.* which are only different Names for the ſame Thing.

[e] See an Inſtance of this in HEYNE, *De Morb. Oſſ. N.* 29.

Proximate Cauſes.

VII. The Proximate Cauſe of this Diſorder is either a Collection or Congeſtion of a viſcid and thick, or of an acrimonious and corroding Humour; or an Inflammation ariſing in the *Medulla*, or in the Subſtance or Cells of the Bone, degenerating into an Abſceſs, and forming *Ichor* or *Pus*. As theſe ſtagnating Fluids can find no Diſcharge from the Bones, eſpecially from their Cavities, they are confined there, till they putrify and become acrimonious, corrode and deſtroy the neighbouring Parts, converting them, particularly the *Medulla*, into a like Kind of Sanies; at length they attack the Bone, and deſtroy that. The Collection of viſcid and pituitary Fluids, with the Expanſion of the Bones, ſometimes happens without Pain, as in the *Pædarthrocaces*[a]: But the Eroſion of the Parts can never happen without the moſt acute Pains, proceeding, as we ſay, from the inmoſt Marrow. But in the Beginning of this Diſorder, when the Miſchief is only in the internal Part of the Bone, the Pain does not increaſe upon external Preſſure: When the Pain increaſes upon Preſſure, the external Parts are brought into Conſent. When this happens, the *Perioſtæum* and Parts that ſurround it, with the Subſtance of the Bone and the *Tunica cellularis* enlarge; from whence a Senſation frequently ariſes, as if the Parts were filled with Air or Wind, and the Diſorder was hence called *Ventoſa Spina*. But when the Tumor is opened, either ſpontaneouſly or by the Knife, if the Bone lays bare, you will frequently find it full of ſmall Eroſions, reſembling a Sponge or Pumice Stone, as it is in a *Caries*. From what has been here delivered, you may learn the near Reſemblance that theſe two Diſorders bear to each other, their Signs, and at the ſame Time ſome material Differences by which they are to be diſtinguiſhed.

Degrees of a *Spina Ventoſa*.

VIII. A *Spina Ventoſa*, ſtrictly ſo called, may very properly be divided into three Degrees. The firſt is, when the Patient complains of a continual grievous Pain in the Bone, which ſeems to him to proceed from the *Medulla*, and torments him ſo that he can have no Sleep. At this Time there is no external Pain or Tumor. In this State the Diſeaſe is confined to the internal Part of the Bone. The ſecond Degree of the Diſeaſe is, when after theſe Pains a red Swelling appears upon the Face of the Bone, either hard, or ſoft, and as it were windy, with external Pain more or leſs. The third Degree is, when after all the former Symptoms, an Abſceſs is formed in the Tumor, which either burſts ſpontaneouſly, or is opened with the Knife, and diſcharges a fetid *Ichor*, or purulent Matter, ſmelling like rank Butter or Lard; and afterwards maintains this Diſcharge in greater or ſmaller Quantities, like a carious Ulcer, and creates, what the Antients uſually called an Ulcer with *Caries* of the Bone. This Species of the Diſorder may be called an *Inveterate Spina Ventoſa*: The other a *recent* or *incipient* one.

Pædarthrocaces.

IX. A *Pædarthrocaces* begins with an Enlargement of the Bone, and generally without any Pain or external Cauſe: But in its Progreſs it is frequently attended with Pain and Inflammation, and at length with Abſceſs, Ulcers, *Caries*, as in the *Spina Ventoſa*, eſpecially about the Joints and Extremities of the Bones; and in ſhort is attended with the ſame Symptoms with the *Caries* and *Spina Ventoſa*. From whence it is evident, that the *Pædarthrocaces* may in ſome meaſure be

[a] MERCKLINUS thinks this cannot happen without Pain, but SEVERINUS and I have often ſeen it.

be looked upon as a distinct Disease, in the Beginning; but if it is not presently relieved, it will at length become a perfect *Spina Ventosa*, differing from the other in nothing but Degree.

X. From considering what has been already delivered, especially what has been taught in the preceding Chapter at *Sect.* VII. concerning the *Prognosis* of a *Caries*, it will be no difficult Matter to form a *Prognosis* of what we are to expect in the Course of Disorders of this Kind. For as it is manifest, that corrupted acrimonious Matter, when it is confined in the Cavity of a Bone, or included in its *Lamellæ* or Cells, cannot be easily discharged, either by Nature or Art; it necessarily follows, that it will, by degrees, corrupt and destroy the Parts that lay near it; till at length the Bone itself, if a timely Remedy is not applied, will be entirely corrupted and destroyed, so as to make it necessary to take off the whole Limb in order to save the Life of the Patient. Nay, what is still worse, if this Disorder arises from a vitiated State of the Blood, when you have taken off one Limb which shall have been affected in this Manner, you shall have it return with equal Fury in another, in the same Manner as it happens in cancerous Cases: Though this is not constantly the Case, especially if you correct this State of the Blood by proper Remedies, and by enjoining a strict Regularity with regard to Diet. In the *Pædarthrocaces*, and frequently in the two first Stages of the *Spina Ventosa*, the Disorder is happily cured by the Administration of proper Remedies. But the Cure will be attended with greater or less Difficulty, in proportion to the Inveteracy of the Disorder, the Progress it has already made, the Strength of the Patient, the Degree of Corruption in the Blood, the Number and Violence of other Symptoms that accompany it; nay, sometimes it will be plainly incurable, unless you proceed to Amputation; and the Strength of the Patient being exhausted, he dies tabid. *Prognosis.*

XI. There are two Methods of treating a *Spina Ventosa*. One is suited to the two Degrees of the Disorder, which we described above, as the milder State: The other to the most violent or third Degree, when the Bones, with the Parts surrounding them, are entirely corroded and destroyed. The best Method that ever I could find for treating the slighter Degrees of this Disorder, is the following. (1.) If the Patient is an Adult, endeavour to correct the Acrimony of his Blood, by prescribing him a Decoction of the Woods, *sc. ex Rad. Sarsaparillæ, Chinæ, Scorzoneræ, Ligno Sassafras, Guaiaci, Juniperi.* Let him drink largely of this every Morning in Bed, as warm as you usually drink Tea or Coffee, giving him from eight Ounces to twelve Ounces at a Time, according to his Strength. In the first Draught let him take *Essent. Lignorum, vel Pimpinell. ad Gts.* 50. *vel* 60. or some other Drops of the same Intention, endeavouring to raise a gentle Sweat. These Medicines will penetrate into the finest Vessels, and even into the bony Fibres, and drive out the noxious Humours, or correct them, greatly promoting the Digestion and Dispersion of stagnating Fluids and Tumors. (2.) This Intention will be greatly forwarded by fumigating the affected Parts with the Steam from Decoctions of resolving or aromatic Herbs. (3.) In the intermediate Times let the Part be rubbed twice in a Day with *Unguentum Mercuriale*, covering it afterwards with the *Emplastrum Mercuriale*. (4.) At the same Time it will be proper to prescribe mercurial Remedies internally, to weak Persons but once, to robust Habits oftner, so as to raise a gentle Salivation: This must be put in Practice, or omitted, according to the Degree *Cure of the Spina Ventosa.*

Degree of the Diſorder, and the Strength of the Patient. I am fully ſatisfied by Experience, that no Good is to be done in this Caſe without the Aſſiſtance of mercurial Remedies, which makes it very ſuſpicious that this Diſorder proceeds from a venereal Taint, or has ſomething very near a kin for its Cauſe. By diligently purſuing this Courſe for ſeveral Weeks (for it will not preſently gain Ground) the firſt and even ſecond Stage of this Diſeaſe, where you have bony Tumors formed, may be cured, and the Tumors diſperſed, or at leaſt brought to that State, that they will not increaſe, but remain as they are, without bringing on any Pain, or other remarkable Inconveniency. This I have frequently ſeen, where I could by no means diſperſe them: Eſpecially where the Patient is regular and moderate in his Diet, living upon ſoft Broths inſtead of ſolid Meats, and drinking the ſmall Runnings of the aforementioned Decoction for his common Drink; or, inſtead of that, the *Decoctum Cornu Cervi*, *Hordei*, *Avenæ*, or any other thin aqueous Liquors.

Cure of the *Pædarthrocaces.*

XII. The ſame Method muſt be uſed in treating the *Pædarthrocaces*, whether attended with Pain or not; giving frequently, at proper Intervals, gently opening Medicines with ſmall Quantities of *Mercurius Dulcis*. If this Diſorder is accompanied with the Rickets, you muſt adminiſter Medicines adapted to this Complaint, and adviſe frequent Exerciſe.

Cure of a bad *Spina Ventoſa.*

XIII. If either of theſe Diſorders ſhould be ſo far advanced, as to be out of the Reach of the Remedies we have already adviſed; if the Pain and bony Tumors increaſe, Abſceſſes are forming, and you have great Reaſon to fear the entire Deſtruction of the Bone: if the Abſceſs does not burſt of itſelf, you muſt not ſtay for its Maturation, but lay the Bone bare with your Knife in the moſt proper Place, which is generally the moſt painful, and deſcending Part, or where it is already burſt. If the Opening is too narrow, you muſt enlarge it; if your Patient dreads the Knife, make your Opening with a Cauſtic, and afterwards make ſeveral [a] ſmall *Foramina* in the Bone with the ſmall Piercer, *Plate* VII. *Fig.* 2. or *Fig.* 7. A. or *Plate* XIV. *Fig.* 8. You muſt pierce down to the Medulla, that there may be Room for a Diſcharge of the confined Matter. But where theſe *Foramina* are not ſufficient for the Diſcharge, you muſt apply a larger Piercer, which they call the Trepan [b], if the Bone will admit of it with Safety; which will not only make greater Room for the Diſcharge of the corrupted Matter, but you will alſo be able to apply your Medicines more conveniently to the Part. Whilſt you are proceeding in this Manner, you muſt inſiſt upon the internal Uſe of the Eſſence and Decoction of the Woods, with antimonial and mild mercurial Medicines: Externally you muſt treat the Ulcer with cleanſing and balſamic Applications, ſuch as *Decoctum Agrimon. Saniculæ, Hyperici vel Ariſtolochiæ, cum Melle Roſar. & Eſſent. Myrrhæ ac Aloes*, which ſhould be injected with a Syringe twice every Day; or a Solution of *Mercurius Dulcis in Aquâ Plantag. vel Aquâ Calcis.* Afterwards you may dreſs with the Eſſences we have juſt mentioned, or *cum Eſſent. Maſtichis aut Succini*, ſpread upon Lint, covering all with a mercurial Plaſter, or with any other that you ſhall think more convenient. This Method is to be continued till the Parts heal.

[a] This has been adviſed by CELSUS, PAREUS, SEVERINUS, SENNERTUS, MARCHETTUS. See MERCKLINI *Not.* p. 483, ſeq. [b] CELSUS has recommended this Method, *Lib.* viii. *Cap.* 2 & 3, and HEYNE, *Lib. de Oſſ. Morb.* p. 68. and PETIT, *Lib. de Morb. Oſſ. Cap. de Exoſtoſi:* And BOERHAAVE, *in Aphoriſm. practic.*

heal. Sometimes the actual Cautery may be uſed to Advantage in this Caſe, to root out the Diſorder, eſpecially when it is only between the [a] *Lamellæ* of the Bone. Raſping or Scraping ſeems to me to be much better ſuited to the *Caries* of the Bone, than to the *Spina Ventoſa*.

Cure of the laſt Stage.

XIV. But when Things are ſtill worſe, and all the Remedies we have hitherto recommended are of no Effect; when the Part is already too much corroded and deſtroyed, ſo that there are no Hopes left of ſaving it, or indeed of ſaving the Patient, but by amputating the diſeaſed Part; you muſt determine on the Operation, which is to be conſidered in two Lights, according to the Difference of the Parts affected. 1. When the Diſorder is ſituated on the ſmall Bones, as on the *Carpus*, *Tarſus*, *Metacarpus*, or *Metatarſus*, or even on the Finger; it will not always be neceſſary to take off the whole Member, that is to ſay, the Finger, Foot, or Hand: But it will frequently ſuffice to remove the corrupted Bone alone. For Inſtance, when the laſt or middle Bone of one of the Fingers has been diſeaſed, I have taken out the foul Bone, and left the ſound Part of the Finger remaining. When the Metatarſal Bone, that ſupports the great Toe, has been diſeaſed [b], I have removed the corrupted Parts from the ſound, and ſaved the Toe. This I did in a Boy of ten Years of Age, and he recovered ſo well, that he walked afterwards as well as before [c]. Where the whole Finger, or only the firſt Bone, has been foul, I have taken off whole Fingers and Thumbs.

When Part only of a large Bone is foul.

XV. In larger Bones, when the whole Bone is not affected, but only a Portion of its external Surface is diſordered with a *Caries* or *Spina Ventoſa*, you muſt by no means take off the whole Limb, but remove that Part of the Bone only which is affected, in the ſame Manner as we taught in the foregoing Chapter on the *Caries*, *Sect.* XVI. But when a large Bone, as the *Os Humeri*, *Tibia*, or *Femur*, or entire Joint of the Arm, Knee, or Foot, is diſeaſed, there is no Remedy but Amputation; making your Wound in the ſound Parts above all that is diſeaſed: But we ſhall treat more fully of this Subject when we write on Chirurgical Operations.

When the whole Bone or Joint is diſeaſed.

Where the Tumor of the Bone will not give way.

XVI. In certain Species of the *Spina Ventoſa*, where the Tumor of the Bone will not yield to the Application which we have adviſed above, and you can come at it with your Hands, PETIT [d] adviſes you to lay the Bone bare by a cruciform Inciſion, and to cut off the extreme Parts of the four Angles of the Skin: And when this is done, to dreſs with dry Lint. On the Day following you are to bore ſeveral Holes in the Tumor, ſo near each other, that it may be pierced like a Sieve: You are then to take the whole off with a Chiſſel and Mallet. The Wound is to be filled with dry Lint: And that the diſeaſed Parts may ſeparate the ſooner from the ſound, he orders the foul Part of the Bone to be dreſſed with a Solution of Mercury in *Aqua Fortis*. This Method is to be continued till you have obtained an Exfoliation. He is very high in the Commendation of this Proceſs, and I think deſervedly prefers it to any other Remedy

[a] SEVERINUS appears to be too fond of the actual Cautery in theſe Caſes, *Cap.* 20. for frequently we cannot get to the Bottom with it, or the Parts are too much corrupted to expect Advantage from it. [b] LE DRAN, in *Obſ.* 112. recites nearly the ſame Caſe, where he took off the *Metatarſus*, Toe, and all; but this ſhould conſtantly be avoided where it is poſſible, for the Toe is of great Advantage in walking.

[c] See in SCULTET. *Obſ.* 90. the Caſe of a Thumb and Hand taken off for a *Spina Ventoſa*.

[d] *Lib. de Morb. Oſſ.* Cap. *de Carie*.

medy in these Cases, even to the actual Cautery, where the Caries has not penetrated too deep.

How to remove an *Exostosis*.

XVII. When an acute Eminence or Excrescence, which is properly called an *Exostosis*, pushes preternaturally above the Bone, creating no Disturbance, Pain, or Deformity, and unaccompanied with *Caries* or *Spina Ventosa*, as I have frequently seen them; in my Judgment it is best to let it alone: For the Remedy will be worse than the Disease, and, by laying the Bone bare, you may bring on a *Caries* or other Inconveniencies. On the other hand, if it occasions any Deformity, impedes any Action, or produces Pain or other Mischiefs, you may take it off in the Manner we have just taught above. You may see various Cases of *Caries*, *Spina Ventosa*, and *Exostosis*, in the Figures of that splendid Work, CHESELDEN's *Osteography*, from *Plate* XLI. to the End: In RUYSCH, *Obs.* p. 94. in his *Thesaur. Anatom.* VIII. *Tab.* 3. and *Thesaur.* X. *Tab.* II. and BIDLOO's *Oper. Anatom. Chirurg.* p. 208. *Tab.* II.

CHAP. X.

Of ULCERS *of the* HEAD.

What these Disorders are.

I. IT remains with me now to say something of Ulcers of the Head, and particularly of those which occupy its hairy Part, and are at this Time called either *Tinea*, *Favus*, or *Achores*: But the Professors of Medicine do not at all agree about the Signification of these Terms. By the Term *Favus*, we commonly understand Ulcers of the Head, that are full of Cavities like a Honey Comb. By *Achores*, those Ulcers which are full of small *Foramina*, which contain a moderately viscid Humour. Many call these Disorders *Tinea*, because, from the Abundance of small *Foramina* in them, they resemble moth-eaten Garments. But for the most Part, the Term *Tinea* at present is applied to a large dry Scab, which Children and Infants are subject to upon the Head, full of thick foul Scales, and very offensive to the Smell: This sometimes extends itself to the Face, in which Case we call it *Crusta Lactea*. This is often benign and of a mild Nature, but sometimes ill-conditioned and dangerous. There is still a worse Kind of *Tinea*, or scabby Head, covering the whole hairy Scalp with an ash-coloured thick Crust, attended with a violent Itching, and stinks grievously: This is generally very difficult of Cure. Persons afflicted with this Complaint, have a very pale unhealthy Countenance. These Disorders are much more frequently met with in Infants and Children than in Adults. They are occasioned either by the Nurse's irregular Course of Life, or by the Child's being used to foul Feeding, from whence foul Blood is made, which produces Ulcers of this Kind. Sometimes they break out in an adult State, resembling a Kind of Leprosy, which is very difficult to cure. In the *Pox* you frequently find both Head and Face, particularly the Forehead, spread with dry Scabs, and scabby Ulcers, which is called a *Venereal Scabies*. Venereal *Gummata* also and *Tophs* of the Head, may be referred to this Class, since they frequently degenerate into Ulcers.

II. Though

II. Though the Ulcers which we have juſt deſcribed, differ from each other in ſome Particulars, yet I ſhall not ſpeak ſeparately of them in this Place, as they are to be cured pretty nearly in the ſame Manner. When they are ſlight, it will be proper to give a gentle Purge now and then, with the Addition of a ſmall Quantity of *Mercurius Dulcis*; adminiſtring between whiles to an adult Patient, Decoctions of the Woods, with edulcorating Pills, Powders, or Eſſences. Infants at the Breaſt may take diaphoretic Powders: But their Nurſes may proſecute this Intention with Powders, Pills, Decoctions, or Eſſences. Externally, you may anoint the Scabs with *Cremor Lactis cum pauca Ceruſſa præparata miſtus*; or with *Oleum Ovorum* alone, or with the Addition of a ſmall Quantity of *Oleum Ceræ*; or with *Unguentum de Enula*, *de Ceruſſa*, *Diapompholygos*, or with any other of the ſame Intention: Obſerving at the ſame Time Regularity in Diet, &c., and defending the Body from the Injuries of the external Air. By this Method, not only Ulcers of the milder Kind are healed, but even thoſe of the more malignant Sort; eſpecially if you give ſmall Quantities of *Mercurius Dulcis* at the ſame Time, or mix *Mercurius Vivus* with your Ointments. But theſe Medicines are to be uſed with Caution. Cure.

III. In the worſe Degrees of this Diſorder, eſpecially where you cannot be perſuaded to uſe Mercurials, you will never ſucceed in your Cure, till you have taken off all the Hair, with which theſe Ulcers have a ſtrong Connection. In ſome Places it is the common Practice to pull out the Hair by the Roots, either by Degrees, or at once, with a *Pitch Plaſter*[a], which is ſpread upon a ſtrong Cloth, or upon Leather, and applied all over the Head, after the Hair has been cut off as far as the Scabs. When it has taken faſt hold, they let it lie on for twelve or twenty-four Hours, and then they tear it off at once, and it brings away with it both the ſcabby Cruſt and the Roots of the Hair: But this cannot be done without great Pain and Effuſion of Blood. When the Plaſter is torn off, they wipe away the Blood with dry Lint, and anoint the Head with ſome *Oleum Laterinum*, with the Addition of a little *Oleum Ceræ* warmed, and cover it with the *Emplaſtrum de Spermate Ranarum pauca Camphora impregnatum*; dreſſing in this Manner every Day, till the injured Parts are clean, and then they heal with *Oleum Ovorum vel Eſſentia Succini*[b]. They preſcribe internal Medicines to correct the Blood, ſuch as you ſaw in *Sect.* II. and adviſe Regularity in Diet. Antimony either alone, or mixt with a ſmall Quantity of *Flores Sulphuris*, is very ſerviceable in this Caſe. You ſhould diligently avoid beginning with the Uſe of Mercurial or Sulphureous Ointments; becauſe they are very apt to repel the noxious Humours, and endanger the Life of the Patient: which Effect they are not obſerved to have, after you have adminiſtred Cleanſers of the Blood for ſome Time internally. Another Method.

IV. In ſcabby Ulcers of the Face, which happen in the Infant State, and are vulgarly called *Cruſta Lactea* or *Achores*, the ſame evacuating and corrective Medicines are to be preſcribed for the Nurſes, which we ordered above, *Sect.* II. The Infants themſelves alſo ſhould be purged frequently, and in the Intervals between Cure of the *Cruſta Lactea*.

[a] JUNCKER in his *Surgery*, p. 280. recommends a Plaſter of Pitch, Scammony, and Reſin.

[b] WEDEL tells us, that the *Tinea* may be cured, and Vermin in the Head at the ſame Time deſtroyed, by waſhing the Head over with liquid Pitch; applying previouſly internal Medicines. *Lib. de Morb. Infant.* p. 61.

between purging, ſhould take diaphoretic Powders prepared *ex Antimonio Diaphoretico, Lapid. Cancrorum, Antimonio crudo, & Flor. Sulphuris.* When they have taken theſe Medicines for ſome Time, you may daub the ſcabby Parts with a Liniment made *ex Cremore Lactis cum Cretâ vel Ceruſſâ*; or in the room of this you may uſe *Oleum Ovorum cum pauco Olei Laterini.* Ointments prepared of Mercury or Sulphur are very dangerous in the Beginning of this Diſorder, or to very weakly Infants. But if Remedies of this Kind ſhould be uſed by unſkilful Perſons, which is frequently the Caſe, to the Detriment of the Patient, you muſt endeavour to ſtrike the Humours out again by preſcribing Sudorifics in different Forms, both to the Infant and its Nurſe, till you have ſatisfied this Intention.

End of the FIRST PART.

INSTITUTIONS OF SURGERY.

THE PREFACE.

HAVING finiſhed the firſt Part of our Inſtitutions of Surgery, which treats profeſſedly of the Five Kinds of Diſorders of the Human Body, which require the Aſſiſtance of the Surgeon; to wit, Wounds, Fractures, Luxations, Tumors, *and* Ulcers; *we ſhall proceed now to the Second Part, which is dedicated to* Chirurgical Operations. *And in this Volume we ſhall take an Opportunity to treat of ſuch Diſorders as remain undeſcribed, either as not properly belonging to any of the foregoing Heads, or ſuch as require particular Contrivances and Machines to be made uſe of in their Cure. In doing this we ſhall conſult Order as far as the Nature of the Subject will admit of it. We ſhall firſt deſcribe thoſe Operations, which may be performed in almoſt all, or at leaſt in various Parts of the Body; as opening a Vein, making Iſſues, applying the actual Cautery, taking off Excreſcences or intire Parts of the Body. We ſhall then proceed to thoſe which have their proper Situations, and happen each to one particular Part of the Body. In performing this Part of our Work, we ſhall begin with thoſe which belong to the Head and each of its Parts, as the Cranium, Eye-lids, Eyes, Ears, Noſe, Lips, Teeth, Gums, Tongue, Palate, Tonſils, Uvula, &c. Then we ſhall deſcribe thoſe Ope-*

rations, which are accommodated to Diſorders of the Neck: From thence we ſhall proceed to the Breaſt, ſo on to the Abdomen, and its neighbouring Parts, to wit, the Anus and Pudenda of both Sexes. Laſtly, we ſhall deſcribe thoſe Operations which are performed on the upper and lower Extremities. Notwithſtanding the great Number of theſe Operations, and the various Methods of performing them, will render this Taſk extremely difficult, yet it ſhall be our principal Care to explain the Nature of each particular Operation, the beſt Method of performing it, and the fitteſt Inſtruments to be made uſe of for that Purpoſe, with all the Clearneſs that the Subject will admit of. By purſuing this Method, we ſhall not only teach the young Beginner the firſt and ſolid Principles of Surgery; but the Surgeon alſo who has already had ſome Experience in his Profeſſion, will, I hope, find ſomething in theſe Inſtitutions, by which he may, in ſome meaſure at leaſt, perfect and adorn his Art.

INSTI-

INSTITUTIONS OF SURGERY.

PART II.

Of OPERATIONS.

SECT. I.

Of GENERAL OPERATIONS *practicable in ſeveral different Parts of the Body.*

CHAP. I.

Of PHLEBOTOMY *in general.*

I. WE begin with the Operation of *Phlebotomy:* Becauſe it is of all the moſt general, performed in moſt Parts of the Body, and by much the moſt frequent in uſe at this preſent Day. By Phlebotomy or Bleeding we here intend the opening a Vein, by a ſharp-edged and pointed Inſtrument of Steel, for extracting a proper Quantity of Blood, either for the Preſervation or Recovery of a Perſon's Health. Phlebotomy what.

II. Veneſection appears to be not only one of the moſt uſeful, but moſt antient Operations in Surgery: Since we find, by the Writings of HIPPOCRATES, CELSUS, and others, that it was even celebrated near three thouſand Years ago. Yet there have not been wanting ſome among the Antients, and Moderns, who have reviled this Practice, as both cruel and fatal to the Healths and Lives of Mankind, as ERASISTRATUS, PARACELSUS, HELMONT, PORTIUS, BONTEKOE, GEHEMA, *&c.* But I think all their Objections too weakly founded to need any Refutation; which might very well be made even only from the daily Experience we have of the great Uſefulneſs of this Operation, in alleviating, preventing, and curing moſt Diſorders of the human Body, eſpecially A moſt antient and uſeful Operation.

thoſe of the acute and inflammatory Kind. The Operation is ſaid to have been firſt hinted to us by the *Hippopotamus*, who, at ſtated Seaſons, uſed to open a Vein with a ſharp-pointed Reed, according to POLYDORE VIRGIL, *De Rer. Inventor.* p. m. 65.

Phlebotomy often difficult.

III. Nor is the Operation in many Caſes practicable with ſo much Eaſe and Safety as is commonly imagined. For though in ſome Patients the Veins lie ſo open and conſpicuous, that even a Novice will find no Difficulty in making their Apertion; yet in others they are either ſo ſmall or deeply ſituated, that the moſt expert Surgeon is ſometimes at a Loſs, and may, by Accident, miſcarry. Add to this, that as the Arteries, Nerves, and Tendons, are frequently very nearly ſeated to the Veins, it is no uneaſy Matter to injure one or other of them with the Inſtrument uſed in Bleeding: which is quickly followed either with a profuſe or fatal Hæmorrhage, an Aneuriſm, violent Pains, Inflammation, Fever, Mortification, or even Death. Phlebotomy therefore ſhould be performed with no leſs Judgment and Caution, than the other important Operations in Surgery: eſpecially as the Reputation of a young Surgeon may ſuffer as much by Neglect or Accidents in this Way, as in many of the other leſs uſual and ſeemingly more difficult Operations.

Qualifications of the Phlebotomiſt.

IV. A good Phlebotomiſt ſhould have a ſteady, nimble, and active Hand, with a ſharp Eye, and undaunted Mind; without which he may either be liable to miſs the Vein, or commit ſome Error that may be injurious or fatal to the Patient and his own Character. For theſe Reaſons it is that Veneſection is leſs readily practiſed by the Surgeon, as he advances in Years: Becauſe old Age is generally accompanied with a weak Eye and a trembling Hand.

Inſtrument for Bleeding.

V. The Inſtrument which is in common Uſe amongſt the Surgeons for opening a Vein is called a *Lancet*. The Shape of this Inſtrument is deſcribed at *Plate* I. A. and at *Plate* XI. *Fig.* 5. The Surgeon ſhould take care to be always provided with a ſufficient Number of theſe, and to have them conſtantly in order, and to have ſome alſo of a larger Size. Thus he will be prepared for Veins in different Subjects. And as this is an Operation that frequently requires to be performed on a ſudden, he will never be at a Loſs. There are many Surgeons in *Germany*, particularly in *Franconia*, *Bavaria*, and *Lower Saxony*, who bleed with a *Fleam*, *Plate* XI. *Fig.* 3. which they uſe in this Manner. They hold one of their Fingers upon the Part B, and applying the Point A to the Vein, they ſtrike the Part C with one of the Fingers of the other Hand, opening the Vein as Farriers do in Horſes. Some of the Surgeons and Bagnio-Men uſe a neater Inſtrument, an *Elaſtic* or *Spring Fleam*, which the *Germans* call *Schnapper*, or *Schnapperlein*, *Fig.* 4. When they have drawn it up, they apply the Point A to the Vein, and then let it go by preſſing upon B. Some again uſe a Lancet in the Form of a Dart, the Figure of which you may ſee in CRONE *de Veneſectione*, p. 33. *Fig.* 4. But ſince the Poſition and Size of the Veins is different in different Subjects, we find that the moſt convenient Inſtrument for our Purpoſe is the *French* Lancet: Though many of our Surgeons are very expert in the Uſe of the *German* Lancet, *Fig.* 3 and 4.

In what Part the Operation is to take place.

VI. Though the Operation of Bleeding is frequently performed in different Parts of the Body, as in the Hand, Foot, Forehead, Temples, Neck, Tongue, Penis, and other Parts; yet it is moſt generally performed in that Vein of the Arm which lies near the Joint of the *Cubit*. Therefore we ſhall begin with teaching

teaching the Method of opening this Vein, and treat more fully of it than of any other.

CHAP. II.

Of OPENING *the* VEINS *of the* ARM.

I. IT is commonly enough known, that the Operation of Bleeding in the Arm is performed on the Veins that lie on the internal Part of the *Cubit*. There are several Things worthy the Surgeon's Notice in this Operation: Some of which regard the Things which are to be done preparatory to Bleeding; some in the Operation itself; others immediately after the Performance of it. Of each of which we shall speak distinctly in their Order. Preparatory to Bleeding you should have in readiness, (1.) *a Linen Fillet*, about a *Paris* Ell in Length, and two Fingers in Breadth, with or without small Strings fastened at each End of it. (2.) Two *small square Bolsters*. (3.) *Porringers* or Vessels to receive the Blood. (4.) *A Sponge* with warm Water. (5.) Some *Vinegar*, *Wine*, or *Hungary Water*, to raise the Patient's Spirits if he should be inclinable to faint. (6.) *Two Assistants*, who must be void of Fear, one to hold the Porringer, the other to reach you any Thing that you shall want. (7.) *A small Wax Candle*, when the Patient is to be blooded at Night, or in a dark Place. (8.) You must place your Patient upon a *Couch*; or, if he is very fearful of the Operation, lay him upon a Bed, lest he should fall into a Swoon. (9.) Lastly, you should take care that no Hair or the Cloaths of the Patient lie in your Way. The Patient himself should take care that nothing should give him any Concern: And he should avoid terrifying himself with recollecting the Mischiefs which have happened by the unskilful Performance of this Operation. Lastly, the Operator should be as expert in bleeding with his left Hand as with his right. For, as you are readier at bleeding in the right Arm with your right Hand, so when you are to open the Veins of the left Arm, you will find it necessary to use your left Hand: And there are some Patients who insist upon being blooded in the left Arm. Preparation for Bleeding.

II. Though the Operation is to be performed at once, with one Puncture, yet many Things are to be observed in order to render it successful. First, it is necessary for the Surgeon to inspect his Patient's Arm diligently, that he may see the Course of the Veins: He must then take hold of the Arm, and extend it towards his Breast, tucking up the Sleeve about a Hand's Breadth above the Bend of the *Cubit*, where he must make his Ligature, rolling the Fillet twice round, and fastening it with a Knot (*Plate* XI. *Fig.* I. D.) The Veins being compressed, and the Blood being stopped in its Return, they will enlarge, and lie fairer to the Eye. The Ligature generally used upon these Occasions is a Slip of fine Scarlet Cloth, but any other Colour will answer the Purpose as well. When you have bound up the Arm in this Manner, you let it go for a small Time till you have taken a Lancet out of your Case: And opening it so that it may make a Sort of an obtuse Angle, you take hold of it with your Teeth about the Joint (A. *Plate* XI. *Fig.* 5.) and hold it some Time till the Veins grow turgid. You are then to lay hold of the Arm again in the same Manner as What is to be done in the Operation.

as we directed before, and extend it to your Breaſt, having an Aſſiſtant ready with the Veſſel in his Hand, at a convenient Diſtance for receiving the Blood.

What Vein is uſually opened.

III. You are now to examine which Vein lies faireſt, and is therefore moſt proper to be opened. For you muſt obſerve that in the Arm there uſually appear three principal Veins: The firſt is called *Vena Cephalica*, and is found in the external Part of the Arm. *See Plate* XI. *Fig.* 1. A. The Second is termed *Baſilica*, and lies on the internal Part of the Arin: In the right Arm it is alſo called *Hepatica*; in the left, *Splenetica*, *See ibid. Letter* B. The Third, which is obliquely ſituated between the former two, is called *Mediana*. *See Letter* C. The Median and Baſilic Veins, as they are larger than the Cephalic, diſcharge a greater Quantity of Blood, but are attended with more Danger in the Operation: For a conſiderable Artery and the Brachial Nerve lie under the Baſilic Vein, and the Tendon of the Biceps Muſcle under the Median. But as they lie fairer to the Eye, and are therefore more frequently the Subjects of the Operation we are treating of, than the Cephalic Vein, it is ſafer and more eligible for the leſs experienced Surgeons to open the Baſilic, or at leaſt the Median Vein. But ſometimes the Veins are ſo ſituated in the Arm, that only one of them will lie expoſed to View, which deprives you of all Choice. Your only Safety in this Caſe depends upon your Choice of a ſkilful and cautious Surgeon.

In what Part the Vein is to be opened.

IV. When you have determined which Vein to open, you are to perform the Operation on that Part which preſents itſelf faireſt to you. But if the Vein has frequently been opened, and the Part which appears largeſt and faireſt is full of Cicatrices, you are not to open above, but below the Cicatrices, by which means the Blood will diſcharge itſelf more freely: For the Part above is generally ſtraitened by the Cicatrix. For this Reaſon, whenever you open a Vein for the firſt Time, begin as high as you can, by which means you will have the more Room to deſcend in repeated Bleedings.

What is to be done immediatcly before Bleeding.

V. Before you apply the Lancet to the Skin, when the Veins are not riſen it will be proper to rub the Arm below the Bandage, which will drive the Blood back towards the Cubit, and render the Veins more turgid. Whilſt this is doing in the right Arm, the Surgeon ſhould take hold of the Patient's Arm in ſuch a Manner that he may lay his Thumb upon the Vein which he intends to open, to prevent the Blood from flowing back, and to keep the Vein from rolling. You are now to fix your Eye upon that Part of the Vein which you intend to open, and taking the Lancet out of your Mouth with your right Hand, ſo placed that the Thumb and firſt Finger may be fixed about the Middle of the Blade: The other Fingers ſhould reſt gently upon the Patient's Arm, to prevent your Hand from ſlipping.

How the Puncture is to be made.

VI. Your Lancet is now to be puſhed lightly and carefully forward by your Thumb and fore-finger, till it has penetrated through the Coats of the Vein; and at that Inſtant to be raiſed a little upwards in order to enlarge the Orifice of the Wound, which will give a freer Paſſage to the Blood. The moſt common and convenient Size of an Orifice is about twice the Breadth of the Back of an Ordinary Knife. You are to keep even between the two Extremes of Raſhneſs and Timidity in making the Puncture. For as in one Caſe you will only divide the common Integuments, and ſo leave your Work undone: So in the other you will run the Riſque of wounding the Artery, Nerve, or Tendon.

The

The Vein may be opened in three Directions. Some open itin a ſtraight Line, *Plate* XJ. *Fig. 2. Letter* A. Others tranſverſely *B:* But moſt Surgeons make an oblique Wound *C. D.* If the Vein is to be opened in the left Arm, the Surgeon muſt change Hands, and do all with his left Hand, which we have directed above to be done with the Right. If you are to bleed with the *German Fleam*, place the Point *A* upon the Vein; and taking hold of the Extremity *B* with your left Hand, drive the Point of the Fleam into the Vein by a Stroke with one of the Fingers of your Right Hand. But if you will phlebotomiſe with the Spring-Fleam, *Fig.* 4. you cock it, by elevating the Hook *c*; and placing the Point *A* upon the Vein, by letting looſe the Spring, it is by a gentle Preſſure plunged into the Veſſel.

VII. Your Aperture being thus made, and the Inſtrument drawn inſtantly back, the Blood will then ruſh forth from the Orifice either in a large or ſmall Stream. Hereupon your Inſtrument muſt be depoſited in the Baſon or Diſh, and not thrown upon the Bed, leſt it ſhould be loſt, or elſe injure the Patient. In the mean time the Blood muſt be permitted to flow as long as it ſhall be judged uſeful or neceſſary: and if it ſhould ſtop too ſoon, as it often may from too great a Stricture of the Bandage on the Arm, it muſt be ſlackened a little, by which means the compreſſed Artery being ſet at Liberty, the Blood will flow from the Orifice as at firſt. If you find the Orifice obſtructed by too great a Tenſion of the Skin, or an Intruſion of the *Membrana Adipoſa*, you ought in that Caſe to return the Bit of Fat, by preſſing with the Finger or a warm Sponge, and to relax the Skin by bending the Arm a little. Laſtly, if the Orifice be obſtructed by thick, grumous, or congealed Blood, that Impediment may be removed by wiping it with a Sponge dipt in warm Water. Treatment after Aperation.

VIII. But that the Patient's Arm may not become painful or languid, by holding it long extended, the Surgeon ſhould ſupport it by the *Cubitus* for a little while; and then give him a Stick, or other cylindric Body, to turn round in his Hand, that by the Contractions of the Flexor and Extenſor Muſcles of the Fingers, the Courſe of the Blood may be accelerated towards the *Cubitus:* which will be ſtill further promoted, if the Patient urges a little voluntary Cough. In the mean time his Attendants ſhould ſtand ready with other empty Cups or Veſſels for receiving the Blood, to carry off ſuch as are full, and adminiſter the Dreſſings for the Deligation, with cordial Water, and other ſuch Neceſſaries. Method of cloſing and binding up the Orifices.

IX. The Quantity of Blood neceſſary to be taken from the Vein at one Bleeding, muſt be determined by the Phyſician, from conſidering the Patient's Diſorder, Strength, Habit, and other Circumſtances. But when the Surgeon attends his Patient without a Phyſician, he may then ſafely proportion this Evacuation himſelf at his own Diſcretion, by reflecting on the Nature of the Patient's Caſe, his Age, Strength, Courſe of Life, and Fulneſs of Habit, *&c.* for he may permit the Patient, that ſhews no Paleneſs of Countenance, nor Diminution of Strength or Spirits, to bleed longer than thoſe that quickly grow faint, *&c.* What muſt be done by the Patient and his Attendants.

X. When there ſeems to be a ſufficient Quantity of Blood diſcharged, the Ligature muſt then be immediately taken off from above the Elbow, and the Skin about the Orifice muſt next be gently ſtroaked or preſſed together by the two fore-fingers of the left-hand: By which means the Lips of the divided The Quantity of Blood to be taken.

Vein are more eaſily compreſſed and cloſed. But while the Surgeon is doing this with his left Hand, he takes the ſmalleſt of the two Compreſſes brought to him by the Servant, and applies it upon the Inciſion with his Right-hand: but ſo as to let what little Blood may remain betwixt the Orifice and the Vein, be diſcharged, before he impoſes the Compreſs. Over the firſt or ſmall Compreſs he ſhould impoſe another that is a little larger, preſſing them both gently on the Orifice with his left Thumb, till the Bandage is laid acroſs. But before the Deligation is performed, according to the Directions we ſhall give for that Purpoſe in the laſt Part of our Surgery, on Bandages, it will be a Piece of Neatneſs and Decency in the Operator, to wipe off what Blood may have adhered to the Arm with a wet Sponge or Napkin, and then to go on with his Bandage. There are indeed many Surgeons who apply but one Compreſs, which they firſt wet in Water, Vinegar, Wine, or its Spirit: Though, in my Opinion, two Compreſſes make the Deligation more firm and ſecure: Though I think it is no great Matter whether they are applied wet or dry; the dry will ſit eaſieſt on the Part.

Treatment after Deligation.

XI. Having applied your Bandage, and drawn down the Patient's Sleeve over his Arm; he ſhould be ordered not to uſe it too early or violently, before the Orifice is well cloſed, which might excite a freſh Hæmorrhage, an Inflammation, Suppuration, or other bad Accident. And if the Patient ſhould faint away ſoon after the Operation, it may be then convenient to wet his Noſtrils with *Hungary* Water or Vinegar, and to ſprinkle ſome of the laſt, or elſe cold Water, in his Face: and eſpecially in Summer-time to let in the freſh and cool Air, by opening the Windows, *&c.* Alſo, if any Wine or cordial Water be at hand, you may give the languiſhing Patient a ſmall Draught thereof; and then the Surgeon will have nothing more to do than waſh his Hands, and the Inſtrument, before he puts it up in his Caſe.

Judgment paſſed on the Blood.

XII. In the next place, it is often cuſtomary to aſk the Opinion of the Surgeon or Phyſician preſent, concerning the healthy or morbid State of the Blood, from its external Appearance. In this Caſe the Surgeon ſhould always make a good Preſage to his Patient and By-ſtanders, even though the Blood ſhould appear bad: for it is not eaſy to expreſs the good Effects that may follow from chearing up the Patient's Mind, which is much better than to leave a heavy Impreſſion on it by a ſevere Prognoſtic. Therefore, if the Blood appear florid, the Surgeon ſhould declare it a Sign that the Patient either is, or ſpeedily will be in good Health. If the Blood appears vitiated, or of a bad Colour, he muſt then pronounce the Bleeding will be extremely ſerviceable to him. If the Patient ſhould, in the mean time, be in a Swoon; the Surgeon ſhould take occaſion even from thence to ſignify the great and ſpeedy Effect the Diſcharge will have towards the Recovery of the Patient's Health. When ſuch or the like encouraging Diſcourſe has been paſſed, the Blood ſhould be ſet by in a cool Place, till the Phyſician or Surgeon renews his Viſit.

Whether the Patient may ſafely drink or ſleep after Bleeding.

XIII. If the Patient ſhould be thirſty after Bleeding, you ought not to deny him the Pleaſure of drinking, eſpecially thin Liquors. The *French* make it a Cuſtom to give the Patient a large Draught of Cold Water after Phlebotomy in inflammatory Diſorders: in which Caſes, if the Patient be of a warm Habit, that Practice may be extremely beneficial. But in cold and weak Habits, it ought not to be encouraged: for then it will be better to give ſome warm Sup-

Suppings of Tea, Coffee, or the like. If any body ſhould propoſe the Queſtion, Whether the Patient may ſleep ſafely after his Bleeding, your Anſwer may be either in the Affirmative or Negative, according to particular Circumſtances. If the Evacuation was made by way of Prevention, or to preſerve the Body in its healthy State, it will be more adviſeable for the Patient to ſhake off his ſleepy Diſpoſition by walking, or engaging in ſome agreeable Sport or Exerciſe: becauſe if Sleep be indulged, the Bandage may get looſe, or ſlip up above the Orifice, and ſometimes thereby occaſion a profuſe and dangerous Hæmorrhage. This Objection ought not, however, to deprive the Patient of a comfortable Repoſe, in caſe of great Weakneſs and Indiſpoſition; eſpecially if he has had no Sleep for a long time before: then it would be denying him a Benefit perhaps greater than the Remedy of Bleeding itſelf. But for the greater Security, it may not be amiſs to let the Nurſe, or ſome body, have a watchful Eye over the Patient during his Repoſe; that in caſe of ſuch an Accident, timely Relief may be had by compreſſing the Vein with one's Finger till the Surgeon can be called.

XIV. When the Surgeon or Phyſician comes again to viſit the Patient, the Blood is uſually ſet out again to have a freſh Judgment paſſed upon it: in which Caſe the Verdict given ought to be ſuch as will exhilarate the Patient, and not depreſs his Spirits, agreeable to what we ſaid before on this Head at *Sect.* XII. The Surgeon muſt in the next place inſpect the Deligation, to ſee if the Bandage be too looſe: and in taking it off, if the Compreſs adheres to the Lips of the Orifice, he ought not to force it away, but to apply his Bandage over it again as before. After waiting a Day or two longer, it will ſpontaneouſly ſeparate, or fall off from the cloſed Orifice, which will by that time be near cicatrized. There are ſome, who being prejudiced in favour of the enthuſiaſtic Doctrine of Sympathy, will have their Blood run into cold Water, or have cold Water poured upon it, in febrile Complaints, thinking by that means to allay the Heat of the Blood. In this Reſpect it may be of ſervice to humour and ſatisfy their Minds, though there may be nothing in the thing itſelf. Behaviour in your After-viſits.

CHAP. III.

Of PHLEBOTOMY *in the* HAND.

I. THERE are two principal Veins in the Hands, which with us in *Germany* are ſometimes opened to bleed the Patient. The one is called *Salvatella*, and runs on the Outerſide of the Back of the Hand towards the Little Finger, being ſometimes denominated *Splenica* by the Antients, who judged its Apertion extremely uſeful in Melancholy, and Diſorders of the Spleen. The other Vein, which is termed *Cephalica*, runs betwixt the Thumb and Forefinger, and was formerly ſo denominated from an Imagination, that bleeding from it was more particularly uſeful than from others in Diſorders of the Head. But we are at preſent convinced thoſe Notions of the Ancients were without Foundation; and though the Patient is bled more difficultly and ſlowly by theſe Veins, yet the Effects will be the ſame as after Phlebotomy in the Arm. Yet it What Veins are opened in the Hands.

it may be ſometimes convenient for the Surgeon to open them either at the particular Requeſt of the Patient; or when the Veins of the Arm are very deeply or obſcurely ſituated, and theſe lie fair and conſpicuous for Inciſion. To which we may add, that Women, in many Parts of *Germany*, pregnant, eſpecially towards the latter End of their Time, generally chooſe to bleed by this Vein, from an Imagination that it occaſions leſs Injury or Weakneſs to the Fœtus.

Method of phlebotomiſing in the Hand.

II. When you are therefore determined from particular Reaſons to phlebotomiſe in the Hand, the Patient muſt firſt hold it in warm Water for ſome time, rubbing it therein well with his other Hand, in order to make the ſmall Veins become turgid and conſpicuous. After this you are to fix a Ligature upon the *Carpus*, that the Veins may continue in that Manner diſtended. When the Hand has been wiped dry with a Napkin, you make an Apertion in the moſt convenient Part of the Vein in the Manner we directed for Veins in the Arm. If the Blood does not flow copiouſly from the Orifice after Inciſion, the Hand ſhould be placed again in hot Water, and taken out when the Diſcharge is judged to have been ſufficient. This done, the Hand is wiped dry with a Napkin, the Orifice defended with two Compreſſes, and your Bandage applied as we ſhall direct in *Part* III. *Chap.* VI. *Sect.* X. on Bandages.

CHAP. IV.

Of BLEEDING *in the* FOOT.

The Veins opened in the Foot.

I. BLEEDING in the Foot is an Operation of very old ſtanding. It having been an Obſervation made by the moſt ancient Phyſicians, that Phlebotomy in this Part proved highly ſerviceable in moſt Diſorders of the Head and Breaſt, and for an Obſtruction of the menſtrual and hæmorrhoidal Flux: upon which Diſcharges greatly depended the healthy State of both Sexes. For theſe Reaſons they therefore denominated thoſe Veins of the Foot, *Saphena* and *Cephalica:* the laſt of which extends itſelf from the internal Ancle to the great Toe; and the firſt, from the external *Malleolus* to the ſmaller Toes. But why one of them ſhould be thought or denominated more *cephalic* than the other, there is not the leaſt Reaſon to be offered; ſince Bleeding from either of them has altogether the very ſame Effect. Therefore, in my Opinion, the Surgeon ſhould always open that which lies faireſt and moſt conſpicuous. But if the Veins upon the *Metatarſus*, or Inſtep of the Foot, do not well appear, it may be then convenient to open one of thoſe at the Ancle, or about the Calf or Ham of the Leg, as I have frequently done myſelf. Nor is the Phlebotomiſt ſo liable to injure any of the Tendons in theſe laſt Parts as he is upon the *Metatarſus*. In the mean time the Operator ſhould, in ſingle Women, expect the Order of ſome prudent Phyſician for his bleeding by theſe Veins: becauſe ſome of them, who are evil-minded, endeavour by this Means to procure a Miſcarriage; which, when known, might make the Phlebotomiſt a Sharer in the ill Report.

II. For the more eaſy and ſucceſsful Apertion of theſe Veins, the Patient muſt firſt waſh both Feet well for ſome time in hot Water: that when the Veins become ſufficiently turgid, the Surgeon may take his Choice of that which preſents faireſt either in the Right or Left Foot, without paying any Deference to the Diſtinction of Right or Left, in any of the forementioned Diſorders. For the Effect, as we obſerved, will be the ſame in either, if they are diſpoſed with equal Advantage for Apertion. Having fixed upon the particular Foot and Vein, your Ligature muſt be applied about two Fingers Breadth above the Ancle; and then the Patient muſt return it into the warm Water, while the Surgeon takes out and prepares his Inſtrument or Lancet. Then kneeling down on one Knee, the Surgeon takes out the Patient's Foot from the warm Water, and having wiped it dry with a Napkin, places it upon his other Knee, or elſe upon a Board laid over the Veſſel of hot Water. He now faſtens or ſecures down the Vein from ſlipping with his Left Hand, as in *Chap.* II. *Sect.* V. *& ſeq.* But if the Veins do not appear well under the Ancles, the Ligature muſt be removed higher, about two Fingers Breadth above where you intend to make the Apertion of the Vein which beſts offers itſelf. 'Tis to be alſo obſerved, with regard to the Surgeon's Poſture, that he may ſeat himſelf on a low Stool or Chair, and place the Patient's Foot in the moſt advantageous Manner upon either Knee. This Method will be preferable to the other in Bleeding with the Spring-fleam, as many do in *Germany:* or the Patient may here ſet the Foot for the Operation upon a low Stool, or any other Support. Manner of opening theſe Veins.

III. The Blood from the Vein thus opened may be received into a Glaſs Cup or Baſon: and if it does not flow freely from the Orifice, the Foot ſhould be returned into the warm Water; which will either prevent or diſſolve the congealing of the Blood that in this Caſe often obſtructs the Aperture. When a ſufficient Quantity of Blood has been thus drawn, which may be known partly from the Time, and partly from the Largeneſs of the Stream, as alſo from the Redneſs of the Water, and Condition or Strength of the Patient; the Orifice is then to be cloſed by the Finger, and, after drying the Foot with a Napkin, to be ſecured by Compreſſes and Bandage. Concerning the Uſefulneſs of Venæſection in the Foot, conſult the Diſſertations of PERDUCIUS, HEREDIA, and STAHLL, who have been oppoſed by HECQUET, *in Lib. ſur la Saignée du Pied.* Pariſ. 1724. The firſt have been again ſeconded by *Jo. Bapt.* SILVA *Medic. Pariſ. in lib. de l'uſage des differentes ſortes des Saignées,* Amſtelod. 1729. Animadverſions againſt this laſt were alſo publiſhed at *Paris* in 1730, by M. CHEVALIER, Phyſician, and QUESNAY, a Surgeon there. Treatment after Apertion.

CHAP. V.

Of BLEEDING *in the Veins of the* FOREHEAD, TEMPLES, *and* OCCIPUT.

I. THERE are many Phyſicians and Surgeons, who think that bleeding by the Veins of the Forehead and Temples is much more ſerviceable and expeditious in relieving all Diſorders of the Head, ſuch as violent Pains, Vertigo, Delirium, Melancholy, and Raving Madneſs, *&c.* than the like Diſcharge by When and how theſe Veins are to be opened.

by Veins more remote from the Parts affected; judging that their Vicinity renders them more capable of evacuating the offending Matter of the Diſeaſe. But, for my own part, I muſt frankly own, that to me there ſeems to be little or no Foundation to expect any conſiderable Difference in the Effects of Bleeding from theſe Veins, in order to a more expeditious Removal of ſuch Diſorders: and this becauſe the external Veins of the Forehead and Temples have little or no Communication with the Brain and internal Parts affected, and do generally yield but a ſmall Quantity of Blood. In my Judgment, Bleeding by the jugular Veins ſeems more likely to anſwer that Intention, as they receive the Contents not only of the forementioned Veins, but alſo of thoſe immediately ſpent on the Brain and Parts affected, and are alſo more large and conſpicuous for Apertion. Yet if the Surgeon be expreſsly ordered by the Phyſician to phlebotomiſe in the Forehead or Temples, in compliance therewith, he ought to obſerve, that before he proceeds to inciſe the Veins, an Handkerchief or Neckcloth ought to be drawn tight round the Neck; that, by compreſſing the jugular Vein, thoſe Branches of it may become more turgid and conſpicuous. The Vein being opened, the Patient muſt hold down his Head, that the Blood may not trickle from his Forehead into his Eyes or Mouth, when the Stream does not ſpin out with ſufficient Force. If the Blood does not ſtop of itſelf after a due Quantity is diſcharged, you muſt compreſs the Orifice with your Finger; and, after wiping the Forehead and Face, apply a Compreſs or two, and then your Bandage.

Phlebotomy in the Occiput.

II. Bleeding from the occipital Veins, which communicate with the lateral Sinuſſes of the *dura Mater*, is both by Reaſon and Experience proved to be ſerviceable in moſt Diſorders of the Brain, where that Part is overcharged with Blood, which may be this Way diverted and evacuated. The celebrated Anatomiſt MORGAGNI[a] eſpecially recommends it, with Scarification and Cupping in thoſe Parts, for all lethargic Diſorders: And ZACUTUS LUSITANUS gives an Inſtance of a deſperate Apoplexy removed by deep Scarification and Cupping upon the *Occiput*, *De Medic. Princip. Hiſt. Lib.* I. *Hiſt.* 33. Theſe occipital Veins are opened by the ſame *Apparatus* as the Vein of the Forehead.

CHAP. VI.

Of BLEEDING *in the* VEINS *of the* INNER CORNERS OF THE EYES.

When and how theſe Veins are to be opened.

IT is well known from Inſpection, and the Writings of Anatomiſts, that there are two Veins which run one on each Side the Noſe through the *Canthi Majores*, or inner Corners of the Eyes; which proceed partly from the Forehead, and partly from the Eyes, and do, like the frontal Vein, diſcharge their Blood down into the external jugular Veins. 'Tis[b] bleeding in theſe canthal Veins that has been univerſally approved by DIONIS and the Generality of Oculiſts for Inflammations and other Diſorders of the Eyes: but upon no better Foundation in my Opinion, than that of bleeding in the Forehead and Temples, (*Chap.* V.) However, when you are to phlebotomiſe in theſe Corners of the Eyes,

[a] *Adverſar. Anat.* VI. *Animadver.* 83.

[b] See his *Surgery*, Edit. 2. p. 494.

Eyes, you must first make a Stricture about the Neck; and, after your Incision, the Patient must incline his Head, that a sufficient Quantity of Blood may be discharged from the Orifice without running into his Mouth: and then you apply a thick triangular Compress with Bandage. As for bleeding in the Veins of the Eyes, we shall consider that in treating of the Disorders incident to that Organ.

CHAP. VII.

Of BLEEDING *in the* JUGULAR VEINS *of the* NECK.

When and which Veins of the Neck are opened.

I. IT has been a very ancient Practice to bleed in the external Jugular Veins of the Neck, for most inflammatory Disorders of the adjacent Parts, for a Quinsey [a], Phrenzy, Madness, Ophthalmia, Apoplexy, inveterate Head-achs, Lethargy, and other Disorders of the Head. Nor are there wanting many among our modern Surgeons and Physicians to encourage the same Practice, and that even from the Authority of Reason and Experience: Since the accumulated and obstructed Blood and Humours may be this Way discharged from the Parts affected, and their bad Consequences prevented. Nor is the Operation at all dangerous; since the Jugular Veins run on each Side the Neck from the Head to the Clavicles, immediately under the Skin, and appearing generally very large, they may be easily perceived and opened: before which you must make a Stricture upon the lower Part of the Neck with a Handkerchief, Neckcloth, or the common Ligature, which must be drawn tight by an Assistant or the Patient, to make the Vein turgid and conspicuous. Or you may place a loose Bandage about the Neck, and let it be drawn downward strongly over the Patient's Breast, either by himself or an Assistant: by which means the Jugular Veins will be compressed on each Side, and become turgid without occluding the *Trachea*, or obstructing Respiration [b].

The Manner of incising the Jugulars.

II. When the Jugular Veins have been by this means rendered turgid and conspicuous, either of them which appears plainest may be secured by the Finger for Incision, either in the Right or Left Side of the Neck indifferently; when the Disorder lies in the whole Head, or in the Neck and Fauces. But when only one Side of the Head, or one Eye is affected, I think the Vein ought to be opened on the disordered Side of the Neck. The requisite Quantity of Blood being taken, the Ligature is next removed, and the Orifice compressed with your Finger, if the Blood does not stop without, while you wipe clean the Neck, and then apply your Compress and circular Bandage. Thus the Blood stops without any Danger of a fresh Hæmorrhage, of which some are without Reason afraid, as I have often experienced. Lastly, it must be acknowledged that the Patient faints away as readily after bleeding in the Neck, as the Jugular Veins are safely and easily opened: but no Danger follows from

[a] GOWER reckons it a specific in this Case. See his *Surgery*, p. 274.

[b] While I am revising these Sheets for the Press, occurs a Woman to whom I prescribe Bleeding in the Jugulars for a violent *Ophthalmia*; but upon applying the Ligature to her Neck, there is no Appearance of the Veins, an Accident I never before met with.

thence. We have an excellent Treatiſe on the Uſefulneſs of bleeding from the Jugulars, publiſhed at *Breſlau* in 8vo. 1735, by TRALLESIUS, a learned Phyſician of that City.

CHAP. VIII.

Of BLEEDING *in the* VEINS, *called* Ranulæ, *under the Tongue.*

IT is very often found of no ſmall Service in a Quinſey, or other inflammatory Diſorder of the Neck to bleed in the two ſmall Veins which run under the Tip or End of the Tongue: eſpecially if a larger Vein has been opened before, either in the Neck, Foot, or Arm, whereby the inſpiſſated and ſtagnated Blood may be gradually evacuted. To bleed in theſe Veins, a Stricture being made upon the Neck as before, you then elevate the *Apex* of the Tongue with your Left Hand, while, with the Lancet in your Right, you circumſpectly open firſt one, and then the other on each Side: becauſe the Apertion of one only will hardly ever diſcharge Blood enough to give any conſiderable Relief. When you judge a ſufficient Quantity of Blood has run out of the Mouth into your Veſſel, remove the Ligature from the Neck: upon which the Flux uſually ſtops of itſelf. But if it ſhould ſtill continue, let the Patient take a little Vinegar, or *Frontiniac* Wine in his Mouth: or elſe you may apply a bit of Vitriol or Allum, or a Compreſs dipt in ſome ſtyptic Liquor, till the Hæmorrhage ceaſes; which can never be dangerous even without ſuch Topics. For if there be not a good large Quantity of Blood diſcharged in the inflammatory Diſorders of theſe Parts, the Apertion of theſe Veins will be of little or no Signification. Yet there are Inſtances where Patients have died for want of ſtopping Blood in theſe Veins. See *Mix. Nat. Cur.* A IV. *Obſ.* 101. and *Ephem. N. Cent.* I. p. 188.

CHAP. IX.

Of PHLEBOTOMY *in the* PENIS.

BLEEDING in the *Vena dorſalis Penis* uſually ſurpaſſes the Benefit of all Remedies whatever in abating inflammatory Diſorders of this Member. This large Vein, which runs along the Back or upper Side of the *Penis*, being generally pretty much diſtended, and conſpicuous in an Inflammation of this Part, may be inciſed about the Middle or back Part of the *Penis*; and kept bleeding till the Member becomes flaccid, and a ſufficient Quantity of Blood be diſcharged proportionable to the Urgency of the Smyptoms. This done, you muſt apply a Compreſs, and the Bandage proper for the *Penis*, as we ſhall direct in the third and laſt Part of our Surgery. But you muſt carefully endeavour to avoid injuring the Arteries or Nerves which enter the *Penis* near this Vein: as alſo not to make your Bandage too ſtrict. For by theſe means the Inflammation and Symptoms may turn out worſe than before.

Of

Of the Symptoms or Accidents which attend Phlebotomy.

CHAP. X.

Of an Ecchymosis.

Ecchymosis, what.

I. MANY are the Accidents which may follow from the Apertion of a Vein. But we shall here only consider the Principal, and begin with an Ecchymosis, or Extravasation of Blood from the Vein betwixt the Flesh and the Skin: Of this there may be various Degrees: so that the Arm hereby often becomes not only much swelled, and of a black-and-blue Colour, but is even sometimes violently inflamed with a most acute Pain, and followed either with a Suppuration, or incipient Mortification in the Limb.

Causes.

II. The Accident we are now treating of frequently proceeds from the Vein having been cut quite asunder by the Phlebotomist: but oftner from the Patient's using his Arm too early after Bleeding, in violent and long Exercises; in which the Contractions of the Muscles make the Veins swell, and force their Blood through the Orifice into the Interstices betwixt the Flesh and Skin, either in a greater or less Quantity, in proportion to the Degree of Violence and Exercise.

Consequences of a slight Ecchymosis.

III. In a slight Ecchymosis or Effusion of Blood under the Skin, there is little or no Danger to be feared, as the stagnant Blood may be generally dispersed without any great Difficulty by the Application of a Compress dipt in Vinegar and Salt, or in Spirit of Wine. Sometimes the Blood suppurates or turns to Matter, which may be much promoted by a *Diachylon* Plaster: and when the Matter is once brought to Maturity, it generally makes its own way through the Integuments, without the Assistance of any Incision: after which, being discharged, the Wound may be healed with a Bit of *Diachylon* Plaster.

Treatment of a large Ecchymosis.

IV. If the Quantity of Blood stagnating in an Ecchymosis be very large and considerable, there is generally but little or no Hopes left to disperse it: but the Disorder too often terminates either in a large Abscess or a Gangrene, after violent Pain and Inflammation have preceded. But to prevent these Consequences, the Surgeon must take his Scalpel, and scarify, or make many little Incisions upon the livid Part to discharge the extravasated Blood; and then apply either a *Diachylon* Plaster, or the Fomentation before recommended for Contusions and Phlegmons, *Part* I. *Book* I. *Chap.* XV. *Sect.* X. & seq. *Book* IV. *Chap.* II. *Sect.* XIV. But if the Arm is already possessed with a violent Inflammation or Gangrene, you ought to scarify it well, and then to invest it with discutient Cataplasms or Fomentations, as we before directed in *Part* I. *Book* IV. *Chap.* XIV. *Sect.* VI. But at the same time in these Cases it is often necessary to bleed in some other Part, and to administer attenuating Medicines internally, till the Inflammation abates, or the Gangrene spreads no farther. You will see more on this Subject, *Chap.* XII. *Sect.* XVI.

CHAP. XI.

Of the PUNCTURE *of a* NERVE *or* TENDON *in Phlebotomy.*

Signs of this Accident.

I. WHAT grievous and cruel Symptoms may ariſe from the pricking a Nerve or Tendon, we have before intimated, in treating of Wounds, *Part* I. *Book* I. *Chap.* I. *Sect.* X. and XI. But you may reaſonably judge, that a Nerve or Tendon has been injured in Bleeding, if the Patient, at the Time of Inciſion, feels a moſt acute Pain, ſo that he can ſcarce refrain from a ſevere Outcry. In a ſhort time after, the excruciating Pains ſtill continuing, the Limb ſwells, becomes inflamed, convulſed, ſtiff, and extended as in the Cramp; which Symptoms, if not timely relieved, threaten Convulſions of the whole Body, a Gangrene of the Part, and Death itſelf.

Treatment.

II. Among the ſeveral Methods of treating theſe Symptoms, from ſuch an Accident, that ſeems to be one of the beſt, which was formerly uſed for the *French* King, CHARLES IX. by his Surgeon AMB. PAREY. For the King had no ſooner declared his intenſe Pain, by crying out aloud, while the Vein was opening, than PAREY imagined, with good Reaſon, that ſome Nerve was injured: and accordingly, the Arm began to ſwell in a little time with excruciating Pains, and at length became quite rigid. Hereupon the King's Phyſicians were immediately called in to a Conſultation with PAREY, and the Treatment agreed on was firſt to bathe the Part injured with warm *Ol. Terebinth. cum Sp. Vin. rect.* and then to inveſt the whole Arm with *Emplaſt. Diachalciteos in Ol. & Acet. Roſar. ſolut.* retained by the expulſive Bandage, which, beginning upon the Hand, aſcends gradually by ſpiral Turns to the Top of the Shoulder. By this Means the Impulſe of the Blood on the Part was not only much abated, but alſo the Pain and Inflammation much diminiſhed. And laſtly, to compleat the Cure, the following Cataplaſm was ordered to be applied to the Arm:

℞. *Farin. Hord. Orob. ana* ℥ ij. *Flor. Chamæmel. Melilot. ana* M ij. *Butyr. recent.* ℥ jſs.

Theſe boiled into a Cataplaſm with Soap-ſuds, were applied to the Arm, till the Pain, and other malignant Symptoms, were totally removed. Notwithſtanding which, the King had a Stiffneſs in moving his Arm for near three Months afterwards: But, by degrees, that went off, and his Arm grew as ſtrong and agile as ever.

A ſecond Method of Cure.

III. Equal Succeſs may alſo be expected from treating the Part with warm *Hungary* Water and *Balſ. Peruv.* for ſeveral Days, till the Pain goes off: And as the *Diachalciteos* Plaſter is ſeldom retained in many of the Apothecaries Shops, you may ſubſtitute *Emplaſt. de Minio vel Saturninum & Diapompholygos.* But great Care muſt be taken, in the mean time, while theſe Medicines are preparing, not to expoſe the Wound open to the Air. Therefore the Wound may be at firſt covered with a Bit of any Sort of Plaſter, and the whole Arm inveſted with a Linen Cloth moiſtened with Oxycrate; which will both abate the Inflammation, and exclude the Injuries of the Air or Duſt from the Part. If the Patient be young, and of a full Habit, it will be alſo proper, at the ſame time,

time, to bleed plentifully in the other Arm. SCULTETUS, *Obſ.* 83. has an Ointment which he much extols for Punctures of the Nerves, as you may there find: where he alſo relates, that he has ſeveral times ſucceſsfully cut through, or totally divided, ſuch punctured Nerves.

CHAP. XII.

Of WOUNDS *of the* ARTERIES *in* PHLEBOTOMY.

Diagnoſis.

I. IN Bleeding, it ſometimes happens, that an Artery is pitched upon and opened inſtead of, or together with, the intended Vein, and this more eſpecially when the Surgeon bleeds in the Baſilic Vein of the Arm; near to which uſually runs the large Brachial Artery[a], an Apertion whereof muſt be followed with a dangerous Hæmorrhage, an Aneuriſm, or even Death. This HILDANUS[b], with myſelf and others, has often obſerved, either from the profuſe Hæmorrhage, or from a Sphacelation of the Limb from the Courſe of the Blood being interrupted. That an Artery is thus accidentally opened inſtead of a Vein, you may diſcover by the Blood's ſpinning very forcibly from the Orifice, by Starts or Leaps, rather than in an even Stream, and extending itſelf into a greater Arch from the Orifice to the Receptacle: The Colour of the Blood is alſo here much more florid, or of a brighter red, than that from a Vein. To which add, that if you here preſs your Finger on the Veſſel below the Orifice, the Blood ſtarts out more violently than before; and quite ſtops, or elſe greatly diminiſheth upon preſſing above the Orifice. The Reverſe of this is found in the Apertion of a Vein.

What the Surgeon muſt do in ſuch a Caſe, when undiſcovered to others.

II. In caſe of ſuch a dangerous Accident, the Surgeon muſt firſt endeavour to keep up his Preſence of Mind, which is very apt to be confuſed by Fear, that thereby the Patient, or his Attendants, may not ſuſpect his Error. In the next place, he muſt carefully obſerve, whether the Blood flows freely from the Orifice, or whether it inſinuates, in a conſiderable Quantity, betwixt the Integuments. If the firſt, he muſt take a large Quantity of Blood, even till the Patient faints, perſuading him and his Attendants, that his Blood appears ſo hot and redundant, as to make ſo large an Evacuation abſolutely neceſſary, after the Example of M. DIONIS[c], when he met with this Accident. When the Patient is in a Deliquium, as the Flux then ceaſes, you may commodiouſly dreſs and bind up the Wound, and by this Precaution hinder a freſh Hæmorrhage, or an Aneuriſm. While the Attendants are otherwiſe employed, the Surgeon muſt place a Farthing, or ſome other Piece of Money, in the Folds of the firſt Compreſs, which being fixed on the Orifice of the Arm wiped clean, he muſt, upon the firſt, place two, three, or more, thick Compreſſes, each larger than the other: and then bending the *Cubitus*, he muſt, for the greater Security, apply two Bandages, in the ſame Manner as after bleeding in a Vein, only a little tighter. It may be next proper to lay a thick, long, and narrow Compreſs

[a] But I have alſo ſometimes obſerved this Artery near the Cephalic Vein.
[b] *Obſ.* 44. *Cent.* III. & LANCISIUS, Lib. *de Cord. & Aneuriſm.*
[c] In his *Surgery*; *Chap.* of *Veneſection.*

preſs upon the Arm, over the Artery, from its Inciſion to the *Axilla*, and to ſecure it in that Poſition by a ſpiral Bandage: That the Brachial Artery being thus compreſſed, the *Impetus* of the Blood on the Wound may be abated; ſignifying to the By-ſtanders, that the Patient's Blood is ſo ardent and rapid, that it cannot well be reſtrained from bleeding again, without this particular Deligation. Thus perhaps his Error may eſcape unſuſpected. Inſtead of the firſt Compreſs with a Piece of Money, you may apply with equal, or more Advantage, a Lump of brown Paper chewed in your Mouth, and then the Moiſture preſſed out of it, ſecure it on the Orifice by ſeveral Compreſſes, and the Bandage as before.

Treatment of the Patient afterwards.

III. The Deligation being compleated, if the Patient does not then recover from the Swoon of himſelf, the uſual Means are to be uſed to recover him, by ſprinkling cold Water in his Face, opening the Windows, applying Volatiles, Vinegar, or *Hungary* Water to his Noſtrils, *&c.* By which means, being brought to himſelf, he muſt be ſtrictly charged to refrain from Exerciſe, to live on a ſpare and thin Diet, and not to uſe his Arm for ſome time, leſt a Relaxation of the Bandage might occaſion a freſh Hæmorrhage, or an Aneuriſm. To avoid this, it may be alſo requiſite to ſuſpend the injured Arm a little bent in a Sling about the Patient's Neck: and to keep it the more ſteady, the Sling may be pinned to the Patient's Clothes, and at Night laid in a convenient Poſture on a ſoft Pillow.

Frequent Viſits neceſſary.

IV. A few Hours after the Deligation, the Surgeon ought to viſit his Patient, and again, at ſhort Intervals, as often as he conveniently can, in order to inſpect the Arm and Bandage, to ſee that the latter ſits tight, and to prevent the Attack of a freſh Hæmorrhage, Pain, Tumor, Inflammation, Gangrene, or other bad Symptoms. If every thing appears right, except only a ſmall, uniform, and ſoft Swelling of the Arm, the Bandage ought nevertheleſs to remain on the Arm, till the fourteenth Day: for ſuch a Swelling does not preſage any thing amiſs, even though it infeſts the whole Arm. But if your Bandage is perceived to get looſe, it ought to be taken off cautiouſly, and re-applied more cloſely. But while the Bandage is taken off from the Arm, the Artery ought to be compreſſed by the Tourniquet, or at leaſt by the Thumb of an Aſſiſtant, graſping the Arm; the Surgeon in the mean time holding his Thumb or Finger preſſed on the Wound, till he re-applies either the ſame or freſh Compreſſes and Bandage. But in this you muſt be careful not to force off the laſt Compreſs or Lump of brown Paper from the Inciſion, if it does not fall off of itſelf, but rather let it remain. However, if it ſhould ſeparate, you may dreſs the Wound with a little *Balſ. Peruvian. vel Capaiv.* till it is well cloſed, and out of Danger, nor liable to a freſh Hæmorrhage. If you come to your Patient, and find his Arm bleeding, the Trunk of the Brachial Artery muſt be immediately compreſſed, either by the Tourniquet, or with the Thumb and Fingers of an Aſſiſtant fixed about the Middle of the Arm: and having provided more or thicker Compreſſes, and a longer Bandage, you then take off the old Dreſſings, waſh clean the Wound with warm Wine, or its Spirit, and next proceed to renew your Deligation more carefully, as we before directed. If the Surgeon meets with the Appearance of a Gangrene from too great a Stricture of the Bandage, he muſt unbind and foment the Arm, or treat it with the Re-

Remedies proper for that Caſe, and, augmenting the Number of his Compreſſes, re-apply his Bandage more cloſely than before. But if the Gangrene proceeds from a Loſs of the Circulation through the Limb, by a Defect of the other arterial Trunk of the Arm, which ſeldom happens, in that Caſe you muſt amputate without delay.

The Patient's Regimen.

V. If the Surgeon meets with none of the forementioned Symptoms, for ſome time after his Deligation, he muſt order the Patient to keep on the Bandage for a Week or a Fortnight longer, keeping his Arm, in the mean time, free from Exerciſe or Motion: leſt the Blood ſhould, by that means, force and extend the (as yet tender) *Cicatrix* into an Aneuriſm. His Diet muſt alſo be all along ſpare and light, as at the Beginning; ſtrictly avoiding all Wines and fermented Liquors, and every thing that will put the Blood into a violent Commotion: in which laſt Caſe the Surgeon will find it neceſſary to bleed in another Part. Thus you may avoid all Danger of an Hæmorrhage, or an Aneuriſm; and the Patient's Arm will become as well as ever, eſpecially, if the Wound be dreſſed with a little *Balſ. Peruv. vel Capaiva, &c.*

What muſt be done if the Error is detected.

VI. Thus far have we deſcribed the Method, in which the Surgeon muſt proceed, when the Error is not diſcovered by the Patient, or his Attendants. But if either of them have ſmelt out the true Caſe, it will be the beſt Way for him to make a free Acknowledgment of his Miſtake or Accident, excuſing the ſame, by aſſuring them, it is no more than what may happen to the moſt expert Surgeon living, in opening ſome Veins: and then promiſing the Patient, that if his Directions are obſerved, he ſhall be perfectly cured, without any Damage: and thus he may compleat his Cure, perhaps better than if his Patient knew nothing of the Matter: for knowing the Caſe to be ſo much more dangerous than that of an inciſed Vein, the Patient will be more ſubmiſſive, and the Surgeon's Orders more punctually obſerved.

Treatment when the Blood inſinuates betwixt the Fleſh and Integuments.

VII. When the Aperture of the Artery, and that of the Integuments, do not exactly correſpond with each other, but the Blood being forced out of the Artery, inſinuates itſelf betwixt the Fleſh and Skin; in that Caſe, which very often happens, the Patient muſt not be bled *ad Deliquium:* for even after that, there may be ſo much Blood extravaſated and retained betwixt the Integuments and Muſcles, as may cauſe a Mortification of the Arm by its Putrefaction, or at leaſt may render the Operation for an Aneuriſm abſolutely neceſſary to be performed. If therefore the Surgeon cannot draw back the Orifice or Inciſion of the Integuments, ſo as to make it correſpond with that of the Artery, and diſcharge the retained extravaſed Blood, he ought immediately to compreſs the Wound with a Lump of chewed Paper, and ſeveral Compreſſes, each larger than the other, which are all to be firmly ſecured on the Part by the Bandage or Deligation before deſcribed at *Sect.* II. of this Chapter; not forgetting the long Compreſs and Bandage, which we recommended for compreſſing the Brachial Artery. Then, after bleeding plentifully ſeveral times, in ſome other Part, the Remainder of the Treatment may be according to *Sect.* III, IV, V, and VI. preceding. But the Patient muſt be viſited again in a little time, to inſpect the Arm: for it often happens, that when you have no apparent Bleeding after Deligation, yet the Blood will inſinuate itſelf betwixt the Muſcles and Integuments, ſo as to diſtend the Arm to an enormous Size. A remarkable In-

ſtance

ſtance of this DIONIS [a] gives from his own Practice. He was once obliged, in this Caſe, to inciſe the Integuments of the whole Arm, whereby he diſcharged four Pounds of Blood, that had been equally diſperſed all around, from the Elbow to the Shoulder. And we alſo meet with a ſimilar Obſervation in RUYSCH [b], in which concreted Blood was lodged almoſt all over the Arm. You may alſo conſult BARTHOLIN. *Epiſt. Med.* 53. *Cent.* III. *Hiſtor. Anatom.* IX. *Cent.* II. and his Hiſtory of an Aneuriſm diſſected, which he ſaw at *Naples*, *An.* 1644.

CHAP. XIII.

Of ANEURISMS.

What an Aneuriſm is.

I. A Throbbing Tumor, diſtended with Blood, and formed by a Dilatation, Wound, or Rupture of an Artery, is by Surgeons uſually denominated an *Aneuriſm:* of which they diſtinguiſh two Kinds, the *true*, and the *ſpurious*. A true Aneuriſm has always a Pulſation, more or leſs, and is formed by a Dilatation only of the Artery, either all around [c], or on one Side of it, much in the ſame Manner as thoſe analogous Tumors of the Veins are formed, which we term *Varices*. So that both Aneuriſms and Varices are a Kind of *Herniæ* of the Arteries and Veins, and accordingly they are by ſome named *Herniæ Arteriarum & Venarum*. But the ſpurious Aneuriſm is, when the Artery, being opened by a Puncture, Wound, Contuſion, Eroſion, or other external Violence, extravaſates the Blood betwixt the Muſcles and Integuments, the Limb itſelf appearing livid, and much ſwelled thereby. A true Aneuriſm may alſo degenerate into one that is ſpurious, by a gradual Dilatation of the Artery, and Extenuation of its Coats, till at length being totally ruptured, the Blood is either extravaſated and retained under the Integuments, or diſcharged freely from the Wound. Hence the Tumor is much larger and leſs prominent, or pointed, in the ſpurious, than in the true Aneuriſm, and is alſo attended with little or no ſenſible Pulſation; but the Putrefaction of the extravaſated Blood very often occaſions a Gangrene and Mortification of the Part, or even Death itſelf, by a profuſe Hæmorrhage. But Aneuriſms may be again diſtinguiſhed from their Circumſtances and Symptoms, into *ſimple* and *complicated*. The firſt are formed without any ill Accidents: the laſt are uſually attended with Immobility, violent Pain, an Abſceſs or Sphacelation of the Part, *&c.* which more

[a] *Chirurg. Operat. Demonſtrat.* VIII. Chap. of Aneuriſms. [b] *Obſ. Anat. Chirurg.* Obſ. 2. pag. 7.

[c] 'Tis a little extraordinary that the learned Dr. FREIND ſhould, in his Hiſtory of Phyſic, contend that all Aneuriſms are formed by a Rupture of the Artery; when we have ſo many Inſtances of their ariſing from a Dilatation only of the arterial Coats, either on one or all Sides. See that deſcribed by me in *Annal. Acad. Juliæ Semeſtri* XII. p. 81. Thoſe in PAREY's Surgery, and RUYSCHII, *Obſ. Chirurg. & Hiſt. Acad. Reg. An.* 1712 & 1721. Alſo LANCISII *Lib. de Corde & Aneuriſmat. & Lib. de Mortib. Subitan. in Schol. Obſ.* 5, § II.

more uſually accompany the ſpurious Aneuriſm [a]. Aneuriſms may be alſo diſtinguiſhed, from the Situation of the Arteries, into *external* and *internal* [b], the firſt being acceſſible, the others not. Another remarkable Difference of them may be taken from their having either a violent or elſe but little or no ſenſible Pulſation [c]. For it is to be obſerved, as we before mentioned, that ſpurious Aneuriſms ſeldom have any conſiderable Pulſation, eſpecially when they are large; whereas the true Aneuriſms, eſpecially the ſmall, have a very ſtrong and ſenſible Pulſation: but in ſome of them the Pulſation increaſes, and in others it diminiſhes, as the Tumor enlarges. See my Account in *Annal. Acad. Juliæ Semeſtri* XII. *pag.* 81.

II. In a true and external Aneuriſm, beſides the forementioned Signs, we obſerve a ſmall Tumor at the Beginning, no larger than a Filbert, which has always a Pulſation. As for the internal Species, as they lie concealed from our Senſes, little or nothing can be ſaid of their Signs, with which, however, the Reader may be ſupplied in LANCISI's Treatiſe on the Subject. The Tumor generally feels ſoft to the Fingers, with a Sort of Fluctuation and Reſiſtance of a Fluid, and is almoſt conſtantly of the ſame Colour with the Skin, having a Pulſation like that of the Artery to which it belongs. Upon preſſing the Finger on the Tumor, as yet ſmall, it diſappears: and upon removing the Finger it returns inſtantly again. But the ſpurious Aneuriſm appears livid, feels hard and turgid, with intenſe Pains: the Tumor is here more plain or equal, and generally without Pulſation, as upon preſſing, it affords a Sort of rumbling or fluctuating Noiſe; and diſtending the whole Limb, or a great Part thereof, to an unuſual Size [d], it very often either degenerates into an Abſceſs or a *Sphacelus*. *Diagnoſis.*

III. Aneuriſms moſt frequently ariſe in the Brachial Artery, from an erroneous Puncture or Injury thereof, in bleeding in the Arm, eſpecially in the Baſilic Vein. For the Artery being in a conſtant Pulſation, will, by urging its Blood againſt the arterial Coats, gradually diſtend them where they make too little Reſiſtance, ſo as at length to form a conſiderable Tumor. If therefore a throbbing or beating Tumor like that deſcribed in the foregoing Paragraph ſhould appear in the Arm a few Days or Weeks after bleeding, it may be certainly depended upon to be an Aneuriſm. But the Origin of Aneuriſms is not from the Lancet alone; nor is their Seat reſtrained to the Arm only [e]; for they *The Seats and Cauſes of Aneuriſms.*

[a] A remarkable Aneuriſm of the ſpurious Kind is deſcribed by BARTHOLIN in a profeſſed Diſſertation, entitled, *Aneuriſmatis Diſſecti Hiſtoria*, Panormi 8vo. 1644. See alſo VAN HORNE *in Epiſt. de Aneuriſmate*; and LANCISIUS, *Lib. de Cord. & Aneuriſm.*

[b] Hiſtories of internal Aneuriſms may be ſeen in PAREY, *Book.* VII. *Chap.* 32. Monſ. BLEGNI, *Zodiac. Med. Gallic. An.* 1681. *p.* 44. RUYSCH. *Obſ. Chirurg.* 37. LANCISII, *Et Annal. Acad. Juliæ locat.*

[c] Of which I have made many Obſervations beſides thoſe in PAREY, *loc. cit.* RUYSCH. *Obſ.* 38. BLEGNI, *l. c. p.* 25. & 42. NUCK *Operat. Chirurg. Experi.* XXIX. LANCISII, *l. c.*

[d] The ſpurious Aneuriſm often acquires an enormous Size, but the true one hardly ever exceeds the Bulk of a Cheſnut, according to GOUEY, *Chirurg. pag.* 231. But that his Opinion is not to be abſolutely depended on, may appear from the ſeveral Accounts we have of large Aneuriſms, particularly one the Size of a Gooſe Egg in HILDANUS, *Obſ.* 44. *Cent.* III. PURMANNUS *Chirurg. curioſa, p.* 212. And in our *Tab.* XI. *Fig.* 6.

[e] AMB. PAREY, *Lib.* IV. *Cap.* 32. aſſerts the Neck to be the Part in which Aneuriſms are moſt frequently formed; but his Opinion is not countenanced by our latter Experience and Obſervations.

they may ariſe from an infinite Number of Cauſes, both external and internal, and may be formed in all Parts where there are any arterial Trunks, or conſiderable Branches diſtributed. Thus we often meet with them from a Wound, Contuſion[a], and Suppuration, and from external Injuries in moſt Parts of the Body. But internally they may ariſe either in the Thorax or Abdomen, from a Diminution of the Strength and Reſiſtance of the external or internal Coats of the large arterial Trunks, from various Cauſes, as an Ulceration, Eroſion, *&c.* agreeable to the Obſervations of FALLOPIUS, (*Lib. de Tumor. Cap.* 14.) SEVERINUS (*Lib. de Abſceſſibus*) RUYSCHI *Obſ.* 37. & 38. LANCISI. (*Lib. de Cord. & Aneuriſmat.*) and our Obſervations in *Annal. Acad. Juliæ Semeſtri* XII. p. 81. We muſt however confeſs, that the Cauſes of internal Aneuriſms are often very doubtful and unſettled: notwithſtanding which, we ought to diſtinguiſh thoſe Cauſes as they occur, into external and internal. They often proceed from a Blow, Fall, or a Fracture of the adjacent Bone, or a violent Straining in lifting great Weights, Jumping, Riding on Horſeback, *&c.* whereby the Blood is accumulated and urged ſo forcibly in the Artery injured, as gradually to diſtend its Coats, and form a Tumor. Sometimes they are owing to an Inflammation, Suppuration, Eroſion, or Ulcer in ſome neighbouring Part, or in a Part of the Artery itſelf: By which the other Coats are ſo weakened, that they are unable to ſupport the Impetus of the Blood. Hence they give way, expand themſelves, and ſwell. In the ſame manner too we often meet with Aneuriſms from a ſlight Puncture, or even barely touching the Coats of an Artery with a Lancet in opening a Vein: in which caſe the exterior Coat of the Artery being divided, and the interior remaining entire, the latter is not alone ſtrong enough to reſiſt the Impulſe of the Blood, but gives way inſenſibly at each Ictus of the Artery, till it at length forms that conſiderable Tumor which we call an *Aneuriſm.* If we therefore conſider that the mechanical Formation of Aneuriſms is in this Manner from a diminiſhed Reſiſtance in the arterial Coats, we ſhall find the Cauſes thereof very numerous, which may weaken an Artery more in one Part than another; ſo as to make it give way to the Force of the Heart, or Impulſe of the Blood and form an Aneuriſm. And this eſpecially, when ſeveral Cauſes concur together, as if violent Straining or Leaping, *&c.* be uſed when the Coats of the Artery are previouſly extenuated or weakened by a Contuſion, Inflammation, Suppuration, *&c.*

Diagnoſis of injured Arteries.

IV. I think we have in the preceding Chapter ſufficiently explained the Manner of enquiring into the great Injuries and Wounds of the Arteries, that may happen in opening a Vein. We ſhall here only enumerate the Signs by which we may diſcover ſlight Punctures, or the ſmaller Injuries of them, which occur in Phlebotomy. But as we are not ſupplied with any certain or characteriſtic Signs indicating ſuch ſlight Accidents, we muſt make the beſt Uſe of a reaſonable Conjecture. If therefore you ſhould perceive a Pulſation againſt the Point of your Lancet, notwithſtanding you have no Hæmorrhage from the Artery, yet you may reaſonable conclude that the external Coat of that Veſſel muſt be in ſome degree injured thereby: and therefore it will be proper to make

[a] Thus FEHRIUS has obſerved an Aneuriſm in a Lad, from a Blow on the Left Side of his Head, which in the Space of eight Days, enlarged ſo as to cover half his Head. *V.* BARTHOLIN. *Epiſt* 53. *Cent.* III.

make your Deligation and Compreſſion to prevent an Aneuriſm, in the Manner we before directed in the preceding Chapter.

V. But if a ſmall beating Tumor ſhould be formed within the Space of a Month after Phlebotomy, either through the Neglect of the Surgeon or Patient, or from leaving off the Deligation too early, it may be pretty ſafely depended on to be one of theſe Aneuriſms from a ſlight Cauſe. But if it be a true Aneuriſm, whilſt it continues recent and ſmall, it gives little or no Uneaſineſs, beſides its Tumor and Pulſation: yet when it has afterwards gradually acquired the Size of an Egg, or one's Fiſt, or even the Bulk of one's Head, as may be ſeen in PURMANNUS *Chirurg. Curioſ.* p. 612. and in our *Tab.* XI. *Fig.* 6. it then occaſions intenſe Pains, Weakneſs, Immobility, and other bad Symptoms in the affected Limb. If then the Help of the Surgeon be not ſpeedily called in, the arterial Coats becoming gradually extenuated, will at length burſt, and be followed by a Train of the worſt Conſequences, if not the Death of the Patient. If the external Integuments ſhould be broke through, a fatal Hæmorrhage muſt follow: and even if they ſhould continue intire, an Abſceſs or Gangrene would deſtroy the Part, as I myſelf have obſerved here in a Patient at *Helmſtadt*. See alſo RUYSCH, *Obſ.* 2. Though the Generality of Aneuriſms afford a dangerous Prognoſis, as [a] BARTHOLIN and [b] HARDER obſerve, yet none are ſo much to be feared as thoſe which are formed internally in the larger arterial Trunks, where there cannot be had a free Acceſs to the Parts, as in the *Aorta*, *Subclavian*, Beginning of the axillary, brachial, and carotid Arteries [c], *&c.* Thoſe Aneuriſms too are generally incurable which are formed in the carotid Arteries of the Neck, in the Subclavian or Axillary near the Shoulder, and in the crural Artery, eſpecially if near the Abdomen. For if the Operation be performed on any of theſe, it muſt be followed either with a profuſe or fatal Hæmorrhage, or elſe a Mortification of the Parts. But thoſe Aneuriſms are much leſs dangerous, and frequently admit of a Cure, which are formed in the external Branches of the Arteries, eſpecially in thoſe running on the *Cranium*, or without the Ribs, and thoſe in the Foot, Hand, or lower Arm. Yet if the Aneuriſm be not recent, though even in the Arm, the Succeſs of the Operation by the Knife will be at leaſt very uncertain, when Deligation and Compreſſion alone will not take their due Effect. For as the arterial Trunk muſt neceſſarily be cloſed or ſhut, it will be almoſt next to impoſſible to prevent the Part, to which the Artery was diſtributed, from waſting away, or elſe from mortifying: ſince the Circulation of the Blood, and their Supplies of Nouriſhment are by this means in a great meaſure, if not totally, cut off; the lateral ſmall Branches of Arteries being incapable of importing a due Quantity of Blood to the Hand and Parts of the *Cubitus*, when one of the larger Branches is wanting [d]. This is therefore a frequent Cauſe of a Mortification in them, ſo as often to oblige the Surgeon

Prognoſis.

[a] See BARTHOLIN. *Epiſt. Med. Cent.* III. *Epiſt.* 53. [b] Jo. JAC. HARDER, in *Apior. Obſ.* 86.

[c] LE DRAN, *Obſ.* 40. *T.* I. relates, that he has found frequent Veneſection of great Service in an Aneuriſm of the *Aorta*. I have experienced the ſame.

[d] That the fellow arterial Branch of the *Cubitus* is not ſo often abſent as Surgeons have imagined, is made apparent, with other juſt Anatomical and Chirurgical Obſervations, in a Medical Diſſertation or Theſis had under me at *Helmſtadt*, by Dr. MORRIUS, *An.* 1730. the Subſtance of which I think to communicate in my Obſervations, which I intend to publiſh ſome time hence by themſelves.

Surgeon to an Amputation, as hath been frequently experienced by myſelf and others[a]; and even Amputation itſelf will very often not ſave the Patient, as may appear from the Caſe in BARTHOLIN, *Epiſt.* 53. *Cent.* I. When an Aneuriſm burſts ſpontaneouſly, the Hæmorrhage is generally ſo profuſe, that the Patient's Life may be loſt in a Minute's Time, if a ſpeedy Compreſſion be not made on the Artery by a ſtrict Ligature, or the Tourniquet, and the Aſſiſtance of an expert Surgeon. And extremely dangerous is the Caſe when the Surgeon, by Neglect or Miſtake, inciſes one of theſe large Tumors inſtead of an Abſceſs, as hath been ſometimes done[c]. Yet it ought to be obſerved here, that ſpurious Aneuriſms are in the general much more dangerous than the true ones. True Aneuriſms are ſometimes tolerable without any great Danger or Uneaſineſs for many Years[d], or as long as the Patient lives; eſpecially if they are defended and ſecured with proper Bandage and Compreſſes: Whereas on the contrary, ſpurious Aneuriſms will not continue many Days without inducing an Hæmorrhage, Abſceſs, and Mortification in the Parts. But both the true and ſpurious Species of Aneuriſm are always the more dangerous and troubleſome as they are larger: Inſomuch, that their Size has deterred the expert and intrepid HILDANUS[e] from performing the Operation on them. And RUYSCH openly declares[f], that, in the vaſt City of *Amſterdam*, no Surgeon had undertaken to perform the Operation for above twenty Years before him. The ſpurious Aneuriſm is alſo more difficult to cure even by the Knife than the true Species: Becauſe the Blood which is extravaſated and concreted all around gives the Surgeon immenſe Trouble to diſcharge it. As for internal Aneuriſms, they not only lie concealed from our Senſes, but are alſo abſolutely deſtitute of any Help or Remedy from Art, becauſe they are inacceſſible to the Hand. But were an internal Aneuriſm to extend and ſhew itſelf externally, it could not be well ſubjected to the Operation, without greatly hazarding the Patient's Life: and therefore the Cure of ſuch have been prudently refuſed by the moſt eminent Surgeons, as FALLOPIUS, PAREY, SEVERINUS, *&c.* cited in BARTHOLIN's *Hiſtoria Aneuriſmatis Diſſecti.* And for the ſame Reaſon we here reſtrain our Doctrine and Treatment of this Diſorder to the external Species of Aneuriſms only. But they who deſire a more particular Account of the Internal, may conſult the learned Treatiſe on the Subject by LANCISI.

Treatment of ſight Aneuriſms.

VI. I ſhall now, for the Information of the younger Surgeon, deſcribe the Method of treating an incipient Aneuriſm, forming itſelf in the Flexure of the *Cubitus*, or Bending of the Arm, where this Diſorder more frequently occurs than in any other Part: and from thence, I think, he may eaſily judge of the Method in which other leſs frequent Aneuriſms are to be treated. Whenever a ſmall Aneuriſm of the true Species begins to form, and ſhew itſelf at the Flexure of the Arm, you are furniſhed with two Methods of relieving it, either by Deligation, or by Inciſion. The firſt of which may be again performed either

[a] V. RUYSCH, *Obſ.* 2. BARTHOLIN. *Epiſt.* & VAN HORN *de Aneuriſmate.*

[b] V. *Phil. Tranſact.* N° 402. *Act. Erud. Lipſ.* Tom. III. *page* 401. PAREY, *Lib.* VI. *Cap.* 32.

[c] V. PAREY, *Lib.* VI. *Cap.* 32. HILDANUS, *Cent.* III. *Obſ.* 43. RUYSCHII, *Obſ.* 38. VAN HORN & LANCISI, *loc. cit.*

[d] Thus SENNERTUS (*Prax. Med. Lib.* V. *Part* I.) gives the Caſe of a Woman who ſuſtained an Aneuriſm the Size of a Walnut on the Flexure of the *Cubitus*, without any Detriment, for the Space of thirty Years.

[e] *Cent.* III. *Obſ.* 44. [f] *Obſ. Chirurg.* 2.

either by Compreſs and Bandage, or by an Inſtrument adapted for the Purpoſe. The Method of relieving and curing this Diſorder by Deligation and Compreſſion, if there be no Extravaſation, ought always to be tried before that by Inciſion, as well in the incipient true as in the ſpurious Aneuriſm: for it would be barbarous to ſubject the Patient to a cruel Operation, for what may be remedied by a milder Treatment. The Patient may be therefore relieved, and the Tumor diminiſhed by Compreſſion, after diſcharging the extravaſated Blood, either with a Compreſs of chewed Paper, or a Bit of aſtringent Plaſter, retained with the other Compreſſes and Bandage we deſcribed in the preceding Chapter. By which means the Diſorder may be conſiderably diminiſhed, if the Deligation be continued on the Limb for ſeveral Weeks or Months: and thus we read of Cures performed as well formerly by HILDANUS (*Cent.* III. *Obſ.* 44.) TULPIUS (*Obſ. Med. Lib.* IV. *Cap.* 17.) ROGERUS, (*Zod. Med. Gall.* 1681. p. 43.) and others of the laſt, as well as of the preſent Century. But if Deligations be found inſufficient, as it was upon the *French* King's Phyſician, M. BOURDELOT (*Zod. Med. loc. cit.*) Recourſe muſt then be had to a particular Machine adapted to the Purpoſe of compreſſing the Aneuriſm; which, if ſmall, may, by the Aſſiſtance of that Inſtrument and a ſtrengthening Plaſter, be compleatly cured. Among the ſeveral Inſtruments contrived for this Purpoſe, we have ſelected the two repreſented in *Tab.* IX. *Fig.* 8. and 9. the Uſe and Application of which may be better underſtood from Inſpection, than a verbal Deſcription. We have alſo, in my Opinion, ſufficiently explained it in our Expoſition of *Tab.* XI[a].

VII. If the Aneuriſm is too large to receive any Benefit from Compreſſure by Deligation, or the preceding Inſtrument; or, if a true Aneuriſm ſhould, by a Rupture of the arterial Coats, degenerate into a ſpurious one, attended with a livid Tumor from the extravaſated Blood, Immobility of the Arm, intenſe Pain, and the Danger threatened from an accidental or profuſe Hæmorrhage; in that Caſe the Patient can have no Relief, but from the Operation by the Knife. Which Operation, however, being attended with much Pain and Danger, ought not to be undertaken without great Care and Circumſpection, and with the Approbation and Advice of other eminent Phyſicians and Surgeons: leſt, if the Succeſs thereby ſhould turn out worſe than expected, it might be raſhly attributed to Imprudence or Miſconduct in the Operator. Treatment of large Aneuriſms.

VIII. There are chiefly two Things required in the Operation: firſt, a Removal of the Tumor or Aneuriſm; and then to conjoin or heal up the Wound in the Artery. In the laſt Century they uſed to amputate the Arm for an Aneuriſm in *Italy*, and then applied an actual Cautery to the divided Artery, as we are told by BARTHOLIN, in his *Hiſtor. Aneuriſmat.* But at preſent we endeavour to preſerve the Patient's Arm, and remove the Aneuriſm by a much milder Treatment. For the ſucceſsful Performance of this Operation, the Surgeon muſt attend chiefly to three Things: firſt, to ſtop the Flux of Blood through the What is required in the Operation

[a] SCULTETUS alſo deſcribes and figures an Inſtrument for this Purpoſe, in his *Armament. Chirurg. Edit.* 4to. *Anno* 1666. *Tab.* XIX. *Fig.* 4. But his does not ſeem ſo well adapted as ours. DIONIS likewiſe mentions the Inſtrument contrived and uſed by Dr. BOURDELOT (deſcribed at large in BLEGNI's *Zod. Med. Gallic.* 1681. *page* 43.) for himſelf, by which *Pouton* or Bridge, he relates, that, within the Space of a Year, he was cured of an Aneuriſm in his Arm, as big as a Pullett's Egg.

the Artery by the Tourniquet, an Inſtrument unknown to the Ancients: ſecondly, to denudate the Artery, and free it from the adjacent Integuments; and, laſtly, to contract or conſtringe the ſame, either by Medicines or Ligature[a]. It will therefore firſt be neceſſary to have all the proper Inſtruments conveniently diſpoſed in Readineſs in a large Plate or Diſh, that there may be no Delays in the Operation. This *Apparatus* muſt take in a *Tourniquet*, to compreſs the brachial Artery, (ſee *Part* I. *Book* I. *Chap.* II. *Sect.* IX. and X. *& ſeq. ad* XV.) a *Scalpel*, *Tab.* I. G. and a Hook, *Tab.* VIII. *Fig.* 2, and 3, to denudate the Artery: to which add a Sponge with ſome warm Wine, or its Spirit, a Pair of obtuſe-pointed Sciſſors, *Tab.* I, C, and D, ſome ſcraped Lint, ſquare Compreſſes of ſeveral Sizes, one narrow Compreſs of a Span in length, with two large Pieces of Linen to inveſt the Arm: and, laſtly, two or three Rollers of two Fingers breadth, and thrice as long, as for Phlebotomy in the Arm. But if the Artery is to be contracted by Aſtringents or Cauſtics, the Succeſs of which is very dubious and uncertain, you muſt then enlarge your *Apparatus* with ſome *Vitriolum Romanum*, *Butyrum Antimonii*, *&c.* Or if you ſecure the Artery by Ligature, which is the ſafeſt and univerſal Practice of the Moderns (becauſe the Eſchar made by Cauſtics has been often obſerved to give way, and excite a fatal Hæmorrhage) inſtead of Aſtringents or Cauſtics, you muſt then provide a crooked Needle armed with ſome ſtrong waxed Thread, twice or thrice doubled; or, inſtead of a Ligature, by a Needle and Thread, you may apply the particular Inſtrument invented by me for this Purpoſe, and repreſented in *Tab.* VII. *Fig.* 4.

How the Patient and Aſſiſtants are to be diſpoſed.

IX. Your *Apparatus* being prepared, the Patient is next to be ſeated in a Chair, leaning back with his Arm extended, in the ſame Manner as for Phlebotomy. Then you muſt place four Aſſiſtants round him, in the moſt advantageous Poſition: and when the Aneuriſm is in the right Arm, it is, in my Opinion, beſt for the Surgeon to ſtand on the right Side of the Patient, placing the moſt expert of the Aſſiſtants next him, to hold the diſordered Arm above the Tumor, together with the Tourniquet applied to it; that he may increaſe or diminiſh his Stricture on the Arm by that Inſtrument, as the Surgeon ſhall direct. One of the other Aſſiſtants ſtanding before the Patient, is to hold the Arm faſt by the *Carpus*, that he may not flinch, or withdraw it in the Operation. A third Aſſiſtant is to ſtand on the left Side, holding the *Apparatus* of Inſtruments. The fourth, or laſt Aſſiſtant, muſt be ready to do any thing the Surgeon may find neceſſary to direct him, during the Operation. But if the Aneuriſm is in the left Arm, the Surgeon and Aſſiſtants are to be diſpoſed in the reverſed Order, as any one may eaſily direct.

Application of the Tourniquet.

X. The firſt Part of the Operation conſiſts in applying the Tourniquet about the Middle or upper Part of the *Humerus*, ſo as thereby gradually to compreſs the brachial Artery, (ſee *Tab.* III. *Fig.* 1. K.) till you can perceive no Pulſation either in the Artery at the *Carpus*, or in the Aneuriſm itſelf. By which means you will be ſure to avoid any conſiderable Hæmorrhage: but you muſt be careful to moderate your Stricture by the Tourniquet, ſo as not to injure the Nerves, or other ſenſible Parts. The Stick, by which the Tourniquet is twiſted,

[a] Surgeons formerly cloſed the Artery, by cauterizing with a red-hot Iron; but that is a Method too cruel, and is, at the ſame time, not ſecure, and often has pernicious Effects.

muſt be held by an Aſſiſtant on the right Side; or if you uſe the Screw Tourniquet, repreſented in *Tab.* V, and VI. that will remain faſt on the Arm, without holding. But it ſometimes happens, as GARENGEOT obſerves in his Surgery, *Chap.* on Aneuriſms, that the Tourniquet cannot be ſafely applied to the Arm in a ſpurious Aneuriſm, by reaſon of the great Extravaſation and Tumor. In that Caſe you may therefore, as the Author directs, apply the Tourniquet over a Ball and Compreſs in the *Axilla*, ſo as to compreſs the Artery, by twiſting the Stick of the Tourniquet above upon the Shoulder.

XI. When the Tourniquet is properly fixed and tightened upon the Arm, there are then three Methods of performing the Operation. The firſt of theſe is, by laying open the true Aneuriſm by a longitudinal Inciſion, continued upward and downward by the Scalpel, according to the Length of the compreſſed Artery: which done, you are to remove the vitiated Blood or Matter therein lodged, either by your Fingers, the Probe, or a Sponge. The Parts being thus cleanſed, you muſt, in the next place, ſlacken the Tourniquet a little, that the ſalient Blood may demonſtrate the upper Orifice of the Artery to you. And in doing this, you need not conſtringe your Tourniquet again immediately, if the Patient be ſtrong, and of a full Habit: but rather permit the Artery to diſcharge a few Ounces of Blood, more or leſs, as may be thought proper. When you have again tightened your Tourniquet, ſo as to exclude the leaſt Hæmorrhage, if your Intention is to treat the Diſorder by Cauſtics and Styptics, you muſt inſert a Bit of blue Vitriol, wrapt up in Cotton or Lint, into the upper Orifice of the Artery; ſecuring it there by ſeveral ſmall Compreſſes, each a little larger than the other, and filling up the reſt of the Space on all Sides with rude Bundles of Lint. You muſt then make a ſtrict Bandage, after preſſing it cloſe with the Fingers and Thumb of your left Hand, over the affected Artery of the diſordered Arm. Inſtead of intruding a Piece of Vitriol into the Orifice of the Artery, you may apply a Doſſil of Lint dipped in, and expreſſed out of the Styptic Liquor of WEBERUS, or in Butter of Antimony: the Effect of which, being ſecured with Compreſſes and Lint as before, will be equal to, if not better than the firſt we propoſed. Over the Dreſſings muſt be applied a ſquare Plaſter, and a large Compreſs of the ſame Form, to be cloſely retained by a Bandage, three or four times as long as is commonly uſed for Phlebotomy in the Arm. M. DIONIS makes his Deligation without the Piece of Vitriol, for which he ſubſtitutes a Lump or two of chewed Paper, or Lint, dipt in ſome Styptic, which he covers with ſeveral ſmall Compreſſes, each larger than the other, and ſecures the whole upon the inciſed Artery by Deligation: which Method of dreſſing may, in many Caſes, be convenient and proper enough.

First Method of Operation.

X. But in order the more effectually to prevent a future Hæmorrhage, it will be neceſſary to apply another Bandage over the former: And, after making ſome circular Rounds with it upon the Part affected, it is to aſcend up the Arm upon the long Compreſs impoſed on the brachial Artery on the Inſide of the Arm, as we directed in the preceding Chapter. That this laſt Bandage may adhere more firmly, it will be neceſſary to paſs it round the *Thorax*, when arrived to the Shoulder, and to faſten it off upon the Arm, diſpoſing the Patient to reſt. When your Dreſſings are thus compleated, and the Tourniquet a little looſened, you muſt obſerve whether any Blood iſſue through the Bandage:

Treatment after Deligation.

dage: and if there be no Appearance of any, it is a Sign your Operation is well performed.

Procedure in Cases of an Hæmorrhage.

XIII. But if you perceive any Blood ooze through the Dreſſings, the Artery muſt be again compreſſed by the Tourniquet, your Dreſſings taken off, and re-applied with more Care and Exactneſs: or elſe a more certain Method muſt be taken to ſecure the End of the Artery, by Ligature, with a crooked Needle and a double waxed Thread, which is the only infallible Means of defending the Patient from a fatal Hæmorrhage, and was formerly propoſed by PAULUS AEGINETA[a], one of the moſt ancient among the *Greek* Phyſicians. But in making this Ligature, the Surgeon muſt have a principal Regard to two Things. He muſt avoid injuring both the Artery itſelf and the adjacent Nerve. In order to which, it will be moſt convenient to make your external Inciſion through the Integuments ſufficiently large, and then carefully to ſeparate the Nerve from the Artery, to which it is attached, by a ſmall Hook: and then to paſs the Head, or obtuſe End of the Needle, foremoſt under the Artery, till you can take hold of the Thread, that its Point may not hurt either that Veſſel or the Nerve. Or elſe, inſtead of a Needle, you may paſs your Ligature under the Artery, by the Inſtrument which I contrived for that Purpoſe in *Tab.* VIII. *Fig.* 4. C. This Inſtrument is to be withdrawn when your Ligature is opened and drawn a ſufficient Length from under the Artery, which is then to be tied with it upon a thin Compreſs of ſcraped Lint, with which you are to defend or inveſt the Artery before the Conſtriction of your Ligature. The Artery being thus ſecurely tied up, you leave about a Hand's Breadth of the Thread or Ligature hanging out of the Wound: in which Manner it is to continue till the Artery is cloſed, and the Ligature comes off ſpontaneouſly. There are ſome Surgeons who alſo direct the lower Orifice of the inciſed Artery to be ſecured by a Ligature as well as the upper: and there are others again who think the ſame to be uſeleſs, or even miſchievous, as indeed it may be, when the Diſorder being in the Flexure of the Arm, the larger Inciſion and Cicatrix this Way made, will, in ſome meaſure, impede or ſtiffen the Motion of the Joint. But if the Aneuriſm be not in the Joint, or in the lower Part of the *Cubitus*, and you perceive Blood to iſſue from the lower Orifice of the divided Artery, then you may, and even ought to make a ſecond Ligature below, as well as above. And thus, after I had tied the upper Orifice in an Aneuriſm of the cubital Artery, upon relaxing the Tourniquet, I perceived Blood ſtart from the lower Orifice, which I therefore ſecured like the other, by tying it with a crooked Needle and ſtrong Thread: ſo that by their Aſſiſtance, with the Application of Balſams, I happily cured the Patient, though a little before in very great Danger of Death. In the ſame Manner you muſt alſo make a Ligature both above and below, even in the Flexure of the *Cubitus*, if you thus find it neceſſary; or at leaſt you muſt compreſs the lower Orifice of the Artery by a proper Bandage and Compreſſes: in which Method I once accompliſhed my

[a] *Lib.* VI. *de Re Medica, Cap.* XXXVII. where he ſays, If a Tumor or Aneuriſm is formed from an Injury of the Artery, we make a longitudinal Inciſion through the Integuments: And dilating the Lips of the Wound by Hooks, we denudate the Artery, under which we paſs a Needle and double Thread, tying it above and below. The intervening Part of the Artery betwixt the Ligatures we lay open by Inciſion, and after diſcharging the Contents, we ſuppurate till the Ligatures are digeſted off.

my Cure of this Diſorder, without making a Ligature below. When the Artery has been thus ſecured by Ligatures, it is a common Practice with ſome Surgeons to divide it tranſverſely a little beneath the Ligature; that the contracting or receding of the Artery into the Fleſh may compreſs its Extremities, and the better prevent a conſequent or dangerous Hæmorrhage. But in my Opinion that Practice is improper, or at leaſt it is unneceſſary; as I have twice ſucceſsfully performed this Operation, and happily cured the Patients of their Aneuriſms without thus dividing the Artery. Laſtly, you are to fill the Wound well with ſcraped Lint, to be firmly ſecured by Compreſſes and a ſtrict Bandage, as we before directed, and as we ſhall more largely explain and demonſtrate in our third and laſt Part of Surgery or Bandages.

Method of preventing an Inflammation.

XIV. In the next place it is a common and no improper Practice with ſome Surgeons to guard againſt an Inflammation by laying Linen Compreſſes dipped in Oxycrate, on each Side the affected Parts of the Arm, to be retained by a ſpiral Bandage; and then to bleed the Patient in another Part: which may be very neceſſary Precautions in Patients of a warm and full Habit. But Phlebotomy with thoſe cooling Applications will be pernicious in ſuch as are of a cold Conſtitution, and have before loſt much Blood in the Operation or otherwiſe; notwithſtanding the *French* recommend that Treatment to be generally followed without any Reſtriction. For I have myſelf cured ſeveral in which I not only omitted Bleeding and the Oxycrate, but even uſed warm Applications of *Sp. Vini Calid. Camphorat. cum Theriaca.* Your Deligation or Dreſſing being thus compleated, the Patient is to be put to Bed, and his Arm laid in an eaſy or a little inflected Poſture upon a Pillow, and the Patient is to be ordered at the ſame time to move himſelf as little as poſſible, in order to reſtrain the Impulſe of the Blood from the Heart on the affected Artery. If you ſhould perceive the Arm to ſwell violently, and threaten an Inflammation, leſt it ſhould be occaſioned by too great a Stricture of your Bandage, you muſt take it off and apply it again as we directed at *N.* XII. preceding. But for a ſmall Tumor or other ſlight Symptoms you ſhould not haſtily remove your Bandage, for fear of a profuſe Hæmorrhage: eſpecially as Experience teaches that even a livid Swelling of the Arm may be ſuſtained in theſe Caſes without any bad Conſequence, provided the Swelling be not over painful or tenſe, nor inveſted with any of the Symptoms of a Gangrene: under which Circumſtances we have directed you to a Method in the preceding Chapter.

How to prevent an Hæmorrhage.

XV. But in order to prevent a fatal Hæmorrhage, when the Cure of an Aneuriſm is attempted by Aſtringents or Cauſtics only, without making a Ligature on the Artery, it may be proper for an Aſſiſtant conſtantly to attend and lie by the Patient, provided with a Tourniquet and the Method of applying it to compreſs the Artery in caſe of ſuch an Accident, till the Surgeon can be called to make a Ligature on the Veſſel by a crooked Needle and double Thread. But ſuch an Accident is, in my Opinion, beſt prevented at firſt by taking up the End of the inciſed Artery with a Needle and Thread, rather than to truſt to the Uncertainty of a Conſtriction or Eſchar made by Cauſtics. 'Tis alſo a prudent Practice of ſome Surgeons to arm their Needle with three Threads; which being paſſed under the Artery, two of them are tied and the other left looſe to be faſtened afterwards by itſelf when the other Threads are relaxed, ſo as to permit a freſh Hæmorrhage.

XVI. With

Management of the Dreſſings.

XVI. With regard to the Bandage and Dreſſings, if they adhere firmly upon the Parts, they ought not to be removed on any ſlight Occaſion, before the third or fourth Day; except a great Inflammation, Tumor, or Hæmorrhage ſhould make it neceſſary to renew the ſame. Then the Surgeon muſt take care that the Tourniquet be duly applied and fixed upon the Arm, or elſe the Artery compreſſed by the Fingers of an Aſſiſtant before he proceeds to take off the Bandage and Dreſſings: and even then he ought not violently to force off the Compreſſes if they adhere, which might bring on a profuſe Hæmorrhage; rather let them remain. When he has cleanſed the Wound as much as poſſible, he ſhould fill it with freſh Lint ſaturated with ſome digeſtive Ointment, leaving ſuch Parts as adhere to be ſpontaneouſly ſeparated in the ſucceeding Dreſſings. In this Diſorder the Dreſſings ought to be repeated as ſeldom as poſſible, eſpecially within the firſt fifteen Days; and then it ſhould be made with all the neceſſary Cautions to prevent a Rupture of the Artery and a profuſe Hæmorrhage.

Treatment of Inflammatory Heats. Fevers, and other Symptoms.

XVII. If within a few Days after the Operation the Patient is ſeized with an Inflammation or Fever, from the intenſe Heat and increaſed Motion of the Blood, threatning an Hæmorrhage or a Gangrene in the affected Arm, he muſt then be inſtantly bled in the other Arm. In the mean time a cooling Regimen and Medicines are to be uſed, and Phlebotomy again repeated in Proportion to the Patient's Habit and the Urgency of the Symytoms. The Diet ſhould be light, ſpare, and cooling, conſiſting chiefly of ſmall Broths and diluent Suppings; induſtriouſly avoiding all hard and ſtimulating or heating Food, as is uſual in large Wounds and other Inflammations.

Agglutination of the Wound.

XVIII. When the Orifice of the Artery is cloſely conſolidated or united, which in common Aneuriſms uſually ſucceeds in ten Days or a Fortnight's Time, your Buſineſs is then to agglutinate or heal up the external Wound in the Integuments, by treating it either with dry Lint or vulnerary Balſams: obſerving in the mean time to make the Patient gently bend and extend his Arm at Intervals. Without this Precaution he may be troubled with an obſtinate Rigidity or Stiffneſs of the Joint and an Incurvation of the Arm: Partly for want of attenuating and diſperſing the Synovia, or Mucilage of the Joint, by repeated Motions; and partly from not ſtretching or extending the Cicatrix as it becomes gradually formed and more indurated.

PURMAN's Method of operating.

XIX. Another Method for curing Aneuriſms is, by fixing the Tourniquet on the Arm, as we before directed: then making an Inciſion through the Integuments, without touching the Aneuriſm, and having freed the diſordered Artery from its Adheſions to the adjacent Nerves, it is then elevated by a Hook ſufficient to paſs a crooked and obtuſe-pointed Needle under it, or our Inſtrument, *Tab.* VIII. *Fig.* 4. armed with a double-waxed Thread. By the tying of which Thread the Artery is conſtringed or cloſed; but in ſuch a Manner that you muſt always place a ſmall Compreſs of Lint upon the Artery under the Knot, leſt it ſhould cut or break through the Coats of that Veſſel. The Artery being thus tied above and below the Aneuriſm, the Tumor is next laid open by Inciſion betwixt the two Ligatures, its Contents diſcharged, and the Wound then treated as we before directed in *N.* XVI. *& ſeq.* And this laſt is the Method PURMANNUS followed in the Cure of that large Aneuriſm which he mentions, *p.* 212. of his *Chirurgia curioſa*, compleating the Cure, and healing up the Wound within the Space of a Month. We have given the Figure of this monſtrous large Aneuriſm in *Tab.* IX. *Fig.* 6. partly for its Uncommonneſs, and to illuſtrate

illuſtrate the Nature of the Diſorder; and partly to refute the Opinion of GOVEIUS [a], *viz.* That a true Aneuriſm never exceeds the Size of a Cheſnut.

XX. The third and laſt Method of performing the Operation for the true Aneuriſm, is by returning or preſſing back the Blood out of the Aneuriſm into its correſponding Artery (which in large Aneuriſms, where the Blood is very much concreted, is a thing impracticable [b]) where this can be effected, the Tourniquet is then applied to the Arm, and a longitudinal Inciſion made through the Integuments as before, without at all injuring the Aneuriſm itſelf by the Scalpel: this done, and the Artery freed from its Adheſions to the Nerve and Parts adjacent, it is then compreſſed by Ligature with a Needle and Thread as before; only without making any Inciſion in the Artery afterwards: by which means the Blood is prevented from returning into the Aneuriſm or diſtended Part of the Artery. You are then to treat the Wound with Digeſtives, as before, till the Ligatures and morbid Part of the Artery are caſt off ſpontaneouſly; after which you may heal and cicatrize as we before directed. This is the Method by which ANELIUS [c] happily cured a very dangerous Aneuriſm within the Space of a Month, at *Rome*. This he prefers, as one may hereby avoid the making a large Wound and Cicatrix, which are the conſtant Attendants of opening the Aneuriſm by Inciſion, and diſcharging its contained Blood, either by the Fingers or Inſtruments, which greatly protracts the Cure of the Diſorder, as well as renders it more painful, and attended with a diſagreeable and uneaſy Scar. After the Operation was performed as above, ANELIUS bled the Patient four times in the oppoſite Arm; and indeed repeated Phlebotomy is recommended by all the other *French* Surgeons who have treated on this Diſorder. But though ſuch repeated Bleeding may be of great Service in abating the Motion and *Impetus* of the Blood, in their warm Climate and Conſtitutions: yet, in our more northern or colder Countries or Conſtitutions, I think it may be very well omitted; as it would too much weaken the Patient, and, as I have happily cured ſeveral Aneuriſms without it.

A third Method by ANELIUS.

XXI. If, as I have ſometimes obſerved, the Coats of the true Aneuriſm ſhould burſt ſpontaneouſly, ſo as to extravaſate the Blood, it then degenerates into a ſpurious Aneuriſm, for which there is no Cure but by the Knife. Here therefore you muſt firſt of all apply the Tourniquet to compreſs the Artery and prevent an Hæmorrhage; you muſt then make an Inciſion through the Integuments ſufficient to diſcharge what concreted Blood may have been extravaſated and intercepted: which done, and the Wound well cleanſed, you muſt ſecure the Artery with a Needle and Thread, as in the true Aneuriſm; dreſſing and healing up the Wound as we have before largely directed.

Treatment of the ſpurious Aneuriſm.

XXII. Whenever you meet with the Brachial, Cubital, or Tibial Artery, wounded either by a Dart, Sword, or other Inſtrument, ſo that the Hæmorrhage thence proceeding cannot be ſuppreſſed, either by Bandage or Remedies, there is then no Method of ſaving the Patient ſo certain and expeditious as this here propoſed for Aneuriſms. You ought firſt to apply the Tourniquet, then denudate the Artery; and, if it be very ſmall, to treat it with Cauſtics or Aſtringents:

The Ligation of other Arteries in the ſame Manner.

[a] See his *Chirug.* pag. 23. 1.

[b] And therefore when the Blood cannot be returned out of the Aneuriſm, this Method will not ſucceed, but one of the former muſt be uſed.

[c] See ANELIUS's *Suite de la nouvelle Methode de guerir les Fiſtules lacrimales*, pag. 257.

gents: but if large, to ſecure it by Ligature with a Needle and Thread, as we before directed: for I may, without boaſting, declare, many are the Patients that have, with my own Hand, been by this means as it were ſnatched from the Jaws of Death. I have even recovered thoſe by Ligature, who have been almoſt ſpent and exhauſted, ſo as to look like Death, through the fruitleſs Attempts of the Surgeons, continued for ten or twelve Days together by Styptics and tight Bandage which had occaſioned their Limbs to ſwell to an enormous Size. But whether or no this Method will ſucceed ſo as to ſave the Limb, in Wounds of the large crural Artery, I have never yet had an Opportunity of experiencing, nor did I ever hear or read of it attempted by others.

Aneuriſms in the Head, Hands, and Feet.

XXIII. In the Method we have here preſcribed, you ought alſo to treat other Aneuriſms, when they are curable: which may be determined, partly from conſidering the Size and Situation of the Artery, and partly from the Size and Nature of the Aneuriſm itſelf. But, for the ſake of Beginners, I ſhall be a little more particular in my Account of other Aneuriſms, and the rather, becauſe it is a Subject of which moſt of our modern Surgeons take little or no Notice. And firſt, an Aneuriſm of the Artery betwixt the Thumb and Fore-finger, occaſioned by a Puncture from a Penknife, was cured by Compreſſion, as we are told by TULPIUS (*Lib.* IV. *Obſ.* 17.) which Compreſſion he made by applying firſt, an aſtringent Plaſter, over that a Plate of Lead, and then by a ſtrict Bandage, having firſt returned the Blood out of the Tumor; the Diſorder was cured within the Space of four Months. The ſame Treatment of Compreſſion may be therefore uſed in moſt other Aneuriſms, eſpecially thoſe which are recent, and not large, after having firſt returned or diſcharged the Blood contained in the Aneuriſm. A Woman ſtruck her Son, of ſeven Years old, ſuch a Blow on the left Side of the Head with a Stick, that, by contuſing the carotid Artery, a throbbing Tumor was inſtantly formed, about the Size of a Hazle Nut; which, in the Space of eight Days Time, grew ſo large as to cover half of his Head, from the ſagittal Suture all over the Temple and Forehead to the Eye: upon her coming for Advice, it was thought proper by the Surgeons to prefer the Operation, though a doubtful Remedy, rather than leave the Patient to the more certain Hazard of his Life. The Tumor was therefore laid open by the Scalpel, the contained Blood diſcharged, and the Wound dreſſed with Aſtringents and tight Bandage: by which means the Patient recovered in a ſhort Time[a]. Thus alſo was cured an Aneuriſm of the Artery behind the Ear, in Proceſs of Time, though with much Difficulty, by the Uſe of Aſtringents and tight Bandage[b]. If an Aneuriſm ſhould ariſe near the Ancle, like that deſcribed by RUYSCH, *Obſ.* XXXVIII. which was opened by an imprudent Operator for an Abſceſs, you ought either to make an Inciſion through the Integuments and Tumor, and to apply Aſtringents with a tight Bandage; or elſe to denudate the Artery, and ſecure it by Ligature with a Needle and Thread, as we directed before. Hence you may be alſo able to treat Aneuriſms formed in any of the other acceſſible Arteries of the Body, where there is any Proſpect of obtaining a Cure. HARDERUS *Apiar. Obſ.* p. 325. takes notice of a Patient's ſudden Death, from opening an Aneuriſm of a carotid Artery in the Neck: And

VAN

[a] See BARTHOLIN. *Epiſt. Med* 53. Cent. III.
[b] *Ephemer. Nat. Curioſ.* Cent. III. Obſ. LXVI. pag. 150.

VAN HORN has obſerved the ſame from an Apertion of an Aneuriſm in the Thigh. *Vid. Epiſt. de Aneuriſmate.*

XXIV. They who deſire a better Idea of the Manner in which the Ligatures are to be made upon the Artery for an Aneuriſm, may inſpect *Fig.* 7. in our ninth Table, where A denotes that Part of the Artery above the Aneuriſm, B the Part below, C the Aneuriſm itſelf, D the ſuperior Ligature, and E the inferior one. But here we may again obſerve, that when the Tumor is on the Flexure of the Arm, the lower Part of the Artery ſhould not be tied with a Ligature, except it be abſolutely neceſſary, for the Reaſons we before alledged. But in what Manner the Circulation of the Blood is carried on through the Hand and lower Parts, after the Operation, I cannot conceive, eſpecially when there is but one Trunk[a] of the brachial Artery near the Elbow; as muſt have been the Caſe with the Patient of ANELIUS, becauſe no Blood returned by the lower Part of the Artery, after its Diviſion, into the Tumor, notwithſtanding he did not ſecure it by Ligature. We muſt therefore defer our Inquiry on this Head, till ſome body may have an Opportunity of examining the Arm of a dead Subject who has undergone this Operation in his Life-time. Dr. WALTER HARRIS, in his eighth Chirurgical Diſſertation, openly condemns this Operation, and calls it dreadful and raſh Butchery: but for what Reaſons himſelf beſt knows. He ſeems, in my Opinion, to have been a very timorous Phyſician, who, out of Fear, or a fooliſh and ill-grounded Compaſſion, is for rejecting ſome of the moſt conſiderable and uſeful Operations in Surgery: without which, it will be impoſſible for the Patient to obtain a Cure, or even to ſurvive any Time.

Some Obſervations on this Diſorder.

CHAP. XIV.

Of injecting Liquors into the VEINS, *and of transfuſing the* Blood *of one Animal into another.*

I. WE treat next of *Injecting* and *Tranſfuſing*, as a Branch of Surgery: becauſe thoſe Operations require the Apertion of a Vein, in the ſame Manner as in Bleeding. The firſt is, the injecting ſome Liquor or Medicine into a Vein opened by Inciſion: and the laſt is, the conveying the arterial Blood of one Man, or Animal, into the Veins of another. Notwithſtanding theſe Operations are ſeldom practiſed by our modern Surgeons, yet they were highly celebrated, and often performed, in the laſt Century, from the Year 1660 to 1680: and therefore we ſhall not think much of our Endeavours here, to give the young Surgeon a clear Notion of the Affair, from whence he may alſo be able to underſtand what Reaſons gave Occaſion for the firſt Invention and Perfor-

The Operations deſcribed.

[a] I have often found two Trunks, or large Branches of this Artery, one ſituated in the inward, and one in the outward Part of the Arm; which below the Flexure of the *Cubitus* communicated again. But moſt Anatomiſts have deſcribed one only.——See my Diſſertation on a dangerous Wound of the crural Artery.——Many Surgeons, unacquainted with this Particular in Anatomy, have taken off the Patient's Arm, where there was no Occaſion, and often endangered his Life.

mance thereof; and what Advantages may be perhaps reaſonably expected from the ſame Operations, even at this Time.

Uſes expected from them.

II. The Generality of Phyſicians, not without Reaſon, attribute moſt Diſorders of the Body to ſome Vice in the Blood. Therefore what Method can be more ready to remove or correct that Vice, than injecting a proper Medicine into the Veins to mix with the Blood itſelf, or the transfuſing the ſound Blood of one Man or Animal into the Veins of another, inſtead of that which is diſeaſed? For by this means the Action of a Medicine on the Blood will be immediate and entire, without being impaired or changed by paſſing the Stomach and Inteſtines, and mixing with various Juices before it arrives to the Veins. But there are even many Caſes which occur, wherein no Medicine at all can be taken by the Mouth, as in Apoplexies, Anginas, the *Hydrophobia*, *&c.* which may poſſibly be this way remedied, when they cannot by any other. And if plentiful Bleeding is ſo ſerviceable in many Diſorders, as the Leproſy, Gout, Epilepſy, Apoplexy, Conſumptions, Scorbutics, Venereal Diſeaſe, malignant Fevers, *&c.* by diſcharging the peccant Matter in the Blood, as it is by many Phyſicians allowed: even the Objections of other Phyſicians againſt it, as weakening the Patient, *&c.* may, by theſe Operations, be obviated or removed. Even old Age may be ſupported, and the very worſt Habits of Body corrected by theſe Means, ſo as to give a firm, juvenile, and healthy Conſtitution. Theſe, and ſuch like, are the vaſt Expectations which have been formed from the preſent Operations by Phyſicians: but the Misfortune is, that they not only meet with Diſappointment in their good Views, but even frequently the Event turns out worſe than the Diſeaſe. For almoſt all the Patients who have been this way treated, have degenerated into a Stupidity, Fooliſhneſs, or a raving or melancholy Madneſs, or elſe have been taken off with a ſudden Death, either in or not long after the Operations. Theſe lamentable and fatal Conſequences have brought the Art of Injections and Transfuſions into Neglect at the preſent: ſo that, being ſuſpected and condemned by proper Judges at *Paris*, where they moſt flouriſhed, we are told they were in a little Time prohibited by a publick Edict of that Parliament.

The Art of Injection deſcribed.

III. Notwithſtanding this, we ſhall give the young Surgeon an Idea of the Manner in which Liquors were formerly, and may now be injected into the Veins of living Men, or other Animals. And firſt, a Vein is to be opened, uſually in the Arm, by your Lancet, as in Bleeding: and having introduced the ſmall Pipe of a Syringe, or a very ſmall Clyſter-pipe with a Bladder (*Tab.* XI. *Fig.* 10.) the contained Liquor is injected or forced into the Vein upwards towards the Heart. Which done, you are to dreſs the Orifice, and make your Deligation upon the Arm in the ſame Manner as after Phlebotomy. But whether or no this Method of injecting proper Medicines into the Blood may ſucceed, eſpecially in deſperate Apoplexies, Anginas, *Hydrophobia*, *&c.* and whether it may not be often uſeful to diſcharge the morbid Blood, and transfuſe ſuch as is ſound, or warm Milk or Broth in its ſtead, ought, in my Opinion, to be determined by future and repeated Experiment. BURMANNUS, in his *Surgery*, (*Part* III. *Cap.* 31.) tells us, that he has not only performed the Operation with Succeſs on others, but alſo very happily upon himſelf, being by this means cured, not only of a troubleſome Itch, but alſo of a ſtubborn Fever. A profeſſed Treatiſe on the Subject has been publiſhed by ELTSHOLTZ, intitled, *Clyſmatica*

tica Nova, ſive Chirurgia infuſoria & transfuſoria, 8vo. 1667. *Editio ſecunda, cum Fig.*

IV. For the Transfuſion of Blood into the Veins, you are firſt to open a Vein in the Patient's Arm or Hand, as at *Fig.* 11. and 12. *Tab.* XI. and then thruſt gently upward into it a ſmall Tube of Silver, Braſs or Ivory. The ſame is to be alſo done with the ſound Perſon; only the Tube muſt here be inſerted downward towards the ſmall End of the Vein. This done, the ſmalleſt of the Tubes is to be inſerted into the other larger one, by which means as much Blood will paſs from the ſound Perſon into the Patient as may be thought proper, and then the inciſed Veins are to be dreſſed or bound up as in Bleeding. But if the Patient does not recover after one Transfuſion, the Operation ſhould be repeated again at convenient Intervals. But before the Patient receives the Blood of the ſound Perſon, he ought to be bled proportionably, that the new Blood, laſt received, may have the freer Circulation. Sometimes a Vein is opened in each Arm of the Patient at the ſame Time, that as much of the vitiated Blood may flow out of one Orifice as he receives of the ſound by the other. For more on this Subject, among others, the Reader may conſult LAMZWARD *in Notis ad* SCULTETUM, and JUNCKEN *Chirurg. Germanica*, pag. 487, where you have Figures of the Operation. If the Blood is to be transfuſed out of ſome Animal into the Patient, then a Calf or a Lamb, for Example, are to be ſecured by Ligatures, and one of their Veins or Arteries opened, either in the Neck, Leg, or Thigh; and the reſt of the Operation managed as before. See *Tab.* XI. *Fig.* 13. and LAMZWARD *in Append. ad* SCULTETI *Armament. Chirurg.* and BURMANNI *Chirurg. P.* III. *Cap.* 31. Laſtly, where Tubes of Metal or Bone were found painful and leſs convenient, for want of being flexible, Operators contrived to faſten an intermediate flexible Pipe betwixt the two others, ſuch as Part of the carotid Artery, or of the Ureter from an Ox, Calf, or Lamb, or the Wind-pipe of a Capon, Duck, &c. by which means the Proceſs becomes much facilitated both to the Operator and Patients.

The Method of Transfuſion.

V. The Contrivance of this Artifice, by which the Blood of one Animal is transfuſed into the Veins of another, is aſſumed by Dr. LOWER, in his Treatiſe *De Corde*, in oppoſition to M. DENIS, who, in his *French* Epiſtle upon this Subject, claims the Invention to himſelf. It is true, the latter made many Experiments in this Way at *Paris*, but with very bad Succeſs. STURMIUS, once a celebrated Profeſſor of the Mathematics at *Altorf*, and VERHUIS, Profeſſor at *Francfort*, attribute the Invention to MAURIT. HOFFMAN: whereas MUYS aſſerts, that LIBAVIUS deſcribed the Proceſs at large in the Year 1615, but without telling us the Book. The firſt Injection of Liquors into the Veins of Animals is generally attributed to the celebrated Sir CHRIST. WREN; but I think we have this Artifice deſcribed before him, by a Profeſſor of Phyſic, in a Treatiſe publiſhed *An.* 1664, in which he explains the Proceſs that had never before been heard of in *Germany*. They who deſire more on this Subject may conſult MAJORIS *Lib. de Chirurgia infuſoria*, ETMULLER *Diſputat. in eod. Argumento conſcript.* ELTSHOLTZ *Clyſmatica nova*, BURMANNI *Chirurgia*, LOWER *De Corde*, SANTINELLUS *in Confuſione Transfuſionis*, MANFREDUS *De Sanguinis Transfuſione*, STURMIUS *in Philoſophia Eclect. Diſſ.* X. MERCKLINUS *De Ortu & Occaſu Transfuſionis Sanguinis*, LAMZWARD *in Appendice ad* SCULTETUM, pag. 29.

The Invention of this Art.

For

For Injections into the Veins in deſperate Diſeaſes, See *Miſc. Nat. Cur. Ann.* IX. and X. *pag.* 144. and LOWTHORP *Phil. Tranſact. Abr.* Vol. III. page 226 to 235.

An EXPLANATION *of the* ELEVENTH PLATE.

Fig. 1. Repreſents an Arm in which a Vein is to be opened: A denotes the Cephalic Vein, B the Baſilic, and C the Median Vein; D the Ligature fixed above the Elbow to make the Vein ſwell.

Fig. 2. Repreſents the ſeveral Forms of inciſing a Vein with the Lancet: A ſhews a longitudinal Inciſion, B a tranſverſe one, and C, D, oblique ones.

Fig. 3. Exhibits the antient *German* Phlebotomus or Fleam for opening a Vein, A the ſharp Point to be fixed on the Vein, B the Handle to be held in one Hand, while the Part C is ſtruck by a Fillip of the Finger of the other Hand, ſo as to drive the Point A into the Vein.

Fig. 4. Is a Spring-Fleam, now in Uſe with ſome. The Part A being fixed on the Vein, and the Part C being elevated, depreſſes the Spring by the End B, which by its Reaction or Elaſticity ſtrikes the End C upon the Fleam A, ſo as to drive it into the Vein. DD is a hollow Caſe of Braſs or Silver, in which the Spring-Part of the Inſtrument B is included.

Fig. 5. Repreſents the *French* Phlebotomus or Lancet, bent ſo as to form an obtuſe Angle, as it ſhould be, for the more convenient holding it in Bleeding.

Fig. 6. Is the great Aneuriſm, as big as one's Head, obſerved by BURMANNUS in an Arm near the Joint or Bend of it.

Fig. 7. Shews the Manner of applying the Ligatures above and below an Aneuriſm, in the Operation for that Diſorder. A B the Artery, C the Aneuriſm, D the upper Ligature, E the lower Ligature.

Fig. 8. Exhibits an Inſtrument contrived both for the Prevention and Cure of Aneuriſms. A A A denote the Plate of Iron or Steel adapted in Form to the Flexure of the Arm, B its Fiſſure, CC Ligatures faſtened to the Ends A A, and extended to DD. E denotes a moveable Steel Plate, joined by the Hinge, I, and covered with a Cuſhion of Cotton or Silk at F, to be fixed upon the Aneuriſm, GG are two ſmall Hooks by which the Inſtrument is faſtened upon the Arm by the Ligatures C C D D, H is a Screw by which the Plate and Cuſhion E F are preſſed down upon the Tumor.

Fig. 9. Repreſents an Inſtrument of the ſame Kind with the former, but of a different Shape. Here the Plate and Cuſhion E F are larger, for bigger Aneuriſms than the former. Its Parts and Explanatory Letters correſpond to thoſe of the preceding Figure.

Fig. 10. Shews the Apparatus with a Bladder and Tube for Injection of Liquors into the Veins: A the Bladder and Tube, B a Vein of the Arm opened, in which the Tube is inſerted.

Fig. 11 and 12 Exhibit the Transfuſion of the Blood from the Veins of one Man into thoſe of another: B denotes the recipient, and A the remittent Arm.

Fig. 13. Shews the Transfuſion of Blood from the crural Artery or Vein of an Animal into the Arm of a Man by the Intervention of the Tube A.

CHAP.




Fig: 1.
Fig: 2.
A B C D
C
Fig: 3.
A
B
Fig: 4
C
D
D
Fig: 5.
A
Fig: 6.
C
C
Fig: 7.
D
C
D
D

Plate II.


Fig: 1.
A
D
C
B
Fig: 2.
A B C D
B
C
Fig: 3.
A
B
Fig: 4
C
A
D
D
Fig: 5.
A
A
Fig: 6.
C
C
B
Fig: 7.
D
E
A
B
C
D
D
A
C
A
C
D




Fig: 8.

Fig: 9.

Fig: 10.





Fig: 11.




Fig: 12.



Fig: 13.





CHAP. XV.

Of INOCULATION *for the* SMALL POX.

The Deſign of this Chapter.

I. THE Art of ingrafting or propagating the Small-pox by Inciſion or Inoculation, has been an Operation equally famous in all Nations with thoſe in the preceding Chapter. Therefore we ſhall, for the Sake of Beginners, deſcribe the Proceſs of it, which under proper Circumſtances may be of great Service to Mankind.

Inoculation deſcribed.

II. The Deſign of this Operation is, to communicate, by Art, a milder Species of the Small Pox to the Infant or adult Patient, than that received by the natural Infection: and this by ingrafting ſome of the variolous Matter. In order to which a ſmall Inciſion[a] is to be firſt made with a Scalpel or Lancet through the Skin of the Arm, and having inſerted a ſmall Particle of the purulent Matter taken from a mild Kind of the Pock, the little Wound is then to be dreſſed with ſome dry Lint, and covered with a Plaſter. After the Operation, the Patient muſt conſtantly keep to his Chamber, the Air of which ſhould be moderately warm, and his Diet regulated by ſome prudent Phyſician, by which means the Diſorder will ſhew itſelf in about ſeven or eight Days, without any malignant Symptons: and, if aſſiſted by a proper Regimen and moderate Warmth, it uſually runs gently through its ſeveral Stages. When the Patient has once had the Diſorder this Way, though never ſo mild, we are aſſured by Experience, that he never has it again: and therefore the Opinion of thoſe ſeems to be well grounded who think the Propagation of the Small Pox by Inoculation might be of general Uſe and Benefit to Mankind, in preſerving the Lives of ſome, and the moſt important Members of others, as the Face, Eyes, Ears, Viſcera, *&c.*

The Operation practiſed by other Nations.

III. Hiſtory informs us, that the Diſorder was this way propagated many hundred Years ago among the *Greeks* and *Turks:* whereas it is but of late Years that the *European* Nations have come into it; among which the *Engliſh* ſeem to have approved and followed it moſt. The Experiment ſucceeded ſo well in the Hands of the *Britiſh* Phyſicians, that the late King *George* himſelf countenanced the ſame in all his Dominions: and from thence the Practice prevailed with Succeſs in *Germany*, particularly at *Hanover*, *Onolſbac*, and *Pyrmont*.

The Objectors againſt this Practice.

IV. It muſt, however, be confeſſed, that there are many, both among the *French* and *Engliſh*, who endeavoured to ſuppreſs and vilify this Practice in their public Libels, condemning it as fatal to Mankind, and unfit to be encouraged among a Chriſtian People; but, I think, all they have objected or advanced has been long ago ſufficiently anſwered and obviated by the learned Dr. JURIN, and other able Phyſicians. They who deſire more particular Accounts, may conſult the Diſſertations publiſhed by the celebrated Phyſician laſt mentioned, as alſo thoſe by PYLARINUS of *Italy*, the celebrated VATERUS of *Vitemberg*, *Act. Erud. Lipſ. Ann.* 1723, 1725, *Act. Natur. Curioſ.* Vol. I. *Obſ.* LXXV. *p.* 133, *&c.* And laſtly, they may conſult Experience, the beſt Phyſician of all.

V. But

[a] But Dr. HARRIS, in his *Chirurgical Diſſertations*, directs only the Cuticle to be braded, and the variolous Matter to be ſpread on the naked Skin.

My own Opinion of it.

V. But, for my own Part, if I may ſpeak freely, I am ſo far from thinking the Practice fatal or miſchievous, that I rather firmly believe it might, under a proper Management, be of the greateſt Uſe and Benefit to the Lives and Healths of Mankind. For, if I think right, the Small Pox ariſes from a peſtilential Virus or Matter lodged in the Blood from the very firſt Day of the Birth, which breaks out almoſt in every Perſon ſooner or later: and the more early, uſually the better. For it is very ſeldom we obſerve the Pock favourable in thoſe more advanced in Years: ſo that the Matter ſeems to multiply itſelf in the Blood, and augment with the Patient's Age. And this, in my Opinion, is the Reaſon why we oftener meet with the Small Pox more mild and favourable in Infants than in Adults. If therefore the Diſorder be procured of a mild Kind by this Operation, and the Blood cleared of its latent Virus, while ſmall in Quantity, and the Infant young, I doubt not but many, and eſpecially the Children of Princes and Nobility, might be thus not only preſerved from Death, but even conducted ſafely through the ſeveral Stages of the Diſeaſe, without the Attacks of its moſt malignant Symptoms. We are convinced by Experience, as well as Reaſon, that the Diſorder which breaks out from a natural Infection is generally more ſevere and fatal than that procured by Art: and no Wonder it ſhould be ſo; ſince in the laſt the Phyſician has an Opportunity of chuſing the moſt favourable Seaſon, and of preparing his Patient beforehand by a proper Regimen, Diet, and Medicines.

CHAP. XVI.

Of SCARIFICATION *and* CUPPING.

The Method of dry Cupping.

SCarification and Cupping was an Operation frequently performed by the moſt ancient Surgeons and Phyſicians[a], notwithſtanding the Moderns have, by their Pride or Neglect, turned the Buſineſs over to thoſe who attend the Baths or Hot-houſes. Yet, as it makes none of the leaſt Operations in Surgery, we ſhall here briefly conſider and explain the ſame. The Operation of Cupping is indeed vague, and not confined to any particular Member of the Body. But whenever the Cupping-glaſs is applied, it is fixed upon the Skin, either entire or ſcarified: and hence we have a twofold Diſtinction of Cupping into *dry* and *gorey*. The Figure of the Cupping-glaſs, for either of theſe Purpoſes, is repreſented in *Tab.* XII. *Fig.* 1. In dry Cupping, the Glaſs adheres to the Skin by expelling or rarifying its included Air by lighted Flax or the Flame of a burning Candle within it, ſo that the Glaſs is preſſed upon the Part with a conſiderable Force by the external Air; in which Artifice our ordinary Cuppers are ſufficiently well verſed. The Uſe of this dry Cupping is twofold; either to make a *Revulſion* of the Blood from ſome particular Parts affected, or elſe to cauſe a *Derivation* of it into the affected Part upon which the Glaſs is applied. Hence we have a Reaſon why HIPPOCRATES[b] orders a large Cupping-glaſs to be applied under the Breaſts of Women who have a too profuſe Diſcharge of their Menſes, intending thereby to cauſe a Revulſion of the Blood upwards from

[a] As we read in HIPPOCRATES, CELSUS, GALEN, *&c.* [b] Sect. V. Aphor. 50.

from the Uterus. And upon the ſame Principle I have myſelf ſucceſsfully cured a profuſe Hæmorrhage at the Noſe, and an Hæmorrhage or Spitting of Blood from the Lungs, by applying Cupping-glaſſes to the Legs and Feet, particularly about the Ancles and Knees. SCULTETUS give us a remarkable Inſtance in *Obſ.* 85. of a Woman, who, by the repeated Application of ſix Cupping-glaſſes (without Scarification) to her Thighs, was not only relieved of the troubleſome Symptoms, cauſed by an Obſtruction of her Menſes, but was alſo thereby freed from the Obſtruction itſelf. Dry Cupping is alſo uſed with Succeſs to make a Revulſion by applying the Glaſſes to the Temples, behind the Ears, or to the Neck and Shoulders, for the Removal of Pains, Vertigo's, and other Diſorders of the Head. [a] They are alſo applied to the Upper and Lower Limbs to derive Blood and Spirits into them when they are paralytic; and, laſtly, to remove the Sciatica and other Pains of the Joints. [b] The Operation is, in theſe Caſes, to be repeated upon the Part till it looks very red, and becomes painful.

II. But Cupping is much oftener joined with Scarification, than uſed alone with us in *Germany*, and in other northern Countries: in which Caſe, the Part is firſt to be dry-cupped till it ſwells and looks red, and the Skin is to be punctured or inciſed by the Scarificator, *Tab.* XII. *Fig.* 2. with which you may make ſixteen or twenty ſmall Wounds in the Skin, cloſe enough to each other to be covered by the Cupping-glaſs, into which the Blood ought to flow from them. (*See Fig.* 3.) In repeating theſe Inciſions, and re-applying the Cupping-glaſs upon freſh Parts of the Skin, the Operator muſt obſerve to begin at the loweſt Part, and thence aſcend gradually, that his Work may not be obſcured by the refluent Blood from above. Having ſcarified the Skin, and applied the Cupping-glaſs with Fire, as before directed, the latter will adhere firmly to the Part, and the Preſſure of the external Air will force a conſiderable Quantity of Blood into it from the Inciſions. But as ſeveral Glaſſes (ſometimes ſix or eight) are often applied at one and the ſame Time, and to different Parts of the Body, the Operator muſt manage his Buſineſs ſo that ſome Glaſſes may be filling, while he is ſcarifying and adapting the others: and in thus ſhifting them alternately, he muſt pour out their Blood into a Pan or Veſſel, waſh them in warm Water, cleanſe the Skin with a Sponge dipped in the ſame Water, and then apply the Glaſſes as before. When the Blood ceaſes to flow faſt enough, you muſt repeat your Inciſions with the Scarificator cloſe by the former, and re-apply your Cupping-glaſſes till a ſufficient Quantity of Blood is drawn, or till it ſtops of its own accord. Your Operation being finiſhed, and the Skin well cleanſed with a Sponge and warm Water, it is next to be rubbed over with a Bit of Deer's Suet to promote the Healing. But if the Blood ſtill continues to flow, which it does but ſeldom, you are then to waſh the Skin with *Sp. Vini*, *Aq. Reg. Hungar.* binding it up with a Compreſs and Bandage.

Cupping with Scarification.

III. The modern Surgeons have, for Conveniency to themſelves and Eaſe to the Patient, contrived a Scarificator, different from the laſt-mentioned, which conſiſts of ſixteen ſmall Lancet-blades fixed in a cubical Braſs Box, with a Steel Spring, as at *Fig.* 4. *Tab.* XII. When the Side of this Inſtrument marked CCCC is applied to the Skin, and the included Spring bent by the Lever A, by depreſſing

The modern Scarificator.

[a] See CELSUS, *Lib.* iv. 6. 2. [b] See DECKER's *Extract.* p. 34.

fing the Button B, it is ſo ſuddenly let looſe as by its Force to ſtrike the Points of the ſixteen Blades out of the Caſe at one Inſtant into the Skin, making as many ſmall Inciſions at once in their regular Order, over which the Cupping-glaſs is to be applied, as we before directed. We meet indeed with the Figure of a Scarificator not much differing from this in PAREY's Surgery, *Book* XI. *Chap.* 5. and after him in LAMSWARD's Notes to the *Armamentarium* of SCULTETUS. But they do not propoſe the Inſtrument for other Uſes than to ſcarify the unſound Parts in an incipient Mortification: whereas this is uſed with good Succeſs by our Cuppers in many other Diſeaſes, as I myſelf have frequently ſeen and experienced. Though M. GARENGEOT [a] condemns it as a bad and uſeleſs Inſtrument: but perhaps that Gentleman never ſaw the Uſe and Effects of it.

Uſes of Scarification.

IV. Cupping with Scarification is uſed in various Parts of the Body, particularly in the Head, Neck, Shoulders, behind or under the Ears, Occiput, Back, and Loins, Legs and Arms, and near the [b] Ancles: and this for making a Derivation, Revulſion, or Evacuation in the various Diſorders incident to plethoric Habits; ſuch as various inflammatory Diſorders in the Head, Eyes, Ears, Tonſils, and Uvula, particularly violent Head-achs, Ophthalmia's, Amauroſes, Suffuſions, *&c.* In all which Caſes it is hardly poſſible to expreſs the general Benefit which may be received from this Operation, eſpecially when timely uſed, and judiciouſly repeated at proper Intervals. Nor is Scarification much leſs beneficial than Phlebotomy in thoſe Patients, whoſe Veins are ſo ſmall or obſcurely ſituated, that it would be dangerous opening them by the Lancet: yet as it is often abſolutely neceſſary to make a Diſcharge of Blood ſome way from them, I have often adviſed this Method to be followed, and with good Succeſs. The excellent Anatomiſt MORGAGNI [c] adviſes Scarification upon the Occiput in Apoplexies, and all ſoporous Affections, as one of the beſt Remedies that can be recommended, either from Reaſon or Experience: becauſe in this Way the heſitating Blood may be diſcharged from the obſtructed Veins of the Brain, which communicate with thoſe of the Occiput, or at leaſt it may, by this means, obtain a more free Motion: but then you ought to ſcarify deep, as he obſerves. Scarification and Cupping upon the Occiput is alſo extremely uſeful in an Ophthalmia, or Inflammation of the Eyes, and a like Diſcharge procured by deep Scarification upon the affected Side in a Pleuriſy, after Phlebotomy premiſed, gives great Relief, according to LANCISI [d]. Laſtly, this Method of evacuating by Scarification and Cupping, makes one of thoſe which are generally repeated at ſtated Seaſons of the Year, like Bleeding and Purging, Spring and Fall, *&c.* which the Patient being once accuſtomed to, ought never to neglect them, for fear of incurring their former, or even worſe Diſorders.

Scarification by ſome rejected and deſpiſed.

V. I muſt indeed own, that there are many among our Phyſicians and Surgeons who contemn this Operation as of little or no Efficacy: and the Reaſon which they offer is, that hereby only that Blood is diſcharged which lodges itſelf betwixt the Fleſh and Skin. But this Judgment ſeems too haſtily formed, and with out

[a] Tract. *De Inſtrument. Chirurg.* Tom. I. pag. 413. [b] Scarification of the Ancles is highly recommended by Jo. TAC. MANN, of *Padua*, in a Treatiſe, publiſhed *A. D.* 1583. [c] *Adverſar. Anatom.* V. pag. 83. & VI. pag. 108. ZACUTUS LUSITANUS alſo mentions a Patient freed from an Apoplexy by repeated Scarification. [d] *Ibid. Adverſ. Anat.*

without a juſt Foundation. For Experience has taught myſelf and many other eminent Phyſicians, that as much and as thick Blood may be diſcharged by Scarification and Cupping, as by Phlebotomy, and conſequently it muſt be little leſs, if not equally beneficial, in all thoſe Diſorders which require Bleeding. But this I can boldly affirm, from my own Reaſon and Experience, that in ſome Caſes Scarification excels Phlebotomy, in as much as the Cupping-glaſs, by firmly adhering to the Skin, not only draws out the Blood, but alſo gives it a greater Impetus or Tendency towards the ſcarified Part; and therefore it conſtantly gives certain and ſpeedy Relief in moſt Diſorders of the Head, Eyes, and Ears, Apoplexies, ſleepy Diſorders, Inflammations of the Tonſils, Hæmorrhages, and Pains of various Kinds, *&c.* ſometimes by Revulſion, and ſometimes by Derivation.

Whether Scarification be dangerous.

VI. There are again other Phyſicians, who imagine Scarification to be not only uſeleſs, but even pernicious: for, ſay they, We have Inſtances of Patients who have been not only violently diſordered, but even killed by the Operation being performed at an improper Time, or with an unclean or infected Inſtrument. [a] Thus a Patient may be in danger of catching ſome foul Diſorder by being ſcarified with an Inſtrument that has not long before been uſed upon one infected with the Leproſy, Pox, Itch, *&c.* for thus the Infection will be inoculated almoſt in the ſame Manner with the ſmall Pox [b]. But if Scarification muſt be condemned and rejected on this Account, ſo muſt alſo Phlebotomy and many other Operations, in which the ſame Inſtrument is applied that has been uſed before. But that the Patient may have no Uneaſineſs from this Quarter, it may not be improper for him to ſee that his Cupper's Scarificator and Apparatus are very clean: or elſe they may keep a Scarificator of their own, which being kept clean and dry, can give no room to make any frivolous Scruples of this Nature.

The Scarification uſed by Surgeons.

VII. There ſtill remains another Sort of Scarification, uſed by Surgeons in violent Inflammations, incipient or confirmed Mortifications, peſtilential Carbuncles, and the like. In which Caſes it has been found highly ſerviceable to diſcharge the ſtagnant and vitiated Blood, by making many ſmall Wounds or Inciſions in the Skin with a Scalpel or Lancet, though without the Aſſiſtance of Cupping-glaſſes. This Kind of Scarification is uſually denominated *Chirurgical* by the Cuppers, in contradiſtinction to theirs: As Surgeons uſe it frequently in Gangrenes and Mortifications, and ſometimes in ſwelled Legs and Dropſies, eſpecially that of the *Scrotum*, and ſometimes for the *Hydrocephalus*. But though it may be ſometimes highly neceſſary to ſcarify the Legs of dropſical Patients, when the Skin is diſtended ſo as almoſt to burſt: yet it ought not to be made indiſcriminately, without abſolute Neceſſity, and a proper Regard to the Patient's Age, Habit, *&c.* Otherwiſe it is even probable, that the ſcarified Part will gangrene or mortify, and deſtroy the Patient. PLINY (*Hiſt. Nat. Lib.* LXXVIII. *Cap.* 1, & 11,) recommends Scarification of the Gums for the Tooth-ach; which, in my Opinion, may not unfrequently be very uſeful.

VIII. Re-

[a] Thus HILDANUS, *Cent.* V. *Obſ.* 71. remarks, that a Palſy aroſe from hence, though it might proceed from a Multitude of different Cauſes.

[b] *Vid.* JORDANUS, *De luc novâ in Moravia.* SPORICHIUS *De Symptom. Crudel. a Scariſicatione.* LIBERIUS *De malitioſâ ſcarificatione, in Obſ.* GREG. HORSTIS, *L.* IV.

The *Egyptian* Scarification.

VIII. Related to Scarification is the Infliction of ſmall Wounds within-ſide the Noſe, Lips, Ears, and Gums, uſed by the *Egyptians*, and recommended by [a] CELSUS and [b] ARETÆUS for abating Inflammations, and relieving various other Diſorders, in which it very often ſucceeds admirably. At which we need the leſs wonder, if we conſider what Relief Nature herſelf often gives the Patient, by making a plentiful Hæmorrhage at the Noſe, in ardent Fevers, Head-achs, &c. Add to this, that the *Egyptians* [c] had a Practice of beating or whipping the Calves of the Legs with Rods, till they looked red, and then ſcarifying, or making Inciſions in the Skin: by which means they procured Relief, and made uſeful Revulſions from the Head and Brain in violent inflammatory Diſorders of thoſe Parts, and in Fevers with Delirium, Watchings, &c. But notwithſtanding the Uſefulneſs of this Practice, it is at preſent hardly ſo much as known among our *European* Nations.

Scarification of the Eyes.

IX. Many of the antient Phyſicians and Surgeons, with HIPPOCRATES, had a Practice of ſcarifying the Inſides of the Eye-lids, and even the Eyes themſelves, with a proper Inſtrument for the Purpoſe, in many of the Diſorders which infeſt that Organ, as is very apparent from the Treatiſe which HIPPOCRATES has left, *De Viſu* [d]. This Operation of ſcarifying the Eyes, though neglected from the Time of HIPPOCRATES, has yet been renewed, or lately introduced again, by the *Engliſh* Oculiſt WOOLHOUSE, at *Paris*: and it has been alſo performed with tolerable Succeſs by ſome others of the preſent Age, as we have Accounts. But for the Inſtruments, and Manner of performing this Operation, we ſhall be more particular in our following Account of the Operations for the Eyes.

CHAP. XVII.

Of BLEEDING *by* LEECHES.

Choice of the beſt Leeches.

I. LEECHES, or *Sanguiſugæ*, are a Species of aquatic Worms or Inſects, of the Shape repreſented in *Tab.* XII. *Fig.* 5. which being applied to any Part of the Body, bite through the Skin, and extract Blood from the ſmall Veins, which frequently conduces much to the Health and Recovery of a Patient: for which Reaſon they have been uſed from the moſt early Times by the antient *Greek* and *Roman* Phyſicians, as may be ſeen in GALEN's profeſſed Diſſertation on this Inſect, commented on by SEBEZIUS. As there are Leeches of different Kinds and Natures, it will firſt be proper to diſtinguiſh and make a due Choice of the beſt. Theſe are always found in clear Brooks or Rivulets: whereas thoſe taken from Lakes, Fiſh-ponds, and ſtagnant Waters generally have ſomething malignant in their Bite, inſomuch as ſometimes to excite great Pain, Inflammation, and Tumor, in the Part, and Uneaſeneſs in the whole Body.

It

[a] *Lib.* IV. *Cap.* 2. where he directs to draw Blood from the Noſe in violent Head-achs.

[b] *De Chron. Morb. Lib.* II. *Cap.* 11. *De Cephalia*, pag. 128.

[c] PROSP. ALPINUS. *Medicina Ægyptior.* p. m. 72. where you have a Figure of this Practice.

[d] CELSUS likewiſe adviſes Scarifications in the Head for many Diſorders of the Eyes, and particularly violent Inflammations, *Lib.* VI. *Cap.* 6. The Succeſs of which I myſelf have experienced.

It is alſo an Obſervation made by ſome of the moſt expert Surgeons, that the beſt Leeches have ſlender and pointed Heads, with greeniſh and yellowiſh Lines or Streaks on their Backs, and their Bellies of a reddiſh yellow: whereas thoſe are the worſt, or moſt malignant, which having a thick and obtuſe Head, and incline from a dark blue to a black Colour on the Back and Sides. But you ought to obſerve it as a neceſſary Caution, never to apply Leeches which have been lately catched in Rivers or foul Water, before they have been kept ſome time in a Glaſs full of clean Water, to be often ſhifted, that they may cleanſe themſelves from what Filth or Venom they may have imbibed: and when they have been thus kept for a few Months, they may be afterwards ſafely uſed, without incurring any bad Accident.

Method of applying them.

II. Before the Leech is applied to the Skin, it ſhould be taken out of the Water to ſtand about an Hour in an empty Cup, or other Veſſel, to drain itſelf; that being thus rendered thirſty and empty, it may both adhere more firmly to the Part, and draw off a larger Quantity of Blood. As for the Part to which they may be applied, that may be on the Temples or behind the Ears, when the Diſorder lies in the Head or Eyes, and eſpecially when the Patient is delirious in a Fever, or overcharged with Blood. But ſometimes they may be commodiouſly enough applied to the Veins of the *Rectum*, in Diſorders proceeding from an Obſtruction of the wonted Evacuation this Way, or in the blind and painful Piles: and by way of Revulſion they will be here uſefully applied in profuſe Hæmorrhages of the Noſe, and ſpitting or vomiting of Blood; in which Caſes they are of incredible Service, eſpecially when the Diſorder ariſes from Obſtructions of the hæmorrhoidal Flux. But before you apply the Leech, the Skin of the Part muſt be firſt well rubbed till it becomes hot and red. Which done, you take hold of the Leech by its Tail with a dry Cloth, or you may place it leaning half way over the Edge of a Cup, and ſo apply it that it may creep out upon the Part; which they are no ſooner fixed upon, but they generally bite and draw the Blood very eagerly. When ſeveral Leeches are to be uſed, you muſt apply each of them to the Part in this Manner ſucceſſively; and if they ſhould refuſe to bite or adhere to the Skin, as they ſometimes do, you may, in that Caſe, put a little Blood of a Pigeon, Chicken, *&c.* upon the Skin. But if that will not allure them, you moſt apply freſh Leeches in their ſtead. The Application of Leeches to the Caruncle in the greater or inner Canthus of the Eye, is found to be extremly uſeful in all inflammatory Diſorders of that Organ, after Phlebotomy has been firſt premiſed.

Treatment after their Application.

III. When the Leeches are diſtended with Blood, they generally ſeparate from the Skin, and leave the Part of themſelves. But if it be neceſſary to draw ſtill a large Quantity of Blood, you muſt either apply freſh Leeches, or elſe cut off the Tails of thoſe which are drawing with a Pair of Sciſſors, by which means the Blood will run through them, and they will draw almoſt as long as you pleaſe. If the Leeches do not ſeparate ſpontaneouſly after a ſufficient Quantity of Blood has been evacuated, upon ſprinkling a little Salt or Aſhes upon the Part, they uſually leave it preſently: which Method ſhould be the rather taken, becauſe forcing or pulling them away often occaſions a Tumor and Inflammation of the Part. The Operation being thus finiſhed, thoſe Leeches which are whole may be returned into the Glaſs again and reſerved for future Uſes: but thoſe die which have had their Tails cut off. The Wound made by this Inſect may be firſt waſhed with warm Wine or Water, and then dreſſed with ſome vulnerary

nerary Plaſter: though there is ſeldom any Occaſion for the latter, as it generally heals up faſt enough of itſelf. They who deſire more upon this Inſect, may conſult GALEN, ALDROVANDUS, GESNERUS, BOTALLUS, PETR. PAUL. MAGNUS, SEBIZIUS, HEUNIUS, CRAUSIUS, SCHRADERUS, STAHLIUS, *&c.* who have wrote thereof more at large.

CHAP. XVIII.

Of ACUPUNCTURATION, *uſed by the* Chineſe *and* Japoneſe.

SOmewhat a-kin to Scarification is the famous Operation of the *Chineſe* and *Japoneſe*, termed *Acupuncturation*. Thoſe Nations rejecting Scarification and Phlebotomy as pernicious, have recourſe to their Acupuncturation and Cauterization, or burning with *Moxa*, as their moſt potent Remedies in almoſt all Diſorders. The firſt of theſe Operations they perform with a large Gold or Silver Needle (*Tab.* XII. *Fig.* 6.) which they ſtrike into the Fleſh, either with their Hand or the little Hammer, *Fig.* 7. It is indeed more than a little ſurpriſing, that ſo deſperate and ſevere an Operation ſhould be ſo much practiſed by a People in other reſpects judicious: and that too, in the Head, Breaſt, Abdomen, Arms, Legs, Thighs, and moſt other Parts of the Body; even in the Abdomen of Women with Child, when the Fœtus is reſtleſs. But I do not know that the Practice has been received by any of our *European* Nations: and therefore, as the Proceſs is ſo much abhorred, we ſhall not here give a prolix Account thereof. They who deſire more, may conſult RHYN *De Artbritide* pag. 145, 183, 190; and KOEMPFER *in Amœnitatibus exoticis*, pag. 582; alſo in his Deſcription of *Japan:* in which Country both theſe Surgeons were Spectators of the Operation.

CHAP. XIX.

Of ISSUES.

The Seat of Iſſues.

I. ISSUES are little Ulcers made deſignedly by the Surgeon in various Parts of the Body, and kept open by the Patient, for the Preſervation or Recovery of his Health. They are by ſome [a] denominated *Cauteria*, but improperly: becauſe by that Term we uſually mean a cauſtic or corroding Medicine. In this Operation the Phyſician endeavours, by Art, to imitate and relieve Nature; who often forms Ulcers in various Parts of the Body of her own accord, for diſcharging pernicious Humours, whereby People are often freed from grievous Diſorders, and enjoy a healthy State. The Parts in which Iſſues are generally made, are either, (1.) the upper Part of the Head; (2.) the Neck; (3.) the Arms, betwixt the Biceps and Deltoid Muſcle, near the Inſertion of the laſt;

[a] CAPIVACCIUS has a Diſſertation *De recta Cauteriorum Adminiſtratione*, in which he treats only of Iſſues, which the *French* alſo term *Cautere*.

laſt; (4.) in the Thighs, eſpecially within-ſide, immediately above the Knee, in a Cavity eaſily felt by the Fingers; and laſtly, (5.) Iſſues are ſometimes made in the Legs, on their interior Side, in a Cavity immediately below the Knee.

The firſt Method of making Iſſues by Inciſion.

II. Though there are ſeveral Methods of making Iſſues, yet none ſeem to be more ready than the following: *viz.* Firſt, to mark the proper Place with Ink; and then elevating the Integument betwixt the Thumb and Fore-finger of the Surgeon and an Aſſiſtant on each Side, you next proceed to make an Inciſion through them, either with the Scalpel or Lancet, big enough to admit a Pea. Which being inſerted and covered with a Plaſter and Compreſs, nothing more is wanting than your Roller to compleat the Operation. Thus by cleanſing and dreſſing the Wound every Morning and Evening with a freſh Pea, it by degrees, in a Day or two, degenerates into a little Ulcer, diſcharging daily a Quantity of purulent Matter; which ſhould be carefully cleanſed or wiped off at every Dreſſing.

A ſecond Method by the actual Cautery.

III. There is a ſecond Method of making Iſſues by wounding the Skin with a red-hot Iron, or actual Cautery; which is uſually included in a Sort of Capſula, or Caſe of Iron, *Tab.* XII. *Fig.* 8. A, to conceal it from terrifying the Patient. When the Caſe B B is fixed upon the proper Part for the Iſſue, the Cautery, or red-hot Iron C, is then preſſed down upon the Integuments, and the Eſchar or Burn, is next to be dreſſed with freſh Butter, or *Ung. Baſilic.* till it at length ſeparates in repeating the Dreſſing every Day. Then the little Ulcer formed is to be filled with a Pea, and dreſſed as before. Though this Method of making Iſſues, according to the Antients, is more ſevere, yet it muſt be equally more efficacious than the other; as the Pain and Cauterization muſt neceſſarily make a conſiderable Revulſion. But there are very few Patients who will ſubmit to it.

A third Method by Cauſtics.

IV. The third and laſt Method of making Iſſues, is, by the Application of potential Cauteries, or corroding Medicines: in order to which, a Piece of Plaſter is firſt perforated, as in *Tab.* XI. *Fig.* 11. and then applied, ſo as its Aperture may cover the Place marked with Ink for the Iſſue. A Piece of the Cauſtic, mentioned *Part.* I. *Book* IV. *Chap.* III. *Sect.* XI. is then impoſed upon the Aperture of the Plaſter, and retained cloſe down upon the Skin with ſome ſcraped Lint, a ſmall Compreſs, and a large Plaſter; and laſtly, with a larger Compreſs and Bandage. The Operation thus far advanced, the Patient is now to be ordered to reſt about ſix or eight Hours, more or leſs, according as the Cauſtic may be in Strength: which Time being elapſed, and the Dreſſings removed, the Eſchar is to be treated as we before directed at *Sect.* III.

V. But in which ever of theſe Methods you make the Iſſue, it muſt be dreſſed at leaſt twice every Day, eſpecially if it runs well, and in the Summer time; and at each Dreſſing you muſt put in a freſh Pea, and cover it with a clean Plaſter, or a Piece of waxed Paper or Silk, or an Ivy-Leaf, retained with Compreſs and Bandage. But the Deligation for Iſſues is much more commodiouſly performed with a leathern Swathe, faſtened by Claſps, as in *Tab.* XII. *Fig.* 9. than by a circular Linen Roller. It is remarkable, that ſome uſe Peas of Silver or Wood to dreſs their Iſſues with, inſtead of the common ones: but the Difference in their Effects is not material. In this manner Iſſues are to be kept open, till the Patient is recovered of the Diſorder for which they were made; and

and in ſome Caſes they ſhould be continued as long as the Patient lives: or if the ſame Diſorder, or ſome other, returns upon drying them up, they muſt be again opened immediately.

Uſe of Iſſues.

VI. Iſſues are uſed chiefly for various Diſorders in the Head, Eyes, Ears, the Teeth, or Sciatica, and other painful Diſorders, which are this way frequently relieved or cured. The Uſe and Advantage of Iſſues is well known, and daily experienced by moſt Surgeons[a], contrary to the Opinion of Helmont, and ſome others, who think they ſerve only to torment and trouble a Patient. However, I muſt frankly own, that a Cure is not to be expected from Iſſues; and though they generally give ſome ſmall Relief, yet in many Caſes I have found it too inconſiderable to be ſenſible: but if, upon Trial, they afford no great Benefit, it is beſt to dry them up again in a little Time. But we muſt not forget to take notice, that it is frequently neceſſary to make two or more Iſſues, to produce any conſiderable Effect in ſtubborn Diſorders, as one in each Arm, or in one Arm and Leg of the ſame Side, *&c.*

Method of drying up Iſſues.

VII. In order to cloſe up an Iſſue, when that ſhall be judged proper or neceſſary for various Reaſons, little more is required than to diſcharge the Pea, and refrain from putting in any more, by which means alone it will cloſe up in a ſhort Time. But if any proud Fleſh ſhould protrude itſelf, it may be amputated, or elſe removed and taken down with *Alum. uſt.* Laſtly, it is obſervable, that when Iſſues of People far advanced in Years ceaſe to make their wonted Diſcharge, and turn of a livid and blackiſh Hue, it is a Sign they are invaded by ſome deſperate Diſorder, and that even Life itſelf is very near its Period. In this Caſe proper topical Remedies ſhould be ſpeedily applied, as *Cantharides, Rad. Irid. Florent. vel Helleb. nigr.*

CHAP. XX.

Of Bliſtering *with* Cantharides.

Bliſtering deſcribed.

I. BY Bliſtering is underſtood an Elevation of the Cuticle, from the Cutis into Veſicles, or Bladders replete with a ſerous Humour, by the Application of external Remedies, and chiefly Cantharides, to the Skin, which may be applied either in form of a Paſte mixed up with Yeaſt; or elſe mixed with ſome Emplaſter, and then ſpread on Linen or Leather, which is the modern Practice: and therefore we conſtantly meet with the *Emp. Veſicator.* ready prepared in the Shops of Apothecaries. Theſe being applied and retained upon the Part with Bandage and Compreſs, in about eight, ten, or twelve Hours Time, will raiſe the Cuticle under the Plaſter in a Bliſter, replete with a thin and acrimonious Lymph. The beforementioned Number of Hours being expired, the Bliſter-plaſter is removed; and the Cuticle, if yet entire, is opened with a Pair of Sciſſors, its Contents being gently abſorbed by Lint or ſoft Linnen. This done,

[a] Vid. Galvani *Trattato delle Fontanelle.* Glandorpii *Gazophylacium, &c. Act. Hafnienſ.* Vol. iii. xii. Muysii *Prax. Med. Obſ.* 2. Schelhamerus, Fred. Hoffmannus, Hilscherus, *&c.* in *Diſſertationibus.*

done, the Part bliſtered is dreſt with ſome ſoft and cooling Plaſter, which Dreſſing is repeated every Morning and Evening, till the Diſcharge ceaſes, and the Part heals. And though it is remarkable, that the Cuticle is ſeparated from the true Skin by this Plaſter, in the ſame manner as it is in Burns; yet it meets with ſo ſudden a Reproduction, as is not a little ſurpriſing. Some make their Dreſſings with Beet or Dock-leaves, ſpread with freſh Butter, inſtead of a Plaſter.

The Size of Bliſter-plaſters.

II. The Size of Bliſter-plaſters varies greatly with the Nature of the Patient's Diſorder, and the Size or Figure of the Parts to which they are to be applied. Thoſe for the Temples and behind the Ears, may be about the Size of a Crown Piece: As may alſo thoſe for the Neck and Arms, Legs and Thighs, and the Top of the Head. But thoſe for the Back and between the *Scapulæ*, may advance to two Hands Breadth.

The Uſe of Bliſter-plaſters.

III. Veſicatories are commonly of very great Benefit, as well as Iſſues, in many of the moſt obſtinate Diſorders: eſpecially when vicious Humours are to be diſcharged from the Blood, or a ſtrong Revulſion to be made from any Part. Thus Veſicatories are of excellent Service behind the Ears, upon the Head, Neck, Arms, *&c.* in all Inflammations of the Eyes, and Suffuſions or incipient Cataracts; as they likewiſe are in all lethargic and paralytic Affections: in which Caſes they give a Stimulus to the Blood and Spirits, and excite thoſe Fluids from a languid to a briſk Motion. Strong Veſicatories are alſo frequently uſed in ardent Fevers attended with a Delirium: in which Diſorders they are properly applied to the lower Extremities, in order to diminiſh the Influx of Blood ſent to the Head and Brain. Laſtly, Bliſters are uſed with great Succeſs in the Small Pox[a], when the Puſtules ſeem to ſtrike in; as alſo in the more obſtinate arthritic and rheumatic Complaints, where they are beſt applied even to the Part in pain, according to the Obſervation of SCULTETUS (*Obſ.* 73.) Bliſters are alſo of great Efficacy when applied to the Legs and Thighs in Aſthmas; and a little below the Elbow for the Tooth-ach.

How to increaſe the Force of Bliſters.

IV. When the Diſeaſe requires a conſiderable Diſcharge this Way, it may be convenient to mix a little Powder of Cantharides with the Ointment or Plaſter, with which the Bliſter is to be conſtantly dreſſed: by which means greater Benefit may be obtained than one would imagine, in many of the moſt obſtinate Diſeaſes.

V. There is another Caſe, in which *Cantharides* are of great Service. When Iſſues, or Ulcers of the Legs, which are of long ſtanding, ſtop, either ſpontaneouſly, or from ſome Diſorder (eſpecially if the Patient is advanced in Years;) in this Caſe ſprinkle *Cantharides* on the Iſſue or Ulcer: let a Piece of Bliſtering-plaſter be made in the Form of a Pea, and be applied to the Part. The Humours that before ceaſed, will generally be provoked by the Stimulus of the *Cantharides*, and be diſcharged again to the great Benefit of the Patient. Add to this, that in ſome Wounds (particularly after the Operation of Lithotomy) where the Lips of the Wound become callous, the Application of *Cantharides* deſtroys the Calloſity; and the Wound heals without Difficulty. See CHESELDEN, and DOUGLASS, *Lib. de Lithot.* But this Application is ſometimes attended with an *Ardor Urinæ*, or great Heat and Pain in making Water; eſpecially if the Bliſters are ſeveral in Number, and ſtronger, or continued longer on the Parts than uſual: in which Caſe the Patient ſuffers the ſame Symptoms as if

Bliſters often accompanied with *Ardor Urinæ*.

[a] See FREIND's excellent *Treatiſe on Fevers* and the *Small Pox*.

if he had taken *Cantharides* internally. But then theſe troubleſome Symptoms are as quickly removed by a frequent and plentiful drinking warm Milk, and amygdalate Emulſions. Laſtly, Bliſters ſhould not haſtily, but with great Caution, be uſed for Patients who are hydropic or cachectic; becauſe they frequently produce an incipient or confirmed Mortification. See BAGLIVI on Bliſters, HOFFMAN, VATER, and others.

CHAP. XXI.

Of INJECTIONS.

Of Injections in general.

MANY Diſorders are very difficultly, if at all, curable, unleſs the Parts affected are injected with ſome proper Liquor, by means of a Syringe and a proper Tube. Which Operation is by Surgeons called *Injection*, and conſiſts chiefly in drawing the Liquor into the Syringe, and forcing it out again into the diſordered Parts. The Method of performing which is too obvious for any body to be ignorant of. But this Obſervation may be neceſſary, To apply the Syringe and Tube to the Parts very carefully, eſpecially in very ſenſible or nervous Parts, to avoid giving the Patient too much Pain: alſo to be mindful, that the Liquor you inject be not too hot or cold. But what Kinds of Liquors and Methods are to be uſed for Abſceſſes and fiſtulous Ulcers, we have before obſerved (in the Book on *Ulcers*, *Chap.* II. *N.* III.)

In Diſorders of the Mouth and Fauces.

II. In Ulcerations and Inflammations of the Tonſils, Uvula, and Fauces, Injections are generally uſeful: but Care is to be taken to preſs down the Tongue with a Spatula (*Tab.* I. *litt.* P.) or the flat End of a Spoon; and having introduced the Syringe two or three Fingers Breadth into the Mouth, the Injection is to be gently thrown in, ſeveral times. A proper Syringe for this Purpoſe is deſcribed by DEKKERUS (*Exercit. Pract.* pag. 242.) furniſhed with a crooked Tube, whoſe Extremity is perforated with ſeveral ſmall Holes, as in *Tab.* VI. *Fig.* 11. This Inſtrument is particularly uſeful, when the Patient's Mouth cannot be eaſily opened by a Spatula, which is often the Caſe.

In Gonorrhœas.

III. Injections are alſo frequently thrown into the Urethra of the Penis, in Men under a Gonorrhœa, in order to waſh out the corrupt Matter, and mitigate the Heat, Acrimony, and Pain. The beſt Syringe for this Purpoſe is that in *Tab.* IV. *Fig.* 10. fitted with a convenient Tube to enter the Penis. Alſo the Syringe in *Tab.* XII. *Fig.* 10. may be very commodiouſly uſed in this Caſe; becauſe the Liquor does not eaſily fly out of it behind. The moſt convenient Liquors for abating the Heat and Pain in this Diſorder, are, warm Milk and Barley-water, ſweetened with Sugar, Honey, or Syrup of Marſhmallows; and after the Uſe of theſe, when we would heal up and ſtrengthen, or gently aſtringe the Parts, we may uſe the following Mixture with Succeſs:

℞. *Aq. Plantag.* ℥iv. *Mell. Roſat.* ℥j. *Sacch. Saturni* ℈j. *M. f. injectio.*

In Diſorders of the Uterus.

If a ſmall Stone ſhould happen to ſtick in the Urethra, its Exit may be very much promoted by injecting Oil of ſweet Almonds or Olives by the Penis. For Diſorders in the Uterus, to expel the After-burthen, when it adheres too ſtrictly to the Womb, or to cure Ulcers in that Part, or cleanſe the *Fluor Albus*,

bus, it is convenient to inject ſome deterging and healing Liquor, by the Syringe which MAURICEAU has deſcribed for that Purpoſe. See *Tab.* VI. *Fig.* 12 and 13. But when this Syringe is uſed, the Surgeon ſhould be careful that its foremoſt high Tube be cautiouſly introduced into the Vagina. To anſwer this End in a ſtubborn *Fluor Albus*, I have experienced the Syringe at *Tab.* XII. *Fig.* 10. to be very convenient.

In Diſorders of the Thorax and Abdomen.

IV. Laſtly, for the Manner in which Liquors are to be injected into the *Thorax* or *Abdomen*, to cure Ulcers or Wounds in thoſe Parts, that has been before deſcribed, when we treated of Wounds. As to the Liquors which are injected by the *Anus*, under the Title of Clyſters, we ſhall conſider them when we come to treat of the Operations proper to that Part.

CHAP. XXII.

Of ACTUAL CAUTERIES.

The ſeveral Sorts of Cauteries.

I. CAUTERIES are by Phyſicians and Surgeons diſtinguiſhed into two Claſſes, actual and potential. By actual Cauteries they intend red-hot Inſtruments, uſually of Iron, which are applied to many Parts and Diſorders. By potential Cauteries we underſtand certain Kinds of corroding Medicines, of which we ſhall ſpeak hereafter in *Chap.* XXIV. Of actual Cauteries, or hot Irons, it is neceſſary for the Surgeon to have a conſiderable Apparatus: inaſmuch as different Diſorders require Cauteries of various Sizes and Figures. Notwithſtanding there are a greater Number of cauteriſing Inſtruments deſcribed and figured by the Writers in Surgery [a], the chief of which we have given you in *Tab.* III. yet it may be neceſſary for the ſkilful Surgeon to invent others, ſuitable to the particular new Diſorders which may ſometimes occur to him.

The Uſe of Cauteries.

II. Cauteries have various and manifold Uſes. For they are not only uſed to deſtroy the dead Parts of carious Bones, in Cancers, to remove *Scirrhi*, Excreſcencies, Carbuncles, and mortified Parts; but alſo to make Iſſues and Setons, to ſtop Hæmorrhages in Wounds and Amputations: and, Laſtly, to remove an Amauroſis, Epilepſy, Sciatica [b], with Pains in the Teeth and other Parts. We are therefore ſo far from condemning the Uſe of Cauteries, as have SEPTATIUS, HELMONT, BONTEKOE, OVERKAMPIO, CRAHN, *&c.* that we rather recommend them as eminently ſerviceable in many of the beforementioned Diſorders. They who are deſirous of ſeeing more upon this Subject, may read ALBUCASIS ARABS, and SEVERINUS concerning the wonderful Effects of cauteriſing, in his elegant Book *De Efficaci Medicina*, *&c.* He may likewiſe conſult Jo. COST. *Prof. Bonon. de igneis Medicinæ Præſidiis*, Venet. 1595. To theſe add FIENUS and BARTHOLINUS *De Cauteriis.*

The Application of Cauteries.

III. For the right Application of Cauteries, various Obſervations are neceſſary. In the firſt place, the Surgeon ſhould ſee that the Size and Figure of the Cautery correſpond to that of the diſordered Part: and while the Patient is preparing for the Operation, to let the Cautery be heating in the fire. After which,

[a] See ALBUCASIS, PARÆUS, ANDREAS A CRUCE, SCULTETUS, *&c.*

[b] SCULTETUS *Obſ.* 72. TULPIUS, *Lib.* iii. *Cap.* 36. DECKER, *Exercit.* pag. 34.

which, it will be neceſſary to ſecure the ſound Parts from the Cautery, to prevent giving more than neceſſary Pain. For this Reaſon it is, that the fleſhy Parts upon a carious Bone are firſt drawn and held aſide by the Fingers of an Aſſiſtant, before the Cautery is applied. When the Inſtrument is ſufficiently hot, it is to be applied and ſtrongly compreſſed upon the diſordered Part, till the Surgeon perceives the Bottom of the diſordered Parts appear ſound. To effect this the more ſpeedily, it will be neceſſary to have ſeveral Cauteries in Readineſs, that if one be inſufficient, he may uſe a ſecond or a third: which Caution is more eſpecially of conſequence to be obſerved in carious Bones and large Hæmorrhages.

The Uſe of Cauteries in Apoplexies.

IV. It may be here not amiſs to take notice, that ſeveral Phyſicians have found by Experience, that Cauteries have ſucceeded in Apoplexies, when all other Remedies have failed. But for the Part to which the Cautery is to be applied, there are various Opinions. SCULTETUS, in *Obſ.* 34. is for having it to be applied to the Occiput: But ZACUTUS LUSITANUS, and RIVERIUS, think it much better to cauterize between the firſt and ſecond Vertebræ of the Neck. Others again pitch upon the Meeting of the coronal and ſagittal Suture: and others prefer different Parts. MISTICHELLIUS, an *Italian* Writer upon the Apoplexy, aſſerts, that no Place can be ſo well pitched upon for Cauterizations in Apoplexies, as the Soles of the Feet. But the Manner in which the Soles of the Feet are to be cauterized in that Diſorder, the forementioned Author has endeavoured to demonſtrate in a particular Table, for which ſee *Tab.* XII. *Fig.* 11. where the Parts to be cauterized are ſignified by the Letters A A, the Cautery by the Letter B. Though that Inſtrument may doubtleſs be of another Figure than a ſquare one. I tried this Practice upon a Perſon in an Apoplexy: but, inſtead of recovering, he died.

CHAP. XXIII.

Of BURNING *with* MOXA.

TO Cauterizations it may not be improper to join burning with Flax and *Moxa*, which latter is a kind of downy Subſtance, ſeparated from the Leaves of a Sort of *Indian* Mugwort, and is uſed by the *Indian* Nations: but the firſt we find was uſed by HIPPOCRATES[a], and the other ancient Phyſicians, to cauterize Parts in pain[b]. Some of the Moderns wonderfully extolled Cauterization with *Moxa*, as the moſt effectual Means to cure, and wholly extirpate the Gout. But for the Art of cauterizing with it, it may be neceſſary to obſerve the following Particulars (*viz.*) In the firſt place to make a ſmall Cone of the Lint or *Moxa*, about a Thumb's Breadth long, (See *Tab.* XII. A B. at the Letter A and B) made much after the ſame Manner as they uſually are for a Suffitus. The Baſis of this Cone is to be ſtuck upon the Part with Gum Arabic, or Gum Tragacanth, and its Point is then to be fired by a Candle, or a burning Coal. By this means not only the Cone will be gradually conſumed, but the painful Part will be at laſt by degrees cauteriſed, and thence the Pains of the Gout

[a] *Lib. de Affect. Cap.* 30. [b] CELSUS, *Lib.* iv. *Cap.* 23.

Gout will frequently have ſome Remiſſion. But if the Pains do not intirely vaniſh at the firſt, a new Cone is to be applied again to the Part, and the Cauterization thus continued till the Pain ceaſes. But, however this Proceſs may have been cried up by many of the *Europeans*, it is at preſent quite in Diſuſe, and that not without Reaſon: for, beſides the acute Pain which it cauſes, it is frequently found to have little or no Effect. But the *Chineſe* and *Japoneſe* have the Operation at this Time in the higheſt Eſteem; inſomuch that it, with their Acupuncturation, makes their chief Remedies.

Theſe Cauterizations are ſaid to be at preſent in uſe among the *Arabians*. More may be ſeen upon this Head in Rhynius *De Arthritide*, pag. 145. Cleyerus *in Medicina Sinica*. Purmannus *in Chirurg. Pars* III. pag. 292. Pechlinus *in Obſ.* pag. 263. Valentini *Polychreſt. Exotic.* pag. 197. and a particular Diſſertation upon *Moxa:* And, laſtly, Koempfer, *in Amœnit. Exotic.* pag. 589. and in his *Hiſtor. Nat. Japon.*

CHAP. XXIV.

Of Caustic *and* Corroding Medicines.

I. CAUSTIC and Corroding Medicines, as they are called by our Surgeons, are thoſe Medicines which, being applied to Parts, conſume, and, as it were, burn them like hot Irons: Whence the *Greeks* gave them the Names of *Cauſtics*; and Celsus[a] denominates them *Adurentia* and *Exedentia*. However they differ in this from actual Cauteries, that they perform their Effects ſlower and with leſs Force and Pain: whereas in the Application actual Cauteries act inſtantaneouſly, and occaſion moſt acute Pain. Potential Cauteries differ among themſelves in various Degrees of Strength, according to their different Subſtance and Preparation: ſo that ſometimes more, ſometimes leſs, is applied to a Part for any Purpoſe. But among the various Kinds of potential Cauteries, the moſt conſiderable and effectual among us is the *Lapis Infernalis*, which is prepared *e Calc. Viv. & Cinerib. Clavellatis*, and which is applied for the opening Abceſſes, as we have before mentioned (*in Part* I. *Book* IV. *Chap.* III. N° XI.) But there are ſome who prefer Lunar Cauſtic, or a Salt prepared from a Calcination of Soap-boilers Lees, or *Ol. Vitriol.* or a Solution of Mercury in *Aq. Fort.* Butter of Antimony, and a Mixture of Soap and Quick-lime; or, laſtly, an arſenical or mercurial Sublimate, mixed with a little Honey. But it ſeems much ſafer to abſtain from the arſenical and mercurial Sublimate, leſt we ſhould occaſion thoſe grievous Diſorders and violent Pains, nay, even Convulſions and Death, which they ſometimes produce[b]. In what Manner potential Cauteries are to be applied for opening Abceſſes, and making Iſſues, we have before declared in *Part* I. *Book* IV. *Chap.* III. N° X. alſo *Part* II. *Sect.* I. *Chap.* XIX. N° IV. for theſe Cauteries are ſaid to be ſtrong enough to remove Warts, Tubercles, Excreſcences, Sarcomas, encyſted Tumors, Wens, and ſcirrhous Tumors, if they are properly applied either ſuperficially, or to the Root of the diſordered Parts. By theſe an Hydrocele may be conveniently opened, and even

The Uſe of Cauſtics.

[a] *Lib.* V. *Cap.* 7 & 8.

[b] This Hildanus mentions, *Cent.* VI. *Obſ.* 22, & 80.

even a whole cancerous Breaſt may be removed. A conſiderable Inſtance of the Succeſs of this Practice in *Germany*, we have from the celebrated SUTORIUS of *Norimberg*, afterwards Surgeon to the Duke of *Brunſwick*. But great Caution is neceſſary in this Kind of Practice, not to irritate ſuch Parts and Diſorders by theſe Medicines, which, if they ſhould prove inflexible, might endanger the Patient's Life. For thus a Scirrhus may often be turned into a Cancer: and if they are applied to the Eyes or Eye-lids, they may hurt Viſion, and may ſometimes occaſion profuſe Hæmorrhages, if applied near large Veins and Arteries. Or, laſtly, they may occaſion Convulſions by injuring the Nerves: though perhaps theſe are not all the bad Conſequences that may attend an injudicious Uſe of potential Cauteries. But for the ſkilful Application of them, we ſhall give ſome Directions hereafter.

CHAP. XXV.

Of Opening ABSCESSES.

THE Methods to be uſed for opening Abſceſſes, I think, have been already deſcribed in *Part* I. *Book* IV. *Chap.* VIII. N°. VIII. Therefore to avoid Tautology, we ſhall refer our Reader thither.

CHAP. XXVI.

Of WARTS.

The various Kinds of Warts.

WARTS are commonly known to be ſmall Excreſcences of the Skin, ſeated in moſt Parts of the Body, but chiefly in the Hands and Face. Their Size and Figure are very various: Some are very broad and flat; ſome again are very ſlender, and others appear in Form of a Pear hanging by its Stalk. Theſe are commonly removed more for the Deformity they occaſion, eſpecially in the Face, Neck, and Breaſts of beautiful Women, than for any Pain or Danger. And, notwithſtanding the great Variety of ſuperſtitious and inſignificant Remedies which are ſometimes uſed by the Populace, and even ſome Phyſicians, for the Removal of Warts, none of them are ſo expeditious and certain as the Means which come from the Surgeon.

Cure by Ligature.

II. To come therefore to the Purpoſe, we ſhall briefly deliver the chief Artifices uſed by Surgeons for the Removal of theſe Excreſcences. And the firſt that offers is that by Ligature: which conſiſts chiefly in this, violently to bind a Horſe hair, or a Piece of fine Thread or Silk about the ſlender and depending Root of the Excreſcence. By this means the nutritious Veſſels being compreſſed, and the uſual Supply of Fluids being cut off, it gradually withers and decays.

Cure by Extirpation.

III. A ſecond Method of removing theſe Excreſcences is, by ſome ſharp Inſtrument, to wit, by taking them up by a Pair of Plyers, and cutting them cautiouſly off with a Pair of Sciſſors. The Wound is to be dreſſed for ſome Time

Time with *Lap. infernalis*, or ſome other cauſtic Medicine, that if there be any of its Root remaining, out of which a freſh Tubercle might ariſe, it may be corroded and deſtroyed.

Removed by Cauſtic.

IV. But if the Excreſcence is of the larger Kind, it is more adviſeable to have recourſe to cauſtic Applications. But to make theſe act the more expeditiouſly, it may be proper to cut off the external and hardeſt Part of the Tubercle, either with a Scalpel, Razor, or Pair of ſharp Sciſſors, and then to dreſs the Wound with Oil of Tartar *per deliquium*, or Spirit of Salt. But if theſe ſeem not ſtrong enough for the Purpoſe, more vehement ones may be uſed; ſuch as Spirit and Oil of Vitriol, *Aq. Fortis*, and Butter of Antimony. On the contrary, the ſofter and ſmaller Kind of Warts may be removed barely by wetting with the Juice of Celandine, or the Milk of Spurge. In the mean time, Care ſhould be taken to prevent any of theſe Applications from getting into the Eyes, when they are uſed about the Eye-lids, which might blind the Patient. To prevent theſe Effects, it may be proper to circumſcribe the Wart with a Ring of Wax, or a perforated Piece of Plaſter, ſo that the Wart may come through; by which means the Wart only will be corroded, and the other Parts remain entire. By the ſame Methods other Tubercles and Spots, which deform a Perſon, may be removed.

Removal by actual Cautery.

V. A fourth Method of removing Warts, is by ſome actual Cautery, accommodated to the Size of the Excreſcence, ſo that it may penetrate to its Root, when applied. (See *Plate* III. *Fig.* 13, 14.) Though there are many violent Means to extirpate Warts, none can equal that of the actual Cautery; which occaſions moſt acute, though uſually but a momentaneous Pain. The Part cauterized may be often dreſſed with *Baſilicon*, or ſome other digeſtive Ointment and cooling Plaſter, ſuch as *de Sperm. Ranar.* This is the moſt certain Method of removing theſe Excreſcences in whatever Part of the Body, the Eyes only excepted; for they never return.

Removal by Evulſion.

VI. The fifth and laſt Method is that uſed by Mountebanks upon the Stage, which conſiſts chiefly in anointing the Tubercle with ſome mollifying Ointment; after which they very violently pull it out by the Nails of their Fore-finger and Thumb. But as this Method of Cure is not very agreeable, ſo it is often found to be alſo ineffectual; for they generally return again from the Remainder of the Root.

Cancerous Warts.

VII. Laſtly, we are here not to omit taking notice of ſome Warts which are livid and blue; which are uſually ſeated in the Face, Lips, and about the Eyes, and are of a cancerous Diſpoſition, much better left to themſelves: for when they are irritated, they frequently degenerate into a Cancer, and miſerably torment the Face, Eyes, and other Parts in which they are ſeated. You will find a remarkable Inſtance of this Kind of Warts in SAVIARD's Obſervations *Obſ.* LXVIII. *p.* 296. which, through an ill-judged Attempt of curing it, was attended with the moſt deplorable Symptoms, and deſtroyed the Patient.

CHAP.

CHAP. XXVII.

The Method of removing EXCRESCENCES, *fleſhy* TUMORS, *and* MARKS *from the* MOTHER.

What an Excreſcence is.

I. EVERY preternatural Tumor, which ariſes upon the Skin, and grows in the Form of a Wart or Tubercle, is called an Excreſcence. They are by the *Greeks* called *Acrothymia*[a]; and if they are born with a Perſon, as they frequently are, they are commonly called *Nævi Materni*, or Marks from the Mother. But if the Tumor is large, ſo as to depend from the Skin like a fleſhy Maſs, it is then uſually called a *Sarcoma*. Theſe Tumors ariſe in all Parts of the Body; more particularly in the Head, Face, Eyebrows, Neck, Breaſt, Abdomen, Anus, Legs, and Arms. But the worſt of theſe Tumors are thoſe which ariſe in the private Parts. The Size and Figure of them are various, ſometimes ariſing to a very conſiderable Bulk, which are deſcribed and figured variouſly by the Writers of Obſervations[b]; with regard to their Colour, ſome reſemble that of the Skin, others are inclined to a black or red. And, laſtly, with regard to their Figure, ſome are like Strawberries, Mulberries, Grapes, Figs, Pears, Mice, and various other Figures.

Their Removal.

II. With regard to the Treatment of theſe, it is to be obſerved, that almoſt the ſame Artifices may here take place as for the Removal of Warts, either by Ligature, the Knife or actual and potential Cauteries, according as their different Sizes, Situations, Figures, the Patient's Habit of Body, his Will, and other Circumſtances may require. For the reſt, when any of theſe Excreſcences have a very large Root, which the *Greeks* call *Mermixia*[c], or if there are large Arteries or Veins near its Root, or if it be firmly joined to any Bone, or have a Tendency to turn cancerous[d], the Surgeon ſhould then remove them with Caution: or, when there is great Danger, he ought wholly to neglect them, to prevent expoſing his Patient to greater Dangers. When theſe Tumors are ſeated near large Veins and Arteries it is often proper to have Styptics, Bandages, and often actual Cauteries in Readineſs to ſtop the Hæmorrhage, eſpecially if they are removed by Abſciſſion.

[a] See CELSUS, *Book* V. *Ch.* 28. *N.* 14.

[b] See SCULTETUS, *Arm. Chirurg.* Plate XXV. ELSHOLST, Hiſtory of the Cure of a *Steatoma*. PECHLIN. *Obſ. Med.* Book III. 46. BARTHOLIN. *Hiſt. Anat.* Cent. 1. 23. PURMAN. *Chirurg. Curioſ.* p. 50, 134, and 370. LAMESWERDE, in his Notes on SCULTETUS, has deſcribed many of a very remarkable Figure. The *Engliſh Philoſoph. Tranſactions* give an Account of a very extraordinary Tumor of this Kind, that was taken off from the Cheek, weighing nineteen Pounds, *N.* 354. And the famous CULME has given a Deſcription of a large *Steatomatous Exoſtoſis* in the Clavicle, with his Treatment of it; publiſhed at *Dantzick*, 1732.

[c] CELSUS, *Book* V.

[d] WEDELIUS ſaw a Caſe, where a *Nævus* turned cancerous on the Application of *Aquafortis*. *De Morb. Infant.* p. 10.

CHAP.

CHAP. XXVIII.

Of ENCYSTED TUMORS, *and especially* SCHIRRI, ATHEROMATA, STEATOMATA, MELICERES, *and others.*

I. WHEN Tumors that arise from different Parts of the Body are contained in certain membranous Coats, they are commonly called *Encysted Tumors*, being sometimes harder or sometimes softer, of a paleish Colour, and usually attended with little or no Pain. These Kinds of Tumors arise almost in all Parts of the Body, from Obstructions, either in the Glands, or adipose Membrane; more especially in the Head, Face, and Neck, where they frequently occasion great Deformity: (See *Plate* XII. *Fig.* 13.) The membranous Coat with which they are invested, is often of a considerable Thickness, and is usually the Coat of the disordered Gland, or some of the adipose Cells. At their Beginning they are usually very small and moveable, being a considerable Time increasing by degrees, till at last they sometimes arrive to an enormous and surprising Bulk. The Consistence of some of these Tumors is soft and fluctuating, of others more hard and consistent. The Figure of them is various, some being like Filberds, Acorns, Bullets, Wallnuts, and Eggs; others again like Pears suspended by a Sort of Stalk, like some of the fleshy Excrescences; some have a very large Root, and resemble one's Fist, Head, or other Figure. The Bulk of some of these Tumors has sometimes increased to that Degree as to weigh many Pounds[a]: others of these Tumors so firmly adhere to the adjacent Parts as to be wholly immoveable, and become of so hard a Consistence as to resemble a bony *Callus*; though many of them always remain soft and moveable. Several Kinds of these encysted Tumors are variously distinguished by their different Consistencies. When their Contents resemble Paste, they are then called *Atheromata*; if they are of the Consistence of Honey, they are termed *Meliceres*; but if they are of a fattish Consistency, like Suet or Lard[b], they then take the Name of *Steatomata:* if they happen in a Gland which becomes indurated, the Tumor is then called *Scirrhous*; and, lastly, when they are of a fleshy Consistence, they are termed *Sarcomata*. Some of these Tumors, as [c]CELSUS has observed, have been found full of Hair. Again, these Tumors are variously distinguished and denominated from their different Situations: For when they are seated under the Scalp, they are called by the name of *Talpa*, *Testudo*, or *Lupia*; when they are seated in the Neck, *Strumæ*, or *Scrophulæ*; but when they are situated in the Hands or Feet, especially near the Tendons of their Muscles, they are usually denominated *Ganglia*. Various Kinds of encysted Tumors.

II. These Encysted Tumors are usually discoverable without much Difficulty by the Eye and Hand: but they are very difficultly discernable from other Tumors barely by their external Signs, if we do not discover their Difference by feeling Diagnosis of encysted Tumors.

[a] PETIT mentions one, that he extirpated, between the *Scapulæ*, of forty-eight Pounds Weight.

[b] I have had some under my Care, that resembled curdled Milk, or new Cheese; for which there is no particular Name. For steatomatous Swellings in the Head consult ROONHUYS, *Obs.* 17 and 20. A Cystic Tumor, that had many membranous Coats, is mentioned by MEEKREN, *Obs. Append.*

[c] *Lib.* VII. *Cap.* 6.

feeling whether they are harder, ſofter, or more or leſs conſiſtent. For as the external Skin receives little or no Alteration in its Colour in the ſeveral Sorts of theſe Tumors, we can therefore form little or no Judgment by it. Nor is it of any great Conſequence to be acquainted with the Nature of the included Matter, except the Hardneſs, before we proceed to the Cure of theſe Tumors: for, whatever Matter they contain, the Manner of Treatment is pretty much the ſame. It is however to be obſerved, that *Scirrhi* or *Sarcomata* are the hardeſt of any of theſe Tumors; next to theſe come *Steatomata*; all the reſt are ſtill ſofter, and may require ſome Variation in their Treatment, according to their Degree of Conſiſtence. Thoſe Tumors ſeated in the Neck, which are called *Strumous* and *Scrophulous*, are commonly thought to be the Glands in the Neck indurated: but I have frequently obſerved *Steatomata* and other encyſted Tumors in the adipoſe Parts of the Neck. For it ſeems ſcarce poſſible that thoſe very ſmall Glands which are ſituated in the Neck, ſhould grow to ſuch a ſtupendous Bulk as ſometimes to hang over the Abdomen: whereas thoſe in the adipoſe Parts may eaſily do ſo. But beſides theſe, there are alſo frequently leſſer Tumors in the Neck, which ſeem to be thoſe Glands indurated and much enlarged, being in fact a Kind of *Scirrhi*.

Prognoſis of encyſted Tumors.

III. When encyſted Tumors are without Pain, are neither too hard, nor too much enlarged, they preſage no great Danger, inſomuch that it is common for poor People, and thoſe who are afraid of the Surgeon's Hand, to bear them, to the End of their Lives, without any great Inconveniency: but when they grow too large, ſo as ſometimes to weigh ten or twenty Pounds; when they cauſe violent Pains, as ſcirrhous Tumors frequently do; then, beſides the intolerable Trouble they give the Patient, they alſo add great Deformity: and, if they are not timely removed, they often occaſion a Conſumption, or Cancer, putting the Patient in imminent Danger of his Life, as we before obſerved in the Chapter of *Scirrhous Tumors*. Every one muſt know, that, in the Treatment of theſe Tumors, for a Cure, the Aſſiſtance of the Knife is conſtantly to be called in: becauſe they will not eaſily digeſt, or be brought to Suppuration, as we have already obſerved in ſcirrhous Tumors. In the mean time, thoſe Tumors are more ſafely and eaſily removed by the Knife, which are of no long ſtanding, are ſoft, ſmall, and moveable: whereas thoſe which are very large and hard, are attended with great Danger; eſpecially if they are ſeated near to large Veins and Arteries, by Nerves, Tendons, or upon the Joints; or, if they happen in feeble and old People. Hence the Surgeon may judge from the Nature of the Tumor, and Circumſtances of the Patient, whether or no, and by what Means, it is curable.

Cure by diſcuſſing.

IV. With regard to the Cure of theſe Tumors, various Methods are proſecuted. I am not inſenſible, that many Surgeons are for an immediate Extirpation of all encyſted Tumors, without any more Delay: but I am rather inclined, with HIPPOCRATES, firſt, to try them with more gentle Methods of Cure. Whenever I meet with theſe Tumors, as yet ſoft, and of no long ſtanding, I think it every Way more proper to diſperſe, or elſe to ſuppurate them, before the Knife is called in for Aſſiſtance. But on the contrary, when theſe Tumors appear to be very hard, and of long ſtanding, it ſeems moſt proper to refrain from topical Remedies: for thoſe Means are ſo far from ſucceeding in the Digeſtion or Diſcuſſion of ſcirrhous and ſteatomatous Tumors, that they very

very often increaſe them, and ſometimes turn them into a Cancer: whereas they might have been tolerable in themſelves for many Years, ſo that under theſe Circumſtances we muſt rely altogether on Extirpation. But if the Patient is afraid of the Knife, and will admit no Means but topical Remedies, it may not be amiſs to uſe ſome of the diſcutient or digeſtive Plaſters; of which Sort are *de Ammoniaco*, *de Galbano*, *de Ranis cum Mercurio*, *Diachylon cum Mercurio*, *de Mellilot. Oxycroc. diaphoreticum* MYNSICHTI, *Diaſapon*, *&c.* SCULTETUS (in *Obſ.* 87.) aſſures us, he has cured various encyſted Tumors of the *Meliceres* kind with *Ceratum diaſinapios:* but before a Plaſter of that is applied, it is generally adviſeable to anoint the Part firſt with *Balſ. Peruv. Ol. Sapon. vel Petroleum*, *&c.* by which means the Tumor frequently diminiſhes by degrees, till it be at length diſperſed. To do this the more effectually, a little mercurial Ointment ſhould be well rubbed into the Tumor every Day before a Fire. See more concerning the Diſperſion of ſcirrhous Tumors in *Part* I. *Book* IV. *Chap.* XVI.

V. If the Tumor does not diminiſh by the Uſe of diſcutient Applications, you muſt endeavour to bring it to Suppuration[a]: and this more eſpecially when it is of the ſofter Kind, like the *Meliceres* or *Atheromata*. For this Purpoſe the Application of a Plaſter of *Diachylon* with the Gums, and the Repetition of warm emollient Cataplaſms to the Tumor, are extremely uſeful: and the more ſo, if you moiſten the middle of the Tumor every Day with a little ſtrong *Sp. Salis Ammoniaci*. When the included Matter has thus acquired a due Softneſs, and the Integuments appear a good deal extenuated, you ought then to open the Tumor by a large Inciſion; and, having diſcharged the Matter with as much as you can of its including Cyſt, the Remainder is to be brought away, by dreſſing with digeſtive or detergent Medicines: for if the Tunics of the Cyſt be not entirely diſcharged, the Tumor generally returns again ſoon after the Wound has been healed. In the Time you are deterging the Abſceſs, it may be proper to apply a *Diachylon* Plaſter externally, to keep the Lips moiſt, and better diſpoſed to unite afterwards. Cure by Suppuration.

VI. But if the Tumor can be neither diſperſed, nor ſuppurated, but continues to enlarge itſelf, 'tis generally in that Caſe moſt adviſeable to make an Extirpation of it, before it grows too large, or degenerates into a cancerous Nature. There are ſeveral Methods in Practice for removing or extirpating theſe Tumors, according to their Size and Nature. Thoſe which are ſmall and hard, or hang by a Root as by a Stalk, are generally beſt removed by Ligature, in the Manner of Warts: by which means they wither, and fall off of themſelves in a few Days. But the moſt ready and expeditious Method is, to cut them off with a Scalpel, and then to heal up the Wound: but if, in removing them this Way, you divide a conſiderable Artery, you may ſtop it by ſome potential, or even the actual Cautery, or elſe by taking it up with a Needle and Thread. Laſtly, theſe Tumors may be often removed by the Application of cauſtic or corroding Medicines retained about the Root by means of Plaſters, Compreſſes, and Bandage: and when you find the Root of the Tumor almoſt corroded through, the reſt may be divided by the Scalpel. Cure by Extirpation.

VII. If the Root of the encyſted Tumor appears too large for it to be conveniently taken off by Ligature, you muſt then remove it either by the Knife or Removal of the larger Kinds.

[a] See SCULTETUS on Suppuration in ſteatomatous Caſes, *Obſ. Chirurg.* 93.

or Cauſtics; though the latter is uſually preferred by moſt Surgeons. In order to extirpate it by the Knife, you muſt firſt make a longitudinal Inciſion upon the Tumor: and if that does not appear ſufficient, make another Inciſion acroſs the former, till you think the Wound large enough for taking out the Tumor. In order to this you next dilate the Integuments, and ſeparate them from the Cyſt of the Tumor, by the Aſſiſtance of your Fingers and the Scalpel, by which Means you are to take it out whole, if poſſible. That you may ſucceed the better in the Operation, it will be proper for an Aſſiſtant to dilate the Lips of the Wound, either with Hooks or his Fingers, and to wipe up the Blood as it flows, with a Sponge, that the Surgeon may not be impeded in his Work. When the Tunic, or Cyſt of the Tumor appears, which is uſually pretty white and hard, the Surgeon is then to take hold of, and elevate the Tumor with the Fingers of his Left Hand, if it be ſmall enough: but if it be too large to be thus held and elevated, it may be done by another Aſſiſtant with the Hook, *Tab.* VIII. or the Forceps, *Tab.* XXIII. *Fig.* 1. or elſe he may paſs a crooked Needle and ſtrong Thread croſs-wiſe under the Tumor, and by that means elevate it, while he circumſpectly frees it from the adjacent Parts; which is generally done with moſt Eaſe in the moveable Kind of theſe Tumors: but in the more fixed, the Taſk is pretty difficult. But in thus freeing the Tumor, the Surgeon muſt be cautious not to injure any important Part that may be contiguous: and if the Tumor, to be extirpated, is either in the upper or lower Extremities, where perhaps a large Artery or Vein is to be divided, in that Caſe the Tourniquet may, and even ought to be firſt fixed upon the Limb. Which Circumſtances being duly obſerved, Tumors of this Nature have been often ſucceſsfully extirpated, of many Pounds Weight, and which have been not only lodged in the fleſhy Parts, but have even adhered to the Bones and Jaws [a].

Treatment after Extirpation.

VIII. The Tumor being thus careſully extracted, if the Wound and Hæmorrhage be ſmall, you may preſs the Lips together with your Fingers; and by covering the ſame with Lint and Compreſſes, retained with a proper Bandage, the Patient is generally cured in a few Days time. But in caſe of a profuſe Hæmorrhage, the Blood is to be ſtopped, either by Ligature, Aſtringents, or the actual or potential Cautery, as we have directed more at large in *Part* I. *Book* I. *Chap.* II.

Removal of Fragments of the Cyſt.

IX. But if by Neglect or Accident the including Cyſt of the Tumor ſhould be broke or wounded in its Extraction, (or induſtriouſly, to ſave the Eye, where it falls on the Eyebrow; or to ſecure the Veins and Arteries in other Parts of the Body) Care muſt be afterwards taken entirely to remove it; otherwiſe the Tumor will ſpeedily return. Indeed if the Tumor be either a *Scirrhus*, *Sarcoma*, *Steatoma*, or in a glandular Part, the Contents are hard enough to make a clean Extirpation of it, notwithſtanding its including Coats be wounded. But when the Matter of the Tumor is ſoft or fluid, by its eſcaping, the Tumor will become flaccid, ſo that it will be hardly poſſible to make a clean Extirpation of the Cyſt without leaving ſome Fragments behind; which muſt, in that Caſe, be brought away, by dreſſing the Abſceſs with Digeſtives, and deterging with *Præcipitat. rub. Alumen uſt. Ung. Ægyptiac. &c.* mixed with your digeſtive

3 Oint-

[a] See ROONHUYSEN, *Obſ.* I. pag. 4. SCULTETUS *cum notis* TILINGII. PECHLIN, *Obſ.* pag. 542. PETIT *apud* GARENGEOT. *Chirur.* Tom. II. Cap. *de Tumor. tunicat.* LE DRAN, *&c.*

Ointments: by which means, having cleared the Sinus, you may incarn and heal, as in other Wounds, without the Danger of a Relapse.

X. If you rather chuse to remove Tumors of this kind by the Use of Caustics, you must, in that Case, apply a Piece of *Lapis infernalis*, *Butyr. Antimon.* &c. upon it, defending the other Parts by a perforated Plaster, as we directed *Chap.* XIX. *Sect.* IV. But, in my Opinion, this is not a safe Practice in those encysted Tumors, which are hard, large, inveterate, and painful, or inclining to be cancerous; for thus you may easily turn a *Scirrhus* into a real Cancer: and even in others, 'tis hardly possible thus to erode them quite away without inducing violent Pains, Fever, Hæmorrhage, and other malignant Symptoms, to the Hazard of the Patient's Life. It is therefore, in the general, much better to have recourse to the Knife for the Removal of these Tumors, when they are large and hard, notwithstanding we now and then meet with an Instance of their being successfully extirpated by Caustics[a]. But if the Tumor appear soft, and yielding, like the *Atheroma* or *Meliceris*, in that Case, I frequently apply a Caustic, so as to make a Way through the Integuments, and Cyst: or else dividing them by an Incision in the Middle, I discharge their Contents, and then deterge and incarn, as in other Abscesses; which last Method I take to be milder than an Incision, and Extirpation of the Cyst by the Scalpel. See Le Dran on the Treatment of a *Meliceris* above the Knee.

Use of Caustics in these Tumors.

CHAP. XXIX.

The Method of extracting foreign Bodies *from* Wounds.

I. We meet with very little in the antient Systems of Physic and Surgery concerning the Extraction of Bullets, which may possibly be, in some measure, owing to their not being so much in use, or at least not so fatal formerly as now. We indeed read in Celsus[b], that leaden Balls were used by Soldiers in War before the Birth of Christ: but then I suppose they were only flung by Slings or Bows, the destructive Powder being at that Time unknown. For the same Reason we also meet with no Directions for extracting Fragments of Bomb or Granade Shells, which are of a later Invention: but then they are more large in the Methods of removing the Ends of Darts, Spikes, Arrows, Swords, and such like Weapons. And though, at this Time of Day, Arrows are hardly ever used but among barbarous Nations, yet it may not be here improper to give brief Directions for their Extraction, if they should chance to come under the Surgeon's Care. In doing this, we shall find that almost the whole Business consists in drawing out the Head, so as that its protuberant Beard or Hooks may not wound and lacerate the contiguous Parts. If it appears to be lodged but superficially under the Integuments, it will be best to draw it out the same Way it entered, provided you first dilate the Wound sufficiently by Incision, rather than give Occasion for any of the adjacent Parts to be lacerated: otherwise it may be thrust forwards, and drawn out in the Direction

[a] Of extirpating scirrhous Tumors in the salival Glands, and those of the Neck and Breast, we shall treat hereafter in particular Chapters.

[b] *Lib.* VII. *Cap.* 5.

rection of its Point in the oppoſite Side, having firſt made an Inciſion to meet it. This laſt Method is moſt eligible, when the Weapon has deſcended very deep, ſo that there is much leſs Space for it to paſs onward, than to be drawn back again; and alſo when it has paſſed beyond any large Blood-veſſels, or Nerves, ſo that it would induce a Laceration of them, to draw it back: And therefore, to avoid them, it muſt be thruſt forward through an Inciſion made in the neareſt and moſt convenient Part of the oppoſite Side. The Method of extracting the Ends of Spikes, Swords, Sticks, or the Fragments of Glaſs, Paper, Clothes, *&c.* you may find in *Part* I. *Book* I. *Cap.* I. *Sect.* XXXIII. and, in the third Chapter following, you will find the Method of extracting Bullets and Grains of Gun-powder, in Gunſhot-wounds. Laſtly, if any of theſe foreign Bodies have ruptured a large Blood-veſſel in the upper or lower Extremities, ſo as to excite a profuſe and dangerous Hæmorrhage, it will, in that Caſe, be immediately neceſſary to apply the Tourniquet upon a convenient Part of the Limb, before you ſearch for the Body; which being extracted, the next Step is to ſecure the ruptured Veſſel, and dreſs the Wound.

CHAP. XXX.

Of SUTURES *of* WOUNDS

The Kinds and Uſes of Sutures.

I. THERE are two Kinds of Sutures uſed by Surgeons in Wounds: the firſt is made with a Needle, and diſtinguiſhed by the Name of the *true* and *bloody Suture*; the other is made by the Application of ſticking Plaſters, and is termed the *dry* or *falſe Suture*. Sutures are not to be uſed indifferently in all Wounds: but in thoſe chiefly, in which the Lips cannot be cloſely approximated by Deligation: as in many of the tranſverſe, oblique, or angular Wounds, which have been lately inflicted, are quite free from any foreign Bodies, and are not attended with any Loſs of Subſtance. In moſt of theſe a Suture will be of great Service, not only by expediting the Healing, or Union of the Wound, but alſo by procuring a ſmaller and neater Cicatrix. The dry Suture is uſed chiefly in ſuch Wounds as are ſuperficial, of no great Depth or Length, and particularly for thoſe inflicted on the Face: though even in theſe there are ſome Surgeons, who prefer and make the true Suture. But I think the different Circumſtances and Diſpoſitions of Wounds may very well direct the Surgeon, ſometimes to one, and ſometimes to the other Kind of Suture: for what need is there for ſtitching up a Wound, whoſe Lips may be well approximated, and retained together by Plaſter and Bandage? I think the Needle ought, in ſuch Caſes, to be ſpared, both for the Eaſe of yourſelf, and the Patient. But on the contrary, in large and deep Wounds, where the Lips cannot be cloſely retained by Plaſter and Bandage, or in thoſe where the Part is almoſt amputated, or hangs by a little bit, as in the Noſe, Ears, Cheeks, Chin, Forehead, Fingers, *&c* there you ought immediately to conjoin the Lips by Suture with Needle and Thread.

Directions for Sutures.

II. As we have already ſufficiently explained the Method of making Sutures in Wounds (in *Part* I. *Book* I. *Chap.* I. *Sect.* XXXIX. *& ſeq.*) we ſhall here only add a few neceſſary Cautions: as, 1. That you ought always to ſhave the Hair of

of the Part clean off, with a Razor, before you attempt to conjoin the Lips of the Wound by dry Suture, with ſticking Plaſters. 2. That when one Plaſter does not well retain the Lips, you muſt apply ſeveral, either by the Side of, or acroſs each other, as in *Tab.* IV. *Fig.* 4, 5, 6. You are alſo to obſerve, 3. That the true Suture with Needle and Thread is of two Kinds, ſimple and compound: the firſt of which comprehends the *knotted*, the *Glovers*, and *circumvoluted*, and the *Suture* of the Tendons. And, among theſe, the firſt is ſo called from its diſtinct Knots, *Tab.* IV. *Fig.* 16. the Glovers from its Reſemblance to the Suture uſed by theſe Artiſts, and by the Surgeon for Wounds of the Inteſtines, *Tab.* IV. *Fig.* 20. The circumvoluted Suture is, when the Thread is wound about the Needle, after it has been paſſed through both Lips of the Wound, as in *Tab.* IV. *Fig.* 21, 22, for the Hare-Lip: In treating of which we ſhall deſcribe it more particularly. The Suture of a Tendon is alſo of a particular Kind, as we ſhall deſcribe in our Chapter of uniting divided Tendons, by this Means, in the End of our Operations. Beſides theſe now mentioned, there were various other Sutures uſed by the antient Surgeons, as the *Sutura Sartoria*, *Sutura Celſiana*, *& Clavata*, the laſt being made upon Quills or cylindrical Sticks, as in *Tab.* IV. *Fig.* 19. But we ſhall not inſiſt upon a particular Deſcription of theſe, which have been long out of Uſe: only we may obſerve, that the *Sutura clavata* has been lately revived, and recommended with a little Variation, by PALFYN and GARENGEOT; who, inſtead of Sticks or Quills, uſe a bit of Silk ſpread with Cerate, and rolled up into a Cylinder. 4. Laſtly, you muſt obſerve, that, in the Suture of deep Wounds, it is frequently neceſſary to introduce a Tent, and leave it at the Bottom of the Wound, till its Fundus appears well deterged, that you may heal it from the Bottom upwards.

CHAP. XXXI.

Of ſeparating ADHESIONS *between the* FINGERS *and* TOES.

I. WE frequently meet with new-born Infants, having ſeveral of their Fingers or Toes cohering, or grown together, either by a ſtrict Adheſion of their Fleſh, or elſe only by looſe Productions of the Skin, as in the Feet of Ducks and Geeſe. Though the ſame Diſorder is alſo ſometimes found in Adults, when their Fingers or Toes have been neglected, after an Excoriation of them in Burns or Wounds: to be freed from which Malady, the Patient is deſirous of invoking the Surgeon's Aid, partly to be rid of the Deformity, but chiefly to recover the proper Uſe of the Fingers. Theſe Adheſions may be ſeparated in a two-fold Manner, according to the Nature of the Diſorder, *i*, *e.* either by cutting out the intermediate Skin with a Scalpel, or Pair of Sciſſars, or elſe barely by dividing them from each other with thoſe Inſtruments when they cloſely adhere. But to prevent their Coheſion again for the future, you muſt inveſt each of the Fingers ſeparately with a ſpiral Bandage about an Inch broad, and dipt in *Aq. Calcis cum Sp. Vini*, according to the Figure in our laſt, or, XXXIXth *Plate*, on Bandages. Adheſions of the Fingers together.

II. Some-

Or with the Palm of the Hand.

II. Sometimes the Fingers, inſtead of adhering to each other, grow to the Palm of the Hand, as I have more than once obſerved from Wounds or Burns; ſo that they cannot by any means be extended, or drawn back to open the Hand. For the Sake of Beginners, I ſhall recite the Method, by which I cured three of theſe Patients. Firſt, I carefully ſeparated the Fingers from their Adheſions with the Palm, without injuring their Tendons, and, after dreſſing them with vulnerary Balſam, and ſcraped Lint, I extended them on a Ferula of thick Paſteboard, in which extended Poſture I treated the wounded Fingers ſeparately, till they were healed. But at every Dreſſing you ought to move the Fingers gently, to prevent a Rigidity, or Stiffneſs of their Joints.

CHAP. XXXII.

Of amputating DISEASED *and* SUPERFLUOUS FINGERS.

INFANTS are ſometimes born with ſupernumerary, miſ-ſhapen, and miſplaced Fingers, of various Kinds, ſome with Nails and Bones, and others without, reſembling fleſhy Excreſcences. When the Deformity or Incumbrance of theſe makes their Amputation neceſſary, it may be conveniently enough performed, either by the Scalpel, or a Pair of Sciſſars; eſpecially when there are no Bones in them: for if there are Bones, you muſt amputate with a ſtronger Pair of Sciſſars for the Purpoſe, able to cut through the Bones. If there are ſeveral of theſe Fingers, and the Infant appears too weak or infirm, to have them all amputated at one Time, it is beſt to take them off at ſeparate and convenient Intervals, intermitting a few Days, ſo as to amputate the next, when the preceding is near well. The Hæmorrhage may be ſtopped with dry Lint and Compreſſes, or ſuch as have been dipt in *Sp. Vini*, and the Wound next healed with ſome vulnerary Balſam, as in others. In the Year 1718, I cured an Infant of three Weeks old, after taking off a ſuperfluous long Finger, which grew to the Thumb, which had a long Bone, and a ſort of Spur like that of a Cock, inſtead of a Nail; ſee *Tab.* XII. *Fig.* 15. I proceeded, firſt, by making an Inciſion through the Skin all round it with a Scalpel, and then cut through the Bone with a ſtrong Pair of Sciſſars. This done, I ſtopt the Hæmorrhage, which was inconſiderable, with Lint dipt in *Sp. Vini*, and a cloſe Bandage; and the Wound was afterwards ſpeedily healed with vulnerary Balſam. I could recite many more Cures of the ſame Kind made by myſelf: but as the Method uſed was the ſame in all, they are not here neceſſary to be mentioned, ſince this alone will ſuffice.

CHAP.

CHAP. XXXIII.

Of amputating ſphacelated FINGERS *and* TOES.

When it is neceſſary to amputate Fingers and Toes.

I. FINGERS and Toes are uſually amputated by the Surgeon, chiefly upon three Accounts; 1. When they are ſo contuſed and ſhattered by Bullets or other Inſtruments, that they cannot be reſtored and preſerved: 2 When they are ſphacelated, or totally mortified, either from Cold, Contuſions, or other Cauſes: and, laſtly, 3. When they become carious, cancerous, or ſcirrhous, ſo as to be curable by no Remedies or Applications whatever, as I have met with frequent Inſtances. Nor is it uncommon for the Fingers of Maſons, Carpenters, and other Labourers, to be accidentally cruſhed, ſo as to make an Amputation of them unavoidable. See ROONHUYS, *Obſ. Chirurg.* 25.

Cautions to be firſt obſerved.

II, Before the Surgeon proceeds to amputate Fingers or Toes, he ought to be firſt well aſſured, that there is no Poſſibility of preſerving them ſound and entire. Therefore if they appear to be but ſlightly cruſhed, or only beginning to be infeſted with a Gangrene, he ought to treat them with diſcutient and ſpirituous Applications, to prevent the Diſorder from ſpreading itſelf; at the ſame time reducing and retaining the bony Fragment by his Fingers, and Deligation, as in other Fractures. But if they are ſo violently cruſhed as to hang but by a little Bit, I know no great Reaſon why they ſhould not be immediately taken off, either by the Sciſſars or Scalpel; as they alſo ſhould when any one Joint is completely ſphacelated; for Delays are, in thoſe Caſes, frequently very dangerous. But if any of the Fingers or Toes ſhould be cut off by any ſharp Inſtrument, ſo as to hang by a Bit, the Wound being recent, though large, you ought not to take off the pendulous Part, but replace it immediately, ſecuring it well by Plaſter and Deligation, and this even when the Part is cut quite off, but obliquely: for I knew an Inſtance of a Butcher's Finger that was cut quite off obliquely, but being immediately fixed, and retained in its proper Place by Deligation with a Linen-rag, it adhered, and became well without any other Medicines. At leaſt, it is always beſt to try, if it will not adhere before you cut it off, and reject it; ſee *Chap.* LXXII. following.

Method of amputating.

III. The Manner of amputating is chiefly threefold: either 1. by a Pair of ſtrong *Sciſſars*[a], or rather ſharp-edged *Pincers*, treating the Wound as we before directed in the preceding Chapter: or, 2. by the *Mallet* and *Chiſſel*, *Tab.* XII. *Fig.* 17. with which the injur'd Parts are taken off at one Blow, as I have frequently done in cancerous Affections with a Caries or *Spina ventoſa* in the Fingers: and ROONHUYS has alſo thus ſucceſsfully amputated the great Toe, being ſcirrhous, notwithſtanding what others may ſay againſt this Method. Or, laſtly, 3. the diſeaſed or mortified Parts are amputated by dividing in the next ſound Joint with a *Scalpel*, leaving or drawing back a large Part of the Skin, to wrap

[a] See PARÆUS. *Book* XVI. *Ch.* 30. SCULTETUS *Plate* LIII. *Fig.* 2. But by this Method the Bones are often ſplintered, or the nervous Parts lacerated; from whence ariſe dangerous Inflammations and Abceſſes, and too frequently a Caries is the Conſequence. DOUGLAS abſolutely rejects the Forceps, the Mallet and Chiſſel, *Operat. Chirurg. Syllab.* p. 45.

wrap over the Stump, that it may heal the ſooner. This laſt Method of amputating, is preferable to the former, in that you are, by this means, certain to avoid any ſupervening Caries, or a ſplintering of the Bone: for which Reaſons I have uſed it with Succeſs for removing Thumbs and Fingers, even of old People, in the Articulation of them with the Metacarpus, when they have been totally deſtroyed by a Caries or Mortification. Some indeed imagine this Method of amputating in the Joint to be not ſo convenient, becauſe a Cicatrix or Skin cannot be induced over the Cartilage. This is, however, an Obſtacle that I never yet met with, and may, at worſt, be eaſily avoided by drawing back and leaving a large Part of the ſound Skin on, and by removing the cartilaginous Extremity of the metacarpal or metatarſal Bone; by which means the Bone and Skin will more intimately unite and adhere. After the Amputation your Dreſſings muſt be made with ſcraped Lint, Compreſs, and Bandage, as we before directed: and, if the Patient be plethoric, in order to prevent Inflammation, or a future Hæmorrhage, it may be proper to take a few Ounces of Blood from a Vein. If any of the two foremoſt Internodes of the Fingers ſhould appear to be carious, and Part of the third, it is better to amputate the injured Part of the laſt by the Mallet and Chiſſel, which will more expedite the Cure, than to take off the whole Finger cloſe to the Metacarpus by the Scalpel. But if the whole Finger or Toe is entirely corrupted, it muſt then be taken off in the Articulation cloſe to the Metacarpus, leaving a good deal of the Skin. See Inſtances of great Toes amputated in LE DRAN, *Obſ.* 112, 113, and 114.

An EXPLANATION *of the* TWELFTH PLATE.

Fig. 1. Repreſents the Cupping-glaſs uſed at preſent in *Germany*, and elſewhere, for dry Cupping, or for extracting Blood after Scarification.

Fig. 2. Is the Scalpel, or Scarificator, commonly uſed by our *German* Cuppers. A the Handle, B the Edge, C the Part which is ſtruck extremely quick by the Finger, ſo as to make the Edge wound the Skin.

Fig. 3. Repreſents the Order or Poſition of the little Inciſions made in the Skin by the Cupper, that they may all be cleanly covered by the Cupping-glaſs, *Fig.* 1.

Fig. 4. Exhibits the modern cubical Scarificator, making ſixteen Inciſions in the Order of *Fig.* 3. by one Stroke upon the Skin, and with very little Pain.

Fig. 5. Gives the Form or Shape of a Leech, for the Information of ſuch as may be ignorant of that Inſect: A the Mouth or Head by which it bites, B the Body and poſterior Parts. But it muſt be obſerved, that one and the ſame Leech may, by differently contracting and expanding itſelf, appear in a hundred Shapes, ſo that its Length and Thickneſs are very uncertain.

Fig. 6. Is the Needle uſed by the Inhabitants of *China* and *Japan* for making their Acupuncturation, which they eſteem in moſt Diſorders, as we do Phlebotomy. A the Handle, B the Point which enters the Fleſh.

Fig. 7. Is the little Hammer uſed to ſtrike in the preceding Needle: A the Head of this Hammer, B its Handle, CC a Caſe in the latter to depoſite the Needle in.

Fig. 8. Repreſents the actual and concealed Cautery, uſed formerly for the making of Iſſues, and is, by ſome, denominated *Capſula Caſſeriana*. A denotes the End of the actual Cautery, or red-hot Iron, protruding itſelf be-

Plate 12. Vol. I. P. 354.






Plate 12.


Fig: 1.
Fig: 3.
Fig: 4.
C
C
C
C
B
Fig: 2.
C
A
B
Fig: 5.
A
A
B
Fig: 7.
A
C
C
B
A
Fig: 6.
B
Fig: 8.
C
B
B
B
B
A
C
A
B
Fig: 9.


Fig: 13.
a
c
b
d
c
Fig: 15
A
Fig: 17.
Fig: 16.
A
B
C
Fig: 12.
B
A
Fig: 10.
B
A
A
C
Fig: 14.
B
A
Fig: 11.
A
A
B
C
B
A
B
B
B

yond the Caſe. B B is the wooden Caſe concealing the red-hot Iron from terrifying the Patient. C the Handle, by depreſſing which, the Cautery is forced into the Skin.

Fig. 9. Is a Machine to be uſed inſtead of Deligation for Iſſues in the Arm, to be made a little longer for thoſe in the Neck, Leg, or Thigh. A A is a leathern Swathe of about two or three Fingers Breadth. C C is a Braſs-plate with ſeveral oblong Apertures, for intercepting the Hook B of the other Plate in the manner of a Claſp.

Fig. 10. Shews the Syringe proper for injecting Liquors into the Urethra of Males, and the Vagina of Females, for various Uſes. A A the Body of the Syringe, B its Extremity, ending with an obtuſe Point inſtead of a ſmall Tube, to prevent the injected Liquor from regurgitating and flying about. C the Ring or Handle of the Sucker, by which the Liquor is drawn into, and forced out of the cylindric Body.

Fig. 11. A A Shews the Parts of the Soles of the Feet, which the *Italian* Phyſician MISTICHELLIUS directs to be cauterized in Apoplexies, B the ſquare Iron Cautery for the Operation, which in that Diſorder, he ſays, is highly ſerviceable.

Fig. 12. Repreſents the Method of burning the Part affected in the Gout with the *Indian Moxa*. A denotes the Cone of *Moxa* not yet fired, and B one that is burning.

Fig. 13. Gives a View of ſeveral encyſted Tumors *a b*; of ſcirrhous Glands in the Neck *c d*; and of a fleſhy Excreſcence or Mark from the Mother, *e*.

Fig. 14. Repreſents the ſmall Scalpel, which I generally uſe for extirpating ſcirrhous Tumors, or Glands in the Neck, Wens, or even ſcirrhous Glands of the Breaſts.

Fig. 15. Repreſents the Hand of an Infant with ſix Fingers: in which A denotes the ſuperfluous Finger with a Nail like a Cock's Spur, which I took off by a Pair of amputating Sciſſars or Pincers. This Inſtrument I alſo uſe in a *Spina ventoſa*, or Caries of the Fingers.

Fig. 16. Is a Hand with a whole Index, A, carious, which I amputate cloſe to the Metacarpus by the Scalpel *Fig.* 14. but then I alſo remove the Head of the firſt Phalanx, that the Wound may heal the ſooner. B denotes a *Spina ventoſa* in the middle Finger, and in the ſecond Internode, which I amputate in the firſt Bone or Phalanx; C is a large Excreſcence or Protuberance at the End of the little Finger, from the ſame Diſorder, which I amputate in the ſecond Bone, both of them by the Mallet and Chiſſel.

Fig. 17. Shews the Method of amputating the great Toe with the Mallet and Chiſſel, uſed by ROONHUYS.

CHAP. XXXIV.

Of amputating the HAND, CUBITUS, *and* HUMERUS.

In what Caſes Amputation is neceſſary.

I. THOUGH the Amputation of Arms and Legs is indeed, with ſome Reaſon, commonly eſteemed one of the moſt terrible and ſevere Operations in Surgery, yet there are many Caſes that occur daily in Practice, in which

the Operation is abſolutely neceſſary and unavoidable, in order to ſave the Life of the Patient: ſuch as, 1. When the Muſcles of the Part or Limb are ſphacelated[a]; 2. Or when the Muſcles and Bones are moſt violently contuſed and ſhattered; 3. When there is an incurable Caries, or *Spina ventoſa*[b]; 4. When the brachial, crural, or other large Artery, is either totally divided[c], or elſe wounded, ſo as to bleed inceſſantly without any Poſſibility of ſtopping the Hæmorrhage but by Ligature; in which Caſe 'tis hardly poſſible to preſerve the Limb from mortifying, or ſave the Patient's Life without Amputation. 5. And, laſtly, this Operation is neceſſary in thoſe Tumors of the Hand and Arm, which ariſe from a *Spina ventoſa*, or ſome other irremediable Cauſe, the Patient being tortured with the moſt excruciating Pains, as deſcribed by M. A. SEVERINUS[d], BIDLOE[e], RUYSCH[f], &c. In the mean time I would adviſe all prudent Surgeons, not to perform this Operation, unleſs there are other ſkilful Surgeons or Phyſicians, who alſo adviſe it, or think it neceſſary; by which means he may avoid many Reflections, which are often unjuſtly thrown upon a Surgeon without ſuch Precaution.

Method of amputating the Hand.

II. To proceed regularly with Amputations in the upper Extremities, we ſhall begin with that of the Hand; which may, on ſome Occaſions, be amputated, in the manner of the antient Surgeons, by one Blow with the Mallet upon a ſharp Chiſſel fixed near the Carpus, as the Operation is repreſented in *Tab.* LIII. of SCULTETUS, *Edit. An.* 1666. But in reality this Method is often found to be not only unſafe, but even of dangerous Conſequence, by violently contuſing or fracturing ſome of the Bones and Parts in the Carpus. It is therefore not without Reaſon that the Moderns reject this Practice for that with the Knife and Saw; with which they take off the Hand more ſlowly indeed, but more ſecurely, provided the Saw be not uſed to the Carpus or Metacarpus: becauſe the numerous Ligaments, Tendons, and ſmall Bones there ſeated, cannot ſafely be divided by the rough Teeth of that Inſtrument. [g] The Practice of the modern Surgeons is therefore here much the beſt, who amputate the Hand by the Knife and Saw, cutting through the Bones of the Cubitus, as will preſently appear.

What is to be obſerved in amputating the Arm.

III. When the Hand, Cubitus, or Humerus, are required to be amputated upon the account of ſome incurable Sphacelus, Caries, or other Diſorder, there are then two Things chiefly neceſſary to be obſerved. The firſt of theſe is the *Place where* the Amputation muſt be made, which muſt at leaſt be one[h] or two Fingers Breadth above the mortified Part, never in the diſeaſed Part itſelf.

[a] See *Part* I. *Book* IV. *Chap.* XIV. preceding.

[b] See *Part* I. *Book* V. *Chap.* VIII, IX. preceding.

[c] I have frequently ſtopped profuſe Hæmorrhages from the brachial Artery by Ligature, and therefore it will not be ſo often neceſſary to amputate the Arm on that account; as many Surgeons imagine and direct.

[d] *Lib de Abceſſ.*

[e] *Exercit. Medic. Chirurg.*

[f] *Epiſt. Anatom.* Problem XIV.

[g] Yet there have been ſome Surgeons who have, in this manner, amputated the Hand by the Saw in the Carpus or Metacarpus, as we learn from SCULTETUS, *loc. cit.* Nor is it impracticable, in my Opinion, to amputate the Hand in its Articulation with the Cubitus by the Scalpel, as in the preceding Chapter; though I muſt acknowledge myſelf to have never yet made the Experiment.

[h] FABRICIUS AB AQUAPEND. L. 1. *Pentateuch. Chirurg. & Operation Chirurg, tit. de Sphacelo,* & SCULTETUS *in Explic. Tab.* LIII. are both of Opinion, in oppoſition to HILDANUS, that the ſphacelated Limb ſhould be taken off in the diſeaſed Part, near the ſound; and what remains unſound, be

ſelf. Nor ought theſe larger Kinds of Amputations to be ever made in the Articulations: for (beſides other Difficulties) there being no Fleſh there to cover the Ends of the Bones, it will be almoſt impoſſible to heal the Stump[i], or prevent a Caries in the Head of the Bone, with other bad Symptoms. The next Thing required after the proper Place for Amputation is aſſigned, according to the uſual Method, is (2.) *the Proviſion and Preparation of the ſeveral neceſſary Inſtruments* and Parts of the Apparatus, which are to be laid in readineſs upon a large Plate, or convenient Part of the Table; yet ſo as that they may be concealed from the Patient's View, who might be not a little terrified and diſheartened by them.

IV. For the ſake of Beginners, we ſhall here enumerate the ſeveral Inſtruments neceſſary to compoſe the Apparatus for this Operation. Theſe are, (1.) the *Tourniquet*, before deſcribed in *Part* I. *Book* II. *Chap.* II. *Sect.* IX. *& ſeq.* (2.) Some Ligatures, or Tapes, of a Finger's Breadth, and about an Ell and an half long. (3.) A middling-ſized Knife (*Tab.* XIII. *Fig.* 1.) for dividing the Skin, to draw it back. (4.) A larger Scalpel, or Knife, of a crooked Figure (*Tab.* XIII. *Fig.* 2.) for dividing the Remainder of the Fleſh. (5.) A Catlin, or double-edged Scalpel (*Fig.* 3.) for dividing the intermediate Fleſh betwixt the Ulna and Radius. (6.) A Piece of Linen Cloth about three Spans long, and ſix Fingers Breadth, ſlit up lengthwiſe about half-way, as in *Tab.* II. *Fig.* 17. (7.) A well-tempered and ſharp Saw[k] (*Tab.* III. Fig. 4.) for dividing the Bones. (8.) A Pair of Pliers, or Forceps to hold the Ends of the Arteries, (*Fig.* 5 and 6.) (9.) Some crooked Needles, armed with ſtrong Thread, and ſome Bits of blue Vitriol wrapped up in Lint or Cotton. (10.) Some ſmall ſquare Compreſſes, (*Tab.* II. *Fig.* 21.) (11.) A large Quantity of ſcraped Lint. (12.) Some aſtringent Powders, to ſtop the Hæmorrhage, or rather, as the former frequently inflames the Parts and impedes the Suppuration, provide ſome *Alcohol Vini & Oleum Terebinthinæ*, in proper Veſſels, though in reality we may well enough omit all of them. (13.) A large Bolſter of fine Tow, of a round Figure, and broad enough to cover the Stump, and retain the other Dreſſings: or, inſtead of this, a Piece of the Fungus called *Lupi Crepitus*, or Puff-ball, of the like Size and Figure. (14.) A Calf's or Swine's Bladder, or elſe a large ſticking Plaſter cut in the form of a *Malta* Croſs, (*Tab.* II. *Fig.* 15.) or three ſeparate Plaſters, two Spans long and two Fingers broad, for inveſting and ſecuring all the other Dreſſings on the Stump. (15.) A Compreſs in form of a *Malta* Croſs, but larger than the Plaſter. (16.) A thick ſquare Compreſs, to inveſt the End of the Limb. (17.) Three other Compreſſes of two Spans long and two Fingers Breadth. (18.) A Roller or Bandage for the Deligation of the whole, of about five Ells long and three Fingers Breadth. Laſtly, (19.) Some Wine, and other cordial Medicines, to aſſiſt and relieve the Patient in caſe of a Deliquium.

The Inſtruments and Apparatus required.

V. The

be removed by the actual Cautery. This they recommend as the ſafeſt and eaſieſt Method. But this Method is now rejected, on many Accounts, by the unanimous Conſent of all modern Practitioners.

[i] But if a ſufficient Portion of the Skin be left on to cover the Stump, it may perhaps heal as readily as the Stumps of Fingers thus amputated.

[k] The Moderns have invented other Saws and Knives for Amputating, as may be ſeen in GARENGEOT's *Tract. de Inſtr. Chirurg.* but theſe here deſcribed being equally as good in all Reſpects, I ſhall not inſiſt on them.

Position of the Patient, Assistants, and Surgeon.

V. The whole necessary Apparatus being thus provided, the next Business is for the Surgeon to dispose the Patient, Assistants, and himself in a proper Posture to begin the Operation. First, therefore, the Patient must be fixed on a low Chair or Stool, in the midst of the Room, the Surgeon standing betwixt his Legs, and fix Assistants at least around him. One of these should stand behind the Patient to hold his Body; another on the Side of the affected Arm, which he is to hold fast by grasping the upper Part of the Cubitus: a third Assistant must hold the Hand, about to be amputated; and a fourth should stand on one Side with the Apparatus of Instruments, to hand them as they may be wanted by the Operator. A fifth Assistant must stand ready with the several Dressings, Compress and Bandage, necessary to complete the Deligation; and the sixth or last should be at liberty to assist the Patient and Operator occasionally, in handing Wine, Cordials, or any other thing they may want.

What must be done immediately before the Amputation.

VI. Things being thus far advanced, the Surgeon, who should have a Napkin before him, to wipe his Hands when there may be Occasion, proceeds to fix the Tourniquet (*Tab.* III. *Fig.* 1. K) moderately tight about the Patient's Arm, in the manner we before directed in (*Part* I. *Book* I. *Chap.* II. *Sect.* IX. *& seq.*) by which means the brachial Artery will be compressed, so as to prevent any profuse Hæmorrhage: and the Nerve being also a Partaker of the same Stricture, will make the Patient less sensible of Pain from the Operation. But to prevent the Tourniquet (*Tab.* III. *Fig.* 1. K) from coming loose, the Turnstick must be held fast by the Assistant standing behind the Patient. But if you apply the Screw-Tourniquet, figured in *Tab.* V. and VI. they will adhere tight upon the Part, without being held by an Assistant. This done, the Assistant holding the upper Part of the Arm, should next draw the Skin strongly upwards, while the Surgeon applies the Tape tight, and circularly about the Part, a little above where it is to be divided, in order to secure the fleshy Parts close to the Bones, that they may be cut through more easily and evenly. Some, as VERDUYN, advise use a leathern Strap with a Clasp, instead of a Tape or Fillet, for this Purpose, which we shall consider in *Chap.* XXXVI. *Sect.* III. following. The Surgeon now encourages his Patient with Expressions of Comfort, and with Wine, or Cordials, before he enters on the Operation.

The Operation itself described.

VII. The Operation itself is next begun by an annular Incision made through the Skin, by the Surgeon, with a small Scalpel, the Arm being extended in a parallel or even Direction, by the Assistants: one of which is then ordered to draw the Skin upward as much as possible. The Surgeon next divides the Flesh, down to the Bones, all round, close by the Margin of the retracted Skin with the larger crooked Scalpel (*Tab.* XIII. *Fig.* 2.) by which Procedure the Skin will wrap over the Stump, and the whole will be healed a vast deal sooner than by the Method formerly used. The Surgeon now takes the Scalpel, with which he divided the Skin, or else the double-edged Catlin, *Fig.* 4. and therewith cuts through the Flesh and Ligaments betwixt the Ulna and Radius; thereby also separating the Periosteum from the Bones where the Teeth of the Saw are to pass, to avoid violent Pain and Inflammation from a Laceration of that nervous Membrane by the rough Teeth of the Instrument. This is no sooner done, but the Assistants draw back the incised Flesh above and below, to open a Passage to the Bones: and that the Flesh above may be drawn up as much as possible, to cut off the Bone higher than the Incision, you must

muſt apply the ſlit Piece of Linen [a] (mentioned before at *N.* IV. 6.) ſo that its Heads being pulled upward by the Aſſiſtant who holds the ſuperior Part of the Arm, he ſtrives to elevate the Fleſh, that the Bone may be taken off as high as poſſible, by which means the Stump will be more eaſily and neatly covered, and the Wound much ſooner healed. The Surgeon muſt fix his Saw in this Operation, ſo that it may work upon both the Bones of the Cubitus at the ſame Time: without which Caution, he will be liable either to cut one of them longer than the other, or elſe occaſion a Fiſſure or Splintering of the ſingle Bone, when it becomes ſo far divided as not to be able to bear the Streſs of the Saw. He muſt alſo move the Saw gently at the beginning, till it is well entered; and then he may go on faſter, but with Diſcretion: and to prevent the Saw from being pinched or obſtructed in it's Motion by the Bones, the Aſſiſtant who holds the ſuperior Part of the Arm ſhould a little elevate the ſame, as the Hand ſhould be a little depreſſed by the other Aſſiſtant, ſo as to make a Space large enough for the Saw to move freely: but this muſt be done gently and cautiouſly, for fear of breaking the Bones. And thus in one Minute or two the Amputation may be completed.

VIII. When the Surgeon has thus amputated the Hand with Part of the Cubitus, his next Buſineſs is, to make a ſtrict Compreſſure and Deligation upon the larger Arteries to ſuppreſs the Hæmorrhage. But the better to diſcover the divided Arteries, the Surgeon muſt order the Aſſiſtant who holds the Tourniquet to relax the ſame a little: or, if it be the Screw Tourniquet, *Tab.* V. or VI. he may looſen it a little himſelf; by which means the Blood ſtarting from the Arteries, will ſhew their divided Orifices. If the Patient be plethoric, the Surgeon may be leſs ſparing of the Blood at this Time, which muſt be received by a proper Veſſel on the Floor: but in caſe of Weakneſſes, the Tourniquet muſt be inſtantly tightened again, to reſtrain the Flux. When the Cubitus is divided very low, near the Carpus, there will not be any great Occaſion to ſecure the Arteries by a Ligature with Needle and Thread: becauſe the two or three Branches which run there, are but ſmall, and may be well enough ſecured by Compreſſes of Lint with ſome Bits of *Vitriol. Roman.* or only by ſquare Linen Compreſſes [b]. But the Fleſh and Ends of the Bones are to be well ſecured and inveſted with Doſſils of dry Lint; over which again fix a large Piece of the Fungus called *Crepitus Lupi*, with or without a large Bolſter of Tow, to be ſecured and retained on the Stump by a wet Bladder, or a Plaſter cut in the Shape of a *Malta* Croſs: or, inſtead of a Plaſter in that Form, you may more advantageouſly apply two or three long and narrow ones acroſs each other in the Form of a Star, upon the Stump; by which the Skin may be drawn down, ſo as to cover the Wound, and procure a ſpeedy Cicatrization [c]: over the Plaſters you are again to place a large Compreſs in form of a *Malta* Croſs, ſo that

Treatment after the Operation.

[a] Some Surgeons uſe a thin Plate of Steel to elevate the Fleſh, inſtead of this Piece of Linen.

[b] M. CHABERT, in his *Obſ. Chirurg. Pariſ.* 1724, aſſerts the Application of Vitriol to be here unneceſſary: ſince the Blood may be ſecurely ſtopped, and the Arteries compreſſed, by properly diſpoſing Linen or Lint formed into Doſſils or Compreſſes about the Ends of the Veſſels; ſecuring them by a cloſe Deligation or Bandage: which, in weak Patients, I have found to ſucceed very well. Others think the Application of Cauſtics both unſafe and injurious, becauſe the Eſchar formed by the Vitriol frequently recedes or ſeparates from the Veſſel, and excites a profuſe Hæmorrhage. *V.* RUYSCH *Epiſt. de nova Methodo Amputandi, &c.*

[c] This Method LE DRAN highly recommends, *Obſ. Chirurg.* Tom. II. p. 309.

that it may closely invest the End of the Limb, where it should be held by an Assistant while the Ends are brought up and applied round the Arm. And lastly, you must fix first, one large square, and then three long and narrow Compresses upon the Stump, so that the last may intersect each other in form of a Star, and come up towards the Humerus; then you finish the Deligation with a long Roller in the manner we shall direct at large, in treating of Bandages for the Arm.

The Hæmorrhage sometimes suppressed by the actual Cautery, or Ligature.

IX. Most of the antient, and not a few of the modern Surgeons, approve of the actual Cautery for restraining the Hæmorrhage from the divided Arteries. This Practice is deservedly rejected by the most expert Surgeons of the present Time; not only for the severe Torture it gives the Patient, but because it is at best very suspicious, and even dangerous; especially in Amputations of the Humerus or Femur: for the Eschar formed by the Cautery very often separates in two or three Days time from the End of the Vessel which is stopped, and thereby occasions a profuse, if not a fatal Hæmorrhage. However the Use of the Cautery will be more likely to succeed in Amputations of the Cubitus or Tibia, than in the Parts beforementioned: but even here it is best to follow the Method at *N.* VIII. preceding, and never to have recourse to the actual Cautery without absolute Necessity. Lastly, if, for the greater Security you are desirous of taking up the Ends of the divided Arteries with Needle and Thread, according to the modern Practice, (which, in my Opinion, is not very necessary in Amputations at the lower End of the Cubitus or Tibia) you are in this Case to take hold of the End of each divided Artery with a Pair of Pliers, termed the Crow's Bill (*Tab.* III. *Fig.* 4. or *Tab.* XIII. *Fig.* 5 and 6.) or some other of a convenient Make: and after passing round your crooked Needle with strong waxed Thread, with the latter you tie up the End of the Vessel.

Amputation of the Humerus.

X. When the Amputation is to be made above the Elbow in the Humerus, the Operation is to be performed almost directly in the same Manner as we prescribed for the Amputation in the Cubitus: except that the brachial Arteries, of which there are sometimes but one, sometimes two or thtee, are to be always taken hold of with a Pair of Pliers, and secured by Ligature with a crooked Needle and waxed Thread, as we just before mentioned in *N.* IX. for in these large Arteries the Use of Styptics or Cauteries are found to be of little or no Efficacy. After the Extremities of the large Arteries are tied up, you must relax the Tourniquet a little, to discover the rest; which are to be also secured in the same manner. Some Surgeons pass a small Needle and Thread through the End of the Artery, whilst held by the Pliers, joining the Thread with that with which they next make the Ligature: which Method, they take, in order to secure the Ligature from slipping off from the End of the Vessel. There are others, who, instead of extending the Ends of the Vessels with a Pair of Pliers, use a very crooked Kind of Needle, with strong waxed Thread, with which they perforate the circumjacent Flesh, first on one Side, and then on the other Side of the Artery, tying up a good deal of the adjacent Flesh together with the End of the Vessel, in order to prevent the Thread from cutting through the arterial Coats: but I think either of these Methods are rather inferior than preferable to the first, in which the Artery is extended with a Pair of Pliers, and then secured by Ligature with a crooked Needle and waxed Thread, passed round the End of the Vessel: for in the two latter Methods there is Danger





Plate 13.







Fig. 3.

Fig. 5.

Plate 13. Vol. I. P. 361.



of paſſing the Needle wide of the Veſſel, or at leaſt the End of the Artery may eaſily fly back, or ſlip out of the Ligature[a].

Treatment after the Dreſſing.

XI. When you have dreſſed the Stump, and compleated the Deligation, according to *N.* VIII. the next Buſineſs is, to give the Patient a Draught of Wine or ſome Cordial; and when he is laid down upon the Bed, the End of the amputated Arm ſhould be compreſſed by the Hands of an Aſſiſtant for ſome Hours, which will not only make the Dreſſings adhere more cloſely, but alſo prevent any conſequent Hæmorrhage. This done, you may, by degrees, relax the Tourniquet ſufficiently to admit of the Blood's Circulation through the Part: and, if upon the Relaxation of it, you meet with no Blood from the Wound, it is a Sign the Operation has been well compleated. In the next place you muſt recommend Reſt to the Patient, and order ſome nouriſhing Emulſion inſtead of common Drink, and paregoric Draughts to be repeated at proper Intervals; that he may hereby recover his loſt Strength, and be eaſed of his Pains by Sleep. The next Day you may again looſen, or elſe totally remove the Tourniquet, and give Orders for a proper Diet and Regimen, ſuch as will abate the febrile Heat and Motion of the Blood, and ſecure the Patient from a freſh Hæmorrhage, as in *Part* I. *Book* I. *Chap.* I. *N.* XLIII. Theſe Accidents may be ſtill better prevented by the Uſe of Phlebotomy at Diſcretion, with cooling Draughts and Powders: but Veneſection muſt be avoided, when the Patient is weak, or has loſt much Blood. If a freſh Hæmorrhage ſhould appear, ſo as not to be ſuppreſſed by the Application of another Compreſs and Bandage, with compreſſing the Stump for ſome Time with the Hands, (which are generally ſufficient) in that Caſe you muſt re-apply the Tourniquet; and, after removing the Dreſſings, make a freſh Ligature upon the Ends of the Arteries: or, if the Ends of the Arteries cannot be taken hold of, you may apply the actual Cautery, and defend the Stump with a larger Quantity of Lint, then ſecure it with an exact Deligation and Compreſſure for ſome time by the Hands, till the Hæmorrhage ceaſes.

When and how to renew the Dreſſings.

XII. The firſt Dreſſings and Bandage ought not to be removed from the Stump before the third or fourth Day, when the Mouths of the divided Veſſels may be ſuppoſed to be well cloſed and united: but in caſe of Accidents, intenſe Pains, Inflammation, Hæmorrhage, or the like, you muſt renew them ſooner. Nor is it amiſs to order a Servant to attend conſtantly for the firſt Week at the Patient's Bed-ſide, provided with a Tourniquet, with which an incidental Hæmorrhage may be ſuppreſſed, till the Surgeon can be called to renew the Deligation. But if every thing ſucceeds well, in renewing your Dreſſings, you ought to remove them one after another very tenderly, and thoſe which are next, or adhere to the Wound, ſhould not be touched at all, much leſs violently forced away, if you are deſirous to avoid irritating the Part, and inducing an Hæmorrhage. 'Tis in this Caſe much the beſt for you to leave the adhering Dreſſings upon the Part for a few Days, and to moiſten them at each Dreſſing with warm Wine or its Spirit, till they become looſe, and ſeparate ſpontaneouſly in the Suppuration, without uſing any Violence. After the firſt Dreſſing, you need not dreſs again above once every other Day,

[a] See DOUGLASS, *Syllabus Operat. Chirurg.* p. 44, 45. where, after the Amputation, he adviſes the Surgeon to unite the Skin croſswiſe.

Day, or every Day at moſt, except your Diſcharge be great, and in the Summer time.

What is to be obſerved in the Dreſſings.

XIII. In renewing your Dreſſings, it is chiefly neceſſary for you to obſerve, that your Wound be well and gently cleanſed from all the foul Matter with Lint, and then to dreſs it with flat Plates or Pledgits of ſcraped Lint; of which that next the Wound ſhould be moiſtened with ſome digeſtive Ointment, and the reſt applied dry. The Pledgits of Lint are to be ſecured and retained upon the Stump by three, four, or ſix ſticking Plaſters of *Emp. Diapalmæ*, or the like, of about a Foot in length, and a Thumb's breadth, croſſing each other upon the Part like a Star: over theſe Plaſters muſt again be fixed a large ſquare Compreſs; and over that three other long and narrow Compreſſes in a ſtellar Poſition, ſecuring the whole by Deligation with your Roller. When your Dreſſings have been thus continued for about a Fortnight, there will not be occaſion for ſo much Lint, nor ſo many Compreſſes as at firſt: nor need you then make your Bandage ſo tight, as there is no Danger of any Hæmorrhage. But in the mean time you muſt continue to treat the Wound with digeſtive Ointments and vulnerary Balſams, retained with Lint, a Plaſter, Compreſs and Bandage, as in other Wounds, till it be healed; which uſually happens in about two Months. For the reſt, it may be here proper to adviſe the Surgeon to apply the Tourniquet, before he removes the firſt Dreſſings; eſpecially in Amputations of the Humerus or Femur, in order to prevent an Hæmorrhage: or at leaſt the brachial Artery ſhould be compreſſed in the Middle of the Arm by the Thumb of an Aſſiſtant.

Treatment of the Fever.

XIV. Laſtly, as Amputations are often followed ſoon after with a Fever, eſpecially in plethoric and ſtrong Habits, it will, in that Caſe, be neceſſary to uſe Phlebotomy with paregoric and cooling Medicines, joined with a proper Regimen and Diet: without which there may be Danger of loſing the Patient, either by the Violence of the vulnerary Fever, as it is termed, by a Sphacelus of the Part, or other bad Accidents.

An EXPLANATION *of the* THIRTEENTH PLATE.

Fig. 1. Exhibits a ſmall-ſized Scalpel, more commodious for dividing the Skin and Fleſh in Amputations than the large crooked one following.

Fig. 2. Is the large crooked or falciform Knife, commonly uſed for dividing the Fleſh to the Bone in Amputations of the upper and lower Extremities, though in moſt Caſes I prefer the ſmall one, *Fig.* 1.

Fig. 3. The Catlin, or double-edged Scalpel, for dividing the Fleſh and Ligament betwixt the Bones of the Cubitus and Tibia; which may be alſo performed by a leſs and ſingle-edged Scalpel, like that in *Tab.* I. G. This Knife is alſo uſed in the Method of amputating the Tibia, which preſerves the Calf.

Fig. 4. Repreſents the Saw uſed for amputating Bones of the Limbs. This Inſtrument is by many delineated as large again as our Figure of it: but a Saw of the ſame Size, or but little larger than our Figure, will perform the Operation as well, and even more commodiouſly than a larger. This and the two preceding Inſtruments are uſually embelliſhed with various Ornaments; which may ſerve to encumber them, and enhance their Price, but can add nothing at all to their Uſefulneſs.

Fig. 5. Repreſents a Pair of Pliers, furniſhed with Teeth at one End, and a Spring at the other, for taking hold of the Ends of divided Arteries, in order to ſecure them by Ligature with ſtrong Thread, and ſtop their bleeding in Amputations of the upper and lower Extremities.

Fig. 6. Is another Pair of Pliers for the ſame Uſe, taken from M. GARENGEOT; which may be alſo made with very flat or no Teeth at the End, to avoid injuring the Coats of the Artery.

CHAP. XXXV.

Of Amputating the FOOT *and* LEG.

The Place for amputating the Tibia.

I. THE antient Surgeons, in amputating the Foot at the Tarſus or Metatarſus, uſed a large Chiſſel and Mallet, and ſometimes a Pair of large cutting Pincers, with which they ſeparated the diſeaſed Parts, and then treated and healed the Wound with Balſams in the uſual Manner: which Practice is confirmed and explained by SCULTETUS, in his *Armament. Chirurg. Tab.* LIV. But as the Tendons and Ligaments, ſeated in thoſe Parts, are, in this Method, violently lacerated and contuſed, the modern Surgeons have therefore juſtly preferred the Amputation of the Toes and Metatarſus by the Scalpel; conducting the Remainder of the Cure as in other Wounds: and in this Manner the Leg may be much better ſupported by the Heel or Stump, than by a wooden Machine*. But becauſe they were afraid of this Practice, from the Difficulty of covering the Bones, and healing up the Wound, they rather followed the more dangerous Method of amputating the Leg about four Fingers Breadth below the Knee, inſtead of taking it off in the lower Part of the Tibia. By this means, though they cut off a large Part of the Leg which was not yet diſordered, they avoided the Deformity and Inconvenience in ſitting down, which the Patient would have met with from preſerving it on: for a long Stump of the Leg can neither be ſtood upon nor well adapted to a wooden Machine: therefore it was thought moſt convenient to amputate it in the upper Part of the Tibia, about a Hand's Breadth below the Patella, to avoid injuring the Tendons of the flexor Muſcles, and the better to adapt the Knee to a Silver or wooden Leg. I am indeed ſenſible that many Surgeons, even at preſent, approve of amputating no higher than the Diſorder has ſpread itſelf, agreeable to the Advice of SOLINGEN, VERDUYN, and DIONIS: but I think their Authorities ought to be but little regarded; not only becauſe of the Difficulty there will be of adapting a wooden Machine to the lower Part of the Leg above the Ancle, but alſo upon the account of the Deformity which the long Stump of the Leg will occaſion, if the wooden Machine is adapted to the Knee.

Obſervations peculiar to Amputations of the Leg.

II. With Regard to the Inſtruments and Dreſſings uſed in this Operation, they are almoſt the ſame which we before deſcribed for amputating the Arm: only it may be here neceſſary to add a few Cautions which relate more particularly

* This GARENGEOT adviſes, *Oper. Chirurg.* Tom. III. p. 417. Edit. 2.

larly to Amputations of the Leg. Theſe are, (1.) To place the Patient upon a low Seat or Bed, ſo that he may lean backward, and extend his Legs[a]. (2.) To ſhave off the Hair with a Razor from the Part where the Amputation is to be made, to prevent the Plaſters, afterwards applied, from adhering to them, ſo as to give the Patient intenſe Pain in removing them. (3.) To ſecure the divided Arteries, which appear in the Stump of the Leg, rather by Ligature, with Needle and Thread, than by Styptics, or actual and potential Cauteries: for though theſe Arteries do not appear very large, yet if they are not ſecured by Ligature, they generally open and bleed profuſely ſoon after the Deligation; eſpecially if the crural Artery be not well ſecured with narrow Compreſſes and Bandage. (4.) The crural Artery is to be compreſſed with the Tourniquet, either of the common Sort, turning with a Stick, or the modern Screw Tourniquet: or elſe you may make a ſtrict Ligature above the Knee with a Bandage twiſted in a cylindrical Form, ſo as to compreſs the Artery deſcending in the Ham, as in *Tab.* XIV. *Fig.* 4. D. yet in my Opinion, it is much better to apply the ſame Ligature higher up upon the Thigh, in order to compreſs the Artery, eſpecially when the Leg is to be amputated near the Knee. See *Tab.* III. *Fig.* I. L. M; by which means the Dreſſings may be more conveniently applied after the Operation, than if the Tourniquet was fixed nearer the Knee.

VERDUYN's Method of Amputating.

III. We have another new Method of amputating the Leg, propoſed by VERDUYN, in a Diſſertation upon the Subject in the Year 1696: which Practice he ſtrongly recommends for the pubiick Good; though he does not pretend to be the original Author of it. There are indeed many who attribute the Honour of inventing this Operation to one SABOURIN of *Geneva*, as GARENGEOT, and ſome other Members of the Royal Academy; who aſſert, that in their Time VERDUYN performed the Operation firſt at *Geneva*, and then at *Paris*. At the ſame time I find the Operation deſcribed and performed by the *Engliſh* Surgeons LOWDHAM and YOUNG, in an *Engliſh* Treatiſe concerning the wonderful Virtues of Oil of Turpentine in Hæmorrhages, together with a new Method of amputating, by JAMES YOUNG, 8vo *Lond.* 1679. The ſame Operation was afterwards improved and deſcribed by my Friend KOENERDINGIUS, Surgeon of the Hoſpital at *Amſterdam*, in his *Dutch* Treatiſe *De Gangræna & Sphacelo, Cruraque amputandi Ratione veteri ac nova*, 8vo *Amſtel.* 1698; which was the ſame Year in which VERDUYN twice performed this new Method of Amputation: a brief Deſcription of which is as follows.

IV. Firſt the *Tendo Achillis* is divided from the Ancle by the Scalpel, *Tab.* XIII. *Fig.* 3. then a longitudinal Inciſion is made upwards, and the Tendon ſeparated from the Bones of the Leg as high as the Part where the Bones are to be amputated by the Saw. See *Tab.* XIV. *Fig.* 4, 5, 6, 7. This done, the Fleſh compoſing the Calf of the Leg. *Fig.* 6. A, is drawn backward with a Cloth towards the Ham, by the Hand of an Aſſiſtant: and then the Integuments and Fleſh upon the Forepart, and betwixt the Bones, are divided in the uſual Manner, by a proper Scalpel, *Tab.* XIII. *Fig.* 1, and 3; and the Bones next amputated by the Saw then the Fleſh is brought over, and adapted to the

[a] HILDANUS in this Caſe places the Patient on the Ground, but his Foot on a low Stool.

the Stump of the Leg, after it has been firſt waſhed with Spirit of Wine: and if there be any unequal and ſuperfluous Parts, they are cut off with a Scalpel; the Remainder being retained in its proper Situation by ſticking Plaſters, or a few Stitches with a Needle and Thread. Laſtly, Compreſſes with a wet Bladder and Bandage are applied in the Manner we before directed, in treating of Amputations in general: or, inſtead of them may be uſed a retentive Machine, figured by VERDUYN and GARENGEOT, for the Purpoſe, being made of Leather, with Straps and Buckles, by which the Stump being ſecured, it is then to be compreſſed for a few Hours by the Hands of an Aſſiſtant, till there is no Danger of an Hæmorrhage. To prevent that, you may alſo apply the Screw Tourniquet, *Tab.* V. *Fig.* 6. or *Tab.* VI. *Fig.* 1. thus the Operation is compleated, the Advantages, of which to the Patient, according to the forementioned Authors, are many: as, (1.) The Calf of the Leg being thus preſerved and adapted to the Stump, cloſes and compreſſes the Mouths of the divided Arteries, ſo as to prevent an Hæmorrhage, without the Uſe of Cauteries, or the Application of Ligatures. (2.) The Ends of the Bones being thus immediately covered with the Fleſh, are not ſo liable to be infeſted with a Caries, as they frequently are in the common Method, which greatly retards, if it does not fruſtrate the Cure. (3.) The Fleſh of the Calf readily unites with the Ends of the divided Bones of the Leg; ſo that by treating the Wound with vulnerary Balſams, in the ſubſequent Dreſſings, the Cure is ſpeedily compleated. Laſtly, (4.) The Fleſh thus adapted to the Ends of the Bones, ſerves as a Pillow ever afterwards to ſupport them; ſo that the Patient may eaſily ſit down, without being obliged to bend the Stump, as he muſt do after the common Method. Add to this, that the Stump may be adapted perpendicularly to a hollow wooden Leg, ſo that the Patient may ſtand or walk upright upon an artificial Leg, as upon his natural one. Every time the Stump is dreſſed, the Portion of Fleſh which wraps over it, muſt be gently ſupported, and preſſed up againſt the Ends of the Bones, that its Weight may not make it ſeparate or ſubſide, ſo as to prevent its uniting. A more particular Account of this Method may be ſeen, illuſtrated with proper Figures, in the forementioned Treatiſe of VERDUYN.

V. Notwithſtanding the before-deſcribed Method had been ſeveral times performed with Succeſs by VERDUYN, and ſome others, yet it met with the Approbation of but few Surgeons: ſo that it was not able to prevail over the common and received Method of amputating the Leg: inſomuch, that it was ſoon after deſerted even by its own Patrons, VERDUYN and KOENERDINGIUS: to which add, that the Patient, upon which SABOURIN performed this Operation at *Paris*, died ſoon after it, as did ſeveral at *Amſterdam*; at which laſt Place ſeveral Patients were troubled with acute Pains, and other bad Accidents, from little Splinters, or the rough Ends of the Bones irritating the Fleſh, even after the Stump was healed up: not to mention the large Quantity of Blood loſt by SABOURIN's Patient, which was even greater than in the common Method of amputating; which, with other Inconveniences, induced KOENERDINGIUS to prefer the common before this new Method, in his Treatiſe on this Subject. Notwithſtanding all this, we find M. GARENGEOT, who ſeems to be ignorant of the forementioned Writings of YOUNG and KOENERDINGIUS on the Subject, endeavouring lately to recommend and re-eſtabliſh this uncommon Method of amputating: as may be ſeen in *Chirurg. Operat. Chap.* of Amputations of the

VERDUYN deſerts his Method.

Leg. M. GARENGEOT there relates, that there were ſeveral Men then living in *France*, who had the Operation happily performed on them in this Manner; ſo that they could not only ſit down eaſily, but alſo leap very nimbly: but if we would reaſonably expect to ſucceed in this Method, the Patient ought to be not only healthy in all other reſpects, but the Cauſe, which requires the Limb to be amputated, ſhould be from ſome external Violence.

The ſame Method practicable in the Humerus.

VI. Laſtly, it is to be obſerved, that the new Method of amputating, which we have been now deſcribing, may, according to the Opinion of our modern Surgeons, be not only performed in the Leg, but alſo in the Cubitus, by preſerving a Quantity of the Fleſh and Integuments, to wrap over the Ends of the Bones. Agreeable to this, the Operation was in the ſame Manner performed with Succeſs by RUYSCH, in the Preſence of VERDUYN and BORTELIUS his Kinſman. See the Treatiſes on this Subject by YOUNG and KOENERDINGIUS; alſo RUYSCHII *Epiſt. problemat.* XIV. *de nova Artuum decurtandorum Methodo.*

CHAP. XXXVI.

Of Amputating the THIGH.

Amputation of the Femur, when neceſſary.

I. THE Surgeon frequently finds it neceſſary to amputate the Leg above the Knee, removing Part of the Thigh itſelf, when a Mortification has reached the Joint, or when the lower Head of the Femur is carious, ſphacelated, cruſhed to pieces, or the large crural Artery irrecoverably wounded. In theſe Caſes the Succeſs of the Operation is very dubious, eſpecially when the Amputation is made very high up in the Thigh. Nor is the Patient in danger of being loſt only from a profuſe Hæmorrhage, from the Diviſion of ſo large an Artery as that of the Femur; but the Quantity of Matter diſcharged daily from ſo large a Wound does often ſo much extenuate and weaken the Patient, that he cannot ſubſiſt till the Cure is completed: therefore whenever the Surgeon finds it neceſſary to amputate in the Femur, he ought to do it as low as poſſible, as near within three Fingers Breadth of the Knee as he can, leaving a good deal of Fleſh, and more of the Skin, to wrap over the End of the Stump: by which means the Cure of the Wound will be much expedited, the Diſcharge of Matter at each Dreſſing rendered leſs profuſe, and the Patient, not being ſo much impaired in his Strength, will be more likely to get happily through the Cure.

Application of the Tourniquet.

II. The Application of the Tourniquet for compreſſing the crural Artery, whether it be the common one with the cylindric Ligature and Turn-ſtick, or the Screw Tourniquet, muſt be made upon the upper and internal Part of the Thigh, as near as you can to the Place where the Head of the internal Vaſtus Muſcle and the Triceps touch each other, as in *Tab.* III. *Fig.* 1. L. M. without this Precaution you may be liable to have ſuch a profuſe Hæmorrhage from the large femoral Artery as will inevitably deſtroy your Patient; which frequently happened to the antient Surgeons before the Invention of the Tourniquet.

III. With

III. With reſpect to Amputations of the Thigh in general, little more need be added to what has been ſaid on this Operation in the Arms and Legs: as in the firſt place, to let the Hair be ſhaved off, and after you have made a circular Inciſion through the Integuments with a ſmall Scalpel, Tab. XIII. *Fig.* 1. to extend or draw them upwards as much as poſſible before you divide the Fleſh, or Muſcles, which laſt you muſt amputate a good deal higher than the circular Inciſion through the Integuments. You may cut through the muſcular Parts at your ſecond Inciſion, either with the Scalpel, with which you divided the Integuments, or with a Knife for amputating Breaſts in *Tab.* XXII. *Fig.* 7. or elſe with the large crooked Knife in *Tab.* XIII. *Fig.* 2. with either of which you muſt cut all round cloſe to the Bone: by which Method of proceeding you will have the Stump of the Bone covered over with Fleſh and Skin in a little time, ſo as to be healed in a few Days; and at the ſame time you avoid the Riſque of a Caries in the Bone from its being expoſed to the Air, as we once before obſerved. For want of this Precaution in Amputations of the Thigh, when the Muſcles have been divided even with the Integuments, the Muſcles have contracted and drawn themſelves up to ſuch a Degree, that I have frequently ſeen the Bone ſtanding out like a Stick for above two or three Fingers Breadth from the Fleſh. In this Caſe the Patient muſt be a long Time, and be much weakened by the Diſcharge of Matter, before the Muſcles can be extended and brought down, ſo as to cover the End of the Bone, without which the Cure can never be completed. With regard to the Hæmorrhage in Amputations of the Thigh, that muſt be always prevented by making an exact Ligature upon the femoral Artery, which is much too large to be ſafely ſecured by any other Method; and, for the ſame Reaſon, your Ligature upon it muſt be very firm and ſecure, by tying it up with a ſtrong Thread paſſed round after the End of the Artery is extended or drawn a little out from the Fleſh with a Pair of Forceps, or a Tenaculum, *Tab.* XIII. *Fig.* 5. and 6[a]. If there appear to be more large Arteries than one divided, they muſt be alſo ſecured by Ligature in the ſame Manner; but for the ſmaller Arteries, it may be ſufficient to cloſe them by Styptics, or Vitriol, and Doſſils of ſcraped Lint without Ligature. The Dreſſings and Deligation are to be much the ſame for an amputated Thigh, as we before directed for an Amputation of the *Humerus:* only the Quantity of Lint, Fungus, Bladder, Compreſſes, *&c.* muſt be proportionably larger, and the Bandages much longer: to which you muſt here add a long, thick, and narrow Compreſs, to be placed all along the Thigh over the crural Artery, and ſecured there by a Bandage peculiar to itſelf: or, inſtead of this, you may fix the Tourniquet, *Tab.* V. *Fig.* 6. or *Tab.* VI. *Fig.* 1. and leave it upon the Limb for ſome time. The Deligation being completed, and the Patient put to Bed, his Thigh muſt be placed in an eaſy elevated Poſture on a Pillow, that the Impetus of the Blood, on the End of the Artery, may be leſs, than in a direct Poſition: which will greatly conduce to the Prevention of a freſh Hæmorrhage. Laſtly, the Stump ſhould be compreſſed for ſome Time by the Hands of an Aſſiſtant, ordering a proper Diet, Regimen, Medicines, *&c.* as we obſerved in Amputations of the Humerus.

The Method of amputating and dreſſing the Femur.

IV. If

[a] PETIT gives a Deſcription of another Ligature, *Mem. Acad. Reg. Pariſ.*

Treatment of Limbs amputated by Gun-ſhot.

IV. If Part of the Leg or Arm ſhould be carried away by a Bullet-ſhot, or Cannon-ball, or be torn off by a Cart-wheel, or Mill, or ſome ſuch other Machine, the firſt Step to be taken by the Surgeon in theſe Caſes, is, 1. immediately to apply the Tourniquet to compreſs the Artery, and ſtop the Hæmorrhage: and then, 2. to cut off the rough End of the Bone by the Saw, or cutting Pincers, that there may be no Points or Splinters to irritate the ſenſible and fleſhy Parts: but if there are no Splinters, or rough Parts, the Surgeon need not cut off any thing: laſtly, 3. to ſecure and cloſe up the Ends of the wounded Arteries, either by Ligature, (when they are large and acceſſible,) or elſe by the Cautery, or by Compreſſure with Lint, Styptics, and Compreſſes, according as particular Circumſtances may indicate to the Surgeon: which being performed, the reſt of the Dreſſings and Deligation are to be completed in the Manner we have before directed for other Amputations[a].

BOTALLUS's Method of amputating rejected.

V. The celebrated *French* Phyſician BOTALLUS formerly invented a very expeditious Method of amputating Limbs in an Inſtant, by letting a ſharp Inſtrument fall down upon them from a certain Height loaded with a great Weight; by which means the Limb is ſtruck off at one Blow, without the Uſe either of Knife or Saw. BOTALLUS has been alſo ſeconded in this Method of amputating by HILDANUS: notwithſtanding which, the Artifice has been reaſonably rejected by almoſt all the prudent Surgeons, who have ſucceeded them: for it is hardly poſſible that a Limb ſhould be taken off in this Manner without ſhattering or ſplintering the Bone. Conſult BOTALLUS in his Treatiſe, *De Vulneribus Sclopetorum.*

The adapting of artificial Limbs.

VI. After the Stump is healed up, the Surgeon may provide an artificial Limb of Silver, for thoſe who can afford it, or of Wood for others; adapted to the Stump, ſo that it may be faſtened on by Straps and Buckles, or by Springs. Of theſe Machines we are furniſhed with various Specimens in AMB. PAREY, HILDANUS, SOLINGEN, *&c.* and by our modern Artiſts, who make theſe kinds of Inſtruments, and other curious Machines: but, for the poorer Sort, it may be ſufficient to ſupply them with a wooden Machine, turned and cut into a proper Shape, with a Hollowneſs or Cavity at the upper End for receiving the Stump of the Knee, that they may, by this means, be enabled to walk, or ſit down, though not in an elegant Manner.

Removal of a Caries.

VII. As a Caries of the Bone is no unfrequent Accident after Amputations, the Surgeon ſhould therefore endeavour to guard againſt it as much as poſſible: even at its very firſt Appearance he ſhould ſtrive to remove it, either by the Application of *Euphorbium*, or the actual Cautery, becauſe it prevents the Progreſs of the Cure; notwithſtanding the Writers in Surgery uſually paſs by this Accident, without taking notice thereof. There ſtill remains a Practice, which, in my Opinion, will very often ſucceed beyond either *Euphorbium*, or the Cautery; and that is, to exfoliate or pare off the diſeaſed Parts of the Bone with a Knife or Raſp till you come to the ſound: by which means the Fleſh will then readily unite with the Bone to complete the Cure, which it cannot while the Caries remains.

CHAP.

[a] PETIT has deſcribed a Machine for ſuppreſſing the Hæmorrhage in the crural Artery after this Amputation, which will effect it without a Ligature on the Artery, or any other Method abovementioned. *Mem. Acad. Reg. Pariſ.* A. 1731. This Machine you will find in my *Plate* XXXIX. with a particular Deſcription of it.

CHAP. XXXVII.

Of Amputating the ARM *in its Articulation with the* SCAPULA.

The Design of this Chapter.

I. THOUGH I never yet attempted to amputate the Humerus in its Articulation with the Scapula, nor ſo much as found it treated of by any of our Chirurgical Writers, except LE DRAN, *Obſ.* 43 and 44. after whom the Operation is deſcribed, without mentioning his Name, by GARENGEOT, (*Chirurg. Operat. Cap.* LIV.) yet, that the Surgeon may not be ignorant of what has been advanced on this Head, I ſhall make it the Buſineſs of this Chapter to give a brief Deſcription thereof.

When the Arm may be thus amputated.

II. According to the two laſt mentioned Authors, there are two Caſes in which it may be neceſſary to amputate the Arm in its Articulation at the Shoulder. The firſt is, when the upper Part of the Humerus is violently contuſed and ſhattered by a Cannon-ball, Bomb, or Granade: the other Caſe is, when the upper Head of the *Os Humeri* is irrecoverably diſeas'd from ſome internal Cauſe, as from an Abceſs, a Caries, or *Spina ventoſa*; to which we may add, a Mortification of the Arm extended to the Shoulder, *&c.*

Previous Requiſites for the Operation.

III. But before you enter on this dangerous and difficult Operation, it will be abſolutely neceſſary to have every Part of your Apparatus of Inſtruments and Dreſſings prepared and diſpoſed each in their proper Order: after which you are to fix the Patient upon a convenient Seat with his Face covered. You muſt next obſerve, that the Tourniquet is not here fixed upon the Arm, as we before deſcribed for the common Amputation of it: but that Inſtrument is, in this Caſe, laid aſide, and the Trunk of the brachial Artery is firſt ſecured by Ligature in the following Manner, before you begin to amputate.

What is to be firſt done in the Operation.

IV. The Patient being properly ſeated, with his Arm extended, and ſecured by an Aſſiſtant, you muſt then carefully ſearch out the true Seat and Courſe of the brachial Artery at the Axilla: in doing which you will be much aſſiſted by being previouſly verſed in the Anatomy of the Part. If the Tumor ſhould be ſo large as to prevent your finding out the Artery, by feeling thro ugh the Integuments, you make a longitudinal Inciſion through them to the Bone, on each ſide the Arm, ſo that you may paſs your Fingers by the Bone, and diſcover the Artery: which done, you muſt then paſs a large Needle[a] with ſix or eight Threads through the Fleſh within two Fingers Breadth of the Cavity in the Axilla, ſo that the Needle may paſs through cloſe to the Bone, and betwixt that and the Artery, without injuring the latter. The Needle and Ligature being thus conveyed betwixt the *Os Humeri* and Artery, the Arm is now let down a little, to relax the Skin, and the Ligature is then tied with a Surgeon's Knot. Your next Buſineſs is, to examine if there be any Pulſe in the Artery below the Ligature as it runs down the Arm: if ſo, your Ligature muſt be drawn tighter till you can perceive no Motion there; and then your Ligature muſt be ſecured from getting looſe by a Knot or two more.

What is to be obſerved in the Operation.

V. There are three Things chiefly neceſſary for you to obſerve in the Operation, after the Artery has been thus ſecured by Ligature to prevent a fatal Hæ-

[a] LE DRAN uſes a ſtraight Needle; but GARENGEOT recommends a crooked one, like that to be ſeen in *Tab.* XIV. *Fig.* 10.

morrhage: theſe are, 1. to leave Skin and Fleſh enough upon the Shoulder; 2. to cut through the muſcular Parts in the moſt convenient Manner; and laſtly, 3. to divide the capſular Ligament which inveſts the Head of the Bone, and connects it to the Scapula, ſo that it may be taken out of the glenoid Cavity in the latter, and be afterwards amputated intirely. To perform each of theſe Intentions with Succeſs and Dexterity, the Surgeon ought previouſly to make himſelf well acquainted with the Nature of the Articulation, with the Poſition of the *Proceſſus Acromion*, and to be careful that a ſufficient Quantity of Skin be preſerved and drawn back to wrap over the Wound; and laſtly, to amputate with his Scalpel two or three Fingers Breadth below the Acromion, ſo as to preſerve a large Portion of the Deltoid Muſcle. This Method will not only fill up the Cavity of the Wound at the Shoulder, ſo as to render it uniform and even, but will alſo much expedite the Cure.

The Manner of amputating.

VII. Every thing being thus far conſidered and advanced, you now take the Scalpel, *Tab.* XIII. *Fig.* 1. or *Tab.* XII. *Fig.* 14. and therewith make your Inciſion through the Integuments, and through the Deltoid Muſcle, as near within the Joint as we before directed: which done, the Arm is then gently elevated, the better to diſcover and divide the Heads of the Biceps Muſcle: and if, in performing this, you divide any conſiderable Arteries or Veins, which bleed ſo as to obſcure your Work, they may be ſtopped for the preſent, either by Compreſſure with the naked Fingers of an Aſſiſtant, or by the Application of Lint and Compreſſes: but if the Hæmorrhage is profuſe, and ariſes from a conſiderable Artery divided, as there frequently is a large Branch here, you muſt, in that Caſe, firſt ſecure it by Ligature, before you proceed farther in your Operation: the next Step is, to divide the Ligament of the Articulation firſt in its upper Part, and then on each Side, but very cautiouſly; moving the Head of the Humerus at the ſame time with your left Hand, that you may only divide the inveſting Ligament of the Articulation without injuring the Artery. Thus you may be ſenſible whether the Arteries are well ſecured: but even afterwards you muſt be very cautious not to wound the Artery, in dividing the reſt of the muſcular Parts between the Articulation. Laſtly, you muſt divide the Skin from the Arm near the Axilla, to leave a triangular Piece with its Corner outermoſt, and its Baſis next the Body, ſo as to be afterwards brought up over the Wound: and thus your Amputation is completed.

What muſt be done after the Amputation.

VII. The Arm being totally removed in the Manner now deſcribed, you muſt next ſearch for the Artery you before ſecured by Ligature, together with a Portion of the Fleſh; and having diſcovered it, you now make another Ligature above the former upon the Veſſel only, by a ſmall crooked Needle, *Tab.* VI. *Fig.* 5. with ſtrong Thread: after which you remove the firſt Ligature from the Fleſh and Veſſel, to prevent it from exciting an Inflammation.

Dreſſings, and Deligation.

VIII. You come now to the Dreſſings of the Stump: which muſt be made with a Pledgit of Lint, with ſmall Linen Compreſſes upon the Ends of the divided Arteries you before ſecured by Ligature: the lower Part of the Skin is then drawn upward, and the upper Part is drawn down together with a Piece of the Deltoid Muſcle: though, in my Opinion, it would be better to apply no Pledgit or Compreſſes to the Arteries or Bone, before you have thus filled the Sinus of the Wound with the adjacent muſcular Parts, and brought the Skin well over; and then you may apply your Pledgit of Lint and Compreſſes: by

which means the Fleſh will more readily unite, and the Wound heal ſooner than if you interpoſed Lint and Compreſſes. In the next place, a large Quantity of Lint muſt be ſecured on the Wound by a ſticking Plaſter, cut in the Shape of a *Malta* Croſs: over which Plaſter you place a large and thick ſquare Compreſs, with a cylindric Compreſs in the Axilla, to reſiſt againſt the Ends of the Arteries, that they may be better able to endure the Impetus of the Blood in their Pulſe. All theſe are to be again retained by a large double Compreſs in the Form of a *Malta* Croſs; and that again inveſted by two other Compreſſes a little more than a Foot long, and four Fingers Breadth. The firſt of theſe is applied obliquely over the Stump, ſo as to let one End come over to the ſound Shoulder, and the other End paſs behind to the ſound Axilla, or about a Hand's Breadth lower: the other and longer Compreſs muſt be placed, ſo as to croſs the former in oppoſite Directions, and to have its Ends croſs each other upon the ſound Shoulder. Laſtly, your Deligation muſt be completed with the Bandage termed *Spica deſcendens*, as we ſhall direct in the laſt Part of our Surgery: but, in making this Bandage, you muſt fix a thick Compreſs, or a ſmall Pillow, in the Axilla, that the Bandage may ſit the neater, and not compreſs the Veins too much which are there ſeated.

An Example of this Method of Amputating.

IX. The abovementioned Operation, as here deſcribed, was performed on a *French* Nobleman for a *Spina ventoſa*, in the upper Head of the *Humerus*, by LE DRAN the elder, with the Conſent and Preſence of the moſt expert Surgeons in *Paris*, as M. MARESCAL, ARNEAU, PETIT, MERY, *&c.* and this he did with Succeſs, the Patient being perfectly cured, as we are told by LE DRAN his Son, and M. GARENGEOT. But the laſt of theſe Authors tells us, in a ſecond Edition of his Operations in Surgery, that the ſaid Nobleman died within ſix Months afterwards of a Plethora*. M. GARENGEOT alſo directs this Operation to be performed for an Abceſs in the Articulation: but whether it would be prudent to perform ſo dangerous and difficult an Operation, for a ſimple Abceſs there, I leave to the Judgment of every experienced Surgeon.

A Method of Amputation not much unlike this, which I recommended in the Arm, after an extraordinary Burn, *A. D.* 1739, you have here as follows: a poor Woman in a neighbouring Town, as ſhe ſat alone at home, being ſeized with a Fit, fell into the Fire; by which Accident her right Arm was burnt to the Bone, from the Hand quite up to the Shoulder, (ſee *Plate* XXXIX. *Fig.* A and B.) there was no Way of preſerving the poor Woman's Life, but by taking off the Arm: but as the Tourniquet could not be applied without giving her exceſſive Pain, (the Skin being burnt to the Neck and Breaſt, and for other Reaſons) I thought it more prudent to take the following Method. I ordered a large Needle, with a Thread of a proper Size, to be paſſed under the Head of the *Os Humeri*, where the brachial Artery deſcends, through the little Fleſh that remained near the Bone. A Ligature was then made with the Thread on the Artery and the remaining Fleſh; then the Fleſh was cut with the Scalpel under the Ligature, and the Bones afterwards ſawed off in the uſual Manner. This was done in the Preſence and under the Direction of my Son, without the Help of the Tourniquet; and with ſuch Succeſs, that the Wound, after the Amputation, bled but little (which is uncommon in theſe

* In his *Operat. Chirurg.* Tom. III. near the End.

Caſes) and healed to our Wiſh, the Woman being ſtill alive and in Health. which ſingular Caſe was publiſhed by my Son the ſame Year; and this Practice there recommended in many other Caſes, particularly in ſtopping Hæmorrhages of the brachial and crural Arteries; as may be ſeen more at large in his Treatiſe on the new Method of amputating the Arm.

An EXPLANATION *of the* FOURTEENTH PLATE.

Fig. 1. Shews the Manner in which the Patient, Surgeon, and Aſſiſtants are to be placed for amputating the Hand, or Arm. A denotes the Patient, B the Surgeon amputating with a Saw; C the Aſſiſtant extending the Hand, D another Aſſiſtant holding the Arm; E the Aſſiſtant who holds the Patient's Body, and takes care of the Tourniquet; F denotes the Diſh or Veſſel placed underneath to receive the Blood.

Fig. 2. Repreſents the Poſition of the Patient, Surgeon, and his Aſſiſtants amputating the Leg. A denotes the Patient ſeated in a Chair; B the Surgeon; C the Aſſiſtant who holds the Foot below the Calf; D the Aſſiſtant who holds the Leg above the Knee: E a Veſſel placed on the Floor, to catch what little Blood may be ſpilt in the Operation.

Fig. 3. Denotes the moſt convenient Part for amputating the Leg at A, and the Thigh at B. but when the Diſorder has extended itſelf higher up in the Thigh, it muſt be amputated proportionably above this Mark, though the Operation is then ſo much the more dangerous.

Fig. 4. Repreſents the Thigh A, with the Leg amputated B, in which may be ſeen the Part for fixing the Tourniquet C D, for amputating the Foot in the Tarſus or Metatarſus. The Tourniquet thus applied may alſo ſerve for amputating the Leg or Thigh, though not ſo conveniently as when placed higher up. In this Figure you have alſo a View of the divided Artery extended a little by the Pliers E, and going to be tied by the Ligature and Knot F. There are ſome indeed who do not approve of this Manner of tying the Ligature: but I have often experienced that it thus anſwers very well[a].

Fig. 5. Deſcribes the Manner of amputating the Leg, ſo as to preſerve the Calf. The Line A B denotes the firſt Inciſion to be made by the Scalpel, *Tab.* XIII. *Fig.* 1. or *Fig.* 3. The Line B C is the Courſe of the ſecond Inciſion, by which the Fleſh of the Calf is ſeparated from the Bones of the Leg. C D the Place where the Bones and reſt of the Leg are amputated. Some reverſe this Courſe of Inciſion, and firſt perforate the Calf with a double-edged Scalpel, *Tab.* XIII. *Fig.* 3. in Line C, and then they direct a Knife in the Courſe B A. But the firſt Method is, in my Opinion, moſt eligible.

Fig. 6. Repreſents the Manner of reflecting back the Calf of the Leg towards the Ham, after it has been ſeparated from the Bones of the Leg by Inciſion: which done, the Surgeon next incides the Integuments, Fleſh and Perioſtæum in the Line B, and then ſaws off the Bones there.

Fig. 7. Denotes a Leg juſt amputated with the Calf A depending, to ſee the Ends of the two Bones; B the Tibia, and C the Fibula.

Fig. 8.

[a] GARENGEOT finds fault with this Method, *de Inſtrument. Chirurg.* Tom. II. p. 219. But LA MOTTE, another of our moſt eminent modern Surgeons, very much approves of it.




D
Fig. 7.
B
Fig. 6.
B
A
Fig. 8.
A
C
B
A
B
Fig. 2.
D
A
G
D
E
E
B
Fig. 9.
A
E
Fig. 10

Plate 14.




Fig. 5.
D
B
C
A
Fig. 7.
B
C
A
Fig. 6.
B
A
Fig. 8.
C
B
A
A
E
D
Fig. 2.
B
D
A
C
E
G
D
E
E
B
Fig. 9.
A
Fig. 10




Fig. 8. Shews the Leg thus amputated, with the Calf A brought over and joined to the Stump B; C denotes Part of the Thigh.

Fig. 9. Represents the Method of applying the Screw Tourniquet (*Tab.* V. *Fig.* 6. or *Tab.* VI. *Fig.* 1.) above the Knee; A denotes the Leg; B the Thigh; CC the Press of the Tourniquet with its subjacent Pillow; D the Place where the leathern or silken Strap E E is fastened by Studs on one Side, and by the Hooks F on the other Side; G the Screw, by turning which, the subjacent Artery is compressed in the Ham.

Fig. 10. Is a large crooked Needle for making a Ligature on the brachial Artery before the Arm is amputated in its Articulation with the Scapula, though the same may be also performed by the straight Needle, *Tab.* XVIII. either of which Needles will also serve for making Setons in the Neck.

PART II. SECT. II.

Of Operations *belonging to the* Head.

CHAP. XXXVIII.

Of making Issues *upon the* Coronal Suture.

I. Issues are sometimes made in the Scalp of the Head upon the Meeting of the coronal with the sagittal Suture: but this Operation is not so frequent in *Germany*, as in *Italy* and *Holland*. Some Physicians think Issues in this Part can be of little or no Service, being not able to discharge any Humours from the internal Parts of the Head; and others again assert them to be of very great Efficacy, for the Truth of which they appeal to daily Experience: so that if we may confide in Experience, and the Authorities of able Physicians, we must readily acknowledge that Issues, made in this Part of the Head, may be highly serviceable in Vertigo's, obstinate Head-achs, Apoplexies [a], Epilepsies [b], Amaurosis [c], Stupidity or Forgetfulness [d], with many other Disorders of the Head, and particularly of the Eyes and Ears, Defluxions, or Catarrhs. Their Uses.

II. To ascertain the proper Place of the Scalp for making these Issues, the antient Physicians [e] direct to shave the Head, and then to measure with two Threads, one extending from the Nose to the Neck, and the other a-cross the first to each Ear: by which means the Point where the Threads touch, or cross each other, will denote the Place where the coronal and sagittal Suture meet; and is therefore the fittest Place for making your Issue, as you may see in the Figures of Scultetus, *Tab.* XXVI. Meekren, *Obs. Cap.* V. and Deckers, *Exercitat.* The proper Seat of these Issues.

[a] Scultetus, *Obs.* 34. [b] See a remarkable Instance in Meekren, *Obs. Chir. Cb.* V.
[c] See Decker's excellent Method in the Amaurosis and Epilepsy. *Exercit. Pract.* p. 109.
[d] Slevogtius Dissert. of an Issue in the coronal Suture, as a Cure for a decayed Memory.
[e] See Celsus, *Lib.* VII. *Cap.* 7. *N.* 15.

Exercitat. Pract. pag. 110. But after all, it must be owned, that this Method of assigning the Place, or Meeting of the Sutures, cannot be certainly relied upon, because of the great Variation there is in this respect in different People. nor is it of any Consequence whether your Issue be made exactly over the Meeting of the Sutures or not; since the Matter discharged by it in both Cases, comes rather from the external Integuments of the Cranium, than through the Sutures from the Brain, as the Antients falsely imagined. It was also a Notion equally wrong, that the Antients entertained of this Part of the Cranium being thinner, and more perspirable, than the rest: though it must be confessed, that Infants, whose Bones are not completely ossified, have this Part soft and membranous, which is usually termed in them, the *Fontanel*, or open Mold: yet in Adults, this upper Part of the Cranium is almost constantly ossified like the rest of the Skull, and frequently the Bones are even harder or thicker here than in other Parts. But these we find were the groundless Reasons, which induced the Antients to make their Issues upon the Meeting of the Sutures. However if the Surgeon will be scrupulously exact in this respect, he may pretty certainly discover the Meeting of the Sutures, without the forementioned *Apparatus*, of measuring by Threads, if he well considers the Course of them in dry Skulls, and feels carefully with his Finger upon the Scalp and Pericranium: for in most Patients the Meeting of the Sutures is sensible to the Touch, either by a small Cavity or Protuberance; upon either of which you may venture to make your Issue.

The Methods of making Issues in the Scalp.

III. To render Issues in this Part more efficacious, they are usually made by the actualy Cautery: in order to which the upper Part of the Scalp is first to be shaved, and then the red-hot Iron is to be strongly pressed by your Hand, so as to burn through the Integuments upon the Part assigned, till you come to the Bones, or naked Cranium. The cauterizing Iron for this Operation may be of two Kinds: the first of which is without a Case, as we have given you a Figure of it in *Tab.* III. *Fig.* 9. taken from Meekren and Deckers; the other, taken from Aquapendens and Scultetus, is furnished with a steel Case, or directing Tube, as we have represented in *Tab.* XV. *Fig.* 1 and 2. but that the Force of the Cautery may not be extinguished by the Integuments before it has reached the Cranium, it may be convenient for the Surgeon, first to make an Incision either longitudinal[a] or cruciform through the Skin, and opening the Lips of the Wound, insert the Tube at *Fig.* 2. that, by pressing the Cautery, *Fig.* 1. through it, you may at the first Time burn into the very Bone[b]. But in whatever manner you make the Issue in this Part, it must be immediately dressed afterwards with a Pea dipt in some digestive Ointment, to be retained by a square Plaster and the four-headed Bandage, applied as we shall direct in treating of Bandages: or the rest, you may consult what has been before said of Issues in general at *Chap.* XIX. preceding. In order to credit the good Effects which many able Physicians affirm they have experienced from this Sort of Remedy, in many obstinate Disorders of the Head, it must be considered, that though there is no immediate Discharge hereby made of pernicious Humours from the Brain, yet the Cauterization makes so strong a Revulsion, and

[a] See Celsus, *Lib.* VII. *Cap.* 7. *N.* 15.

[b] Thus Meekren, in his Figure, expresses an Incision before the Application of the Cautery; but says nothing of it in the Description.

and the Pain it excites gives ſo ſtrong a Stimulus to the Veſſels, as frequently to remove Obſtructions, and the inveterate Pains they have occaſioned, even in one Inſtant. For more concerning the Uſes of Iſſues in this Part, the Reader may conſult, beſides the forementioned Authors, MARC. DONATUS, *Lib.* II. *Hiſt. Miral. Cap.* 4. M. A. SEVERINUS, *Pyrotech. Chirurg. Lib.* II. *Part.* 1. *Cap.* 6. RIVERIUS, *Cent.* II. *Obſ.* 93. AQUAPENDENS, *Oper. Chirurg. Cap.* 1. CLAUDINI *Reſponſ. de Cauterio in Sutura Coronali*, &c.

CHAP. XXXIX.

Of ARTERIOTOMY *in the* TEMPLES.

I. ARTERIOTOMY, as the Word imports, is the Apertion of an Artery with a ſharp Inſtrument, in order to extract a proper Quantity of Blood, for the Recovery of a Patient; performed almoſt in the ſame Manner as bleeding in a Vein. Though this Operation is not ſo often performed at preſent with us, as it was formerly among the antient Surgeons, for fear of the profuſe Bleeding, or an Aneuriſm, which may be occaſioned by wounding this Veſſel; yet, if it be well adapted to the Patient's Diſorder, and ſkilfully performed, it may be very often of the greateſt Service, and yet not attended with any bad Conſequences. We read of the Apertion of Arteries made by the antient Phyſicians in various Parts of the Body; as in the Forehead, Temples, behind the Ears, in the Occiput, betwixt the Thumb and Fore-finger, &c. where-ever the ſmaller Arteries lie fair for Inciſion, ſo that their Pulſation may be perceived by the Finger through the Skin. But among the modern Phyſicians and Surgeons we hardly ever meet with this Operation performed in any other Parts but the Temporal Arteries, which may be opened by the Lancet without much Difficulty or Danger, as they lie very near the Skin, ſo as generally to be very perceptible to the Touch; and, being reſiſted by the *Os frontis* on which they are incumbent, they may be very eaſily compreſſed, to prevent any profuſe Hæmorrhage, or dangerous Aneuriſm. But even here every prudent Surgeon muſt own, it is much more difficult to make a fair Apertion of an Artery, than of a Vein; becauſe they ſeldom appear viſible through the Skin, and then you have no other Guide but their Vibration on the Finger. We ſhall not here enlarge upon the extraordinary Artifices which we read to have been uſed for Arteriotomy by the antient Surgeons, becauſe they are now obſolete. We ſhall, in this Place therefore, only deſcribe the Operation with its Dreſſings and Uſes, as they at preſent obtain among our modern Surgeons and Phyſicians. Arteriotomy deſcribed.

II. Firſt, the Patient muſt be ſeated conveniently with his Head inclined to either Side againſt the Light, that the Surgeon may the better diſcover the Artery, in order to which, he had beſt place the two foremoſt Fingers of his left Hand upon the Artery, at a little Diſtance from each other, as he will be directed by its Pulſation; and obſerving well the Courſe or Direction of it within that Space, to dip the End of the Lancet carefully into it betwixt his two Fingers: but it will be here neceſſary to inciſe deeper, as the Veſſel lies lower, than in Phlebotomy. You muſt alſo inlarge your Inciſion more, by elevating the The Method of Operating.

the Point of your Lancet as you draw it out: nor need you be afraid of cutting the Artery quite in two; for it will not be attended with any bad Conſequences after Compreſſure and Deligation. If now the Blood follows your Lancet in a very florid and ſalient Stream, ſtarting at every Pulſation of the Veſſel, you may be ſatisfied the Artery is well opened; otherwiſe you muſt repeat your Inciſion, till your Lancet has either divided or opened the Artery, which you may know by the forementioned Signs: but as the ſmall and thin Point of the ordinary Lancet may be eaſily broke off againſt the Bone, I have experienced the Scalpel, *Tab.* I. *Fig.* G. to be more convenient; eſpecially if your Inciſion be downward and not upward. To do the Patient any conſiderable Service by this Evacuation, you ſhould bleed him plentifully; that is, to take about a Pound of Blood, or a Pound and half, or more, if he be plethoric, otherwiſe your Operation will be of little or no Benefit: therefore we need the leſs wonder at the Practice of the Antients, whoſe Method was to bleed the Patient in this Manner till he fainted. If you are deſirous of opening an Artery in the Occiput, or behind the Ears, rather than in the Temples, your Operation may then be conducted in the Manner we have now deſcribed.

Deligation. III. When a ſufficient Quantity of Blood has been taken, your Deligation muſt be made with three ſquare Compreſſes, each larger than the other; laying on the ſmalleſt firſt, in which muſt be included a Farthing, a Bit of Lead, or a Pellet of chewed Paper, to compreſs the wounded Artery againſt the ſubjacent Bone: your other two Compreſſes being laid over the ſmalleſt according to their Size, they muſt be there firmly retained and ſecured by the *Faſcia nodoſa*, which we ſhall deſcribe at large when we come to treat of Bandages at the latter End of our Surgery. The Head thus properly inveſted with your Bandage, muſt continue ſo at leaſt a Week or eight Days before you take it off, to prevent a profuſe Bleeding, or an Aneuriſm: and if the Deligation ſhould, within that Time, get too looſe, it muſt be tightened again, and continued till the Cure is completed.

The Uſes of Arteriotomy. IV. The Uſes of Arteriotomy are ſo many and conſiderable, that not a few Phyſicians recommend it as the laſt Reſource in many Diſeaſes of the Eyes, and the moſt obſtinate Diſorders in the Head, from whence the Patient will often find Relief when all other Means have been tried in vain; eſpecially when they are cauſed by too great a Fulneſs of Blood. Experience can beſt teſtify the good Effects of Arteriotomy in Vertigo's, obſtinate Head-achs[a], Epilepſies, Suffuſions, and Inflammations of the Eyes, and moſt of the other plethoric Symptoms which attack theſe Parts: but particularly in Apoplexies, it has been lately demonſtrated, in a profeſſed Treatiſe on the Subject[b], to be the moſt effectual and expeditious Method of relieving the Patient. I ſhall therefore leave the prudent Reader either to countenance or condemn the Opinion of thoſe who think *Arteriotomy*

[a] LANZONUS confirms the good Effects of Arteriotomy in obſtinate Head-achs, Madneſs, Epilepſies and Diſeaſes of the Eyes, *Ephem. Nat. Cur. Cent.* III. p. 142. It is alſo approved by BARBETTE, and much extolled by SEVERINUS, particularly in the Epilepſy. In a moſt obſtinate Cephalæa I have found it effectual.

[b] By CATHERWOOD, entitled, *A new Method of curing Apoplexies*; notwithſtanding which, the Operation has been twice performed by me on two apoplectic Patients the one an old and the other a young Man, but without the expected Succeſs; for they both died ſoon after, though the Operation was made in the Beginning of the Diſorder, and aſſiſted with other proper Remedies; from whence we ſee, that *Arteriotomy* will not always cure Apoplexies.

teriotomy too dangerous to be put in practice, and even then of no more Uſe than Veneſection; ſince the Uſes and Effects of it are atteſted by the Obſervations and Experience of our beſt Phyſicians, and the Danger of it may be totally removed by proper Compreſſes and Deligation: yet, I muſt own, that, with regard to the Reputation and Character of a young Phyſician or Surgeon, it may be generally adviſeable to defer this for the laſt Help, in Caſes which will admit of Delay. After all, it will be equally neceſſary to aſſiſt this, as well as many other Operations in Surgery, by ordering a proper Diet, Regimen, and Medicines adapted to the Patient's Diſorder, if we expect to make any conſiderable Cure.

CHAP. XL.

Of the HYDROCEPHALUS.

I. HYDROCEPHALUS is a preternatural Diſtention of the Patient's Head to an uncommon Size, by a Stagnation and Extravaſation of the Lymph: which, when collected within-ſide the Bones of the Cranium, the *Hydrocephalus* is then termed *internal*; as it is *external*, when retained betwixt the common Integuments and the Cranium. The firſt Kind of the Diſorder is ſeldom to be met with but in Infants, who contract it whilſt they are in the Womb, or in a difficult Birth. Among others, the Reader may conſult, on this Head, WEDELIUS, *De Morb. Infant.* pag. 47. and RUYSCH, *in Theſaur. Anat.* II. *Tab.* III. which laſt has given a very ample Account of this Diſorder. It generally appears in the Infant whilſt young and new-born: but if it has advanced to any great Degree, it is a dangerous Caſe, and generally incurable. For if you make a Paracenteſis in the Head, to diſcharge the Lymph, your Operation is no ſooner performed, but the Infant dies, as Phyſicians have been too often well aſſured by Experience. If the Diſorder be in its firſt Stage, and but beginning to ſhew itſelf, it will be moſt adviſeable to try what may be done by Medicines; ſuch as gentle and repeated Purges, to draw the Humours downward with corroborating Medicines internally. Externally you may apply to a good Purpoſe a large Compreſs round the Head, dipt in warm *Aq. Calcis & Sp. Lavend. vel Aq. Reg. Hungar.* This Compreſs muſt be retained by a proper Bandage, termed the *Reflex Capeline*, which is deſcribed in the *Third Part* of our Syſtem treating on Bandages. The internal Hydrocephalus.

II. In the external *Hydrocephalus*, as we obſerved, the Humours are lodged betwixt the external Integuments and the Cranium. Hence you may diſtinguiſh this Species by the Softneſs of the Head and Skin externally. But in the internal *Hydrocephalus* the Head feels as hard as uſual, though it is much more diſtended and enlarged: the Reaſon of which Appearances is manifeſt from what we ſaid in the laſt Paragraph. Though the external *Hydrocephalus* is not without Danger, yet it may be much more readily cured than the internal Species; but the more difficultly as it is of a longer ſtanding. The Cure muſt be attempted as well by internal as external Remedies at the ſame Time. Apply Cathartics, Diaphoretics, Diuretics, attenuating and ſtrengthening Medicines for internal Uſe: and externally, a Compreſs dipt in the Fomentation beforementioned for the internal *Hydrocephalus*. Or, you may apply thoſe Waters and The external Hydrocephalus.

Spirits to the Head, together with diſcutient Caps or Bags filled with the Tops of dry *Marjorana*, *Origanum*, *Serpillum*, *Pulegium*, *Chamomilla*, *Salvia*, *Roriſmarina*, *Lavendula*, *&c.* warming them before you ſecure them on the Head by the proper Bandage. HILDANUS writes, that he happily cured an *Hydrocephalus* barely with the repeated Application of *Aqua Calcis* as a Fomentation, by means of a Sponge. To the forementioned Remedies we may add an Errhine, or cephalic Snuff, compoſed *ex ſummit. Marjoranæ*, *Lil.conval. Mari veri*, *Hippocaſtan*, *Nicotiana*, *&c.* Add to theſe the repeated Chewing of Tobacco in the Mouth, to diſcharge the Seroſities from the Head by Spitting. Laſtly, ſome foment the Head with the Fumes of burning Spirit of Wine highly rectified. But if all theſe Means prove unſucceſsful, Recourſe muſt then be had to chirurgical Helps: among which, you ought, firſt, to try a large Bliſter, applied behind the Ears, on the Occiput and Neck: if this does not altogether anſwer your Intention, you may add Scarification and Cupping upon the ſame Parts. PISO relates, that he cured a Man of an *Hydrocephalus*, by making Iſſues in the Neck: and therefore Setons, one of which will effect as much as two Iſſues, may be here alſo highly ſerviceable. When all other Means have been uſed in vain, ſome of the Antients adviſe a deep tranſverſe Inciſion to be made at the Bottom of the Head or Occiput: which I cannot approve of, as it may eaſily wound, or even totally divide, the Blood-veſſels and Muſcles there ſeated. But as this Danger may be avoided by deep Scarification and Cupping upon the ſame Parts, the Diſcharge that way may be equally ſerviceable, and much more commodious. The Parts ſcarified are to be afterwards dreſſed with Lint, ſpread with ſome digeſtive Ointment: adding ſometimes a little *Præcipitatum rubrum*, to keep up the Diſcharge. When the Diſorder is thus removed, you heal it up with ſome vulnerary Balſam, keeping the Patient, for a conſiderable Time, in a Courſe of proper internal Medicines, and under a ſuitable Diet and Regimen. Hiſtories of this Diſorder are given in PAREY, LUSITANUS, KERKRINGIUS, and others: but particularly VESALIUS relates, that he found nine Pounds of Serum in the Ventricles of the Brain, in a Subject who died with an *Hydrocephalus*, *Anat.* Lib. I. Cap. 5.

CHAP. XLI.

Of TREPANNING *the* CRANIUM.

When the Trepan is neceſſary.

I. TREPANNING is univerſally underſtood to be a Perforation, or Opening made in the Bones of the Cranium by a kind of *Terebra*, or round Saw, which has its Name from the *Greek* Word τρίω, and by the *Latins* called *Modiolus*. This Operation was performed by the Antients, not only in Fractures and Depreſſions of the Cranium, but alſo in thoſe other obſtinate Diſorders of the Head and Brain, which could not be relieved by internal Medicines, and the Uſe of Iſſues upon the coronal Suture: by which means they thought to give a more immediate Vent to the offending Humours. But the modern Surgeons never uſe the Trepan at preſent for internal Diſorders of the

Head[a]; though they ſeldom neglect it in Fractures and Depreſſions of the Cranium, cauſed by Blows, Falls, Bullets, and other external Injuries. They alſo frequently apply it in Fractures and Fiſſures of the Cranium, to diſcharge extravaſated Humours, which, by injuring the Brain, might occaſion the Death of the Patient. The Trepan is therefore uſeful, not only in theſe Caſes, to elevate the depreſſed Parts of a fractured Bone in the Cranium, for which you may conſult *Part* I. *Book* I. *Chap.* XIV. but alſo the moſt fatal Symptoms, and Death itſelf are avoided, by diſcharging the extravaſated Blood through an Aperture made by this Inſtrument. It is well known, that the Bones of the Cranium are often fiſſured, and the adjacent Blood-veſſels, lacerated by external Injuries, without any apparent Fracture or Depreſſure of them; ſo that if the extravaſated Blood be not removed by the Trepan, by preſſing on the Brain it will greatly injure, if not totally deſtroy its ſeveral Functions. The Conſequences of neglecting this Inſtrument in ſuch Caſes will be Reſtleſſneſs, Delirium, Convulſions, Vertigo, Apoplexies, Stupidity, with a Loſs of the Senſes, Speech, and voluntary Motion, and at laſt Death itſelf. Sometimes only the milder of theſe Symptoms appear, and in but ſmall Degree, when the Head has been injured by external Violence: but in ſome time afterwards, when the Blood or Humours have been accumulated, the moſt fatal Symptoms do then gradually approach, and even threaten the Life of the Patient. But if Death is not the immediate Conſequence, as there is no natural Vent for the extravaſated Blood or Lymph, it muſt conſequently putrify, and, by corroding the Brain and its Membranes, will inevitably deſtroy the Patient in a little Time, if it be not prevented by a judicious Application of the Trepan, for diſcharging the offending Matter. This Inſtrument therefore ought never to be neglected in urgent Caſes of this Nature.

II. The leſs Time you loſe, the better, before the Application of the Trepan: but in the Operation itſelf you muſt go on ſlowly and carefully. For it is extremely difficult, if not impoſſible, for you to take out a Piece of the Cranium by this Inſtrument, without injuring the ſubjacent *Dura Mater*, to which it is moſt intimately attached, ſo as to be often in ſome Degree wounded, though you uſe the greateſt Circumſpection. For this Reaſon I am induced to condemn the Advice of thoſe[b], as very unſafe, who direct to trepan the Cranium immediately upon every ſlight Diſorder of it. I ſhould therefore adviſe you, with CELSUS, and moſt of the Moderns[c], to try firſt the Uſe of other Remedies, both external and internal, as Phlebotomy, Purging, Clyſters, diſcutient Bags, *&c.* rather than immediately to ſubject the Patient to the Trepan, before you are convinced it is abſolutely neceſſary[d]. But you may ſee more upon this Head in *Part* I. *Book* I. *Chap.* XIV. *Sect.* XXXVI, *& ſeq.* where we treat of Wounds in the Head. On the other hand, there are many Caſes, in which Delay

The Trepan not to be uſed haſtily.

[a] Though indeed there is an Inſtance related in the *Mem. Acad. Pariſ. Chirurg.* where an obſtinate Head-ach was cured by Trepanning, *Tom.* I. p. 226. But there follows another, p. 227. where the Operation failed in a ſimilar Caſe.

[b] See FIENUS *De Trepanatione*, and BOHNIUS *De Trepanationis Difficultatibus*. Likewiſe ROONHUYS, *Obſ.* I. p. 1.

[c] Among which are CÆSAR MAGATUS, Lib. II. *De Vulneribus*, Cap. 41. and DIONIS in *Chirurg. Operat* —CELSUS, Lib, VIII. Cap. 4.

[d] See ſome excellent Obſervations on the Uſe of the Trepan in dangerous Caſes, *Mem. Acad. Chirurg. Gallic.* Tom. I. p. 188.

Delay may be of the moſt fatal Conſequences; where, being convinced of the Inſufficiency of other Remedies, you ought immediately to have recourſe to the Trepan, in order to elevate or remove the depreſſed or fractured Parts of the Cranium, and to diſcharge the Humours extravaſated internally.

The Event of the Operation doubtful.

III. The Surgeon can hardly ever be certain of the Succeſs of this Operation: becauſe he cannot be previouſly aſſured in what Manner or Degree the Brain, and its including Membranes, are injured, the Diſorder generally turning out worſe than its Symptoms indicated: therefore we need the leſs wonder that moſt Patients miſcarry after the Uſe of the Trepan, not from the Operation, but the Violence of their Diſorder, or the Injury received: and ſome there are, who, being much better after the Operation, appear ſeemingly in a fair Way for Recovery, and yet miſcarry contrary to Expectation. Upon Enquiry made after the Cauſes of this unexpected Diſappointment, and ſudden Death of the Patient, they appear chiefly to be two: either from, 1. an Inflammation or Suppuration of the Brain and its Membranes, from the Putrefaction of ſome Blood or Matter that could not be diſcovered or diſcharged; or, 2. from ſome Attack of the Blood on the Parts affected, by Irregularities committed by the Patient in the Non-naturals, either in Drinking, and bad Diet, (or by an unwholeſome Air[a]) or by Frights, Anger, Venery, or other intenſe Paſſions, *&c.*

What Parts of the Cranium may be trepanned, and what not.

IV. But before we proceed to acquaint the young Surgeon with the Method of performing this Operation, it will be previouſly neceſſary to point out to him, upon which Part of the Cranium it may be convenient for him to apply the Trepan. And, in general, the Place where the Fiſſure appears, will be moſt convenient for the Trepan, if nothing contra-indicates: but, in Fractures, it will be proper to trepan a little below the injured Part, that the extravaſated Humours may be more eaſily diſcharged: yet, if the Fragments of the Bone can be removed, ſo as to make Way for the Extraction of the Blood and Splinters which injure the Brain, the Uſe of the Trepan may be in that Caſe neglected. It muſt be next obſerved, that there are ſeveral Places in the Cranium, which ought not to be in any Caſe trepanned: as, 1. upon the Sutures where the Bones meet with each other, eſpecially upon the ſagittal Suture, as HIPPOCRATES has long before obſerved; becauſe, in theſe Parts, the *Dura Mater* is more ſtrongly attached to the Cranium, and under the ſagittal Suture runs the longitudinal Sinus of the *Dura Mater*, which, by trepanning in this Place, might eaſily be injured, to the Hazard of the Patient's Life. Yet, in Caſes of urgent Neceſſity, the Trepan may be uſed upon the coronal Suture, and ſometimes upon others; Inſtances of which may be ſeen in CARPUS, *Lib. de Fract. Cranii*, HILDANUS, *Obſ.* I. *Cent.* 2. 2. It is equally dangerous to trepan the Cranium in the Middle of the *Os Frontis*, eſpecially in that Part which forms the Fontanel: becauſe under theſe is ſeated the forementioned Sinus of the *Dura Mater*, which might eaſily be wounded by the Inſtrument. 3. The Trepan muſt not be applied upon any of the Sinuſes of the *Os Frontis*. 4. nor ought it to be uſed where any large Vein or Artery ſpreads itſelf. 5. If the fractured Part of the Bone, upon which you fix the Trepan, is looſe or carious, you might then injure the Brain by this Inſtrument. 6. It has been judged improper to trepan in

[a] It has been obſerved by the *French* Phyſicians, that Trepanning has failed in many Hoſpitals, on account of the Impurity of the Air. *Mem. Acad. Chirurg.* Tom. I. GUILLEAM. alſo relates, that this Operation generally proved fatal in *Paris*, though ſucceſsful in other Parts of *France*.

in the lower Parts or Baſis of the Cranium, which are inveſted with Muſcles, as about the Occiput and Temples: though the Moderns find that the Trepan may be very well uſed, and even applied upon the lower Parts of the Cranium, and upon the temporal Bones, after the Muſcles have been freed from them[a]. 7. Laſtly, it will be improper to trepan upon the cruciform Eminence of the *Os Occipitale*. Notwithſtanding theſe Rules or Cautions, if a violent Fracture ſhould happen in or near the forementioned Places, you ought to trepan as near to the affected Part as poſſible: and if the Fracture has paſſed a-croſs the Sutures, you muſt trepan within a Finger's Breadth of the Suture on each Side. Sometimes it is impoſſible to diſcover the particular Part of the Cranium, which is injured; the Patient, in the mean time, being afflicted with the moſt urgent and dangerous Symptoms, ſuch as Vomiting, Drowſineſs, Convulſions, Fever, Bleeding at the Noſe and Mouth, with the Loſs of his Senſes and Speech. In theſe Caſes, it will be neceſſary to trepan firſt on the Right Side, then on the Left, afterwards upon the Forehead, and laſtly, uppon the Occiput, and ſo round till you meet with the Seat of the Diſorder: for it is much better, in theſe deſperate Caſes, to try a doubtful Reemedy, than none at all, as CELSUS[b] rightly adviſes, that the Surgeon may not be accuſed of having neglected any thing which might conduce to the Recovery of the Patient. You muſt not think it a new or uncommon Practice to make ſeveral Perforations in the Cranium after one another by the Trepan: for, in many Caſes, we meet with extravaſated Blood or Splinters of the Bone, which require the Uſe of the Trepan in other Parts, beſides where the Wound itſelf manifeſtly appears. Therefore the Operation muſt be repeated, till you can diſcover and remove the Cauſe of the Diſorder: ſo that it is no wonder to meet with three or four, nay ſeven or twelve Perforations in the Cranium, made by the Trepan in the ſame Patient; of which we are furniſhed with many Inſtances, particularly in SCULTETUS, *Obſ.* 7. GLANDORPIUS *Speculum Chirurg. Obſ.* 3. p. 46. to which add DIONIS in his Operations, and many others: but what is more, we read of the Trepan being applied twenty-ſeven different Times with Succeſs upon a Count of NASSAU, in STALPART, VANDER WIEL, *Cent.* 1. *Obſ.* 8.

V. After having pitched upon the Part to be trepanned, your next Buſineſs, is, to ſhave the Scalp, and make an Inciſion through the Integuments, to lay bare the Cranium, except it ſhould have been already done to your Hand by the Wound. The Inciſion of the Integuments may be made in the Form of a Croſs +, or in the Figure of the Letter X, V, or T, large enough to admit the Crown of the Trepan upon the Bone. After your Inciſion is thus made, you muſt elevate and ſeparate the Integuments and Perioſteum from the Cranium by the Edge and Handle of the Scalpel: and having wiped off the Blood, you muſt inſert a large Quantity of ſcraped Lint, to dilate the Wound, and compreſs the divided Veſſels, in order to diminiſh the Hæmorrhage, which indeed, though profuſe, may, in many Patients, be ſerviceable. A Compreſs muſt be next applied, dipped in *Sp. Vin. Aq. Calc.* or *Sp. Vin. Camphorat. calid.* to be retained by the Kerchief Bandage: thus the Patient is to be left, if the Diſorder

Proviſion for the Operation.

[a] See ROUHALT *De Vuln. Cap.* p. 91. & ſeq. and SAVIARD, *Obſ.* 27. p. 136.

[b] In *Lib.* II. *Cap.* 10. To which we may add the Sentence of HIPPOCRATES in *Aphor.* vi. Sect. I. *Deſperate Diſorders require deſperate Remedies.*

order will permit, for a few Hours, that the Blood may be ſtopped before you apply the Trepan: otherwiſe the Work will be ſo much obſcured, that you cannot ſee what you are about: yet if any Delay will be dangerous, you ought to apply the Inſtrument immediately: before which, if the Hæmorrhage be great, you may ſecure the Ends of the divided Arteries by Ligature with a crooked Needle and Thread: but if you are in great haſte, the Hæmorrhage muſt be ſuppreſſed for the preſent by the Fingers of an Aſſiſtant preſſed upon the Part.

Apparatus of Inſtruments and Dreſſings.

VI. We come now to the Apparatus of Inſtruments and Dreſſings, which muſt be provided before you enter upon the Operation. The firſt and principal is the Trepan or Terebra, with its Crown, *Tab.* XV. *Fig.* 3. Some of the Antients uſed a Trepan made in the Shape of a common Gimlet, according to the Figures of FABRICIUS AB AQUAPENDENTE, ANDREAS A CRUCE, and SCULTETUS (*in Officina Chirurg.* pag. 14, & ſeq.) *Tab.* II. *Fig.* 7, &c. which inſtrument they applied with one Hand; from whence it was uſually denominated the Hand Trepan. But, as this Inſtrument labours under many Defects, which renders the Application of it leſs commodious, the Moderns, at preſent, uſe a [a] Trepan like that repreſented in *Tab.* XV. *Fig.* 3. with a Handle turning round, like that uſed by Coopers, which is much more commodious than the antient one; eſpecially if the Crown of it be not made cylindrical, but broader above than below, in the Shape of an inverted Cone, as it is repreſented in *Fig.* 3. A. by which means the Inſtrument, meeting with more Reſiſtance as it deſcends further through the Bone, is not ſo liable to ruſh in upon and wound the Brain. The Inſtrument contrived in this Manner is, by ſome, termed the Trepan of HILDANUS, though it was known and deſcribed by CELSUS[b], and others of the Antients, long before HILDANUS. The Crown of this Inſtrument, marked A, is joined to the lower Part of the Handle B, by a Screw, ſo that it may be taken off and put on at Pleaſure: or elſe, that a Crown of another Size may be ſcrewed in its Place, ſince it will be neceſſary for the Surgeon to be provided with Crowns of different Sizes. The Connection of the Crown, with its Handle, is, by ſome of our modern Surgeons, made in a different Manner from that here repreſented, but with no great Advantage, in my [c] Opinion; ſince that of the Make here repreſented, is found to anſwer moſt Purpoſes conveniently enough. The Trepan is diſtinguiſhed into Male and Female: in the firſt of which the Crown is furniſhed with a ſharp Point or Pyramid A. but when the ſaid Point or Pyramid, *Fig.* 4. is taken out by the Winch, *Fig.* 5. the Trepan is then termed Female. You muſt next be alſo provided with a Scalpel of a particular Make, with a round and flat Head, as repreſented at *Fig.* 6. which is, by ſome, denominated the lenticular Scalpel: to which add another Inſtrument for gradually depreſſing the *Dura Mater*, of the Shape repreſented at *Fig.* 7. You muſt be alſo provided with a perforating Inſtrument, *Fig.* 8. which muſt be ſcrewed into the Cavity B of the Handle *Fig.* 3. alſo a Hair Bruſh, like that repreſented at *Fig.* 9. with a ſmaller Terebra or Wimble, like that in *Tab.* VII. *Fig.* 7. a Lancet, an Elevator, *Tab.* VII. *Fig.* 7, 8, and 14. a Tooth-pick made of a Quill,

[a] See alſo AMB. PAREY, *Lib.* IX. *Cap.* 18. where he gives ſuch a Figure of the Crown of this Inſtrument.

[b] *Lib.* VIII. *Cap.* 3.

[c] *Vid.* GARENGEOT *Tract. de Inſtrument.* Tom. I. pag. 115.

[a] Quill, a Probe with a ſharp Point, ſome Doſſils of Lint: and laſtly a Veſſel with ſome *Spirit. Vin. rect.* all which are to be placed in order in a large Diſh or Plate, that they may be ready to the Surgeon's Hand in performing his Operation. The Apparatus of Dreſſings and Bandage to be applied after the Operation, conſiſts of a Doſſil of Lint of an orbicular Figure, which muſt be tied round the Middle with a Piece of Thread, about a Span long, the Form of which is repreſented in *Tab.* XV. *Fig.* 11. beſides which, there muſt be added another round Bundle of Lint of a convenient Size, ſecured by a Thread like the preceding, as repreſented at *Fig.* 12. You muſt alſo have ſome Pledgits of Lint, *Fig.* 13. for covering the other Dreſſings, and filling up the Cavity in the Cranium. To theſe add ſome *Mel. Roſar. & Tinct. Succin. vel Maſtich.* ſome ſcraped Lint, a ſquare Compreſs: and laſtly, a large Napkin, or ſquare Piece of Linen, to make the Kerchief or Bandage for the Head. All which Particulars are to be diſpoſed in order upon one or two large Plates, that they may be readily found, and handed to the Surgeon as he wants them.

VII. The Apparatus being thus provided, we come next to the Operation itſelf. To perform it with a greater Readineſs and Exactneſs, the Patient muſt be diſpoſed in a convenient Poſture upon a Couch, or ſome other low Seat, in ſuch a Manner, that the Surgeon and Aſſiſtants may have free Acceſs to perform each their Part. This done, and the Dreſſings removed, the Wound is next to be cleanſed from the extravaſated Blood, or other Foulneſs: after which, you place the Head in a convenient Manner upon a Pillow, to be held faſt by an Aſſiſtant. The Surgeon now takes the perforating Trepan, *Fig.* 8. which he adapts to the Handle B, inſtead of the Crown A, *Fig.* 3. ſo that by turning round the Handle D, he makes a ſmall Entrance or Aperture with his Inſtrument, and then applies the male Trepan with a Crown, *Fig.* 3. A. Upon the Top of the Handle C C, the Surgeon fixes his left Hand, upon which he places his Chin or Forehead[a], while with his Right Hand he ſlowly and carefully turns round the Handle, till the Crown of the Trepan, with its Spindle, have made a circular Entrance deep enough in the Cranium: then he removes the Spindle, and continues his Work carefully with the Crown of the Trepan only, as long as he ſees convenient, all the Saw-duſt being firſt bruſhed off from the Cranium and the Teeth of his Inſtrument with Bruſhes of Hog's Briſtles. He now continues to uſe the Trepan till the Saw-duſt becomes Bloody, which denotes, that he has penetrated the Diploë, or intervening ſpongy Part of the Cranium: but it is to be obſerved, that he will not always meet with this Sign, becauſe in ſome Sculls the Diploë is wanting in the Part trepanned. However, when his Saw-duſt becomes bloody, the Inſtrument muſt be directly laid aſide, and, after waſhing away the Blood with a Sponge dipt in *Sp. Vin.* he then ſcrews the ſmall Terebra, *Tab.* VII. *Fig.* 7. B, by two or three Turns, into the ſmall Aperture in the Middle of the trepanned Piece of Bone, and then takes it out again, making two or three more Turns with the Crown of his Trepan. then

The Method of Trepanning.

[a] Moſt Surgeons formerly placed their Forehead upon their left Hand, on the Inſtrument; but it ſeems to be a better Practice to lean the Chin as M. PETIT and GARENGEOT direct; becauſe then the Operator has a better View of his Work.

then he examines with a Probe or Tooth-pick, whether the Plates of the Cranium are ſufficiently ſawed through; which cannot be better known, than by carefully attending to the Colour of the circular Groove or Diviſion: for when that appears of a blue or grey Colour, which was before white, it is a Sign, that you have penetrated ſo far through the lower Plate of the Bone, as to render the *Dura Mater* almoſt conſpicuous through it: the Trepan muſt therefore now be applied with greater Circumſpection, leſt the Saw-Teeth of its Crown ſhould ruſh in upon and wound the *Dura Mater*, which might be attended with violent Inflammation and the moſt malignant Symptoms. But if the bony Plate appears livid in one Part of the circular Groove, and white in another, it is a Sign that the Trepan has not cut equally through; and therefore it muſt be inclined and preſſed a little harder upon the whiteſt Parts, moving round the Handle ſlowly and carefully, till the Saw-Teeth of the Crown have cut deep enough to make the round Piece of Bone looſe or moveable. In that Caſe it will not be convenient to cut totally through the Bone with the Saw-Teeth of the Trepan. To avoid wounding the ſubjacent *Dura Mater*, you ſhould rather ſcrew in the Terebra again, *Tab.* VII. *Fig.* 7. B, or ſome ſuch Inſtrument, till you find that by pulling this upward with the Aſſiſtance of an Elevator, you can totally remove the round Piece of Bone.

The Treatment after Trepanning.

VIII. Having thus extracted the round Piece of the Cranium, the Blood uſually follows it: which being wiped off, the Surgeon is carefully to examine, whether there are any Fragments or rough Parts remaining to be extracted, or Depreſſions to be raiſed: if there are, you muſt do it immediately: if not, you muſt ſmooth the rough Parts about the lower Margin of the Aperture, by applying the headed Scalpel, *Fig.* 6. to prevent the *Dura Mater* from being pricked and injured by any of the ſharp Splinters. This done, the Blood will more readily diſcharge itſelf: but to promote its Exit, you may gently incline the Patient's Head on one Side and another, tenderly and carefully preſſing the *Dura Mater* itſelf, either by the Head of the Scalpel, *Fig.* 6. or the Depreſſor, Fig. 7. by which means the Patient is no ſooner relieved from the Weight or Preſſure of the extravaſated Blood on his Brain, but he inſtantly begins to recover his loſt Senſes, either ſuddenly or by degrees, like one juſt awoke out of a deep Sleep. When the Patient has thus recovered his Senſes, and the Blood notwithſtanding is in ſome meaſure retained, the Surgeon ſhould direct him to fetch a deep Breath, and hold it with a Strain, like one that has a hard Stool. Others rather recommend violent Sneezing, provoked by Sternutatories, in order to force out the extravaſated Blood: the Succeſs of which in my Opinion, muſt be very precarious, if not ſometimes fatal.

When extravaſated Blood, or a bony Fragment or Splinter are lodged under the *Dura Mater*.

IX. If the *Dura Mater* appears diſtended or elevated, and of a blackiſh blue Colour at the trepanned Aperture of the Cranium, it is uſually a Sign that Blood or Matter are retained underneath it. Therefore there remains but one and a doubtful Remedy for it; which is, to make a Perforation through the *Dura Mater* (as alſo the *Pia Mater* when the Matter lies ſo low) with a Lancet or Scalpel, to give Vent to the retained Blood or Matter, which will otherwiſe certainly prove fatal to the Patient, by eroding ſome of the larger Blood-veſſels. I know there are ſome, who think the *Dura* and *Pia Mater* cannot be perforated without deſtroying the Patient, and therefore they forbid

it. But the Succeſs of this Practice, if you avoid the larger Arteries and Veins, is confirmed, not only from my own Experience, but likewiſe the Authorities of [a] PAREY, [b] GLANDORP, [c] COITER, [d] FALLOPIUS, [e] MAGATUS, [f] MARCHETTI, [g] ROHAULT, [h] BLANCARD, and other creditable Writers, who teſtify, that many have had this Operation performed without Danger. If you meet with any bony Fragments or Splinters which irritate and wound the Brain, they muſt be carefully extracted, either by your Fingers or the Pliers: or if any Parts of the Bone are depreſſed only, you muſt raiſe them by your Fingers, a Lever, or an Elevator adapted to the Purpoſe. When a Splinter is inſinuated betwixt the *Dura Mater* and the *Cranium*, ſo that you cannot extract it by the firſt Aperture you made with the Trepan, a ſecond or third Perforation muſt be made by the ſame Inſtrument, till you have removed every thing injurious to the Brain and its Meninges. Sometimes it will be neceſſary to cut off or remove the bony Fragments, by making a ſecond Perforation into the firſt, like a half Moon, by the Trepan when the Fragments are ſtrong, or by the ſmall Saw (*Tab.* VII. *Fig.* 9.) by a Pair of cutting Forceps, or laſtly, by the Mallet and Chiſſel, to be ſeen in the ſaid *Tab.* VII. But when the Fragments are thin and weak, you may remove them by the lenticular Scalpel, *Tab.* XV. *Fig.* 6. that you may afterwards extract or remove the vellicating Splinters. When there is a long Fiſſure in the Cranium, you may trepan upon each End of it: but when the Fiſſure runs in ſeveral Directions, you muſt trepan upon each, becauſe every one of them has uſually extravaſated Blood or Matter lodged underneath [i].

X. Having deſcribed the Method of perforating the Cranium by the Trepan, and of diſcharging the extravaſated Blood, Matter, and bony Fragments, we next proceed to the Dreſſings and Deligation. Theſe are made, firſt, with a round Pledgit of dry Lint, *Fig.* 11. to be laid next the *Dura Mater*, with a Thread faſtened to it, and hanging out of the Aperture, that it may be placed under and drawn out from beneath the Cranium: upon which Pledgit of Lint is afterwards poured ſome *Mel. Roſar.* diluted with a little *Sp. Vini*; though there are ſome who recommend the Application of *Tinct. Maſtich. Succin. &c.* which are, in my Opinion, too ſtrong and acrid, becauſe they often moleſt the Patient with violent Pain. You then lay on a like Pledgit of Lint, furniſhed with a String, as in *Fig.* 12. with other Doſſils, till the Cavity is replete. In the next place, the Cranium and Wound itſelf muſt be dreſſed with Lint, ſpread with ſome mild digeſtive Ointment, or *Mel. Roſar.* upon which add a ſquare Compreſs dipt in warm *Sp. Vini*, or *Sp. Vini Camphorat. cum Aq. Calc.* and then you ſecure the whole, without a Plaſter, by the Capeline or Head-Bandage, deſcribed in the End of our Surgery. Deligation and Dreſſings.

XI. In the ſubſequent Dreſſings, which muſt be repeated once or twice every Day, you muſt ſtrictly avoid fat and oily Applications; which will deſtroy the Membranes and foul the Bones: inſtead of ſuch, you muſt apply balſamic and healing Topics, eſpecially *Mel. Roſar. cum pauco Sp. Vini*, *Tinct. Maſtich. &c.* Of renewing the Dreſſings.

[a] Lib. ix. Cap. 21. [b] *Obſ. Chirurg.* 4. [c] *Obſ. Anatom. & Chirurg.* [d] *De vuln. Capit.* Cap. 48. [e] *De vuln.* Lib. ii. Cap. 42. [f] *Obſ.* 14. [g] *Pag.* 83. 116. [h] *Obſ. Med. Phyſ.* Cent. 1. *Obſ.* 27.

[i] The *French* Surgeons have deſcribed, and illuſtrated with Obſervations, a great Variety of Caſes, where a frequent Repetition of the Trepan was requiſite. *Mem. Acad. Reg. Chirurg.* Tom. I. p. 251 *& ſeq.*

The Wound being thus conſtantly dreſſed and attended, you will have an Exfoliation of a thin Plate from the trepanned Margin of the Bones, uſually within forty or fifty Days, which ought not to be pulled away by Force. Your Exfoliation being obtained, there will then appear new Fleſh and Callus, ſhooting up from the clean Bone and *Dura Mater*, ſo as at length to fill up the whole Cavity. By that Time you find the Cavity about half filled, you muſt moderately compreſs the ſprouting Fleſh and Callus by ſcraped Lint and Bandage, to prevent it from being too ſoft and lax: and when it is arrived even with the Surface of the Bones of the Cranium, you muſt endeavour to conjoin and extend the Integuments over it, by the Aſſiſtance of Sticking-Plaſters; that the new-formed Subſtance may intimately unite with the ſuperinduced Skin. This new-formed Subſtance, with which the Cavity in the Cranium is filled, becomes gradually more and more indurated: but ſo as even at laſt to reſemble rather a Cartilage than a Bone, which, upon boiling the Cranium, ſeparates, and falls out from the other bones: and it is from the weaker Reſiſtance of theſe cartilaginous Places that ſuch as have been trepanned are ſubject to Diſorders and Pains in their Heads, upon a Change made in the Weight and Temper of the Atmoſphere: though that Inconvenience may be partly remedied by conſtantly keeping the Place armed with a Plate of Silver.

The Removal of Accidents.

XII. If a Vein ſhould open itſelf ſo as to bleed profuſely after the Operation has been performed with the Trepan, then you muſt ſprinkle on ſome *Pulv. ex Bolo Armeno, Sang. Dracon. Thure & Colophon. &c.* compreſſing the Part for ſome time with Lint: but if the Brain or *Dura Mater* ſhould be inflamed, you muſt apply diſcutient and cooling Topics externally, *Aq. Flor. Samb. cum pauc. Gutt. Sp. Nitri dulc.* the Patient muſt alſo uſe Abſtinence, with Phlebotomy, and cooling diluent Medicines internally. Even ſome (as ROHAULT, p. 123.) recommend Scarification of the *Dura Mater* itſelf, before the laſt preſcribed Mixture is applied: but, if a Suppuration ſhould follow, ſo as exactly to form an Exulceration, the Surgeon muſt cleanſe away the Matter, or Sordes, with ſcraped Lint, or by an Injection mixed with *Mel. Roſ. Sp. Vini & Tinct. Maſtich. Succin. vel Elix. prop. ſine alcali vel acido.* If, after the Patient has been once trepanned, he perceives great Uneaſineſs and Diſorder in ſome other Part of the Head, it is a Sign there ſtill remains ſome foreign Body to be removed: therefore the Trepan muſt be again applied upon the aſſigned Place. If any ſpongy Excreſcence, or proud Fleſh ſhould riſe up above the Level of the Wound upon the Cranium, it may be removed by ſome of the following Methods; either by ſtrong Depreſſion with Lint dipt in *Sp. Vin. vel Tinct. Maſtich.* and a tight Bandage, or by applying the round Piece of Lead, *Fig.* 14. contrived by BELLOSTE [a], which is, by ſome, made perforated, and furniſhed with Handles, as at *Fig.* 15. which is to be put into the Aperture of the Cranium, and well covered with round Pledgits of Lint: but you will ſeldom have Occaſion for this Inſtrument, if the firſt Method be uſed. Or, laſtly, if the Excreſcence has already ſurmounted the Surface of the Cranium, it may be cut off, either by tying it round with a Thread, or with a Pair of Sciſſars, and the reſt may be taken down with *Vitriol. Cærul. Pulv. Sabin. vel Alum. uſt.* and for the future you muſt make a ſtricter Compreſſure and Deligation with more compact

[a] See his Treatiſe, entitled, *Le Chirurgien d'Hoſpital.*

pact Doſſils of Lint: by which means the ſprouting Excreſcence will be not only compreſſed and reduced, but the Wound itſelf will readily heal in a little Time. We have an Account of the Trepan being repeated to the eighth Time for a Caries in the Cranium, *Mem. Chirurg.* Tom. I. p. 262. There is another, of a ſucceſsful Operation on the coronal Suture, and the Repetition of it, p. 255. but in p. 244. it is adviſed to omit it. LE DRAN alſo has many curious Obſervations on Trepanning. As to the Abuſe of it, conſult ROONHUYS in his Obſervations.

CHAP. XLII.

Of Extracting Bodies fallen into the EYES.

Things to be extracted from the Eyes.

I. IT is no uncommon Thing for the Eyes to be moleſted with a Bit of Glaſs or Sand, a Splinter of Wood, or from off a Quill, or the Toe or Finger-Nails, and ſometimes by little Inſects, or cauſtic and pricking Bodies of various Kinds; which, by ſlipping into this tender Organ, we daily experience will produce excruciating Pain and Inflammation. To remove theſe, and prevent their bad Conſequences, the Surgeon's Aid is often required; whoſe chief Buſineſs is, to diſcharge the foreign Body as ſoon as poſſible, by ſome of the Means we ſhall hereafter preſcribe.

Methods of Expulſion.

II. The firſt and moſt eaſy Method of diſcharging theſe Subſtances, is, by agitating and extending the Eye-lid with one's Fingers, holding the Head down at the ſame time; by which means the increaſed Flux of Tears, excited by the vellicating Body, very often waſhes the ſame out of the Eye, without much Difficulty. But if this Method does not ſucceed, the next Remedy is, to blow ſome levigated Pearl or Crabs-claws through a Quill under the Eye-lid; that, as theſe are waſhed out by the Tears, they may alſo take away the foreign Body with them. Otherwiſe the Surgeon muſt take the ſmall round Head of a ſlender Probe, or a little Pair of Pliers, the End of a Tooth-pick, *&c.* and extending the Eye-lids gently from the Eye, carefully ſearch for, and tenderly extract the offending Body. There ſtill remains a very eaſy and certain Method for removing theſe injurious Subſtances from the Eyes, by dipping a Pencil-bruſh of ſoft Feathers, or a Bit of fine Sponge faſtened in a Quill, in warm Water, by which you may bruſh them out from betwixt the Eye and its Lid. Lime, or any acrid Salt, and ſuch like Subſtances, may be waſhed out by warm Water, or Milk, either by injecting them, or with a Feather or Bit of Sponge. When the foreign Body is removed, the Surgeon muſt furniſh his Patient with a cooling anodyne Collyrium *ex Aq. Roſar. Damaſc. cum albumine ovi conquaſſata, & pauxillo Saccbar. Saturni, vel Lap. Tutiæ præparat.* with which the Eye is to be frequently waſhed; not neglecting to bleed the Patient at the ſame time, if there be any conſiderable Inflammation.

CHAP. XLIII.

Of TUBERCLES *and* EXCRESCENCES *on the* EYE-LIDS.

Kinds. I. THE preternatural Tubercles, which we frequently meet with upon the Eye-lids, are of various Sorts and Sizes. If the Tubercle be ſmall, hard, red, immoveable, and ſeated upon the Eye-lid above the *Cilia*, or Range of Hairs, it is then denominated by the *Greeks*, *Crithe*, and by the *Latins*, *Hordeolum*, from its ſuppoſed Reſemblance to a Barley-corn. This Tumor is included in a Kind of Cyſt, which, by Inflammation, degenerates into a thickiſh Matter: from whence frequently proceed intenſe Pains and various other Diſorders of the Sight. The Seat of the *Hordeolum* varies, being ſometimes immediately next to the Skin, and ſometimes within-ſide the Eye-lid, under its Muſcle. When the Tubercle is moveable, 'tis uſually denominated *Chalazium*, or a Stithe. Some are termed *Grandines*, as being like Hail: others are named *Hydatides*, being Veſicles replete with watery Humour. Sometimes ſeveral Species of the encyſted Tumors are formed upon the Eye-lids, as the *Atheroma*, *Steatoma*, and *Meliceris*; of which we have already treated in *Chap.* XXVIII. preceding. It may be here obſerved once for all, that almoſt all the Tubercles on the Eye-lids are of the encyſted Kind, ſome having a ſmall depending Baſis, and others a broad one, as may be ſeen in *Tab.* XV. *Fig.* 16, 17, 18.

Prognoſis. II. We are, from the Importance and Obviouſneſs of this Organ, obliged to undertake the Cure and Removal of many of theſe Tubercles: which, in other Parts of the Body, might be very well neglected. Yet we ought not, even here, to call in the Aſſiſtance of the Knife, when they are very ſmall, and not troubleſome to the Sight; for they are often tolerable without Danger, though they may perhaps give a little Deformity. 'Tis remarkable, that theſe Tubercles ſeldom give way to topical Remedies: nor ſhould you be over forward with the Uſe of emollient Cataplaſms, which are recommended by ſome; becauſe the Eye itſelf may be injured by them, and therefore Extirpation is to be preferred.

Cure. III. Almoſt all Tubercles of the Eye-lids, which do not hang pendulous by a ſmall Root, are removed by making an Inciſion through the Integuments by the Scalpel, ſo as to avoid wounding the Tumor, in order to take it clean out, as we before directed for encyſted Tumors in *Chap.* XXVIII. foregoing. But if the Coats of the Tumor are wounded, or adhere very firmly to the adjacent Fleſh, ſo that it cannot well be extirpated whole by the Scalpel, it may be cut out as far as you well can by a Pair of ſmall Sciſſars; and the Remainder eroded and caſt off by dreſſing with *Ægyptiacum*, or ſome other digeſtive Ointment, mixed with *Præcipitat. rub. vel Lap. infernal.* after which you may complete the Cure with Balſams, as in other Wounds. In ſome Caſes, when I think the Tumor cannot be totally extirpated, I make an Inciſion through its including Cyſt, together with the common Integuments; and, after expelling or diſcharging its Contents, deſtroy the reſt with Digeſtives and Cauſtics, as I directed for encyſted Tumors: but here you muſt be very careful to prevent any of the Cauſtic from falling into the Eye, which might greatly injure, if not

deſtroy

deſtroy its Sight. But we are furniſhed with a much more ready and eaſy Way of removing thoſe Tubercles of the Eye-lids, which hang pendulous by a ſmall Root, as at *Fig.* 17 and 18, which is, either to cut them off inſtantly by a Pair of Sciſſars, or elſe gradually by a Ligature with a Silk Thread. But another Method muſt be taken with the *Hordeolum*; becauſe that, contrary to moſt encyſted Tumors, is uſually attended with Pain and Inflammation. Therefore in theſe laſt, it will be proper, firſt, to try to diſperſe them by diſcutient Applications; and if that will not ſucceed, to bring them to Suppuration before they are inciſed. It will greatly conduce to diſperſe and eaſe the Pain of an incipient *Hordeolum*, if the Patient frequently foments it with his faſting Saliva, or elſe with a Mucilage *ex Sem. Cydonior.* or the Pulp of a roaſted Apple mixed with a little Saffron and Camphire. If none of theſe ſucceed, but the Tumor holds on its Inflammation, and begins to turn yellow, you may ripen and break it with a *Diachylon* Plaſter, or a Mixture of Honey and Meal. But the Cure of it will be ſooner completed, if you invert the Eye-lid, by Inciſion with a Scalpel a-croſs the Tumor, ſo as to ſeparate the Skin of the Eye-lid, and extract the Cyſt entire, if it be hard; otherwiſe you may open the Cyſt, and diſcharge its included Matter, and deſtroy the Remainder by Digeſtives; by which means you will avoid an unſightly Scar in the Eye-lid, and the Wound itſelf will heal without the Application of other Medicines.

CHAP. XLIV.

Of WARTS *on the* EYE-LIDS.

I. THE Eye-lids are frequently moleſted as well with Warts as the foremen-tioned Tumors, which often both obſtruct the Sight, and disfigure the Eye; for which Reaſons the Patient is deſirous of their Removal. Theſe Warts adhere to the Eye-lids, either by a broad or ſlender Baſis; and may be extirpated either by the Knife, Ligature, or Cauſtics, in the Manner we directed for Warts in general, in *Chap.* XXVI. preceding. You muſt never apply the actual Cautery to deſtroy theſe Warts, as you may for thoſe in other Parts of the Body: nor ſhould you apply Cauſtics but with the greateſt Circumſpection; leſt, if any Part ſhould ſlip into the Eye, it might greatly injure, or deſtroy the Patient's Sight[a]. If a Wart on the Eye-lid appears blackiſh, or livid, you will generally have Reaſon to fear its turning cancerous, as it will do, eſpecially if irritated with Inſtruments or Medicines: for this Reaſon, theſe are uſually termed, *Noli me tangere*, by the moſt expert Oculiſts; ſo that it is beſt to leave this Species of Warts to themſelves. I happily removed a large Wart from the upper Eye-lid by Ligature, which had no broad Root, but impeded the opening of the Eye-lids; the Figure of which Wart you may ſee in *Tab.* XV. *Fig.* 17. A.

CHAP.

[a] Thus TIMÆUS à GULDENKLE, *Lib.* I. *de Affect. Capit. Cap.* XXI. relates the Caſe of a Surgeon, who blinded a Woman by endeavouring to remove a Wart from her Eye-lid by the cauſtic Juice of Spurge.

CHAP. XLV.

Of Relaxation and Tumor of the EYE-LIDS, *termed* Phalangoſis *and* Ptoſis.

Nature of the Diſorder.

I. WE frequently meet with the Eye-lids either tumified, or relaxed to ſuch a Degree, as greatly deforms the Eye, and impedes its Viſion. Sometimes the relaxed Eye-lid ſubſides in the Manner repreſented by *Fig.* 19. *Tab.* XV. occaſioned either from a Palſy of the Muſcles, which ſuſtain and elevate the Eye-lids, or from a Relaxation of the *Cutis* above, from various Cauſes. Sometimes an œdematous or aqueous Tumor is formed on the Eye-lids, ſo as almoſt entirely to exclude Viſion: which laſt Caſe ſhould be well diſtinguiſhed from the former, and may be remedied, without much Difficulty, by the Uſe of internal and topical Medicines. Such are Purges with Diuretics and Sudorifics inwardly; and a Compreſs dipped in warm *Sp. Vin. Camph. & Aq. Calc.* externally. But, in the paralytic or relaxed Caſe, you muſt uſe nervous and cardiac Medicines, and apply a little *Balſ. Peruv. cum Aq. Reg. Hungar.* &c. If theſe Medicines ſhould all miſcarry, the beſt and moſt expeditious Method is, to extirpate a ſufficient Quantity of the relaxed *Cutis*; and, after healing up the Wound, the Remainder may become ſufficiently ſhortened.

The antient Method of Cure.

II. The Antients contracted the Skin thus relaxed, by extirpating Part of it with the Aſſiſtance of a Ligature with a Needle and Thread. Having firſt carefully ſecured it by Ligature, by paſſing the Needle through the Bottom of the Skin, they then cut it off cloſe to the Ligature: which, in many Caſes ſucceeded very well. Sometimes they firſt amputated Part of the relaxed Skin by the Sciſſars or Scalpel, and then ſecured the Wound, either by Ligature or Suture, with a Needle and Thread, as we read in HIPPOCRATES, *Lib. de Vict. acut. Sect.* 66.) CELSUS *Lib.* VII. *Cap.* 7. *N.* 8.) and PAULUS ÆGINETA *Lib.* VI. *Cap.* 8. But the Hæmorrhage frequently proves ſo large in this laſt Method, as to obſcure the Wound, and render it impoſſible to make a neat Suture, or Ligature: to avoid which Inconvenience, the famous *German* Oculiſt, BARTISCHIUS, formerly contrived a wooden Inſtrument, *Tab.* XV. *Fig.* 19. B B. to intercept the redundant Part of the *Cutis*, and compreſſing it by turning the Screw D D, ſo as to obſtruct the Blood-veſſels, and hinder the Circulation, the intercepted Part mortified in a few Day's time, and caſt itſelf off.

The modern Treatment.

III. But as the laſt mentioned Practice of BARTISCHIUS was attended with great Pain, Inflammation, and other Inconveniences; VERDUYN has much improved upon him, by making almoſt a ſimilar Inſtrument of Braſs, but with Perforations in its upper and lower Plates, as in *Tab.* XV. *Fig.* 21. By which Inſtrument the redundant *Cutis* is not only compreſſed, but alſo ſecured with a Ligature, by paſſing a Needle and Thread through the Apertures, and leaving about four or five Inches of the Thread hanging down on each Side; you then amputate the redundant Skin, cloſe to the Edge of the Inſtrument, with a Scalpel, or Pair of Sciſſars: after which you remove the Inſtrument, and make a Ligature with the Threads. Having performed your Operation, the Wound is, for the firſt Time, to be dreſſed with ſome vulnerary Balſam and ſcraped Lint: but, in the ſubſequent Dreſſings, you may ſpread your Lint with ſome

digeſtive Ointment, to be retained with Compreſs and Bandage, as we directed in other Wounds of this Part. After a few Days, when the Lips of the Wound appear to be pretty well cloſed or conjoined, you may then cut the Ligature, and carefully extract the Threads, removing them, not all at once, but one at a Time, in each Dreſſing, and compleating the Cure with ſome vulnerary Balſam and Plaſter. You may cauterize the Wound before the Removal of the Inſtrument, which will not only ſuppreſs the Hæmorrhage, and render the Diſorder leſs liable to return again, but may perhaps, at the ſame time, ſave you the Trouble of making a Ligature or Suture. Sometimes this Diſorder is ſo great, as to deſtroy the Figure of the Eye, or ſo obſtinate and inveterate as to return again, after a repeated Performance of the Operation; which renders the Caſe incurable. Laſtly, we may obſerve, that RAW invented an Inſtrument, not much differing from the former in its Make and Uſes; (ſee *Fig.* 22.) but you may ſee the original Invention of this Inſtrument highly controverted between him and RUYSCH[a], who rather attributes it to ADRIANSONIUS.

CHAP. XLVI.

Of the Trichiaſis, *or Inverſion of the* EYE-LIDS; *in which the Hairs irritate the Eyes.*

I. THE *Cilia*, or Margins of the Eye-lids are ſometimes inverted, ſo as greatly to irritate the ſenſible Coats of the Eye, and bring on intenſe Pains and Inflammation; which, without timely Aſſiſtance, may greatly injure, if not totally deſtroy the Sight. This Diſorder is, by the *Greeks*, termed *Trichiaſis*, or *Diſtrichiaſis*, *hairy*; and ſometimes *Entropion*, *Inverſion*, becauſe herein the Lids and their *Cilia*, or Hairs, are inverted, ſo as to offend the Eye. The Diſorder is generally occaſioned from an irregular Cicatrix formed from a Burn, the Small-Pox, an Ulceration, or Wound from ſome external Injury. Sometimes a Relaxation of the Skin, and a paralytic Diſorder of the Eye-lids, deſcribed in the preceding Chapter, make one of the chief Cauſes of a *Trichiaſis*, Nor is the Cure of a *Trichiaſis* to be effected without much Difficulty, eſpecially when the Diſorder is become inveterate. Cauſe of the Diſorder.

II. 'Tis hardly poſſible for the Surgeon to remove this Diſorder, ſo as to prevent its returning, without extirpating the offending Hairs: which every one muſt allow to be no eaſy Operation, that has ſeen anything of the Diſorder. for if you cut the Hairs cloſe off, it will be to no Purpoſe, becauſe the rigid and ſharp-pointed Stumps of the Hairs will ſhoot up and irritate the Eye worſe than the Hairs did before. Some indeed endeavour to cure the Diſorder, without extirpating the Hairs, by clearing them out from the Eye, and keeping them folded back, or paſted on the Outſide of the upper and lower Eye-lids by ſome ſticking Plaſter. But this Practice is not often attended with the deſired Effect; becauſe the Motion of the Eye-lids looſens the Hairs, and they become again inverted, ſo as to offend the Eyes, as before. In this Caſe therefore the Practice of ſome is conformable to the Advice of CELSUS (*Lib.* VII. *Cap.* 7. *N.* 8.) who directs to burn out the Roots of the Hairs, one by one, with a ſlender, Method of Cure.

[a] See RUYSCH *Epiſt. Anat.* XIII. and RAVIUS *in Tract. de Septo Scroti.*

ſlender, but broad-pointed Needle of Steel, in the Shape of a *Spatula*, heated red-hot. But ÆGINETA (*Lib.* VI. *Cap.* 13.) directs to extract each Hair firſt with a Pair of Pliers, before the Cauterization of their Roots; which is an Operation ſo painful, that the Patient will hardly ſubmit to it. Therefore ſome chuſe to fill up the Cavities at the Roots of the Hairs, after their Extraction with *Lap. infernal.* or ſome other Cauſtic, taking great Care that no Part of it ſlips into the Eye. Or it will be better to touch their Cavities with a ſmall Pencil-bruſh dipt in *Sp. Salis Ammoniaci cum Sp. Vini rectificatiſſ.* by which means they will cicatrize and cloſe up, without producing any more Hairs. When there are many injurious Hairs to be thus extracted; it will be better to remove them at ſeveral Times, than all at once; otherwiſe you may induce too great Pain and Inflammation on the Eye, whoſe *Cornea* ſhould be alſo defended from the Cauſtic or Cautery here uſed by a ſmooth hollow Plate of Lead, Wax, or Horn, adapted in the ſame Manner as for artificial Eyes. If the Diſorder ſhould ariſe from a Relaxation of the Eye-lids, it will be neceſſary to treat it in the ſame Manner we directed in the preceding Chapter.

Other Methods. III. But if all the Hairs of the Eye-lids are thus inverted, and the Patient will not permit them to be extracted by the Roots, and to be afterwards treated with Cauſtics; there then remains but one, and a lamentable Method of removing the Diſorder, by amputating the *Cilia*, or cartilaginous Margins of the Eye-lids themſelves; which the Patient had better ſubmit to, notwithſtanding the Deformity it may occaſion, rather than be blind. After the Operation, a *Collyrium* ſhould be made, and applied *ex Aq. Roſar. Alb. Ovor. & pauc. Sacchari Saturni, vel Aquâ & Sp. Vini ana*; and the Wound muſt be treated in the ſubſequent Dreſſings with ſome Balſam till it be healed. But lately CORTUMIUS, in a profeſſed Diſſertation de *Trichiaſi*, under Profeſſor GOELICKE, 1724, has propoſed a new Method of removing the *Cilia*, rather by Cauſtics with *Lap. infernal.* than by Amputation. When the Patient is laid on his Back, he directs firſt to arm and defend the Eye with Lint or Leather; and then to rub the *Cilia* with ſtrong *Lapis infernalis*, till the cartilaginous Margins of the Eye-lids with their Hairs, are eroded and removed; after which you are to dreſs firſt with dry Lint, and then with a *Collyrium ex Aq. Roſar. & Alb. Ovor.* to be often renewed. The next Day you muſt remove the Lint, or leathern Defenſative from the Eye, to avoid an Inflammation from it; and, if any ſmall Eſchar ſhould be formed underneath the ſame, it may be removed by ſome digeſtive Ointment: by which means, if you clear the Eye well from the Lint, he aſſerts that the Wound will be cured generally within the Space of ſix or eight Days.

CHAP. XLVII.

Of the Anchyloblepharon, *or* CONCRETION *of the* EYE-LIDS.

Deſcription. I. THE Diſeaſe termed *Anchyloblepharon*, is when the Eye-lids cohere, or grow to each other, or to the Eye itſelf. It is eaſily diſtinguiſhable from the glewing up of the Eye-lids in the Small-Pox and Inflammations, by an Inſpiſſation of the Juices and glutinous Matter, by which they are ſtrongly

faſtened together for ſome Time, but without intimately concreting, becauſe they ſeparate again ſpontaneouſly in a little Time afterwards.

II. Sometimes the Eye-lids cohere, ſo that they cannot be opened, to admit the Light for Viſion, either in one or both of the Eyes, as in *Tab.* XV. *Fig.* 23. AA. Sometimes again the Eye-lids grow to the Globe of the Eye itſelf, either to its *Tunica cornea*, *albuginea*, or both: which Accidents generally ariſe from violent *Ophthalmias*, Burns with Gunpowder or other Fire, the Small-Pox, cauſtic Remedies, or an Ulceration of the Parts from many other Cauſes. 'Tis true, this Diſorder is ſometimes born with the Infant; and may ſometimes ariſe in Adults from a fleſhy Excreſcence in the Angles of the Eyes growing to the Eye-lids, as I had once an Inſtance myſelf. See *Miſcell. Nat. Cur. Dec.* II. *Ann.* 8. *pag.* 135. Cauſes.

III. The Cure of all the ſeveral Species of this Diſorder is, in ſome meaſure, both doubtful and dangerous, but of none more than that in which the Eye-lids are conjoined to the *Cornea*; for in that Caſe it will hardly be poſſible to free them without blinding, or at leaſt injuring the Patient's Sight. Nor is there leſs Difficulty to free the Eye-lids from each other, when they cohere from a Burn. Therefore in all Burns and Ulcerations of the Eye-lids, great Care ſhould be taken to treat them with emollient and cooling Topics, and to keep them free from Adheſions, to which all inflamed and excoriated Parts are extremely ſubject. When the Eye-lids grow together in the Small-Pox, they generally adhere at the ſame time to the *Cornea*, from whence they cannot eaſily be ſeparated without injuring the Sight; for, after the adhering Parts have been freed from each other with the greateſt Judgment and Caution, there are almoſt conſtantly ſome little Scars or Specks left upon the *Cornea*, which greatly impede the Sight for the future, and which it will be almoſt impoſſible to remove. Prognoſt.

IV. From what has been ſaid concerning the Nature of the Diſorder, you will readily conclude, that the Cure muſt conſiſt in a ſkilful Separation of the conjoined Parts. In order to which, the Patient is firſt to be placed on a Bed or Chair againſt the Light, in the moſt convenient Poſition for the Operator; who is firſt to examine whether the Eyelids are totally conjoined, or whether there may not be ſome ſmall Interſtice left, which you will generally meet with in the greater or internal *Canthus* of the Eye next the Noſe. If the Eye-lids are ſtrictly conjoined in every Part, you may then begin to make your Diviſion in either of the *Canthi*, or Angles, which appears to be moſt convenient; but with a ſoft Hand, and great Circumſpection, to avoid wounding the *Cornea*, or Eye itſelf. When you have made a ſmall Aperture, a Pair of Sciſſars, or Scalpel, with a blunt Point, are to be introduced, with which (*Tab.* XV. *Fig.* 25.) you gradually and carefully divide the Lids from each other. But if there is naturally left a ſmall Aperture betwixt the Eye-lids, where they do not adhere, you may then immediately introduce one of the forementioned obtuſe-pointed Inſtruments, and proceed to make your Inciſion; or, if you have none that are obtuſe-pointed, introduce a ſmall grooved Director, *Tab.* XV. *Fig.* 24. and then you may ſafely divide with the common Sort of Sciſſars, Scalpel, or a Lancet. Cure.

V. When the Eye-lids have been carefully ſeparated from each other, you muſt then examine with a Probe, whether they adhere to the Eye itſelf; if they do, you muſt again free them cautiouſly with an obtuſe-pointed Scalpel or Lan- Adheſions of the Globe.

cet: but when the whole Globe, or the greater Part of the Eye is firmly attached to the Lids, the Operation is both difficult and dangerous; as it will be almoſt impoſſible to free the *Cornea* without injuring the Sight: which Accident may be avoided and the Cure more eaſily obtained, when the Lids adhere only to the *albuginea Tunica* of the Eye. Even Wounds of the laſt mentioned Tunic are of ſo little Conſequence, that I would always chuſe rather to cut off Part of that in dividing them, than to leave Part of the internal Membrane of the Eye-lid adhering to it; for the internal Membrane of the Eye-lids cannot be amputated without inducing great Injuries on the lacrymal Gland and Duct: therefore it is highly neceſſary, for this Operation to be performed by an expert and ſteady Hand.

Treatment after the Operation.

VI. When the Lids have been freed from the Globe of the Eye, the next Buſineſs is, to prevent them from joining again; which they will certainly do, if not prevented by interpoſing ſome Lint, or a thin Plate of Lead, Wax, Leather or a Bit of Gold-beater's Skin, cut in the Shape of a Half Moon, and moiſtened with *Ol. Amygd. dulc.* Either of theſe are to be left ſeveral Days in the Eye, till there is no Danger of future Adheſions; and if they ſhould fall, or be taken out, they muſt be again replaced in a ſhort Time. If the Patient cannot bear the Interpoſition of the forementioned Plates, as is ſometimes the Caſe, he muſt then frequently agitate and work round his Eye-lid, at Intervals, after having uſed a Collyrium *ex Aq. Plantag. Lap. Tutiæ pp. & Sacc. Saturni*, or a Powder prepared *ex Saccharo, Margaritis & Lap. Cancror.* And, laſtly, the Surgeon himſelf muſt ſometimes paſs the obtuſe End of a Probe betwixt the Lids and the Globe of the Eye, to free and keep them from Adheſions.

Adheſions in the Small-Pox.

VII. When the Eye-lids are glued together by a gummoſe and inſpiſſated Matter in the Small-Pox, and Inflammations of that Organ, ſo that they cannot eaſily be opened; they ſhould never be forcibly pulled aſunder, but firſt moiſtened a conſiderable Time with warm Milk, or other emollient Topics: by which means the Patient will generally be able to open the Eye himſelf ſoon after.

CHAP. XLVIII.

Of the EVERSION *and* GAPING *of the* EYE-LIDS, *termed* Ectropium *and* Lagopthalmia.

Origin of the Diſorder.

I. WHEN the Eye-lids are everted or retracted, ſo as to ſhew their internal or red Surface, and cannot ſufficiently cover the Eye, the Diſorder is then denominated *Ectropium* and *Everſio Palpebrarum*, by the *Greeks* and *Latins*. When the upper Eye-lid only is thus diſordered, it is then denominated *Lagopthalmus*, *Oculus leporinus*, or Hare eyed. Some indeed will have the *Lagophthalmia* a Retraction of the upper Eye-lid without any Everſion, ſo that it cannot cover the Eye; which Accident does alſo happen to the lower Eye-lid, as I have often obſerved, without any Everſion, though it is not mentioned by others as a Species of the *Ectropium*. Sometimes this is a ſimple, or original Diſorder; and ſometimes only a Symptom or Conſequence of another, as an Inflammation, Sarcoma, Tumor, *&c.* When the Diſorder is ſimple, or original, it generally ariſes from a Contraction of the Skin of the Eye-lid by the

Scar of a Wound, Ulcer, Burn, &c. or from an Induration and Contraction of the Skin, after an Inflammation; and sometimes it may proceed in a great measure, from the Use of astringent *Collyria*, injudiciously applied in Disorders of the Eyes.

Cure by Medicines.

II. The Cure of this Disorder consists in relaxing and elongating the external Skin of the Eye-lid, so as to cover the Eye; which is often no easy Task to perform, especially when the Disorder is become inveterate. When the Disorder is recent, it will be best to try the Application of Emollients; such as the Vapours of hot Milk or Water, Oil of Almonds, or Olives, Mucilage of Quince-Seeds, Hare's Fat, *Ung. Dialthææ*, &c. to be continued for several Days on the Scar or contracted Skin of the Eye-lid; which must be often extended either upwards or downwards, according as the Disorder is either in the upper or lower Lid. When the Patient goes to Bed, it will be proper to bring the Eye-lids close to each other, and to restrain them so by Plaster, Compress, and Bandage, to be repeated or renewed every Night. But if none of these Means take effect, you must then have recourse to the Operation, when you judge the Case curable; which is performed in the following Manner.

Cure by the Operation.

III. First, you make a semilunar Incision in the external Skin of the Eye-lid, next its Tarsus, or cartilaginous Margin, making the Angles of the Incision downward in the upper Lid, and upward in the lower Lid (as in *Tab.* XV. *Fig.* 26. AA.) that, by this means, the Skin may be elongated. If the Skin does not appear to be let out enough by one Incision, you must make two or three more, running parallel with the first, and about the Distance of a small Packthread from each other. When the Eye-lid is thus sufficiently elongated, you must dress the Wound first with dry Lint stuffed into the Incisions; and then with Lint, moistened with some vulnerary Unguent, which will both prevent the old Skin from uniting again, and at the same time cause new Flesh to sprout up in the Incisions, to elongate the Skin. Lastly, to forward the Extension and Cure, a Piece of sticking Plaster should be fastened to the Margin of the Eye-lid, to keep it extended either up or down; which Method is to be continued till the Eye-lids will shut close.

When the Disorder is from an Inflammation or a Sarcoma.

IV. When the Disorder arises from an Inflammation, or fleshy Excrescence within-side the Lid; you must, in that Case, first remove the Inflammation by the Remedies we have elsewhere described for that Purpose; and then, after arming the Eye with a defensative Plate, remove the Excrescence by *Lapis infernalis*[a]: and thus, by removing the Impediments, the Eye will recover its former Action. When the Disorder proceeds from an *Encanthis*, *Hypersarcosis*, or *Sarcoma*, as in *Fig.* 27, 28, 29, *Tab.* XV. you may remove it by the Directions we shall presently give in the two following Chapters.

When the Disorder is incurable.

V. When the Skin of the Eye-lid has continued violently distorted or contracted from the Patient's Birth, there is seldom any Hope of curing it: and it is still more impossible to obtain a Cure, when the lower Eye-lid is everted through a Weakness of the orbicular Muscle in old People, without any Appearance of a Scar; in which Case the Operation will be to no Purpose. If any good can be done, it will be most likely by corroborating and spirituous Medi-

[a] See IVESIUS on this Subject, *Lib. de Morb. Ocul.* See likewise ROONHUYS, *Obs.* 18. & BIDLOO *Exercit.* p. 153.

cines both external and internal: but in general, this Diſorder is always the more obſtinate and difficult to cure, as it is more inveterate, or of longer Standing. We have a learned Diſſertation *de Eſtropio* by KECKIUS, *ſub Præſidio* ZELLERI, *Tubing. An.* 1733.

CHAP. XLIX.

Of the ENCANTHIS.

I. WE ſometimes meet with a Tubercle, formed in the greater or internal *Canthus* of the Eye, growing out either from the *Caruncula lachrymalis*, or from the adjacent red Skin; which Tumor is ſometimes large enough not only to obſtruct the *Puncta lachrymalia*, but alſo Part of the Sight, or *Pupillæ* of the Eye itſelf[a]. In this Diſorder the Tears continually run down the Cheek, which greatly deforms the Eye and Face, and gives Riſe to an *Opthalmia*. See *Tab.* XV. *Fig.* 27. A. This Tubercle, denominated *Encanthis* by the *Greeks*, is of two Kinds; the mildeſt of which is that without Hardneſs and Pain: but the moſt obſtinate and malignant Species is livid, and very painful, tending, in ſome meaſure, to a cancerous Nature.

Treatment of the firſt Species.

II. In the Beginning of the mild Species of the *Encanthis*, it will be highly uſeful to ſcarify firſt, and then to apply ſome mild eſcharotic or cauſtic Medicine; of which the moſt innocent is a Powder of *Saccbar. Canarienſ. & Vitriol. alb. aut Alum. uſt.* in the Proportion of five Parts of the firſt to one of either of the laſt. A little of this Powder being carefully ſprinkled upon the Tumor, is afterwards to be waſhed out of the Eye with warm Water. If this proves inſufficient, you may ſometimes touch the Tubercle with *Lapis infernalis*, but with great Caution. But to turn off the Humours from the Eyes, and prevent a Relapſe of the Diſorder, you muſt have recourſe to Iſſues or Setons, with Phlebotomy, and cooling Purges. If you find, that the Application of the Medicines takes no effect, or if the Tubercle is of the malignant Species, you then draw it out either with a Hook, *Tab.* XV. *Fig.* 30, 31. or a Pair of Pliers, or elſe, when it is very large, with a Needle and Thread paſſed through it, and tied together like a Sling for a Handle; by which you muſt gradually and carefully extend and draw up the Tubercle, in order to avoid wounding the Eye itſelf, or the lachrymal Caruncle, which would be attended with very bad Conſequences: for as the lachrymal Caruncle in the greater *Canthus* of the Eye, ſtops and prevents the Tears from overflowing, and running down upon the Cheek, if you was to cut off Part from it, the Conſequence would be a watery Eye, or conſtant Flux of Tears over the Cheek. It is therefore rather better to leave Part of the morbid Tubercle, then cut off any Part of the lachrymal Caruncle; becauſe any Remains of the firſt may be afterwards cleared away, by degrees, with Eſcharotics, if you cannot take it off with a Pair of Sciſſars. After an Extirpation of the Tubercle, you muſt apply deterging and healing Medicines, or a Collyrium *ex Lap. Tutiæ, Myrrhæ, &c.* till the Wound is healed.

Treatment of a malignant Encanthis.

III. In a malignant Encanthis, inclining to be cancerous, being hard, livid, and very painful, 'tis generally better to let it alone, and to mitigate its Uneaſineſs with cooling and lenient Collyria, then to exaſperate it by the Operation,

[a] See a Figure of a large *Encanthis* in PURMANNUS's *Chirurgia Curioſa*, pag. 134.





Plate 15.


Fig. 1.
B
A
Fig. 2.
C
C
Fig. 3.
D
Fig. 4.
B
A
E
Fig. 5.
Fig. 6.
Fig. 7.
Fig. 8.
B
A
Fig. 9.
B
B
A
Fig. 10
Fig. 11.
Fig. 12.
Fig. 13.
Fig. 14.
Fig. 15.
Fig. 16.
A
B
D
C
D
Fig. 23.
A
Fig. 26.
A
C

Vol. I. P. 397.

Fig. 17.



Fig. 18.







Fig. 19.








Fig. 23.







Fig. 26.



Fig. 27.



Fig. 28.



Fig. 29.








Plate 15.

Vol. I. P. 397.



tion, or by eſchoratic Medicines; otherwiſe you may perhaps bring on Symptoms worſe than the original Diſeaſe, as is frequently done in cancerous Diſorders by improper Treatment. We have an extraordinary Cure of this Diſorder related by PURMANNUS in his *Chirurgia Curioſa*; in which, after having extirpated the very large Tubercle by Ligature, he applied an actual Cautery to its Root with Succeſs.

CHAP. L.

Of the Sarcoma *and* Hyperſarcoſis, *or* Excreſcence *formed betwixt the* EYE *and its* LIDS.

I. RELATED to the foregoing Diſorder are thoſe Tubercles, or fleſhy Excreſcences, on the inner Surface of the Eye-lids, termed by the *Greeks*, *Sarcomata* and *Hyperſarcoſes*; (See *Tab.* XV. *Fig.* 28, 29.) which, in the Beginning, as uſually very ſmall; but by degrees advance to a conſiderable Bulk. Some of them are ſmooth and even-ſurfaced; and ſome again are rough and unequal, like the Raſberry or Mulberry; of which Excreſcences I have ſeen and cured ſeveral. Deſcription.

II. I generally remove theſe Tubercles, firſt, by carefully extracting them with a ſmall Hook, *Tab.* XV. *Fig.* 30, 31. and then cutting down to the Root with a Pair of ſmall Sciſſars. After letting it bleed a while, I order the Patient frequently to waſh his Eye with a Collyrium *ex Lap. Tutiæ, Aloeæ & Sacc. Saturn. in Aq. Roſ. ſolut.* till the Wound is healed. Inſtead of a Hook you may alſo extend the Tubercle, by paſſing a Needle and Thread through it. Some endeavour to remove theſe Tubercles by Eſcharotics, and *Lap. infernalis*; but I think Inciſion to be much ſafer, as well as more expeditious, and leſs painful. Cure.

An EXPLANATION *of the* FIFTEENTH PLATE.

Fig. 1. Is an Iron Cautery to make Iſſues in the Head: A the Handle, B the Cautery.

Fig. 2. A denotes the Cannula to receive and direct the Cautery, *Fig.* 1.

Fig. 3. The *Trepan* which I uſe. A denotes its *Crown*, B the Place where the Crown is ſcrewed on. CC the upper Part of the Handle, upon which the Hand is laid in the Operation. D the Arch of the Handle by which the Inſtrument is moved round; E a Spike in the Crown. The Moderns have a Method of faſtening the Crown on the Trepan, otherwiſe than by ſcrewing; but this is my Way.

Fig. 4. Repreſents the Spike taken out of the Crown.

Fig. 5. Is the Key or Winch, by which the Spike is taken hold of and ſcrewed into the Crown.

Fig. 6. A lenticular Scalpel, with which the rough Edge of the Bone is ſmoothed after the Uſe of the Trepan.

Fig. 7. Is a Steel Inſtrument, commonly called a *Depreſſor*, with a flat Button at its End, to preſs down the *Dura Mater*, and diſcharge the latent Blood. The ſame Inſtrument is alſo by ſome termed *Meningophylax*.

Fig. 8. Is a Kind of *Terebra* to faſten to the Handle at B, *Fig.* 3. after having taken off the Crown, being uſed to make the firſt Entrance for the Spike of the Trepan, and to perforate Bones in the *Spina Ventoſa*; whence it is alſo ſometimes named the *perforating Trepan*; A denotes its Point, B the Screw to faſten to the Handle.

Fig. 9. Is a Hair-bruſh to cleanſe the Teeth in the Crown of the Trepan.

Fig. 10. Is the exfoliating Trepan, which is ſometimes uſed to pare away a carious Part in a Bone. A its Point; BB the Wings which ſcrape the Bone, when the Inſtrument is turned round.

Fig. 11. A Doſſil of Lint, furniſhed with a Thread, for dreſſing the trepanned Cranium.

Fig. 12. A Pledgit, or round Compreſs of ſcraped Lint ſecured with a Thread.

Fig. 13. Is another Pledgit of Lint without a Thread, to fill the Aperture of the Cranium.

Fig. 14. Is the leaden Plate of BELLOSTE, to defend the Aperture and Dreſſings.

Fig. 15. Denotes the Shape in which the ſaid Plate is to be firſt bent.

Fig. 16. A denotes an encyſted Tumor, or *Atheroma*, in the upper Eye-lid; and B is another in the lower Eye-lid.

Fig. 17. A large flat Wart on the upper-lid, having a ſlender Root, ſo as to fit it for Removal by Ligature with a Piece of Silk.

Fig. 18. Is a *Sarcoma* or Excreſcence on the Outſide of the Eye-lid, with a ſmall Root.

Fig. 19. Repreſents the *Phalangoſis* and *Ptoſis*, or Tumor, and Relaxation of the Eye-lids. A denotes the Diſorder in the left Eye: BB an Inſtrument contrived by BARTISCHIUS, adapted to remove this Diſorder in the right Eye: DD a Screw by which the two Arms of the Inſtrument are approximated, or brought together.

Fig. 20. Is an Inſtrument like the firſt, but improved by VERDUYN, and as it is figured by RUYSCH, *in Epiſt. Anat.* XIII. AA and BB denote the two Arms of the Inſtruments without any Perforations, to remove various Tubercles by approximating them by the Screw CC, and moving the Hinge D, by which they are connected.

Fig. 21. Denotes the ſame Inſtrument of VERDUYN, only a little larger, and perforated with many ſmall Holes *aaaaa*, to make a Suture for this Diſorder of the Eyes.

Fig. 22. Is an Inſtrument for the ſame Uſe corrected by RAW, and taken from his *Epiſt. de Septo Scroti*, being made more crooked, and ſhutting differently. A the Manner of paſſing the Needle through its Apertures: B the Thread drawn through to conjoin the Wound of the Eye-lid.

Fig. 23 Exhibits an Eye with the *Ancyloblepharon*, or Concretion of the Eye-lids, marked AA.

Fig. 24. Is a ſmall grooved Director, ſometimes uſeful to divide Concretions of the Eye-lids.

Fig. 25.

Fig. 25. A ſmall Scalpel with an obtuſe Point, uſed in ſeveral Diſorders of the Eyes.

Fig. 26. Repreſents the Manner of inciſing the lower Eye-lid in the *Ectropium*, or *Lagophthalmia*, or Everſion and Retraction of the Eye-lids.

Fig. 27. Repreſents an *Encanthis*, or Excreſence in the Corner of the Eye near the Noſe.

Fig. 28 and 29. Denote a *Sarcoma* and *Hyperſarcoſis*, or fleſhy Excreſcence within-ſide the Eye-lid; that marked A belonging to the lower Eye-lid, and that at B to the upper Lid.

Fig. 30. Repreſents a ſmall Hook, for elevating and extending thoſe Tubercles, to extirpate them; the crooked Point of which may be made either ſingle or double, as you may ſee by removing the Gripe B in *Fig.* 31. where CC denote the two Prongs, D D the Handle.

CHAP. LI.

Of BLEEDING *in the* EYES.

I. THOUGH Blood-letting in the Eyes has been, a few Years ago advanced by the *Engliſh* Oculiſt Mr. WOOLHOUSE, as an Invention of his own; yet it manifeſtly appears, from various Treatiſes, that the Operation was both known, deſcribed, and practiſed above an hundred Years before among the *German* Phyſicians[a]. This Operation is cried up by Mr. WOOLHOUSE, as of greater Conſequence than any other Diſcovery in Phyſic; he even thinks it preferable to the celebrated Philoſopher's Stone[b]. Not a new Diſcovery.

II. Blood-letting may be ſucceſsfully uſed in the Eyes: 1. Whenever thoſe Organs are inflamed; that is, when the Blood-veſſels, ſpent on the White of the Eye, appear much larger and more numerous than uſual; wherein it will often ſucceed, when other Medicines, and even Phlebotomy, have been tried without their due Effects, and when the Inflammation runs to ſuch a Height as to endanger the Sight. 2. It may be uſed to Advantage when the *Cornea* is infeſted with Specks or Abceſſes; for, after dividing the Veſſels which ſupply the Diſorder, it may be much more eaſily removed. 3. It may be uſed when a red Coat or Film grows upon the Eye; for the oftener the Veſſels are inciſed, which nouriſh the Film, the ſooner it will ſhrink, and diſappear. Laſtly, 4. it may be uſed by way of Prevention, when the foreſaid Diſorders have been removed, and threaten a Return, by the Intumeſcence of the Veſſels in the White of the Eye; in which Caſe you therefore ought to inciſe the turgid Veins, and foment them. In what Caſes uſeful.

III. There

[a] See MAUCHART, *in Diſſert. de Opthalmoxyſi*, pag. 18. FELIX PLATERUS *Prax. Med.* 8. Lib. i. *Tit. de Vitiis Læſ.* 1609. pag. 280. *& 4to Baſil.* 1656. pag. 238. He is again cited on this Head by M. A. SEVERINUS, *in Medicina Efficaci, Anno* 1682. *edit.* pag. 50. Cap. x. which treats of letting Blood in the Eyes.

[b] See the *Diſſertations ſcavantes & critiques de M.* WOOLHOUSE, pag. 310. and *Diſſert. Opthalm.* pag. 224.

Method of Operating.

III. There are ſeveral Ways of performing this Operation, of which we ſhall here only relate the chief. Firſt, the Patient is to be ſeated conveniently on the Bed-ſide, or on a Chair, with his Head held in a proper Poſture by an Aſſiſtant: which done, the Surgeon makes a tranſverſe Inciſion with a Lancet upon the turgid ſmall Veins in the Corners of the Eye, ſo as to open them, or cut them quite aſunder. Some uſe a ſmall Pair of Sciſſars, inſtead of a Lancet, to divide the Veſſels; but, in uſing either of them, the Eye-lids muſt be held apart from each other by the Fingers of one Hand, while the Veſſels are inciſed by thoſe of the other. Some, again, elevate the ſmall turgid Veins with a crooked Needle before they divide them, the Eye-lids being, in the mean time, held aſunder by an Aſſiſtant[a]. But it would be ſtill better to have theſe crooked Needles made thin and double-edged, ſo that they may divide the Veſſels, of themſelves in the Elevation, without the Uſe of Lancet or Sciſſars. Laſtly, there is no material Objection, why this Operation may not be almoſt as advantageouſly performed by the ſcarifying Inſtrument we ſhall deſcribe in the following Chapter.

What is to be done after the Operation.

IV. The ſmall Veins being thus inciſed or divided, their Diſcharge of Blood ſhould be promoted by Fomentations of warm Water, or a Decoction *ex Euphraſia Hyſſop. Veronica, &c.* frequently applied to the Eye by means of a Sponge, or ſoft Linen Rags: for this Operation will be more ſerviceable, as the Diſcharge procured is more copious. But if once performing it does not ſuffice to remove the Swelling and Inflammation, it may be ſafely repeated two or three Times more; aſſiſting it in the mean time with the Uſe of a proper Regimen, Diet, and Medicines, both external and internal. I muſt indeed confeſs, that after having performed this Operation myſelf, on ſeveral Patients, firſt at *Altorf*, and ſince at *Helmſtadt* in *Germany*, I could not poſſibly prevail on them to have it repeated, and it was with the greateſt Difficulty that they were perſuaded to it at all; ſome being deterred from it by Fear of loſing their Eye-ſight, and others upon the Account of the great Pain which it muſt neceſſarily inflict on this tender Organ. The Reaſon of its being ſeldom performed on Infants, is the Difficulty of perſuading them to hold their Head and Eyes ſteady; and the Danger of applying a Lancet, or other ſharp Inſtrument, when thoſe Parts are in Agitation, is very apparent to every one.

Another Method.

V. To this Operation is related that by Inciſion, propoſed in a Diſſertation under CAMERARIUS at *Tubingen*, *Ann.* 1734. for a venereal Ophthalmia: in the moſt violent Symptoms of which Diſorder it is adviſed to make a circular Inciſion in the White of the Eye round the *Cornea*, to diſcharge the ſtagnant Blood, or other Matter diſtending that Membrane, and obſtructing its Veſſels. But whether this is a ſafe and uſeful Practice, or whether it may not be uſed with Succeſs in other violent Ophthalmias, as well as the venereal, can be only aſcertained by the beſt of Teachers, Time, and Experience.

CHAP.

[a] This is the Method preferred by M. ST. YVES, in Lib. *De Morb. Oculor.* pag. 195.

CHAP. LII.

Of SCARIFYING *the* EYES.

I. SCARIFICATION of the Eyes agrees, in many Reſpects, ſo much with the Bleeding of them, deſcribed in the laſt Chapter, that it is no great Wonder Mr. WOOLHOUSE, though a famous Oculiſt, ſhould confound them one with the other[a]. But I think there is a manifeſt Difference, at leaſt enough for any one to diſtinguiſh betwixt them, becauſe the Parts are different; for the interior Surface of the Eye-lids is here the Subject of Scarification, as well as the White of the Eye, to which the foregoing Operation is confined: and then again they are each of them performed by different Inſtruments, as will preſently appear. Coincides much with the laſt Operation.

II. That Scarification of the Eyes is no modern Invention, is apparent from its having been deſcribed and performed by HIPPOCRATES[b], CELSUS[c], ÆGINETA[d], and others among the antient Phyſicians. But there are ſeveral Reaſons to be offered for its having come into Diſuſe with the Phyſicians of the ſucceeding Ages. It might be owing partly to its ſeeming a difficult, dangerous, and very painful Operation, and partly from their judging it to be of little or no Efficacy, as we find by many of their Writings. However, the firſt that revived the Practice among the Moderns, after it had lain neglected for ſo many Ages, was the celebrated *Engliſh* Oculiſt Mr. WOOLHOUSE. Not a new Operation.

III. To ſcarify the Patient's Eye, he muſt be firſt ſeated on his Chair or Bed in an advantageous Poſture againſt the Light, with his Head ſecured from moving by an Aſſiſtant: after which the Operator preſſes his Thumb and Fore-finger on the Eye-lids, ſo as to elevate, or open, and turn them outward, that their interior red Surface may come into View; which may be done with moſt Eaſe in the lower Eye-lid. He now takes his ſcarifying Inſtrument in the other Hand, and rubs it backward and forward with great Swiftneſs upon the internal Surface of the Lid, and upon the White of the Eye itſelf if he thinks proper, and ſometimes even upon the *Cornea*, moving from one Corner of the Eye to the other, ſo as to lacerate the ſmall turgid Veins, and make them bleed plentifully. But this in general is an Operation much ſooner learned from Inſpection, than a verbal Deſcription. Method of Operating.

IV. The Diſcharge of Blood from the ſcarified Veſſels ſhould be promoted as much as poſſible by the Applications propoſed for that Uſe in the preceding Chapter, at *Sect.* IV. which will alſo cleanſe the Eye, and abate its Inflammation at the ſame Time. But, in order to prevent the ſcarified Parts from adhering to each other, they ſhould not be bound up, at leaſt in the Day-time, but the Lids ought to be frequently agitated by the Patient: and if they are bound up at Night, you ought, firſt, to interpoſe a Bit of Gold-beaters Skin, or ſome ſuch Subſtance, to keep them aſunder. Mr. WOOLHOUSE recommends the Interpoſition of three or four Seeds of Clary for this Purpoſe, or rather a Bit of Gold-beater's Skin anointed with ſome Eye-ſalve: for without ſome ſuch Precaution, you will hardly avoid Concretion or Adheſion of the Parts ſcarified. Treatment after the Operation.

[a] See MAUCHART *De Ophthalmoxyſi*, pag. 17. [b] Lib. *De Viſione*. [c] Lib. VI. Cap. 6. N. 26.
[d] Lib. III. Cap. 22. *De Trachomate*.

How long the Scarification muſt be continued, or how often repeated, will belong to the prudent Phyſician to determine, from the particular Circumſtances of the Caſe: but in the mean time it will be highly neceſſary to call in the Aſſiſtance of a proper Regimen, Diet, and Exhibition of both external and internal Medicines; for, by neglecting theſe Helps, your Operation may not only prove ineffectual, but perhaps induce a worſe Diſorder on the Eye. Conſult PLATNERUS's Diſſertation *De Scarificatione Oculorum*, pag. 36, *& ſeq.*

The Inſtruments to be uſed,

V. The Inſtruments uſed by different Authors for this Operation, are various. HIPPOCRATES ſeems to have uſed a ſort of prickly Thiſtle, like the *Atractylis*[a]. Some of the antient Phyſicians ſcarified with a ſmall Steel Raſp in the ſhape of a Spoon: See *Tab.* XVI. *Fig.* 5. with which they rubbed the internal Surface of the Eye-lid till it bled, as we read in CELSUS (*Lib.* VI. *Cap.* 6. N° 26. and ÆGINETA *Lib.* III. *Cap.* 22.) the firſt of which Authors calls it *Specillum aſperatum*, and the laſt *Blepharoxyſton*. Others uſe the rough Plant, named by Botaniſts *Equiſetum magis nudum*, which ſeems to be very well adapted to the Intention. Others again recommend the Pumice-ſtone, *Os Sepiæ*, *&c.*

The lateſt and beſt Inſtrument,

VI. But the lateſt and beſt Inſtrument for this Operation is found to be the Beards of Barley or Rye, which are furniſhed with Rows of ſmall Teeth or Hooks denoted by A in *Fig.* 3. *Tab.* XVI. Ten, twelve, or fifteen of theſe Beards are to be cut and tied together by a String, ſo as to reſemble a Sort of Bruſh for Clothes, as in *Tab.* XVI. *Fig.* 4. the Teeth of each Beard or Spike being turned outward all round, their ſlender Ends form a Sort of Handle A, to be held and worked round and acroſs by the Fingers, to ſcarify the Inſide of the Eye-lids, and the Eye itſelf with the Part B. Hence this Scarification of the Eyes is, by the modern Surgeons, not improperly called *Ophthalmoxyſis* or *Blepharoxyſis.*

Its inventor.

VII. The firſt Contriver of this Bruſh for the Eyes appears to be Mr. WOOLHOUSE, the Oculiſt: who, though he preached up the great Uſes of his Inſtrument to his Pupils, yet ſtudiouſly endeavoured to conceal it, and its Application, from them; till in 1726, M. MAUCHART (preſent Profeſſor at *Tubingen*, and Archiater to the Duke of WIRTEMBERG) his *quondam* Pupil, publiſhed both his Inſtrument and its Uſes, with the Method of applying it. About two Years afterwards, the celebrated PLATNERUS of *Leipſic* explained the whole Buſineſs more at large, in a Treatiſe *De Scarificatione Oculorum*; in which we have the Figure of the Eye-bruſh uſed by Mr. WOOLHOUSE, as you find it repreſented by me in *Tab.* XVI. *Fig.* 4.

Uſes of the Eye-bruſh.

VIII. This Eye-bruſh, or Scarificator, is ſaid by the Author, Mr. WOOLHOUSE to be very uſeful in all Diſorders of the Eyes which require Bleeding: as when the ſmall Veſſels are obſtructed, and the whole Eye inflamed, whether from external or internal Cauſes, as a Blow, Wound, Cataract, Pterygium, Hypopyon, Staphiloma, or the like. In all which Caſes, the internal Surface of the Eye-lids ſhould be chiefly ſcarified, in order to diſcharge the heſitating Blood: and, if I may credit Mr. WOOLHOUSE, this Practice is more effectual in removing Inflammations, induced by external Cauſes, or a Chirurgical Operation, than in

[a] See MAUCHART, lib. c. pag. 6, *& ſeq.* PLATNER, l. c. pag. 25.

in original Ophthalmias or Inflammations of the Eyes: but in the *Chemoſis*, or moſt violent Inflammation of this Organ, it will be neceſſary to ſcarify the Eye itſelf with this Bruſh, as well as the internal Surface of its Lids. 2. He aſſigns the Uſe of his Bruſh to be for the Removal of the Pterygium, Abceſſes, and white or other coloured Specks and Films on the Eye: for, by ſcarifying the *Tunica albuginea* of the Eye, and ſometimes the *Cornea* itſelf, or rather the *Pterygium* upon the *Cornea*, the Veſſels which ſupply thoſe Impediments and Blemiſhes of the Sight are lacerated, and, with the Uſe of other Medicines, deſtroyed; and conſequently, they muſt, in a little Time, dwindle and diſappear. 3. He judges his Inſtrument highly ſerviceable in ſtrengthening and recovering a weak or impaired Sight; or even to remove an *Amauroſis*, or Cataract, which are not of any long ſtanding: for, by the ſtrong Stimulus of this Operation, the ſtagnant Humours are put into Motion, the obſtructed or compreſſed Nerves and Blood-veſſels are again opened, and rendered pervious, and the Eye, by that Means, reſtored to its priſtine Vigour. 4. The *Ophthalmoxyſis*, or bruſhing up of the Eye, is very ſerviceable for the Cure of an *Atrophe*, or *Tabes* of that Organ; as it occaſions a greater Influx of Juices to the Parts, which are therefore ſupplied with more Nouriſhment. 5. This Operation may contribute to the Cure of an *Hypopyon*, or *Hypohæma*, that is, a Collection of Blood or Matter under the *Cornea*, occaſioned by ſome Blow, or other external Violence, which muſt be diſperſed, in order to clear the Sight. 6. This is no deſpicable Remedy for eaſing and removing intenſe Pains, termed by the Antients *Ophthalmoponia*, and when the Light itſelf is intolerable to them: for this being an internal Inflammation of the Eye, cauſed by an Obſtruction and Diſtention of the Veſſels near the *Retina*, the Blood diſcharged by ſcarifying with this Bruſh muſt certainly draw off what is ſuperfluous, and greatly eaſe this ſenſible Part. And laſtly, 7. The Bruſh will be often found very uſeful and neceſſary in Palſies, incipient Mortifications, and many other Diſorders of the Eye-lids, as well as of the Eyes themſelves. See PLATNERUS *De Scarificatione Oculorum*, pag. 37, *& ſeq.*

When Scarification is improper.

IX. But it is not to be imagined this Inſtrument will be uſeful in all Diſorders of the Eyes indiſcriminately, as PLATNERUS, WOOLHOUSE's Pupil, obſerves. For it will be improper, 1. in a dry *Lippitudo*, or *Xerophthalmia*, where the Eye is hot, dry, itches, and the Patient cannot look at the Light without great Pain. It will be alſo equally improper, 2. in Diſorders of the Eyes from a *Venereal* or *Scorbutic* Cauſe; for, unleſs the Vices of the Juices be firſt corrected and removed, as this Operation augments their Influx upon the Parts, it may increaſe, rather than relieve the Diſorder. Nor will it be to any Purpoſe to try the Bruſh, 3. in an old *Cataract*, *Gutta ſerena*, or *Hypopyon*, where the Diſorder is become fixed and incorrigible by Length of Time. And, laſtly, you muſt not expect it to cure, 4. an *Ectropium*, *Trichiaſis*, *Anchyloſis*, and many other Diſorders of the Eye-lids, for which it is not deſigned.

Concerning the Bruſh.

X. With regard to the Eye-bruſh before deſcribed, it is to be obſerved, that a ſmall Force will blunt it, and therefore it cannot well be uſed more than once: a new Bruſh muſt be provided againſt every Operation. 'Tis to be likewiſe obſerved, that the Beards of old Barley are not ſo proper as thoſe of new, which is not altogether full ripe: becauſe the firſt, being very brittle, will be apt to ſhatter, and leave ſome of its Teeth behind in the Coats of the Eye,

which may be followed with bad Conſequences. For the ſame Reaſons alſo it ſhould not be the Product of too rich a Soil, nor have paſſed under the Action of the Flail in thraſhing the Grain.

My Opinion of the Operation.

XI. After all, I muſt confeſs, that, upon Trial, I never could experience any great Effects from this Operation, which I have frequently performed in moſt Diſorders of the Eyes: and, what is more, I have known many Patients afflicted with various Diſorders of the Eyes, which have been reported by WOOLHOUSE, and his Pupils, to be cured by this Practice, when the only real Advantage they received from it, was the Abatement of their Pain; which I take notice of thus openly, leſt it might be imagined, I did not ſucceed for want of operating as I ought, in the Manner of Mr. WOOLHOUSE. I muſt indeed own, that it makes an uſeful Evacuation in *Ophthalmias*, and that I have often experienced its good Effects in many inflammatory Diſorders of the Eyes, eſpecially when aſſiſted with Phlebotomy and Bliſters: and thus I make no doubt but its Author and his Followers may have cured many Diſeaſes of the Eyes. But it may, in general, be queſtioned, whether thoſe Diſorders would not have gone off as readily by Bleeding, Purging, Bliſters, and Scarification in other Parts, as by this Practice; at leaſt the Difference will hardly countervail the extraordinary Pain it gives. We know, that Diſorders of the Eyes were very well cured before the Diſcovery of this Practice by Mr. WOOLHOUSE, and may perhaps be better removed at preſent by ſome, who are ignorant of his *Apparatus*: at leaſt this I may venture to ſay, that if, with Difficulty and much Perſuaſion, you draw in the Patient to ſubmit once to ſo rough an Operation upon ſo tender an Organ, you will not find it practicable to allure him to it a ſecond Time. Children in particular, who yet are more ſubject to Diſorders of the Eyes than Adults, are ſcarce ever prevailed on to undergo the Operation; and female Patients are extremely averſe to it. Nor ſhould I inſiſt upon the ill Conſequences attending the Teeth of the Inſtrument's being left ſticking behind in the Coats of the Eye, and the wounding of the *Cornea*, *&c.* from the intenſe Pain obliging the Patient to move his Head and Eye, which may cauſe an Inflammation even worſe than the original. Even the moſt prudent Oculiſts are obliged to own, that the Practice is beſet with many Inconveniencies in the very Diſorders to which it is moſt adapted; nor can we meet with Examples enough of its good Effects to over-balance the Danger and excruciating Pain that attends it. I would therefore adviſe the young Surgeon not to be over-fond of his new Eye-bruſh, nor bring it into his Practice but in Caſes of the laſt Neceſſity, when all other Means are ineffectual. It is alſo remarkable, that among the modern *French* Surgeons and Oculiſts, none take any Notice of this Practice but ST. YVES, notwithſtanding it made ſo much Noiſe at firſt. In general, the *French* Surgeons are very ſcanty and defective in treating on Diſorders of the Eyes.

CHAP.

CHAP. LIII.

Of the EPIPHORA, *or* WATERY EYE.

I. THE *Epiphora*, or watery Eye, is a Diſorder, in which the Tears being obſtructed from paſſing through the lacrymal Ducts into the Noſe, are forced to run down over the Cheek with Deformity and Uneaſineſs to the Patient. There are ſome indeed who confound this Diſorder with the *Fiſtula lacrymalis*; but unjuſtly; becauſe in the laſt the Tears are not ſincere, but mixed with a purulent Matter flowing from an Ulcer in the lacrymal Sac. But, that the Nature of both theſe Diſorders may be the better underſtood, it will be proper to give you an Idea of the Courſe and Figure of the lacrymal Ducts, as you will find them repreſented in *Tab.* XVI. *Fig.* 6. where *a a* denote the *Puncta lacrymalia* in the Eye-lid, *b* the *Caruncula lacrymalis*. *Fig.* 7 and 8. repreſent the lacrymal Ducts of each Eye ſeparated and here entire: *a a* denote the *Saccus lacrymalis*, as it is called; *b b* the *Puncta lacrymalia*, with their ſmall Tubes or Ducts, *c c*, leading into the lacrymal Sac. The Letters *d d* denote the *Canalis naſalis*, opening into the Noſe by the Aperture *e e*. In *Fig.* 9. you have a View of theſe Ducts annexed to the Eye, where the lacrymal Points are marked *a a*; the Caruncle *b*; the Ducts from the *Puncta lacrymalia c c*, leading into the *Saccus lacrymalis d*, thence into the *Canalis naſalis e*, and by that into the Noſe through the Aperture *f*[a]. Nature of the Diſorder.

II. This Diſorder of the Eye may proceed from many Cauſes, which impede or obſtruct the Paſſage of the Tears into the Noſe through the before deſcribed Parts. Thus, if the *Puncta lacrymalia* are ſtopped up, it will produce an *Epiphora*, or watery Eye: but as long as the Paſſages into the Noſe are clear, that Humour, which is ſeparated by the lacrymal Gland, to moiſten and cleanſe the Eye, will be drank in by the lacrymal Points, conveyed from thence into the Sac, and from thence it will, by degrees, paſs into the Cavity of the Noſe itſelf. The *Epiphora* may therefore proceed, (1.) From ſome hard Tumor or Tubercle, as the *Encanthis*, in the greater Canthus or Angle of the Eye, obſtructing the *Puncta lacrymalia*. (2.) From a Contraction or Concretion of the *Puncta*, after a Wound, Ulcer, or Burn of the Eye-lid: and (3.) From the ſame Cauſes, or from an Obſtruction of the *Canalis naſalis*; as may frequently happen in an Inflammation, from an inſpiſſated or gummy Matter. (4.) It may be cauſed by a Polypus, Caruncle, or Excreſcence in the Noſe, compreſſing and occluding the lacrymal Duct internally. (5.) From a *Fiſtula lacrymalis*. (6.) An *Ectropium*, or Inverſion of the Eye-lids. (7.) From an Eroſion, or Loſs of the *Caruncula lacrymalis*. And, laſtly, (8.) from a Wound in the lacrymal Duct, blocking up the ſame with an ill-formed Cicatrix. Cauſes.

III. The Diſorder itſelf may be readily diſcovered both from the Looks and Relation of the Patient: but, to find out its immediate Cauſe, requires much more Attention. When it ariſes from a Loſs of the lacrymal Caruncle, a Diſ- Diagnoſis.

3 tortion

[a] This Paſſage of the Tears is by many thought to be a modern Diſcovery: but the celebrated Anatomiſt MORGAGNI, in his firſt and ſixth *Adverſaria Anatomica*, has demonſtrated the Courſe to have been known and obſerved by GALEN, VEGETIUS, BERENGARIUS, FALLOPIUS, CARCANUS, STENO, &c. After MORGAGNI this Part has been explained at large by ANELIUS, *in Lib. De Fiſtula lacrymali*, and MEIBOMIUS, *in Epiſt. De Vaſis Palpebrarum novis.*

tortion of the Eye-lids, an Encanthis, or a Polypus, in the Noſe, the Cauſe is generally obvious enough: but when it is from a Concretion of the *Puncta*, the Cauſe can only be known by Inſpection, and conſidering whether there has been any Wound or Burn, &c. When the *Puncta* remain open, and the naſal Canal is concreted or obſtructed, the Tears have a ready Admittance into the Sac, but not into the Noſe; which therefore diſtend or dilate the Sac, from whence the Diſorder is ſometimes named a *Hernia lacrymalis*; and by ANELIUS it is termed a *Hydrops Sacci lacrymalis*[a]. In this Caſe, upon preſſing the Finger on the lacrymal Sac, it does not diſcharge its Contents into the Noſe as it ought, but the Tears return again through the *Puncta* into the Eye. See *Tab.* XVI. *Fig.* 10. A. Sometimes the lacrymal Sac is thus dilated, ſo as to form a very conſpicuous Tumor externally; which, by Preſſure with the Finger, will for the preſent be greatly diminiſhed, or elſe totally diſappear. If the Diſorder is at the ſame time accompanied with a *Fiſtula lacrymalis*, the aforeſaid Preſſure will diſcharge a purulent Matter along with the ſerous Humour; whereas in the ſimple Epiphora, it will appear quite limpid and aqueous.

Prognoſis and Cure.

IV. The Prognoſis and Treatment of this Diſorder will turn out various, according to the particular Cauſe and Circumſtances. When accompanied with an Encanthis, Polypus in the Noſe, a Diſtortion of the Eye-lids, or a *Fiſtula lacrymalis*, the Epiphora cannot be cured, till you have firſt removed thoſe Symptoms which cauſe it. When it ariſes from a Concretion of the *Puncta lacrymalia*, you ſhould carefully examine whether the Ducts leading into the *Saccus*, marked *c c*, *Fig.* 7 and 8, are all along cloſed and concreted, or whether their Orifices only are occluded with a thin Film: for, if they are all the Way concreted, whether from a Cicatrix, Wound, or Burn, there will be no Poſſibility of a Cure; whereas the thin Skin occluding their Orifices, may be eaſily perforated with a ſmall Needle, and kept open, till they are healed, with a Briſtle, or Silver Wire, dipped in *Ol. Ovor.* as at *Fig.* 11, 12, 13.

Obſtruction of the Duct.

V. If the *Puncta* appear to be pervious, and in their natural State, you may conclude the *Canalis naſalis* to be obſtructed; which being uſually occaſioned by a glutinous Matter, may be generally removed, ſo as to cure the Diſorder, if it has not been too long neglected. To diſperſe and remove the Matter, Diſcutients muſt be often applied, with repeated Preſſure by the Finger, to expel the ſtagnant Humours, that they may not become acrimonious, erode the Membranes, and bring on a *Fiſtula lacrymalis*. One of the beſt Diſcutients for this Purpoſe is a Tincture of Aloës diluted in ſome Eye-water, or an Infuſion of Hyſſop, Betony[b], or ſome mineral Waters, or the Salts extracted from them mixed with an Eye-water, &c. In the mean time ſhould be ſometimes uſed a Sternutatory *ex Majoran. Lil. Conval. Mar. Majoran. Hellebor. &c.* and if theſe Means prove ineffectual, you may treat the Patient in ANELIUS's new Method of curing a *Fiſtula lacrymalis*, by paſſing a ſmall Silver Probe, *Tab.* XVI. *Fig.* 11, 12, 13, into the *Puncta*, and through the lacrymal Duct and Sac into the *Canalis naſalis*, and ſo into the Noſe. But this is an Operation that ought not to be attempted by every one, who is not an expert Operator, and well verſed in the Structure of theſe Parts: otherwiſe you not only miſcarry in your Operation,

[a] In *Diſſert. ſur la nouvelle Decouverte de l' Hydropiſie du Conduit lacrymal.* Paris 1716.

[b] This Infuſion is highly commended by SCHOBINGERUS, for a *Fiſtula lacrymalis*, in his Treatiſe on that Subject. p. 20.

Operation, but greatly injure the Patient[a]. The Paſſages are to be thus cleared by the ſlender Probe every Morning and Evening, for ſeveral Days, injecting afterwards ſome of the beforementioned Liquors by a ſmall Silver Syringe, *Tab.* XVI. *Fig.* 14. the ſlender Tube of which is to be inſerted into the lower *Punctum lacrymale*, as we ſhall more particularly direct in the following Chapter: and thus, by the repeated Uſe of Injections, the Diſorder will be either removed, or elſe degenerate into a *Fiſtula lacrymalis*, and muſt then be treated accordingly. Laſtly, when this Diſorder ariſes from a Loſs of Subſtance in, or an Eroſion of, the lacrymal Caruncle, it will be to no Purpoſe to uſe Remedies, becauſe the Caſe is incurable. *Vid.* HEBENSTREIT. *Diſſert. De Oculi lacrym.* Lipſiæ 1743.

CHAP. LIV.

Of the FISTULA LACRYMALIS, *and of the Diſorders related to it.*

The *Fiſtula lacrymalis* deſcribed.

I. THE *Fiſtula lacrymalis* is generally underſtood to be a little Ulcer in the greater or internal Canthus of the Eye next the Noſe, which either of itſelf, or by Preſſure, diſcharges a purulent Matter. The Seat of this Ulcuſcle is in the *Sacculus lacrymalis*, or Paſſage for the Tears into the Noſe: therefore the *Fiſtula lacrymalis* is more or leſs dangerous, in proportion to the Size and Condition of the Ulcer, which ſometimes lies concealed only in the Sacculus, and diſcharges its Matter through the *Puncta lacrymalia*; but ſometimes again it not only erodes the Sacculus, but alſo the external Skin, and the adjacent Bone. If the Skin is not eroded through, the Fiſtula is thence denominated *imperfect*; as it is termed *perfect* after having made its Way through the Integuments[b]: but when it has alſo eat through the adjacent Bone, or rendered it carious, it is then uſually termed a *complicated Fiſtula lacrymalis*. It is remarkable, that the generality of Phyſicians and Surgeons had a wrong Notion of the Nature and Treatment of this Diſorder, till the Beginning of the preſent Century. Their Error might be owing partly (1.) To the Multiplicity of Diſeaſes to which this Part of the Eye is ſubject, and the Number of different Names which are frequently given to each of them. (2.) To the real Nature of the Diſorder, having been examined into by very few Surgeons and Anatomiſts; for moſt of them imagined the Seat of the Ulcuſcle to be either in or under the lacrymal Caruncle; whereas the more accurate of the Moderns diſcovered, that the purulent Matter was diſcharged neither from nor behind the Caruncle, but rather out of the *Sacculus lacrymalis* through the *Puncta*[c]. Having acquired a wrong Idea of the Diſorder, they were conſequently led by that into a wrong Practice

[a] Yet this Operation is far from being impracticable, as many Surgeons not well verſed in theſe Diſorders (and among the reſt GARENGEOT) would fain perſuade us. For to ſay nothing of ANEL, I myſelf have often and often performed it upon Numbers of Patients.

[b] This Species of the *Fiſtula* is what CELSUS (*Lib.* vii. *N.* 7.) ſeems to term *Ægilops*, but he does not ſpeak very intelligibly of it in this Place.

[c] FALLOPIUS was perhaps the firſt Anatomiſt that obſerved this, in *Tom.* II. *p.* 224. See alſo MORGAGNI, *Adverſ. Anat.* VI. 64.

Practice: both which the Moderns have endeavoured to correct, and not without Succeſs.

The different Names and Kinds of the Diſorder.

II. But that our Reader may be a better Judge of the falſe Opinions which have been entertained and advanced concerning this Diſorder by the principal Writers in Surgery, we ſhall endeavour briefly to relate them: and firſt, ſome of them have by the Name of *Fiſtula lacrymalis* underſtood that Kind of Diſorder which we term *Epiphora*, or the watery Eye, and have deſcribed in the preceding Chapter. (2.) Others ſeem to uſe the Term *Fiſtula lacrymalis*, *Anchylops*, and *Aegilops*, as ſynonymous; ſo that there is no Poſſibility of knowing their Meaning, till we are furniſhed with a proper Diſtinction and Explanation of thoſe Diſorders ſeparately; for the *Anchylops* is, by the generality of the modern Writers, uſed to ſignify a Tubercle in the greater Canthus of the Eye next the Noſe, whether it be ſeated in or near the lacrymal Sac, or whether it be with or without an Inflammation accompanying it. It ought to be here obſerved, that the *Sacculus lacrymalis*, as well as other Parts, is ſubject to encyſted Tumors, Inflammation, and Abceſs, and very often to a Diſtenſion or Rupture, now termed a *Hernia lacrymalis*; (ſee *Tab.* XXVI. *Fig.* 10. A B. and *Fig.* 16 and 17.) in which laſt, upon preſſing the Finger on the Tumor, it ſubſides more or leſs, and the ſerous Humour diſcharges itſelf either through the *Puncta lacrymalia* at the Eye, or into the Cavity of the Noſe, or both Ways. We define an *Aegilops* to be a ſmall Tumor formed after an Inflammation or Abceſs in the greater Canthus of the Eye, near the *Sacculus lacrymalis*; which in Time, by the Acrimony of its purulent Matter, erodes the external Skin and lacrymal Ducts, ſometimes eats away the Fat round the Globe of the Eye, and ſometimes renders the *Oſſa plana*, and other Bones near the Noſe, carious to a dangerous Degree. Sometimes the upper, lower, or both of the lacrymal Ducts, are ſo eroded, as to diſcharge large Quantities of purulent Matter through the *Puncta* in the greater *Canthus*; and then it forms the *Fiſtula lacrymalis*, whoſe Characteriſtic is a purulent Matter. But, when the diſcharged Humour is quite limpid and aqueous, the Diſorder ought then to be denominated an *Epiphora*, as we obſerved in the preceding Chapter. (See *Fig.* 18. *lit. a* and *b.*) From what we have here advanced, I think it will not be difficult for any one to diſtinguiſh the different Diſorders of this Part; which, from their Affinity, are very often confounded by Phyſicians and Surgeons.

Cauſes.

III. An *Anchylops* may proceed from many Cauſes; and, among others, an Inflammation or encyſted Tumor may produce this Diſorder, as well as occaſion a ſimple *Fiſtula lacrymalis*, or an *Aegilops:* yet the firſt ariſes ſtill more frequently from a Relaxation and Diſtenſion of the lacrymal Sac; ſo that we generally meet with an *Aegilops* and *Fiſtula lacrymalis* fixed in the greater Canthus of the Eye at one and the ſame time: this ſeems to ariſe from an Obſtruction of the Paſſage of the Tears, or purulent Matter, into the Noſe; the Conſequence of which muſt be an Extenuation and Tumor of the lacrymal Sac. An *Aegilops* is generally cauſed by a previous Inflammation or Abceſs, which frequently erode the lacrymal Ducts and the external Skin, and even produce a *Fiſtula lacrymalis* in its worſt Degree. But though there are many more Cauſes beſides Inflammation, which may produce a *Fiſtula lacrymalis*, yet there is no Cauſe ſo frequent or immediate as an Exulceration of the lacrymal Sac, or of the adjacent Membranes. But when once the lacrymal Ducts are eroded, the Matter finds

finds an immediate Paſſage into the ſubjacent *Sacculus*, as at *Fig.* 18. A *Fiſtula lacrymalis* may alſo frequently proceed from an Obſtruction of the inferior lacrymal Duct, termed the *Canalis naſalis*, *d d*, *Fig.* 7 and 8. from whatever Cauſe that Obſtruction may ariſe: for no Obſtruction can be formed, without inducing a Stagnation of the Humour, which will therefore become acrid, diſtend the Duct, and either erode, or totally deſtroy, its Membranes. And in this Manner the Diſorder is frequently occaſioned in many Patients who have had an Inflammation in their Eyes, in the Membranes of their Noſe, or in theſe Ducts themſelves, or when thoſe Parts have been injured by the Small-pox, as I have frequently obſerved: though it muſt be confeſſed, that the Diſorder ſometimes ariſes ſpontaneouſly, without the Aſſiſtance of any of the beforementioned Cauſes.

Kinds of the Fiſtula lacrymalis.

IV. There are various Species of theſe *Fiſtulæ*. The firſt Diſtinction of them is, (1.) Into *perfect* and *imperfect*: the former of which is, when the purulent Matter flows out through an Eroſion of the Skin in the Canthus; and the latter, when the Matter is diſcharged through the *Puncta lacrymalia*, the Skin remaining entire: which laſt Kind is generally accompanied with a Tumor of the lacrymal Sac. You may have an Idea of the perfect Kind, from conſulting *Tab.* XVI. *Fig.* 19. *a. b.* Some of theſe *Fiſtulæ* are again diſtinguiſhed into (2.) *ſimple* and *compound*; the laſt of which is, when a Calloſity, Caries, or the like, attend. Some again are, (3.) *mild* and *recent*; others *old* and *malignant*. (4.) Some *intermitting* and *periodical*; others *continual*. Still more Diſtinctions of the ſeveral Species of this Diſorder may be ſeen in *p.* 8. of our profeſſed Diſſertation on the Subject in 4to 1716, at *Altorf*. We have ſtill another Diſtinction of theſe *Fiſtulæ* into *true* and *falſe*, made by M. GARENGEOT: by the *true*, he underſtands an Ulceration of the lacrymal Ducts; and by the *falſe*, he intends an Ulceration in the adjacent Parts only, which we term an *Aegilops*. Some[a] will have a Calloſity eſſentially neceſſary to the Formation of a *Fiſtula lacrymalis*; becauſe a Callus is conſtantly found in moſt other *Fiſtulæ*: but this is not the common and received Notion of a *Fiſtula lacrymalis*, as we are taught by the Authorities of CELSUS, FALLOPIUS, CARDAN, WOOLHOUSE, and MORGAGNI, *adverſ. Anat.* VI. *p.* 82. and from daily Experience. M. ST. YVES[b], the late famous Oculiſt at *Paris*, aſſerts that he ſeldom found a Callus in theſe *Fiſtulæ:* and I myſelf have obſerved a great many, and thoſe inveterate lacrymal *Fiſtulæ*, which have yet had no Calloſity. There are ſome Surgeons again, who imagine that there never can ariſe a *Fiſtula lacrymalis*, without an Obſtruction of the *Canalis naſalis* at the ſame time, becauſe ſuch an Obſtruction muſt be the Occaſion of the *Fiſtula:* but even this Opinion is without Foundation, as hath been long ago evinced by the Authorities of the beſt Writers, and as I have been frequently aſſured by Experience: for I have often obſerved, and am now acquainted with ſome of theſe *Fiſtulæ*, in which the purulent Matter has a free Exit from the lacrymal Sac through the *Puncta lacrymalia*, if you preſs it with the Finger every Day; and at the ſame time the *Canalis naſalis* appears to be open, becauſe the purulent Matter is alſo diſcharged through it into the Noſe[c].

V. Having

[a] As SIGNOROTTUS and PLATNERUS, in *Diſſ. de Fiſt. lacrymali*, Sect. 1, 2, 3.

[b] See his *Traité des Maladies des Yeux*, pag. 59. and SCHOBINGERI *Diſſ. de Fiſtul. lacrym.* p. 3.

[c] Some will have it, that the purulent Matter flows only through the upper, and others only through the lower *Punctum lacrymale*; but it has generally a Paſſage through both, though often more is diſcharged through one than the other.

Signs of the Fiſtula lacrymalis.

V. Having in general deſcribed and explained the ſeveral Kinds of theſe *Fiſtulæ*, and the Diſorders related to them, we ſhall now proceed to the Signs by which they are diſcovered. And firſt, you may be pretty well aſſured, that the Patient has a lacrymal *Fiſtula*, if he complains of the Tears being more copious than uſual, and running over his Cheek, and that a Quantity of purulent Matter is found collected in the Eye, in a Morning chiefly; and at the ſame time you obſerve no Appearance of Inflammation: but if you preſs the lacrymal Sac with your Finger, it diſcharges a Quantity of purulent Matter by the *Puncta lacrymalia*. This appears to me the moſt certain Sign of a *Fiſtula lacrymalis*: and with me FALLOPIUS, WOOLHOUSE, and ANELIUS concur. You may judge whether there be any Caries from the ill Smell, and from the livid or blackiſh Colour of the Part, with the Diſcharge of purulent Matter; and eſpecially, if the Bone appears bare or eroded to the Eye or Probe, in open *Fiſtulæ*. The Colour of the Matter diſcharged is ſo far from giving a ſure Indication, whether or no the Bone is carious, that I have often found it of a good Colour, when at the ſame Time the Bone appeared rough and eroded to the Probe: but you may be generally aſſured, there is a Caries of the Bone, if the *Fiſtula* has been of very long ſtanding, and diſcharges a large Quantity of Matter. But the Seat of the Caries is not always the ſame, being ſometimes in the *Os lacrymale*, ſometimes in the *Os planum*, and in the *Os maxillare ſuperius*. You may diſcover whether the *Canalis naſalis* be obſtructed, from little or none of the purulent Matter, or injected Liquor, being able to make its Way into the Noſe, but all returning through one of the *Puncta lacrymalia*[a]. A Callus in theſe *Fiſtulæ* may be diſcovered by the unuſual Hardneſs or Reſiſtance which the Parts give to the Finger; but this is not a frequent Symptom in lacrymal *Fiſtulæ*, as hath been often obſerved by ST. YVES, M. GARENGEOT, and myſelf. If theſe Parts are infeſted with an encyſted Tumor, they appear preternaturally enlarged, and harder than uſual, nor does the Tumor ſubſide by preſſing it with the Finger; and there appears no Sign of Inflammation. But if the Tumor ſubſides by Preſſure with the Finger, you may conclude there is a *Hernia lacrymalis*, or Dilatation of the lacrymal Sac. Laſtly, an *Ægilops* is diſcovered by the Appearance of an Exulceration in the greater Canthus of the Eye next the Noſe, without affecting the lacrymal Ducts.

Prognoſis.

VI. The ſeveral Diſorders before enumerated uſually terminate differently, according to particular Circumſtances. But as the Eye itſelf, and the ſpongy Bones of its Orbit, are ſo nearly ſituated, it is hardly poſſible the Patient ſhould eſcape a Caries in the laſt, with many grievous Symptoms in that Organ itſelf. An *Anchylops* or *Ægilops* may very eaſily degenerate into a Fiſtula; and a ſlight Fiſtula may become obſtinate, malignant, and even cancerous; which having deſtroyed the Bones, there are then but little Hopes of obtaining a Cure. Theſe Diſorders are in general more or leſs malignant, according as the Patient is of a good or bad Habit of Body, as the Matter of the Fiſtula is more or leſs acrimonious,

[a] I obſerved an uncommon Species of the *Fiſtula lacrymalis* here in a Student, *Anno* 1726; in which, though the Diſorder had been of eight Years ſtanding, yet no Matter could be diſcharged by preſſing with the Finger. The Tears conſtantly iſſued down upon his Cheeks, and after Sleep the Eye was found replete with a purulent Matter: but when a Quantity of Liquor was injected at either *Punctum*, it ran out with ſome purulent Matter through the other. There was no Tumor of the lacrymal Sac; but, upon inciſing the Integuments, the lacrymal Bone was found carious.

monious, and as the Patient is more or leſs regular in his Diet and Courſe of Life. If the Patient is in other Reſpects well, the Diſorder recent, and without a Caries, Callus, or other bad Symptoms, there is no great Danger: but the Diſorder may be cured, by the Method of ANELIUS, in a few Days time. The perfect or compleat Fiſtula which has eroded through the Skin, is generally attended with a Caries; and is therefore hardly, if at all, curable, before the carious Bones are removed. Alſo a Callus muſt be firſt removed before you can cure thoſe *Fiſtulæ* in which it is found: but if both Calloſity and Caries are abſent, a Cure may be obtained with much more Eaſe and Expedition. again, in general, the older or more inveterate the Fiſtula, the more difficult it is to cure; becauſe in them the Bones are commonly infeſted with a Caries: and if that is not perfectly removed, though you ſhould, in Appearance, cure the Diſorder, it will quickly return again. But what is more than a little ſurpriſing, there are ſome Surgeons who write, that ſeveral of theſe *Fiſtulæ* which, have been accompanied both with a Callus and a Caries, have been cured barely by leaving the Diſorder to Nature[a]. Unleſs the *Canalis naſalis* be rendered pervious, and kept open, the Cure cannot be compleated: for though you remove the Callus and Caries by the Knife or Cautery, the Patient will be afterwards troubled with a watery Eye, in which the Tears run down over the Cheeks. The compreſſing Inſtruments formerly uſed to relieve this Complaint, do little more than moleſt the Patient, or frequently turn a mild into a malignant Fiſtula. But the Practice of the modern Surgeons is greatly to be preferred before that of the Antients in this Diſorder: for the firſt being reformed by the Authority and Example of ANELIUS, about the Year 1712, have ever ſince continued to cure recent *Fiſtulæ* of this Species after his Manner, without either the Uſe of the Scalpel, Terebra, or Cautery, provided there is no Callus or Caries in it, notwithſtanding what others may ſay to the contrary; whereas formerly they hardly ever cured a *Fiſtula lacrymalis* of any Kind without the Uſe of one of thoſe ſevere Remedies[b].

Treatment of the Anchylops.

VII. If the Patient is troubled with an *Anchylops*, or Tumor or Inflammation in the greater Angle of the Eye next the Noſe, the Surgeon muſt in that Caſe, uſe his Endeavours firſt to diſperſe it, to prevent the Tumor from degenerating into an Abceſs or Fiſtula. This Intention may be beſt anſwered towards the Beginning of the Diſorder, by moiſtening the Part with a little *Sp. Vitriol.* dipping a ſmall Bruſh, or the End of the Finger therein, ſeveral Times in a Day, as in treating upon Tumors we directed for the Furuncle: but in this Practice you muſt be very careful to avoid injuring the Eye itſelf; upon which Account it may in ſome Caſes, be ſafer to uſe a Liniment of *Mel. Roſar.* acidulated with *Sp. Vitriol.* covering the Part afterwards with a Diachylon Plaſter. In moſt Caſes, a Cure may be almoſt as readily obtained by frequent fomenting with Compreſſes dipped in warm *Sp. Vini Camph.* and a Cataplaſm *ex Pomis coctis, vel aſſatis Camphoroque miſt.* to be continued till the Tumor ſubſides, and the Inflammation is diſperſed. If the Tumor ſhould appear to be of the encyſted Kind, you may treat it as we have directed in *Chap.* XXVIII. *Sect.* I. *N.* VI, and VII. foregoing: by which Method I happily extirpated a large encyſted Tumor by the Scalpel, which was very deeply ſituated in the

[a] This does but very ſeldom happen. See more in MAITRE-JEAN, *in Lib. De Morbis Oculorum, in Cap. de Fiſtula lacrymali.*

[b] See my Diſſertation on this Subject.

Orbit of the Eye of a certain Maid. Laſtly, when the Tumor ariſes from a Diſtention of the lacrymal Sac, you muſt treat the Diſorder by the Methods we ſhall preſently direct at *N.* X. following.

Treatment of the Ægilops.

VIII. If the laſt mentioned Tumor or Inflammation rather tends to Suppuration than to be diſperſed by the preceding Treatment, it will then be proper to forward its Maturation or Converſion into Matter as much as poſſible, leſt an obſtinate Fiſtula, or worſe Conſequences, ſhould be the Effects of too long Delay. The Suppuration of it may be conveniently promoted by a Diachylon Plaſter with the Gums, or an emollient Cataplaſm frequently applied warm. As ſoon as you can diſcover that the Matter is ſuppurated, you are to open the moſt depending Part of the Tumor, either with a Lancet or Scalpel, to diſcharge and preſs out the Matter, that it may not eat through its including Cyſt, or the adjacent thin Bones. That being thus diſcharged, the Abceſs or Ulcer muſt be next deterged by dreſſing with digeſtive Ointments, or *Mel. Roſarum cum Myrrha, vel Ung. Ægyptiac. ſeu Præcipitat. rub. Portiuncula permiſt.* after which it may be healed with vulnerary Balſams, in the Manner we directed for Abceſſes in general. If the Abceſs in this Diſorder ſhould break of its own accord, as I have frequently known it to do, and its Aperture or Orifice appears too narrow to give a free Diſcharge to the Matter, it may be afterwards dilated with a Tent, prepared Sponge, Gentian Root, or rather by the Scalpel, and then treat it as before. If the Bone appears foul, it will be neceſſary to apply ſome ſcraped Lint, with a few Drops of *Sp. Sulph. aut Vitriol.* or a little *Pulv. Euphorb.* laying over it a Compreſs dipped in *Aq. Calcis*; by which Means having removed the Caries, the Wound will be diſpoſed to heal. Sometimes it will be found neceſſary to exfoliate or ſcrape the foul Bone with the Raſp, repreſented in *Tab.* VII. *Fig.* 3, 4, 5, or *Tab.* XVIII. *Fig.* 9. Some Surgeons think it a more ready Method of Cure, to cauteriſe the Bone with red-hot Irons, adapted to a Tube or Caſe, as in *Tab.* XVI. *Fig.* 21 and 22. compleating the reſt of the Cure with Balſams or vulnerary Medicines, in the Manner we ſhall explain more at large in treating of this Diſorder at *N.* XII. following.

Treatment of ſlight Fiſtulæ.

IX. The Treatment of the true Species of lacrymal *Fiſtulæ*, in which there is an Ulceration of the lacrymal Paſſages, is various, according to the different Nature, Degree, and Circumſtances of the Diſorder. For when the Fiſtula is recent, the Patient of a good Habit, the Skin intire, and the Ducts not ulcerated or obſtructed, but diſcharging freely a mucous, and not a purulent Matter into the Noſe; you ought not in theſe Circumſtances, to have immediate Recourſe to the Knife, Terebra, or Cautery, but firſt endeavour to cure the Fiſtula by the mildeſt Methods of Treatment, before you try the ſeverer Operations of Surgery. In this Caſe, you ſhould frequently expreſs the Matter included in the lacrymal Sac by your Fingers; leſt it become ſo acrid, as to erode the adjacent Parts by its too long Stay: and, in the Intervals, you ſhould ſtrive to cleanſe or deterge the Parts by the repeated Uſe of the mundifying Remedies, which we adviſed for the watery Eye in *Chap.* LIII. *N.* V. At the ſame time, too, you muſt call in the Aid of Phlebotomy, Purges, Scarification, Bliſters, Diet, and Regimen, according to the Patient's particular Habit and Circumſtances.

Cure by Compreſſion.

X. M. DIONIS tells us, in his *Surgery*, that he has cured many of theſe recent *Fiſtulæ*, particularly in Infants, barely by Compreſſion in a proper Manner: and GARENGEOT alſo affirms the ſame to have been done formerly at *Paris* by that

that eminent Surgeon M. ARNEAU. By the firſt of theſe the Compreſſion was made in the following Manner: 1. Firſt of all he impoſed a Piece of *Emplaſt. de Minio* upon the Tubercle or Fiſtula of the lacrymal Sac: then 2. he applied a ſmall triangular Compreſs of about the Thickneſs of one's Finger, or, inſtead of the one thick Compreſs, ſeveral thinner ones upon each other, in order to fill up exactly the Cavity in the Angle of the Eye next the Noſe. In the next Place, 3. he adapted another Compreſs over the former, dipping both of them firſt in ſome *Aq. Calc.* or *Sp. Vini.* Laſtly, 4. he firmly ſecured and preſſed down the Compreſſes upon the Tumor by a ſtrict Deligation with a circular Bandage; that by this Means, none of the vitiated Humours might be collected or retained, and that the relaxed *Sacculus* might, by degrees, recover its former Tone and Dimenſions. But, according to M. DIONIS, this Treatment muſt be continued for ſeveral entire Months to cure the Patient. It is to be obſerved, that ſome uſe a peculiar Inſtrument for compreſſing the Parts diſordered, inſtead of Compreſſes and Bandage: of which Inſtrument there are ſeveral Kinds propoſed by FABRIC. AB AQUAPENDENTE, SCULTETUS, PALFYN, and myſelf, in *Tab.* XVI. *Fig.* 20. taken from PLATNERUS. But, after all this Method by Compreſſure will be to no Purpoſe when the lacrymal Ducts are concreted or obſtructed: for the Advantage of this Practice can only take place when there is an Abceſs near the lacrymal Sac, as in *Fig.* 18. or at leaſt when the lacrymal Ducts are found pervious.

Cure by Inciſion.

XI. When the Diſorder is become ſo malignant or inveterate as not to be relieved by the preceding Method of Compreſſion, the general Practice of Surgeons in that Caſe was formerly, and now is, to lay open the Tubercle, or diſtended lacrymal Sac almoſt in the Middle, betwixt the internal Canthus and the Noſe: and this either by Cauſtic, or rather by Inciſion with a Scalpel or a Lancet; but with great Circumſpection, to avoid wounding the lacrymal Ducts and *Puncta*, which lead to the Sac, or the Ligament which faſtens one Eye-lid to the other, which would greatly deform the Eye. 'Tis generally adviſed to make this Inciſion obliquely: as, for Example, from *d* towards *e* or *c*. *Fig.* 9. *Tab.* XVI. or in *Fig.* 10. from B towards A; for which ſome prefer the ſtraight, and others the crooked Scalpel: but either of them will do, in my Opinion; for I have ſucceſsfully performed the Operation with both. Your Inciſion muſt be continued downward, till you have penetrated into the Cavity of the lacrymal Sac, enlarging it afterwards both upward and downward in the aforeſaid Direction from the Top of the Sac down to the *Canalis oſſeus*. The Wound is not to be dilated by filling it with Lint (though PLATNERUS and GARENGEOT recommend a particular Inſtrument for this Uſe) and laſtly the Dreſſings are to be ſecured with Compreſs and Bandage. There are others again who rather approve of making this Inciſion in a ſemicircular Form like an Arch, whoſe Convexity muſt be towards the Noſe, and Concavity towards the Eye; beginning the Inciſion at the lower Part of the *Apophyſis naſalis* of the *Os Frontis*, where that Bone meets the *Os maxillare* and *lacrymale*, and continuing your Inciſion from thence, in the Form of an Arch, to the Meeting of the internal *Apophyſis* of the *Os jugale*, as we have repreſented by the dotted Line *c b Fig.* 19. *Tab.* XVI. When your Inciſion is ſufficiently enlarged by the Knife, you muſt dilate it further with Lint, as before: by which means you have an Opportunity the next Day of obſerving, whether the Bones be carious, and in what

Part

Part or Manner it will be beſt to perforate them. If the Wound ſhould bleed much, you may apply a Pledgit of Lint dipt in *Sp. Vini rectificatiſſ.* to be retained on the Part with a Compreſs, and a little ſtricter Bandage. In the ſubſequent Dreſſings you muſt uſe *Eſſent. Succin. Ol. later.* and other detergent Applications, as we before directed for the *Aegilops*, at *N.* VIII. When the Parts are well cleanſed, you may finiſh the Cure with ſome vulnerary Balſam and deſiccative Plaſter, retained with a thick triangular Compreſs, as we directed at *N.* X. and thus the Wound gradually heals. Others again apply the compreſſing Inſtrument beforementioned upon the Wound over the Compreſs and Plaſter; but not very often with the deſired Succeſs, becauſe the *Canalis naſalis* is generally hereby obſtructed.

The antient Treatment of a callous Fiſtula with Caries.

XII. In a callous *Fiſtula lacrymalis* the Method of Treatment uſed by the antient Surgeons was, to open the Ulcer firſt, and then to dreſs it with *Trochiſc. de Minio, Præcipit. rub. Ung. Ægyptiac. Lap. infernal. &c.* with which they removed the Calloſity, and then finiſhed the Cure in the Manner we before directed. But if a Caries alſo accompanied it, they applied *Pulv. ex Euphorbio*, or *Sp. Sulphur. Vitriol. &c.* with ſcraped Lint. If theſe did not anſwer, they then raſped or ſcraped the vitiated Bone, as we directed at *N.* IX. or elſe applied the actual Cautery ſeveral times, according as the Caſe required. The cauterizing Inſtruments uſed in this Diſorder, were of various Figures, as the Surgeon beſt fancied; as you may ſee by thoſe figured in AQUAPENDENS, SCULTETUS, SOLINGEN, PALFYN, DIONIS, GARENGEOT, PLATNER, *&c.* Some were uſed naked without any Tube, as thoſe we have repreſented in our *Tab.* III. *Fig.* 14 and 16. Others again were furniſhed with a Tube, which was firſt placed in the Wound cloſe to the Bone, and then the Cautery was conveyed through it, to avoid burning the Skin and Lips of the Wound: ſee *Tab.* XVI. *Fig.* 21, 22. The Eſchars formed by the Cautery were afterwards ſeparated by ſome digeſtive Ointment, and the Wound then healed with vulnerary Balſams, as we directed before. But in performing this Operation you ſhould firſt not only bind up the Patient's ſound Eye, that he may not be terrified at the Sight of the Cautery, but you ſhould alſo ſecure the diſordered Eye by an Inſtrument in the ſhape of a Spoon, *Tab.* XVI. *Fig.* 23. that it may not be touched by the Cautery. It will be alſo previouſly neceſſary to dry the Bone well with Lint before you apply the Cautery, which will otherwiſe be too ſoon extinguiſhed. But, after all, this Treatment, in order to cleanſe the Fiſtula by the Cautery, will be to little or no Purpoſe, ſo long as the *Canalis naſalis* remains obſtructed: nor can the Tears be diſcharged into the Noſe, unleſs a new Paſſage be made for them by perforating the Bones with the Cautery; otherwiſe the Patient will be continually moleſted with a watery Eye after the Fiſtula is cured: ſo that this Method of Cure will, in my Opinion, ſucceed beſt when the *Canalis naſalis* remains pervious and entire, or when there is a Suppuration without-ſide the lacrymal Sac. Therefore it will be highly neceſſary to diſtinguiſh thoſe *Fiſtulæ*, in which the *Canalis naſalis* is occluded, or ſhut up, from thoſe in which it is not.

Cure by perforating the Os lacrymale.

XIII. To remove the laſt mentioned Symptom, the watery Eye, in the Cure of theſe *Fiſtulæ*, ſome Surgeons have propoſed the following Method: *viz.* After opening the lacrymal Sac, as we directed before at *N.* XI. the next Day they perforated the *Os Unguis* with a ſharp-pointed Inſtrument for the Purpoſe,

(*Tab.* XVI. *Fig.* 24. or *Tab.* VII. *Fig.* 7. or *Tab.* XXIV. *Fig.* 2.) which is carefully paſſed obliquely through the upper and lower Part of the *Os ſpongioſum* into the Cavity of the Noſe: after which they introduce and leave a ſmall Tent in the Wound, which is frequently cleared and opened with a Probe; till being healed, it forms an artificial lacrymal Duct. Some remove the Caries, and make an artificial lacrymal Duct, at the ſame time, by the forementioned Inſtruments, or by a Director, without any actual Cautery: which laſt is, however, uſed by ſome like that at *Fig.* 21. with the Tube *Fig.* 22. [a] with which the Bones are perforated, and a Paſſage made for the Tears into the Noſe as before. Though theſe Methods of Cure are very troubleſome and painful to the Patient, yet they are at preſent uſed as the beſt we are acquainted with. And St. YVES, the famous Oculiſt of *Paris*, treated his Patients in the ſame Method, as he informs us in his Treatiſe on *Diſorders of the Eyes*.

ANELIUS's new Method of curing lacrymal Fiſtulæ.

XIV. But, in conſideration of the great Difficulty there is to perſuade timorous Patients, eſpecially thoſe of higher Rank, to undergo the Severity and Fatigue of the forementioned Operations of Inciſion, boring, cauterizing, *&c.* ANELIUS, in the Year 1712, endeavoured to contrive a more ſafe and eaſy Method of curing theſe *Fiſtulæ*, in favour of the Duke of SAVOY, who was then troubled with the Diſorder; which Method ſucceeded ſo well, as to cure not only recent, but even inveterate *Fiſtulæ*, if not accompanied with Callus or Caries, and that even without the Severity of the Knife, Cautery, or Compreſſion, in the following Manner.

The Uſe of ANELIUS's Probe.

XV. He firſt provided himſelf with a ſlender Probe, in the Form of an Arch, made of ſmall Silver-wire, as in *Tab.* XVI. *Fig.* 11, 12, 13. then placing the Patient in a convenient Poſture againſt the Light, he opens the upper Eye-lid with the Fingers of one Hand, while, with thoſe of the other, he introduces the crooked Probe through the upper *Punctum lacrymale* into the Sac; which may be done with more or leſs Difficulty, according as the Surgeon has before conſidered the Figure, or Poſition, and anatomical Structure of the Parts. After having introduced the Probe into the Sac, he gently agitates and preſſes it downwards, and towards the Noſe, with a certain Slight, into the obſtructed *Canalis naſalis*, which is by this means, opened. Theſe Ducts are much more eaſily opened by this Artifice, when they are only obſtructed by Matter, or ſome glutinous Humour, than when they are totally cloſed and concreted, as is frequently obſerved in theſe *Fiſtulæ* which are inveterate: for the laſt ſometimes require the Probe to be preſſed into them ſo forcibly, as to excite ſome Pain, and often ſet the Noſe a bleeding a little [b]. But to prevent the newly-opened Duct from cloſing again, M. ANELE thinks it neceſſary to inject ſome Liquor every Night and Morning, or oftener; and then to repeat the Introduction of the Probe as often as it may be found neceſſary, till no more Matter iſſues from the *Puncta lacrymalia:* which denotes the Ulcer to be cleanſed, and the Ducts to have recovered their natural State.

XVI. To

[a] See SOLINGENIUS, PALFYNUS, and GARENGEOT.

[b] M. GARENGEOT appears to be ignorant of the Uſe of theſe Probes, when he thinks they cannot open the Ducts, but only ſerve to ſearch out the lacrymal Sac. See *N.* XXV. following. STAHLIUS was the firſt, who paſſed the Briſtle through the *Puncta lacrymalia* into the lacrymal Sac; but not with the View of opening the naſal Duct.

Uſe of M. ANELE's Syringe.

XVI. To inject theſe Parts, I muſt recommend the Syringe, contrived by ANELIUS, and repreſented in *Tab.* XVI. *Fig.* 14. or elſe ſome other like it. The Tube A, in the anterior Part of this Inſtrument, is about the thickneſs of a Hog's Briſtle, and is to be inſerted into the *Punctum lacrymale* of the lower Eye-lid, as being leſs moveable: in which manner you force the healing Injection ſeveral times into and through the lacrymal Sac, in order to waſh out the Sordes, and render the Ducts pervious[a]. To perform this Operation the more eaſily, your Patient ought to be placed againſt the Light, with his Head either erect, or a little inclined backward: and, if the Diſorder be in the right Eye, the Surgeon ſhould ſtand on the right Side of the Patient, and having filled the Syringe with a ſuitable Injection, he then places his left Ring-finger under the *Punctum lacrymale* of the lower Eye-lid, near the lacrymal Sac, and thereby draws down the Eye-lid, to bring the *Punctum lacrymale* into View; and thus he more eaſily inſerts the Tube of the Syringe, and, at the ſame time, his Finger ſerves as a Fulcrum, or Support, to the others which move the Syringe. Having in this manner, ſecured the Eye-lid, the Surgeon next takes the Syringe by its Hinder-part C, betwixt the Fore and Middle-finger of his Right Hand, and carefully inſerts the Tube A, in the lower End of the Syringe D, into the lower *Punctum lacrymale:* after which he preſſes the Handle of the Sucker B into the Syringe by the Thumb of the ſame Hand, ſo as to force the Liquor through the lacrymal Duct, Sac, and *Canalis naſalis* into the Noſe; from whence it will run into the Fauces, and ſome Part of it will eſcape through the upper *Punctum lacrymale.* But to ſay Truth, the whole of this Method is much better and eaſier demonſtrated by Practice, than expreſſed by Words. If the Diſorder be in the left Eye, the Surgeon muſt then ſtand on the right Side of the Patient, and manage the reſt of his Operation as before. If the Surgeon pleaſes, he may, for Variety, inſert his Syringe, and inject by the upper *Punctum lacrymale*, after having turned it upward and downward by his Finger. But to inject by either of them as he ought, he ſhould be provided with good ſharp Eyes, and a dextrous Hand: though he will find it the moſt eaſy of the two, to inject by the lower *Punctum lacrymale.*

What more muſt be done.

XVII. Theſe two Operations of Probing and Injecting muſt be continued, or repeated every Day, till you find, 1. that the Injection will paſs freely into the Noſe without the Aſſiſtance of the Probe; and, 2. that there is no purulent Matter diſcharged either ſpontaneouſly, or by Preſſure from the lacrymal Sac into the greater Canthus of the Eye: and then you may conclude, from theſe two circumſtances, that the Cure is completed: which however is not always performed within the ſame time, but ſooner or later according to the Nature and Degree of the Diſorder. When mild, it is ſometimes cured within four, eight, fourteen, or twenty Days; and ſometimes longer. But there is hardly any lacrymal Fiſtula ſo bad, but it may, by this means, be cured in time, provided it be free from Callus and Caries. I have myſelf often cured theſe *Fiſtulæ* in ſo ſhort a Space as three Days, by this Practice: and have even found, by Experience, that this Method of ANELIUS will not prove altogether unſucceſsful, even in thoſe *Fiſtulæ* which have a ſlight Caries. By this Method I cured a Girl of ten Years old, in the Year 1727, of an inveterate *Fiſtula lacrymalis*, with a

2 ſlight

[a] M. GARENGEOT (in Cap. *De Fiſt. Lacrym.*) adviſes the Tube of the Syringe to be agitated, till you have introduced it into the lacrymal Sac; but this is not neceſſary; it is ſufficient you inſert it into the *Punctum lacrymale*, or the Beginning of the Duct.

ſlighter Caries, which I injected every Day for ſix Months: the Patient is at this Day well, and married.

Treatment of complete or open Fiſtulæ.

XVIII. In the perfect or complete Species of the *Fiſtula lacrymalis*, in which the external Skin is eroded or ulcerated, you may much more eaſily open the Paſſage of the occluded naſal Canal, than in the other Kind. For in this Diſorder you may readily paſs the forementioned Probe of ANELIUS, immediately through the *Canalis naſalis* right down into the Noſe, and that even with its largeſt End foremoſt, marked *b*, in *Fig.* 12. I have even ſeveral times opened the naſal Canal readily in this Species of the Diſorder, by the Probe marked K. in *Tab.* I. For deterging the Ulcer, and compleating the Cure, you muſt follow the Methods we have before propoſed; only inſtead of a Tent of Lint, you ſhould uſe one of Lead or Wax, and touch the *Canalis naſalis* every other Day cautiouſly with a conical Bit of *Lapis infernalis*; and, after healing up the external Lips of the Wound, uſe the Injections adapted to keep open the naſal Canal for a conſiderable Time. M. PETIT has ſometimes ſucceſsfully uſed a thick waxed Thread, to keep open the naſal Canal, inſtead of a Tent, as we are informed by M. GARENGEOT, in his Chapter on this Diſorder. But when you find the *Os Unguis* foul or vitiated, you muſt enlarge the Opening of your Ulcer, and remove the Caries, or perforate the Bone, as we before propoſed.

Fiſtulæ without Obſtruction of the naſal Canal.

XIX. In thoſe lacrymal *Fiſtulæ*, which have no Obſtruction of the naſal Canals, inſtead of probing, you muſt more frequently waſh out the offending Sordes by Injection. The beſt Injections in this Caſe are of the Decoctions of vulnerary Herbs, all mineral or medicated Waters; or *Aqua Calcis*. When you perceive the lacrymal Sac too much relaxed or diſtended, you muſt endeavour to recover its Tone by topical Remedies, as *Hungary* Water, *&c.* and the Lips of the Wound muſt be touched frequently with the *Lapis infernalis*; by which the relaxed Skin will be greatly ſtrengthened: you ſhould alſo apply the compreſſing Inſtrument repreſented in *Tab.* XVI. *Fig.* 20. or ſome other figured for the ſame Purpoſe by AQUAPENDENS, SCULTETUS, or PALFYN.

Callous and carious Fiſtulæ.

XX. But it muſt not be imagined, that the Method of probing and injecting, contrived by ANELIUS, will cure all lacrymal *Fiſtulæ* whatever: for in ſuch as are inveterate, and attended with an obdurate Callus, or a ſpreading Caries, this Practice will be to no Purpoſe: nor are we as yet furniſhed with Remedies ſufficient for the Cure of ſuch *Fiſtulæ*; though I can acquaint you, that *Archiater* BRUNNERUS aſſures me in a Letter, that he cured a lacrymal Fiſtula of the very worſt Kind by a mercurial Injection. It very often happens too, that the Flux of purulent Matter in this Diſorder cannot be leſſened, nor the naſal Canal kept open by Injection, ſo as to make a Paſſage into the Noſe, even though it may ſeem pervious to the Probe; of which I have known various Inſtances, without being able to account for the Cauſe. In theſe Caſes, therefore, if the Patient preſſes for a Cure, there remains but one Method of relieving him, and that is, by removing the Callus and Caries, and by making a new Paſſage, or an artificial naſal Canal into the Noſe. See N° XII. and XIII. preceding. Sometimes the Caries penetrates ſo far into the *Oſſa ſpongioſa* of the Noſe, that it is impoſſible for you to extirpate the ſame either by Remedies or the Cautery; though I muſt confeſs this to be a Caſe that never occurred in my own Practice. But even in the very worſt Caſes, the Diſorder may be pal-

liated, and the Patient much relieved, by making a Paſſage for the purulent Matter to run into the Noſe, which before diſcharged itſelf with great Uneaſineſs at the Corner of his Eye: and in theſe Caſes too you will find Injections of the greateſt Service.

Other new Methods of Cure.

XXI. We before obſerved, that, in imperfect *Fiſtulæ*, where the Skin is not eroded, you ought firſt to make an Inciſion through the Integuments before you perforate the *Os Unguis*. But, to render the Operation leſs formidable and ſevere, a certain Surgeon of *Hamburg* thought it beſt to perforate the Skin, Sacculus, and Bone at once, with an Inſtrument contrived for that Purpoſe, repreſented in *Tab.* XVI. *Fig.* 24. keeping open the new-formed lacrymal Ducts by a Tent, till the Wound was healed externally. Laſtly, as ſome of the Moderns have found, that the new naſal Canal formed by perforating the *Os Unguis*, does frequently fill up, or grow together, they have endeavoured to prevent it (by WOOLHOUSE's Direction) by inſerting a ſmall Tube of Lead, Silver, or Gold, *Tab.* XVI. *Fig.* 25. which is left there ever after, and the external Wound healed up over it, that the Paſſage may not afterwards cloſe up. In this Practice I have ſeveral times ſucceeded myſelf; but then I uſed a Tube a little larger than the common, as at *Fig.* 26. that the Tears might have a free Paſſage; healing up the Wound afterwards over the Tube.

LEMORIERE's Method.

XXII. We have ſtill another new Method of curing lacrymal *Fiſtulæ*, propoſed to the Royal Academy at *Paris*, by M. LEMORIERE[a]. He firſt opens the lacrymal Sac in the uſual Manner by a Scalpel, and then inſerts a particular Kind of ſharp-pointed and crooked Forceps, *Tab.* XVI. *Fig.* 29. A, with the Beak of which he breaks through the *Os lacrymale* into the Cavity of the Noſe. In the next place, he dilates the Perforation with the Forceps, *Fig.* 30. with which he further lacerates and breaks the *Os lacrymale*, and Membrane of the Noſe, to enlarge the Duct, ſo that it may not eaſily cloſe up again, which it is otherwiſe very apt to do. After removing the Forceps, he dreſſes the Wound for the firſt Days with Lint, and ſome digeſtive Ointment; but, on the third or fourth Day, he introduces a Bit of Wax-Candle into the new-formed Duct inſtead of a Tent, which ſhould be about the Thickneſs of a Straw, or one Line at leaſt in Diameter, made a little crooked, and armed with a ſmall Head, as at *Fig.* 31. A, B; this he continues in the Duct for the Space of thirty or forty Days, till the Parts are well formed; after which he removes the Candle, and heals the Wound: by which Method, he aſſerts, the Duct may be certainly kept open without any Danger of Concretions.

M. ST. YVES's Method.

XXIII. We have alſo another Method of curing theſe *Fiſtulæ*, given us by the famous Oculiſt ST. YVES of *Paris*, and deſcribed by SCHOBINGERUS, in a Treatiſe *De Fiſtula lacrymali Baſil. Ann.* 1730, as follows: firſt, he gently elevates and ſtretches the Skin at the greater Canthus of the Eye, as in opening a Vein, and then makes an oblique Inciſion with a Lancet, through the Integuments, and lacrymal Sac from the Eye-lids towards the Tendon of their orbicular Muſcle[b]; he next dilates the Wound by inſerting a Tent of prepared Sponge, and defends it with a Piece of Plaſter: the next Day, after removing the Dreſſings, he

[a] In *Memoir. Acad. Reg.* An. 1729. pag. 590. Edit. *Amſtel.*

[b] I ſuppoſe the Inciſions muſt be made from below upward; but it does not appear from this Deſcription.

he examines the State of the Wound and *Os Unguis* with a Probe, and by Injection; and is particularly careful in his Enquiry, whether the Bone be carious. This done, he ſupports the Patient's Head in a reclined Poſture with one Hand, while, with the other, he cautiouſly and obliquely perforatest he *Os Unguis* towards the Noſe with a triangular Probe, by the *French* called *Troicar*; in doing of which, great Care muſt be taken not to miſtake the *Os planum* for the *Os Unguis*, leſt, by perforating the firſt, you ſhould run into the *Antrum Highmorianum*, or elſe upon the *Apophyſis naſalis* of the *Os maxillare*. Add to this, that when the Apex of the Trocar has entered obliquely through the *Os Unguis*, you muſt then direct it betwixt the two *Laminæ* of the *Os ſpongioſum* in the Middle of the Noſe, that you may avoid injuring thoſe *Laminæ*, or any of the adjacent Parts. The Perforation thus made, the Surgeon now directs the Patient to breathe deep, and blow out the Air forcibly through his Noſe; that by the Exit of the Air and Blood through the Wound, he may judge whether the Perforation be rightly made. To dilate and keep the Paſſage open, he at firſt inſerts a Bit of Wood like a Wedge, and covers it with a Bit of Plaſter: but for the ſame Purpoſe, he afterwards dreſſes with Tents of Lint dipt in Cerate, which Tents he renews every third Day, gradually enlarging them, but never exceeding the Thickneſs of a Gooſe-quill; and afterwards he gradually diminiſhes the Thickneſs of the Tents before the Wound is quite healed [a]; by which means he aſſerts, that the foul Bones will caſt off and ſeparate ſpontaneouſly, without the Help either of actual or potential Cautery, and a new Paſſage will be formed for the Tears from the lacrymal Sac to the Noſe. If any Splinters or Aſperities of Bones offer themſelves in the Cure, they muſt be removed, Sinuoſities muſt be opened, and Ulcerations in the *Membrana Schneideriana* and lacrymal Sac deterged with *Lap. infernalis*, or other Eſcharotics. At every Dreſſing the Patient muſt cloſe his Noſtrils, and endeavour to force the Air through the new-formed Duct, to diſcharge the Sordes, and clear the Paſſage, which muſt be afterwards filled with a Tent dipt in Oil [b], and covered with a Plaſter; and when the Sides of this artificial *Canalis naſalis* appear conſolidated, the Tent is omitted, and the Plaſter only uſed till the external Wound is alſo cicatriſed, which, he ſays will generally be within the Space of ſix or eight Weeks; and, laſtly, towards the End of the Cure, when the Parts are near cicatriſed, you may inject ſome proper Liquor through the *Punctum lacrymale*, which, by paſſing into the Noſe, will demonſtrate whether you have rightly ſucceeded.

XXIV. With regard to the Method of curing lacrymal *Fiſtulæ* by probing and injecting, propoſed by ANELIUS, SCHOBINGERUS, in pag. 22. of his Diſſertation on this Subject, writes, that it is almoſt univerſally rejected, or forgot, becauſe it requires an uncommon Dexterity or Slight in the Adminiſtration thereof. I grant, indeed, it may be rejected, or forgot, by thoſe who are ignorant of the *Encheireſis* of the Operation, and Anatomy of the Parts: but, for my own part, it is my general Practice, and I find no Difficulty in it; though one would imagine, from the Deſcription SCHOBINGERUS gives of it, that he could ſcarce at all perform it, not being ſufficiently verſed in its *Encheireſis*. An Obſervation on the Method of ANELIUS.

XXV. It

[a] I queſtion whether it be abſolutely neceſſary to obſerve all theſe Circumſtances minutely.

[b] 'Tis the general Advice of Surgeons, never to apply Oil or Fat to injured Bones: and, as I can ſee no Reaſon why it ſhould be applied to theſe tender ones, I think it is ſafer to uſe a Tent dipt in *Sp. Vini rect.* or ſome Tincture, rather than Oil.

Some Errors of M. GARENGEOT.

XXV. It is alſo remarkable, that M. GARENGEOT, in his *Operations*, paſſes by this Method of ANELIUS with little or no Mention of it, as a Thing of no Conſequence; and, in his Treatiſe of *Inſtructions*, he deſcribes it ſo lamely, that one may be ſatisfied he never attempted or performed it. The Probe, too, which he figures for this Operation, is ſo ſlender and weak, and ſo ill-ſhaped towards its upper End, that one can never be able to open the naſal Canal by it. He likewiſe repreſents the End of the Tube for the Syringe ſo ſlender, that it muſt be impoſſible for it to have any Perforation or Cavity as it ought; beſides which, it will be apt, like a Needle to run into the Eye-lid itſelf inſtead of the Duct. Laſtly, he directs to uſe a *Speculum Oculi*, inſtead of the Fingers, to ſecure the Eye-lids in this Operation, which *Speculum* he figures double, ſo that the Operator will be more obſtructed than aſſiſted by the Inſtrument; when the whole Buſineſs may be performed with the greateſt Eaſe by the Fingers only, according to the Directions given by myſelf, and ANELIUS, for above theſe twenty Years paſt, and as I have above an hundred times performed it. In the next place, M. GARENGEOT writes, that the lacrymal Probe cannot be conducted into the naſal Canal, becauſe ("*le Detour eſt trop grand*") of the great Incurvation of the Paſſage to it; whereas the Probe may be thus conducted without Difficulty by one verſed in the Artifice, and acquainted with the Courſe of the Ducts. And ſo far is the Thing from being almoſt impoſſible, as he aſſerts it to be, that I readily performed it above twenty Years ago, barely after the Reading of ANELIUS's Account of it, without ſeeing it done by another. I muſt indeed own, that ſeveral Surgeons have, at Times, applied themſelves from *Hamburg*, aud other remote Parts, to me at *Helmſtadt*, to inſtruct them in the *Encheireſis* of this Operation, which they before thought impracticable, becauſe they had ſeveral Times miſcarried in it; but, after they had been ſhewn the Artifice a few Times by me, they found no Difficulty in performing it themſelves. I had once a Student in Divinity under my Care for a lacrymal Fiſtula, who, after having ſeen me paſs the Probe every Day for ſome Time through the *Punctum lacrymale* and naſal Canal into his Noſe, could, upon trying, eaſily perform the ſame himſelf by looking in a Glaſs; and became, at length, ſo expert in it, as to paſs it with more Nimbleneſs and Dexterity than I could myſelf: for by that Time you would imagine the Probe entering the lacrymal Punctum and Duct, he had ſlipt it alſo inſtantly through the lacrymal Sac and naſal Canal into his Noſe; which Proceſs he would repeat ſeveral times in an Hour, without any Difficulty or Uneaſineſs, and there leave the Probe, to keep the Paſſage open. I have been the more prolix on this Artifice, to refute the Impoſſibility of it, and demonſtrate M. GARENGEOT not only unſkilled in the Operation, but even ignorant of the chief Uſe of the Probes which he repreſents, when he ſays, they ſerve only to ſearch out the lacrymal Sac; whereas the chief Deſign of them is to open the obſtructed Cavity of the naſal Canal, in the watery Eye and lacrymal Fiſtula. Nor does the aforeſaid Gentleman ſo much as mention the Name of ANELIUS, the Inventor of theſe lacrymal Probes and Syringe; for what Reaſon I muſt leave others to judge. Conſult MORGAGNI, in *Adverſar. Anatom.* VI. 64.

XXVI. Nor muſt I omit mentioning here a Method of PETIT's; which is this: he makes an Inciſion in the lacrymal Sac; into which he introduces a grooved Probe, paſſes it into the Noſe, and, by this Means, opens the Canal.

Through

through the Groove in the Probe he admits a Wax candle (Fr. *Bougie*) to keep the Duct open: this Candle he changes once a Day, till he thinks the internal Surface of the Canal is perfectly cicatrised; and he uses it no longer. The Tears now pass, as usual, from the Eyes into the Nose, and the external Wound is closed in two or three Days: but I know by Experience, that this Method does not always succeed.

Authors divided as to the Treatment of these Fistulæ.

XXVII. From what has been said in this Chapter, it will manifestly appear that there are various Methods of treating lacrymal *Fistulæ*, according to different Authors, and the several Species of the Disorder; insomuch that there is not any one Operation in Surgery besides, in which Surgeons are less uniform or more unsettled in their Practice. You will find this Disorder considered more largely, with many other different, but less considerable Methods of treating it, in our professed Dissertation *De Fistula lacrymali*, *Altorf.* 1716.

The Author's Method of treating lacrymal Fistulæ.

XXVIII. It now remains for me to acquaint the Reader briefly with the Methods in which I myself usually treat these *Fistulæ*. And first, in the Beginning of the milder Species, I approve of the Method of probing and injecting, contrived by ANELIUS; which I usually continue for the Space of several Days or Weeks, according to the Nature of the Disorder, and especially when I perceive it diminish by this Practice. But when I find little Benefit result from it, I have Recourse to the Knife, with which I carefully lay open the Skin and lacrymal Sac, by an oblique or semi lunar Incision; then waiting till the Hæmorrhage ceases, the next Day I perforate the *Os Unguis* into the Nose, by the Instrument for this Purpose in *Tab.* XVI. *Fig.* 24. or *Tab.* XXIV. *Fig.* 2. In performing which, I observe the several necessary Circumstances, as I have before directed. After washing the Wound with warm Wine, I first fill the new-formed Duct with a Tent, and a Day or two after with a Piece of Wax-candle, or a leaden Plummet, about the Thickness of the Instrument at *Fig.* 21. A, dipped in some Balsam or some mineral Water, till the Canal is completely formed; to effect which the sooner, I now and then touch the Surface with a Stick of *Lap. infernal.* after the Tent or Candle is extracted; and in this Method I continue three or four Weeks, or longer. I next insert a small Cannula of Lead, Silver, or Gold, *Tab.* XVI. *Fig.* 15. from PLATNERUS, and heal up the Wound over it: but as the Bore of that Cannula often proves too small to transmit the viscid Juices of these Parts freely into the Nose, I generally prefer one that is a little larger, as at *Fig.* 26. which I insert, and heal up the Wound over it as before. The Tube thus left in the new-formed nasal Canal, is generally so far from being uneasy to the Patient, that I have known many who could not tell whether the Tube was left in or not, after their Cure was compleated. But to prevent any Obstructions, or other Accidents, towards the End of the Cure, the Day after I have closed the Lips of the Wound, I inject some *Decoct. Veronicæ* (or some mineral Water) several times every Day through the *Puncta lacrymalia* by the Syringe of ANELIUS, that the Tears may have a clear Passage to the Tube. I must indeed confess, that though these Tubes will generally very well suffice to convey the Humours into the Nose, yet, in some malignant *Fistulæ*, when the Tubes are not large, they do not answer their Intention, but leave the Patient molested with a watery Eye. I never yet used the actual Cautery for the Cure of these *Fistulæ*, and I really think it is hardly ever necessary, notwithstanding many Authors lay so great a Stress upon it: but on the contrary

contrary, I imagine the Baſis of the Cure to conſiſt in making an artificial naſal Canal ſufficiently large, by the Method here preſcribed; ſo that it may not eaſily be again cloſed or obſtructed. Even if you meet with a Caries in the *Os Unguis*, it may be very well removed without the actual Cautery. And laſtly, you may from hence conclude, thoſe perforating Inſtruments and *Canulæ*, which are too ſmall to make an ample Paſſage through the *Os Unguis* into the Noſe, not well adapted to ſucceed in this Operation.

Cautions. XXIX. I think it will not be improper to cloſe this Chapter, by giving the young Surgeon a few Cautions with regard to our preſent Subject. And firſt it will be neceſſary for him to keep the Patient's Body open with lenient Purges, eſpecially when he is to call in the Aſſiſtance of the Knife; not neglecting to open a Vein in plethoric Subjects, and to repeat it upon the Approach of inflammatory Symptoms after the Operation. 2. In Patients of an ill Habit, afflicted with theſe *Fiſtulæ*, the Juices muſt be corrected by the Uſe of alterative and evacuating Medicines before and after the Operation, eſpecially a Decoction of the Woods, and a mercurial Purge now and then. (3) If the lacrymal Fiſtula be attended with ſome other Diſorder, a Regard muſt be had to treat the latter with proper Medicines ſeparately. (4.) With regard to the Surgeon's Poſture for performing this Operation, I uſually do it ſtanding; but PLATNERUS performs it ſitting, almoſt in the Manner of couching a Cataract. *Diſſ. de Fiſt. lacrym.* pag. 41. (5.) The ſame Author directs (pag. 43.) to remove the *Perioſtæum* from the Bone in this Operation, alſo to divide and extirpate the lacrymal Sac by a tranſverſe Inciſion, after ſeparating it from the *Os Unguis:* but as I can ſee no Reaſon for this Practice, I never came into it, and yet I cured my Patients equally well; and therefore of two Evils, the leaſt is to be choſen. (6.) In order to cure the *Hernia* of the lacrymal Sac, PLATNERUS adviſes to open it with the Scalpel, and afterwards to heal it with *Balſ. de Mecha*, that the Sac may be contracted, and rendered firmer by the Cicatrix. I myſelf have ſucceeded in this Practice: but then, a few Days after the Inciſion, I touched the Lips of the Wound every Day with *Lapis infernalis*, and injected afterwards a Decoction of *Veronica cum pauxillo Sp. Vini.* (7.) In a Caries of the *Os Unguis*, PLATNERUS adviſes not to perforate it, but to burn it through into the Noſe by the actual Cautery, according to the antient Practice. But as this ſevere Practice is not attended with any Advantage, and as the Caries of the Bone may be removed by perforating it without Fire, I prefer the milder Method. (8.) In cutting theſe *Fiſtulæ*, M. GARENGEOT adviſes to divide the *obliquus inferior* Muſcle of the Eye, if it appears bare of its Fat: but as he gives no Reaſon for this Practice, which may be followed with dangerous Conſequences to the Eye, I think it ought to be rejected. (9.) The ſame Author aſſerts, that the new Perforation into the Noſe cannot be kept open, and that therefore the Tears will not have a Paſſage thither after the Operation: alſo, that the *Puncta lacrymalia* will be uſeleſs after the Operation: but, if this be compared with what has been here advanced, and tried by the Experience of myſelf and others, the Reader muſt naturally conclude that Gentleman to be but little verſed in Diſorders of the Eyes, which is alſo proved from his not mentioning what has been propoſed on this Subject by ST. YVES, WOOLHOUSE, and LEMORIERE.





Plate 16.


Fig. 1.
A
B
Fig. 2.
A
B
Fig. 3.
A
Fig. 4.
B
A
Fig. 5.
B
A
Fig. 6.
a
b
d
Fig. 7 & 8.
Fig. 9.
Fig. 10.
A
Fig. 16.
Fig. 17.
Fig. 18.
a
b
Fig. 19.
a
b
Fig. 13.
a
b
Fig. 11.
a
b
Fig. 12.
A.
Fig. 15.
B
D
Fig. 14.
A
Fig. 24.
F
F
F
E
C
B
Fig. 20.
A



Fig. 21.
Fig. 22.
Fig. 23.
Fig. 24.
Fig. 25.
Fig. 26.
Fig. 27.
Fig. 28.
Fig. 29.
Fig. 30.
Fig. 31.
Fig. 20.
A
B
C
D
F
a
b
c

Plate 16.



An EPLANATION *of the* SIXTEENTH PLATE.

Fig. 1. Repreſents an obtuſe pointed Hook, to draw the Eye-lids aſunder in ſome Operations: it was ſent me under the *french* Name *Hameçon plat*, or the flat Hook. A is the flat End; B the Handle.

Fig. 2. Repreſents the Needle A, fixed in a Handle B, for elevating and diſſecting the ſmall Blood-veſſels on the *Conjunctiva* and White of the Eye; as alſo to elevate and diſſect a *Pterygium*.

Fig. 3. Denotes a Beard of Rye or Barley, to make the Bruſh or Scarificator: in which A denotes the ſmall Hooks and Points which ſcarify the Blood-veſſels of the Eye.

Fig. 4. Is an Eye-bruſh compoſed of twelve or fifteen of the foregoing Beards: A the Handle; B the Part which ſcarifies.

Fig. 5. Is the Eye-raſp of CELSUS and AEGINETA, made in Shape almoſt like a Spoon: A the Handle; B the rough and convex Part, with which the Antients ſcarified the Eye-lids. This I received from M. MAUCHART. We have another a little different from this repreſented by PLATNERUS, *in Diſſert. de Scarif. Oculor.*

Fig. 6. Repreſents the left Eye; whoſe two *Puncta lacrymalia* are denoted by *a a*, and the lacrymal Caruncle betwixt them is marked *b*.

Fig. 7 and 8. Exhibit a View of the lacrymal Ducts, as they paſs from each Eye into the Noſe: *a a* the lacrymal Sac; *b b* the *Puncta lacrymalia*; *c c* the Ducts which lead from the two *Puncta* into the Sac; *d d* the naſal Canal; *e e* the Opening of the ſame Canal into the Noſe.

Fig. 9. Shews the Manner in which the before-deſcribed Ducts are ſituated and diſpoſed with regard to the Eye: *a a* the *Puncta lacrymalia*; *b* the lacrymal Caruncle; *c c* the Ducts which lead from the *Puncta* to the lacrymal Sac: *d* the ſaid Sacculus; *e* the *Canalis naſalis*; *f* the Aperture of it into the Noſe.

Fig. 10. Shews an *Anchylops*, and a *Hernia* or Diſtenſion of the lacrymal Sac.

Fig. 11. Is a very ſlender Probe of ſilver Wire, a little crooked, and armed with a ſmall Head or round Point, for opening and clearing the lacrymal Ducts and naſal Canal, when they are obſtructed in *Fiſtulæ*, or a watery Eye, as propoſed by ANELIUS.

Fig. 12. Is another Probe of the ſame Kind, and for the ſame Uſe, but ſtronger; which I uſe in more obdurate Obſtructions of theſe Parts.

Fig. 13. Is another Kind of Probe, which I now uſe for the ſame Intentions, but more conveniently, as it is ſhorter.

Fig. 14. Is a ſmall ſilver Syringe, as deſcribed by ANELIUS, to inject Liquors through the *Puncta lacrymalia*: A the Tube which enters the lacrymal *Punctum* and Duct; B the Handle of the Sucker; C, D, the hollow Cylinder.

Fig. 15. Is another ſmall Tube of a different Make, which may be adapted to the End of the Syringe by the Screw B.

Fig. 16 and 17. Demonſtrate the ſeveral Ways in which the lacrymal Sac may be diſtended or relaxed.

Fig. 18.

Fig. 18. Shews how an Abceſs or Tubercle may be formed, ſo as to deſtroy the lacrymal Duct; *a* that upon the upper Duct, *b* one upon the lower Duct, like that which I ſaw in the Duke of SAVOY.

Fig. 19. Repreſents a complete lacrymal Fiſtula: *a* one with a pretty large Opening; *b* one with a narrow Opening; the Line *b c* denotes the Courſe for Inciſion in theſe *Fiſtulæ*.

Fig. 20. Is a ſteel Inſtrument for compreſſing the lacrymal Sac, from PLATNERUS: A the Bolſter which is impoſed on the lacrymal Sac; B the Hinge; C the Screw which preſſes the Bolſter on the Sac; D the upper Part which goes over the Forehead; E a Hook which goes into the Holes of the Strop, to ſecure the whole upon the Head.

Fig. 21. Is an Iron Cautery, for perforating the *Os lacrymale*.

Fig. 22. A Cannula adapted to the preceding Cautery, to be fixed upon the Bone before the Cautery is applied.

Fig. 23. Repreſents an Inſtrument made of Silver or Braſs; which in the Part marked *a* is made hollow like a Spoon, to cover and ſecure the Eye, while the Cautery is paſſed through the Aperture *b* to the carious Bone; *c* the Part which ſerves for a Handle. This may alſo ſerve to cover the Eye when you cut for the *Fiſtula lacrymalis*.

Fig. 24. Repreſents an Inſtrument for perforating the Integuments, lacrymal Sac, and Bone, at the ſame time; or you may only perforate the Bone with it, after the lacrymal Sac is opened by Inciſion. A the Point; B the Handle.

Fig. 25. A B denote ſmall Tubes to be inſerted into the Perforation of the *Os Unguis*, according to WOOLHOUSE and PLATNERUS, and to heal up the Wound over it.

Fig. 26. Is a Tube of the ſame Kind, but a little larger; which I uſe for the ſame Purpoſe, and may be beſt made of Lead or Gold.

Fig. 27, 28. Are ſilver Tubes uſed by PLATNERUS, to keep open the new-made Paſſage to the Noſe, till it is become callous or cicatriſed.

Fig. 29. Repreſents the Forceps of LEMORIERE: A the ſharp-pointed and crooked Beak, which perforates the *Os Unguis*; B B its Handles, by which you open and ſhut its Beak.

Fig. 30. Repreſents the Head only of the ſame Forceps, opened as it is when you dilate the Parts, after perforating the *Os lacrymale*.

Fig. 31. Denotes the Shape of the Piece of Wax-candle, which LEMORIERE uſes inſtead of a Tent, to keep open the Perforation to the Noſe: A its Head; B that End which goes into the Noſe.

Of

Of OPERATIONS on the EYES.

CHAP. LVI[a].

Of SUFFUSIONS *or* CATARACTS.

I. AFTER having conſidered the Diſorders of the Parts adjacent, we come now to thoſe of the Eye itſelf; the chief of which is that termed a *Suffuſion* by the Antients, and a *Cataract* by the Moderns. The *Greeks* call it *Hypochyma* and *Hypochyſis*; the Deſcription of which Diſorder has been very imperfect, till of late. We deſcribe a Cataract or Suffuſion, with the Generality of Oculiſts, to be a Diſorder of the Humours in the Eye; by which the *Pupilla*, which ought to appear tranſparent and black, looks opaque, and of ſome other Colour, as inclining to white, grey, blue, brown, *&c.* And thus Viſion is variouſly impeded, or totally deſtroyed. A Cataract deſcribed.

II. It is remarkable that the Generality, and even the moſt eminent Surgeons and Phyſicians, have been all along greatly deceived, till within the preſent Century, both as to the Seat and Cauſes of the Cataract; moſt of them believing it to be a Pellicle, or membranous Subſtance, formed always in the aqueous Humour: whereas the moſt expert Surgeons and Oculiſts have of late Years found, that, by repeated Diſſections of the Eye thus diſordered, there is hardly ever any white Membrane or other foreign Subſtance to be found in the aqueous Humour; but that it is almoſt conſtantly an Opacity in the cryſtalline Lens; and therefore the true and common Cauſe of a Cataract is, according to myſelf and the reſt of the Moderns, an Opacity of the Cryſtalline, and not any thing in the aqueous Humour, as the Antients ſuppoſed. Indeed the Antients might have been led into this Error very eaſily, from the Appearance which the Diſorder affords, without diſſecting the Eye: for, by barely inſpecting that diſeaſed Organ, the opake Cryſtalline looks like a Membrane in the aqueous Humour; by couching or depreſſing which, with a proper Inſtrument, the Eye recovers its former Viſion. This is confirmed by various Obſervations and Experiments made by ſeveral eminent Members of the Royal Societies at *London* and *Paris*, and the *Commerc. Literar. Norimberg*; and may be ſeen, conſidered more at large, in our profeſſed Treatiſe *De Cataracta, Glaucomate,* Cauſes according to the Antients.

[a] M. GARENGEOT here is much to be commended; that is, his Treatiſe of *Chirurg. Inſtruments, Tom.* I. *Cap.* XIV. *p.* 414. He laments and inveighs againſt the Negligence of Surgeons in leaving the Operations on the Eye wholly to Mountebanks and Strollers; and exhorts them earneſtly to vindicate this moſt noble Branch of Surgery to themſelves (though at preſent few are furniſhed even with proper Inſtruments) and to beſtow due Pains and Attention upon it. Yet I cannot but wonder, that amongſt all his *Chirurgical Operations*, this very Gentleman ſhould deſcribe in his Writings, or point out to his Pupils, but one relating to the Eye; which is the Operation performed in the *Fiſtula lacrymalis*. This confirms me in what I hinted in the foregoing Chapter, that he was not very converſant in the Cure of theſe Diſorders.

Glaucomate, & Amauroſi, An. 1713. and in our Apology for, and our Vindication of, the ſame, *An.* 1717 and 1719[a].

The firſt Diſcoveries of the true Cauſe.

III. It is almoſt eighty Years ſince the preceding Error of the Antients, with regard to the Cauſe of Cataracts, began to be publickly remarked by M. QUARE, ROLFINCKIUS, GASSENDUS, RAUHAULT, BORELLI, and others. But theſe Gentlemen having but few Obſervations to eſtabliſh their truer Notion of the Diſorder, their Obſervations were not only thought, by the Generality, to be anomalous, but even the old Error, of Cataracts being conſtantly formed by a Membrane, ſtill prevailed; and the rather, becauſe there were few or none who took the Pains to diſſect any Eyes affected with this Diſeaſe. But at length M. BRISAC and MAITRE-JEAN, by new Experiments and Diſſections of Eyes thus affected, demonſtrated apparently, that Cataracts aroſe not from any Membrane, but an Opacity of the cryſtalline Lens[b]. But though theſe laſt Gentlemen were much miſtaken, in thinking themſelves the firſt Propoſers of this Diſcovery; yet their Merit is not inconſiderable, for having more carefully proved, and demonſtrated by inconteſtable Obſervations and Experiments, what had been ſtarted by their Predeceſſors, and at that Time almoſt buried again in Oblivion. For, to ſay nothing of myſelf, the whole Drift of the Eſſays and Obſervations on this Subject, given us by the Learned in *France*, *England*, and *Italy*, tends largely to prove, that the ordinary and *moſt common Cauſe* of Cataracts is from an Opacity of the cryſtalline Lens.

The Author's Opinion.

IV. I ſay only the moſt common Cauſe of Cataracts is from an Opacity of the Cryſtalline, without abſolutely denying, as ſome do[c], that a membranous Subſtance may be ſometimes formed in the Eye, ſo as to cauſe the like Diſorder. I rather recommend this Point to be decided by farther Obſervation and Experiments. For though when I firſt wrote on the Cataract I was furniſhed with five Obſervations of my own, beſides thoſe of BRISAC and MAITRE-JEAN, in which an Opacity of the Cryſtalline appeared to be the ſole Cauſe; yet I even then entertained an Opinion, and afterwards declared it, that I thought a Membrane, or other ſolid Body, floating in the aqueous Humour, might ſometimes alſo cauſe a Cataract, as I once obſerved in diſſecting a recent Subject. Nor has this Caution of mine turned out uſeleſs to others, ſince I received a Letter from Profeſſor WIDEMANNUS, Director of the *Acad. Natur. Curioſ.* which aſſures me, he found and demonſtrated ſuch a Membrane to ſeveral eminent Phyſicians of *Norimberg*, as LOCHNER, THOMAS, and GOCKELIUS, in both the Eyes of a Woman who had Cataracts: but then he at the ſame time obſerved in one Part of the Cryſtalline an incipient, and in the other Part a complete Opacity. After the Operation, which was performed three Years before the Woman died, ſhe became quite blind of that Eye whoſe Cryſtalline was wholly opake; and with the other Eye, whoſe Cryſtalline began to be obſcured, ſhe could only diſcern and diſtinguiſh large Objects. A Caſe much like this LANCISI tells me

[a] PLATNER, a Pupil and Friend of WOOLHOUSE (in *Program. Anatom.* 1736. relates that in the Eye of a Female Patient who had a Cataract) he found no Pellicle or membranous Subſtance; but the Cryſtalline Lens was opake, and leſs than uſual. So that even WOOLHOUSE's Pupils agree with me, that this Diſorder is not properly called a *Glaucoma*, but a *Suffuſion*.

[b] This is the Opinion received and defended by the preſent *Engliſh* Oculiſt, TAYLOR, in his Pamphlet on the Cataract, *Lond. An.* 1736.

[c] Among theſe I am reckoned as one by TAYLOR, in *Page* 5. of his ſaid Pamphlet; but unjuſtly, ſince my Writings on the Subject demonſtrate the contrary.

me he obſerved in GARELLI, *Archiater* to the Emperor, upon diſſecting whoſe Eyes he found a whitiſh Membrane in each, floating in the aqueous Humours: but then here again the Cryſtallines were yellowiſh and ſomething obſcure, though his Eyes had never undergone any Operation while he lived. Thus theſe Membranes ſeem generally attended with a Diſorder of the Cryſtalline. From theſe and a few of the like Obſervations, it appears, that a Cataract may ſometimes be cauſed by a Membrane in the aqueous Humour, though generally and moſt frequently from an Opacity of the cryſtalline Lens.

Aſſertors of membranous Cataracts refuted.

V. Though an Opacity of the cryſtalline Lens appears, from Obſervation and Experiment, to be the common and moſt frequent Cauſe of Cataracts; yet it had been denied by ſeveral[a], many of which have no other Reaſon to offer, than that they think it very extraordinary, and almoſt impoſſible, that ſo many eminent Phyſicians, and profeſſed Oculiſts, ſhould have been thus miſtaken, for ſo many Ages, in judging it to proceed from a Membrane. Others think the Method of curing this Diſorder, by couching or depreſſing the cryſtalline Lens, is ſo ſevere and dangerous an Operation, that it muſt inevitably deſtroy the whole Sight of the Eye, becauſe they judge the Cryſtalline to be abſolutely neceſſary for Viſion. But how egregiouſly theſe are miſtaken, may appear from the ſingle Inſtance of the expert Anatomiſt WENCKERUS; who found both the Cryſtallines at the Bottom of the Eyes many Years after he had couched; the Patient, in the mean time, enjoying his Sight very well, eſpecially with one Eye, even to his Death, when they were diſſected. A like Obſervation we have, given us by BENEVOLI, firſt ſeparately, *Florent. Anno* 1722, and afterwards joined to a Treatiſe *De Caruncula in Urethra*; to which add the ſeveral Experiments made by the *French*, mentioned long ago in my Treatiſe on the Cataract. There are ſome again who, being fond of cavilling about Words, contend that ſuch an Opacity of the Cryſtalline ought rather to be called a *Glaucoma* than a Cataract; but with no more Reaſon on their Side than the former. This Diſorder of the cryſtalline Lens affords the ſame diagnoſtic Symptoms, and is cured by the ſame Practice with what has all along obtained among the Antients in their Suffuſion or Cataract; and therefore this Diſorder really is, or at leaſt deſerves the Name of their Cataract. On the contrary, we find that a *Glaucoma* is all along deſcribed by the moſt expert Surgeons and Phyſicians, as a Diſeaſe which very ſeldom happens, and which is wholly incurable. There are other frivolous Objections ſtarted, which the Reader may ſee refuted more at large in our Treatiſe on the Subject, with the Apology for the Vindication of it. We therefore aſſert, that a Cataract is hardly ever cauſed by any Membrane, or other Body floating in the aqueous Humour; becauſe it appears from Experience, that out of fifteen Patients, you ſhall hardly find one Cataract cauſed by a Membrane, all the reſt proceeding from an Opacity in the cryſtalline Lens. And conſequently we may depend on what has been advanced by the moſt expert Surgeons

[a] We have a Diſſertation *De Cataracta*, publiſhed in 1721, at *Straſburg*, by FREYTAGIUS; in which he aſſerts the general Cauſe of Cataracts to be a Membrane in the aqueous Humour; but, inſtead of proving it anatomically, he would perſuade us, he had ſeen his Father extract ſuch Membranes with a Hook above an hundred Times: but few will believe him, who know any thing of the Diſorder, and what has been advanced concerning it by others.

geons in *France*[a], *England*[b], and *Italy*[c], viz. *that the common Cauſe of Cataracts is not any Membrane, but an Opacity of the Cryſtalline*, notwithſtanding what others may ſay to the contrary.

Diagnoſis. VI. From what has been ſaid, it will be no difficult Matter to diſtinguiſh a Cataract from the reſt of the Diſorders of the ſame Organ. For, 1. it differs from an Amauroſis, or *Gutta Serena*, which ſome call the black Cataract; becauſe in this laſt the Eye loſes the Sight without any viſible Diſorder in the Eye, or any Change in the Appearance of its Pupilla. 2. An *Albugo*, or white Speck in the Eye, is not behind the Cornea and Uvea, as is the Cataract, but in the *Cornea* itſelf. 3. The *Ungula*, or *Pterygium*, is a preternatural Tunic without-ſide the *Cornea*. 4. The *Hypopyum* is indeed ſeated behind the *Cornea* in the aqueous Humour; but then it conſiſts of a purulent and fluctuating Matter; whereas the Cataract is a ſolid Subſtance. 5. A *Glaucoma* does indeed appear in a great meaſure like a Cataract, ſo as to deceive many, if they do not conſider that though both of them are ſeated behind the Pupilla, yet the *Glaucoma* being in the vitreous Humour, lies deeper than the Cataract, whoſe ſeat is in the Cryſtalline: therefore the firſt will generally appear of a darker blue, or a grey Colour, as its Name imports; whereas the Cataract uſually appears of a Pearl Colour, and ſeated immediately behind the Pupilla: add to this, that it has been conſtantly obſerved by Phyſicians, that the *Glaucoma* very rarely happens in compariſon with the Cataract; and when once it is formed, there is no Poſſibility of removing it, which cannot be ſaid of the opake Cryſtalline.

Species of Cataracts. VII. Cataracts have been diſtinguiſhed by Surgeons and Oculiſts into various Species. As, 1. By the Time of their ſtanding, into *recent* and *inveterate*. 2. By their Growth, into *incipient* and *confirmed*. 3. Into *mature*, when the Pupilla is totally obſcured; and *immature*, when the Pupilla being but partly obſcured, the Patient is as yet capable of perceiving Objects. Some Cataracts never come to Maturity, or at leaſt but very ſlowly. 4. According to the Symptoms, Cataracts, are again diſtinguiſhed into *ſimple* and *complicated*; the latter being when the Cornea, Uvea, or vitreous Humour are alſo affected, or when the Pupilla is immoveable, too much contracted, or adheres to the adjacent Parts. Sometimes there is a Tabes of the Eye attending it; and at other Times it is joined with ſome Diſorder of the Retina, or optic Nerve. 5. Cataracts are generally immoveable, but ſometimes they tremble or fluctuate upon touching the Eye with the Finger, being then called a *ſhaking Cataract*. 6. Almoſt all of them are of different Shades, though they approach nearly to the ſame Colour, to wit, that of Pearl, whitiſh, or grey, and are accordingly denominated *white* or *grey Cataracts*. We do not frequently meet with Cataracts of a yellow or greeniſh Colour, and ſeldom with any marbled, or looking like Cheeſe, or like a glowing Iron. 7. In ſome Cataracts the cryſtalline Lens degenerates into a milky Fluid, and in others into a purulent Matter, like that of Abceſſes; and in couching theſe, the Matter will eſcape, and confuſe the Humours of the Eye upon breaking the Capſule of the Cryſtalline with the Needle. And hence again we have a Diſtinction of Cataracts into *milky* and *purulent*.

[a] M. PETIT and MORAND, in *Hiſt. Acad. Reg. An.* 1722, 1723, and ST. YVES of *Paris* in his Book on *Diſeaſes of the Eyes*, Chap. *on the Cataract*.

[b] Mr. CHESELDEN, and others, in *Phil. Tranſact.*

[c] As MORGAGNI, SANTORINI, COCCHUS, BENEVOLUS, &c.

purulent. 8. Cataracts are again uſually diſtinguiſhed by Oculiſts into *true* and *ſpurious:* by the firſt, we mean one in which the Opacity appears immediately behind the Pupilla; but the ſpurious is, when the Opacity ſeems to be ſeated otherwiſe. Laſtly, 9. Cataracts are not undeſervedly diſtinguiſhed into curable and incurable; for thoſe of a grey or whitiſh Colour are the moſt eaſily cured. To theſe we may add ſuch as have no Colour, the Patient being ſenſible of Light and Darkneſs; alſo thoſe, in which the Pupil does not adhere, but can contract and dilate itſelf. On the contrary, you can have no great Hopes of curing complicated or fluctuating Cataracts, in which the Patient can neither perceive Light nor Darkneſs, and in which the Pupilla or Uvea adheres, is immoveable, and either contracted or dilated; or when it appears of the unuſual Colours at 6. and 7. preceding. We find ſome again diſtinguiſhing Cataracts into *common* and *uncommon.* By the firſt, they intend ſuch Opacities of the Cryſtalline as appear of a whitiſh Colour; and by the laſt, they mean thoſe of any other Colour; which indeed differ very remarkably from the former, in appearing not convex, like them, but flat or concave, as we have lately obſerved ſome, and as I find it alſo remarked by the accurate Oculiſt M. ST. YVES, in his Treatiſe on the Diſeaſes of this Organ.

VIII. We before demonſtrated, that the common and uſual Cauſe of Cataracts is an Opacity of the cryſtalline Lens, and hardly ever a looſe Membrane. But to explain the Manner in which the Cryſtalline becomes thus obſcured, we muſt conſider, that when the Juices are too thick and glutinous to paſs freely through the very minute ſerous Veſſels of this Body, they ſtagnate and obſtruct thoſe Veſſels, which become afterwards contracted and dried. Thus it may be formed in various inflammatory Diſorders of the Head and Eyes; and particularly after ſome external Violence has injured that Organ, as a Fall, Blow, Burn, *&c.* or expoſing the Eyes too much to the Heat of the Summer Sun, or an intenſe Fire. *Cauſes.*

IX. The principal Sign of a Cataract is, therefore, a ſmall Cloud, or whitiſh Opacity of the Cryſtalline. To ſatisfy your Patient whether it may be cured by couching, you ought to be firſt well aſſured, whether it be of the mature or immature Kind; for if it be of the latter, the Operation will be abſolutely miſchievous. The Signs of a *mature* Cataract, fit for couching, are, when the Pupil having loſt its native Blackneſs appears moveable, and equally of a duſky Hue, the Patient being ſenſible of Light and Darkneſs, but incapable of diſtinguiſhing Colours. On the contrary, you may judge it to be *immature*, if the Opacity is not equally ſpread behind the Pupil, the Patient being as yet able to ſee Objects imperfectly, eſpecially upon turning his Back to the Light. But, if the Patient can neither diſcern Light nor Darkneſs, it is a Sign the Retina or optic Nerve is greatly affected, and that the Diſorder is an *Amauroſis*, or *Gutta Serena*, for which no Cure can well be expected. You may alſo diſcover whether the Pupil adheres to the Cataract, and is become rigid, by obſerving whether it contracts or dilates itſelf in a ſtrong Light, or in the Dark; alſo if it does not move upon rubbing or touching the Eye with your Finger. If any ſmall Specks appear behind the Pupil, ſome Parts of the Cryſtalline are either inſpiſſated, or elſe ſome minute Pellicles are ſprouting from the Uvea, as I remember to have ſeen, and which may poſſibly unite into a Membrane. Sometimes only the Middle, the Margin, or elſe one half of the Cryſtalline is become opake; and *Diagnoſis.*

in the firſt Caſe, Objects will ſeem to the Patient to be perforated in the Middle. If any Tunic appears plain or convex within-ſide the Pupil, it denotes the Surface of the Cataract, as ST. YVES obſerves.

Prognoſis. X. There is ſcarce any Diſorder, the Event of which is more uncertain than that of the Cataract: which will ſometimes admit of a Cure, and ſometimes not. But, to ſay the Truth, Medicines will generally have little or no Effect, when the Diſorder is confirmed or inveterate; notwithſtanding what ſome may boaſt of their wonderful *Arcana* for this Purpoſe[a]. Almoſt the ſole Relief is therefore to be had from the Surgeon's Hands and Inſtruments. We very rarely meet with Inſtances of this Diſorder being cured by leaving it to Nature alone. And yet, by the Operation itſelf, a Cataract that bids faireſt for Recovery, though treated in the moſt judicious Method, ſhall frequently be the worſe for it; when one that ſeemed to be irrecoverable, ſhall be cured by the ſame Treatment, beyond all Expectation. However, a Cataract is much milder and more tolerable to the Patient than many other Diſorders which we eſteem deſperate and incurable: becauſe neither the Diſeaſe nor the Operation are uſually accompanied with intenſe Pain, nor Hazard to the Patient's Life. But, in the general, thoſe Cataracts are moſt likely to be cured, which are mature and not complicated, the Patient being capable of diſtinguiſhing Light and Darkneſs, and the Pupil retaining its natural and free Motion. But there can be little Hopes of ſucceeding in thoſe where the Pupil is rigidly contracted, the Uvea firmly attached to the Cataract; or where the Pupil, having loſt its natural round Figure, is lacerated, angular, and variouſly diſtorted. The Succeſs of the Operation is rendered ſtill more doubtful, if the Patient is weak, aged, or afflicted with a violent Head-ach, or when the Eye is too much ſhrunk up, or enlarged and ſwelled. The Cataract is alſo the worſe, as it degenerates more from the Pearl Colour; for the moſt unuſual Colours always proceed from and denote the worſt Affections of the Eyes. Yet even many of theſe are often cured by the Operation beyond Expectation, when the Eye is free from other Diſorders. For the milky and purulent Cataracts, though there is Danger of the opake Matter mixing with the aqueous Humour in the Operation, ſo as to render the Succeſs of it doubtful; yet it has been often obſerved by the moſt expert Oculiſts, that this Matter will ſubſide to the Bottom of the Eye, and the Humours recover their former Clearneſs[b]. It is indeed difficult to couch a variegated or marbled Cataract, as being too ſoft, and not yet arrived to a due Conſiſtence. Therefore when this Species does not give way to Remedies, you ought to defer the Operation till the whole Pupil appears opake, which denotes the Cataract to be ſufficiently mature. The Diſorder has been judged the more difficult to cure, as it is more inveterate, by the antient Surgeons and Phyſicians; and yet it has been obſerved by ſome of the modern Oculiſts, that Cataracts, without other Diſorders in the Eyes, may be often cured, though of twelve, eighteen, or even thirty Years ſtanding, if the Eye is free from other Diſorders. If the Patient cannot diſtinguiſh Light and Darkneſs, the Operation

[a] HOVIUS audaciouſly aſſerts (*in Lib. De circulari Humor. in Oculis Motu*, pag. 122.) that he can thus, at any Time, cure all Sorts of Cataracts, whether recent or inveterate. But, upon the ſtricteſt Enquiry into the Truth of the Matter, I can meet with no Inſtances of his Succeſs.

[b] See my Treatiſe on the Cataract, *p.* 255. See MAITRE-JEAN, Lib. *De Morb. Oculor.* Cap. *De Cataracta.*

tion will be but of little more Service than for removing the Deformity of the Eye; becauſe then the Cataract is accompanied with an *Amauroſis*, or *Gutta Serena*. In Infants the Operation is generally leſs ſafe, and more impracticable, than in Adults, by reaſon of their Impatience and Strugglings. Nor ſhould the Operation be performed on thoſe who have a Cough, Catarrh, Defluxions, and Vomiting, before thoſe Diſorders are firſt removed; leſt, by the Patient's being diſturbed in the Operation by thoſe Symptoms, his Eye might be irrecoverably injured and ſpoiled for the future. In thoſe Cataracts which move or fluctuate from one Side to the other, there is generally little or no Hope of the Operation ſucceeding; but when the opake Body appears before the Pupil, it may then be ſometimes extracted through an Inciſion in the Cornea.

Treatment of doubtful Cataracts.

XI. When the Cataract appears even deſperate or incurable, I think it is better to attempt to reſtore the Patient's Sight by the Operation, though in vain, rather than leave him to certain Blindneſs without uſing the beſt Means; and this the rather, becauſe the Operation may be performed, without inducing intenſe Pains, or endangering the Patient's Life; which are indeed Reaſons ſufficient to deter moſt People from Lithotomy, and the more ſevere chirurgical Operations. When the Patient is blinded by the Cataract, he cannot be blinded again by the Operation, if it does not ſucceed. The leſs Proſpect there is of curing the Diſorder, the more Honour and Fame will the Operator acquire, by recovering the Patient's Sight beyond all Expectation.

Of the Glaucoma and Gutta Serena.

XII. Surgery can be of little or no Service towards the curing of a *Gutta Serena*, as hath been hitherto univerſally allowed; till of late, the *Engliſh* Oculiſt TAYLOR has given out, that he can cure it by an Operation. The Falſity of this, Experience has demonſtrated. The Diſorder we now ſpeak of, is not ſeated in the anterior or middle Part of the Eye, but either in the Retina, the optic Nerve, or in the Brain itſelf, to which Parts no Operation can be extended. If there is any Room left to expect a Cure, it will be more reaſonable to attempt it by ſuch internal Medicines as will raiſe a Salivation, and purge; adding at the ſame time Phlebotomy, Scarification, and Setons or Iſſues, eſpecially thoſe on the coronal Suture, or in the Neck. What we have ſaid of the *Amauroſis*, or *Gutta Serena*, holds true in a worſe Degree of the *Glaucoma*; which being an Opacity of the vitreous Humour, is univerſally allowed, both by the antient and modern Surgeons, to be incurable by any Operation whatever. It is remarkable, that this vitreous Humour is ſometimes ſo much indurated, as well as diſcoloured, that it reſembles a Cartilage; as appears from an Obſervation formerly communicated to me by the celebrated Anatomiſt and *Archiater* LANCISI.

The two Methods of curing Cataracts.

XIII. There are chiefly two Methods of curing Cataracts; either by couching with the Needle, or by the Uſe of internal and external Remedies. It is true, there are ſome who reject all Methods of treating Cataracts by Medicines as uſeleſs and trifling; yet I think there are ſome Caſes in this Diſorder which ought to be recommended to the Care of the Phyſician. Nor are there Inſtances wanting, as well among the Moderns as Antients[a], of Patients, who, by the Help of Nature, aſſiſted with Medicines, have been freed from Cataracts beyond all Expectation; eſpecially when the Diſorder is incipient, and not firmly rooted or fixed in the cryſtalline Lens. But leaving the Phyſician to direct a proper

[a] *Vide* CELSUS *Lib.* VI. *Cap.* 6. and the modern Writers on the Diſorder.

proper Regimen and Courſe of Phyſic adapted to the Patient's Habit, Age, and other Circumſtances, we ſhall here proceed immediately to deſcribe the Methods of curing Cataracts chirurgically, by the Help of the Hands and convenient Inſtruments; of which CELSUS has treated with great Accuracy.

Surgeons adviſed to be diligent in learning this Operation.

XIV. But firſt it may be proper for us to admoniſh Surgeons to make themſelves better acquainted with the Operation for couching Cataracts, and to be more converſant in the Practice thereof, and not to leave the Buſineſs to Quacks and itinerant Pretenders, as we have ſeen it done but too much of late[a]. If the Practice is, as we ſee often, well enough executed by theſe boaſting Pretenders, what might we not expect from the Hands of the more prudent and regular Surgeon, were he to engage more in this Practice? which is, in reality, attended with leſs Danger or Hazard than the common Operation of Phlebotomy: for, in couching a Cataract, you run no Riſque of wounding a Nerve, Tendon, or Artery, as you do in opening a Vein. But leſt our Reader ſhould think we are recommending the Operation, for its Eaſineſs, to the Practice of every one, though ever ſo unſkilful; we ſhall here enumerate the ſeveral neceſſary Qualifications for an Oculiſt, whom we may venture to truſt in the Cure of this Diſorder. 1. He muſt be very well verſed in the anatomical Structure, and in the Functions of the ſeveral conſtituent Parts of the Eye, that he may avoid injuring any of them ignorantly. 2. He muſt be well acquainted with the beſt Inſtruments and Methods of operating, to be learned from a frequent and cloſe Attention to the Practice of ſome expert Maſter. 3. His Mind muſt be intrepid, his Hand ſteddy, and his Eye ſharp and quick-ſighted. 4. He ſhould be equally ready with his Left as with his Right Hand; that he may couch the left Eye with his Right-Hand, and the Right Eye with his Left Hand. 5. He muſt have made himſelf previouſly expert in the Practice, by repeated Trials upon the Eyes of Brutes, and of dead Men, before he ventures to couch the Eyes of the Living.

The Time of Couching; and previous Preparation of the Patient.

XV. But, in order to the more ſucceſsful and eaſy Performance of this Operation, it will be previouſly neceſſary for the Surgeon to appoint the moſt convenient Time, and to prepare his Patient in the beſt Manner, by a proper Regimen and Medicines. With regard to the firſt, ſuch a Seaſon ſhould be choſen, in which the Air is pretty temperate as to Heat and Cold, as in Spring and Autumn. The Day appointed for the Operation ſhould eſpecially be ſerene and clear, and the Hour generally in the Forenoon; not but the Afternoon will do very well, and may be, in ſome Caſes, preferable for weak and timorous Patients, who are uſually in better Spirits after a moderate Dinner. The Apartment for couching the Patient in will be fitter as it is lighter, provided the Sun does not ſhine in upon you; for ſo ſtrong a Light as the Sun's Rays will cauſe the Pupil to contract itſelf, ſo that you cannot have ſo large a View of the Parts and Inſtrument within the Eye. As for the Preparation of the Patient, he ſhould not only obſerve a proper Regimen and Diet a few Days before the Operation, but he ſhould alſo in that Time take ſome alterative and evacuating Medicines, with the Uſe of Phlebotomy, to prevent the Eye from being moleſted by intenſe Pain, Inflammation, Suppuration, and perhaps a Loſs of the whole,

[a] It is a little extraordinary, that M. GARENGEOT ſhould take no Notice of this Operation in his Treatiſe, as if it made no Part of Surgery.

whole, after the Operation has been performed[a]. It may alſo be generally convenient to give the Patient a Clyſter, if he has not eaſed himſelf lately; and, that his Courage may not fail him, the Operator ſhould take Care that he may have ſome Gravy Soop, or other ſtrengthening Sippings in the Morning, before he begins his Operation. Laſtly, nothing can more conduce to the Patient's Recovery, and the Prevention of Accidents, after the Operation, than to procure him a ſound Sleep afterwards by an anodyne Draught or Emulſion; by which the Faculties both of his Body and Mind will be recruited, and the lately ſuppreſſed Cataract will not be apt to aſcend again.

Of the Aſſiſtants and Needles.

XVI. The Surgeon ought never to undertake the Operation by himſelf, but to provide two Aſſiſtants, one to hold the Patient's Head (as in *Tab.* XVII. *Fig.* 1. A.) and the other, to adminiſter the Needle and other Neceſſaries. But he muſt be more particularly provided with *couching Needles*, and with a *Speculum Oculi*. Of the *Speculum* you have two Forms at *Fig.* 15 and 16, and of the couching Needles there are a great many Kinds, the chief of which are repreſented in *Tab.* XVII. *Fig.* 2, 3, 4, 5, 6, 7, 8, 9, 10, and 11. The beſt of them are, in my Judgment, thoſe at *Fig.* 5, 6, and 10. All have a little broad and ſharp Point like a Tongue or like a Barley-corn, but flatter; and that at *Fig.* 6. with a Sulcus in its Point, ſeems better adapted to couch the Cataract, than any of thoſe which have either a narrower or a broader Point. For thoſe with too ſlender a Point, as in *Fig.* 2 and 4. do eaſily lacerate the Cararact; and thoſe with a more obtuſe Point, as in *Fig.* 8. meet with Difficulty in perforating the Coats of the Eye. For theſe Reaſons many Surgeons uſe two Needles in this Operation, one with a ſharp Point, (*Fig.* 7 and 9) to perforate the Coats of the Eye, and the other with a broader or more obtuſe Point (*Fig.* 8.) to depreſs or couch the opake cryſtalline Lens. But it is much eaſier to write of the Advantage of uſing two Needles, than to experience it in Practice. But which ever Sort you chuſe, Care muſt be taken, that it be firſt well poliſhed with Cloth or Leather, before you uſe it to the Eye; that neither its Roughneſs, nor any Particles of Ruſt, may injure that very tender Organ. Mr. FREYTAGE beforementioned greatly recommends a Needle ſhaped like a Hook, for extracting membranous Cataracts out of the Eye; but if this ſucceeds ſo well, why did he not give us the Figure of it?

Apparatus of Dreſſings, deſcribed.

XVII. That there may be no Obſtruction, nor any Time loſt in the Operation, it will be neceſſary to provide every thing in Order which may be wanted for the Dreſſings, after the Couching is performed. Such as, 1. a cooling Collyrium *ex Aq. Plantag. cum Ovi Alb. ſubact. & cum Aluminis, vel Tutiæ, vel Croci, aut Camphoræ portiuncula.* Others uſe common *Sp. Vini* for a Collyrium. ST. YVES uſes a Mixture of ten Parts Water, and one *Sp. Vini*, which he recommends as the beſt. 2. A large Compreſs of ſoft Linen, ſufficient to cover the diſeaſed Eye. 3. A Bandage of about three Ells long, and two Fingers broad; or elſe an Handkerchief folded together in Form of a Triangle, to retain the Compreſs and Dreſſings on the Eye. Laſtly, 4. you muſt provide ſome *Aq. Reg. Hungar. vel Acetum, vel Sp. Cl &c.* to rub the Patient's Noſtrils, if he ſhould faint in, or ſoon after the Operation.

[a] Such a Caſe as this is deſcribed by my Son, in his Account of the Operation for a Cataract, performed by TAYLOR at *Amſterdam*, in 1735, upon one of our Friends.

Poſition of the Patient.

XVIII. There now remains but one more Pre-requiſite before the Surgeon enters on his Work; and that is, to fix and ſecure the Patient in the moſt convenient and advantageous Poſture. He therefore muſt be placed againſt the Light on a much lower Seat than that of the Operator, as you may ſee in *Tab.* XVII. *Fig.* 1. E. the Surgeon himſelf C, being ſeated on a much higher Chair D. If the Patient can ſee either perfectly, or but in Part with the Eye, which is not couched, it muſt be firſt covered or blindfolded with a Handkerchief or Bandage; leſt, by ſeeing the Inſtrument approach, he ſhould move his Eye, and diſturb the Operation: upon which Account it may be alſo proper to admoniſh the Patient, that if his Eye ſhould recover its Sight very ſuddenly in the Operation, as is not unfrequent, he may not ſtir, or make any Exclamations of Joy till it is over; leſt, by a ſmall irregular Motion, the whole Cure ſhould be fruſtrated, and his Sight loſt for ever. The Patient ſhould fix his Hands on the Surgeon's Thighs, and his Legs alſo betwixt thoſe of the Operator. Sometimes it may be proper for an Aſſiſtant to hold up his Feet, that he may not riſe out of the Chair before the Operation is finiſhed. Behind the Patient muſt ſtand the Aſſiſtant A, ſecuring the Head, when the left Eye is to be couched, with his Left Hand on the Forehead, and his Right Hand upon the Chin, which he muſt preſs cloſe to his Breaſt, ſo as to hold the Head firm and ſteady; becauſe a very ſmall Motion of the Head may cauſe perpetual Blindneſs, as we are aſſured by ſad Experience.

Method of Operating.

XIX. Every thing being thus prepared in Readineſs, the Patient is ordered to open his Eye-lids as wide as poſſible, and to turn his Eye inwards towards his Noſe, that a ſufficient Portion of the White of the Eye may appear in the leſſer Angle of the Orbit towards the Temple. (See *Plate* XVII. *Fig.* 14. A.) The Operator now divaricates the Eye-lids with the Fore-finger and Thumb of his left Hand, when it is the left Eye, and of his right Hand when it is the right Eye he couches; and thus he at the ſame time firmly ſecures the Eye from moving; ſee *Fig.* 1, and *Fig.* 14. Some there are who uſe the *Speculum Oculi*, *Fig.* 15 or 16, for this Purpoſe, which, in my Opinion, will more impede than aſſiſt the Operator; but I ſhall not adviſe thoſe to reject it, who are fond of uſing it. The Oculiſt next takes the couching Needle, handed to him by an Aſſiſtant, betwixt the Thumb, Fore and Middle-finger of his right Hand, in the Manner we uſually hold a Pen in Writing, as you may ſee in *Fig.* 1. and *Fig.* 14. he then places the two lower Fingers of the ſame Hand upon the Patient's Cheek, to ſupport thoſe which guide the Needle, and that they move freely, as in Writing: then he carefully enters the Needle almoſt in the Middle of the White of the Eye* betwixt the *Cornea* and external Angle of the Orbit; proceeding, not obliquely, but ſtraight, through the Coats of the Eye, over-againſt the Cataract, to avoid wounding the Blood-veſſels; ſee *Fig.* 14. A. As ſoon as the Needle is perceived to be through the Coats of the Eye, which may be known by your loſing the Reſiſtance, its Point is then inclined towards the Cataract; (ſee *Fig.* 14. B.) which being entered by the End of your Inſtrument, you thereby endeavour to depreſs it gently below the Pupil to the Fundus of the

* The true Place for perforating the Coats of the Eye by the couching Needle, has been largely and elegantly treated of in *Mem. Acad. Reg. Pariſ. An.* 1726, pag. 370. *Edit. Amſtel.* by M. PITIT, who aſſigns the Place to be two Lines Diſtance from the *Cornea*. The Place approved of by TAYLOR we ſhall conſider hereafter.

the Eye, whether it be a Membrane or an Opacity in the cryſtalline Lens; for we are not as yet furniſhed with diſtinguiſhing Marks ſufficient to know one Caſe from the other by their external Appearance, except the Obſervations of ST. YVES. If you perceive the Cataract deſcend with the Point of the Inſtrument below the Pupilla, which it will do the firſt Time, when mature and conſiſtent enough, you are then to continue it there a little while, that it may afterwards ſtay at the Fundus of the Eye. If, upon elevating your Inſtrument again, the Cataract does not riſe above the Pupil, your Operation is well performed; and therefore the Needle is now to be drawn out of the Eye in a ſtraight Line as it entered. If the Cataract riſes again afterwards above the Pupil, as it frequently does, you muſt again couch it with the ſame Needle, as before, keeping it down a longer Time, till it remains ſuppreſſed below the Pupil. M. FREYTAGE indeed adviſes to extract the Cataract, which he thinks is always a Pellicle, by a Hook through the *Cornea*; as, he ſays, he has frequently ſeen done by his Father: but as he neither deſcribes the Hook, nor the Method of Extraction, and as I much doubt whether this Hook would not alſo extract or lacerate the *Retina*, *Choroides* and *Sclerotica*, 'tis, in my Opinion, beſt to neglect his Advice.

Reflections on the Operation.

XX. When the Cataract adheres firmly to any of the Coats of the Eye, 'tis often a very difficult Taſk to couch or depreſs it entire; and therefore in this Caſe you may firſt divide it with the Needle, and then couch or depreſs each Part ſeparately. The ſame muſt be done if you happen to lacerate, or break the Cataract in Pieces in the Operation; and, by this means, the Patient has often recovered his Sight, as we read in CELSUS, GUILLEMEAU, PAREY, BARBET, BRISSAC, and as I have twice obſerved myſelf. If the Cataract adheres ſo firmly to the Uvea, that it can hardly be thence ſeparated, it is often convenient to perforate it in the Middle; by which Means the Rays of Light, paſſing through the Perforation to the *Retina*, the Patient can ſometimes ſee tolerably well afterwards: which Practice may perhaps ſucceed beſt when the Cryſtalline is very thin; for I once found it ſo diminiſhed in Thickneſs in a dead Subject, that it was ſcarce thicker than one's Thumb-nail, and firmly adhered at the ſame Time to the Uvea. But when the Cataract appears to be yet too ſoft, it is adviſeable with BRISSAC, to withdraw the Needle, and defer the Operation till it becomes more conſiſtent, rather than deſtroy the Patient's Sight by confuſing the Humours. When both Eyes are to be couched, 'tis beſt not to perform the Operation on both at one Time, but to intermit a few Days, that the Patient may the better endure the ſame without too violent Symptoms. If you couch the right Eye, the Operation muſt be reverſed; that is, you muſt hold open the Eye-lids with the Thumb and Fingers of your right Hand, and couch the Cataract by the Needle with thoſe of your left; becauſe the Vicinity of the Noſe to the greater Canthus of the right Eye, will impede the Action of the right Hand for this Operation. Though in *Tab.* XVII. *Fig.* 17. you have the Figure of a Needle contrived and ſent me by a Friend, with which you may couch the right Eye with the right Hand. A the Needle, B the Handle, C the Incurvation which reſts on the Noſe.

Treatment after the Operation.

XXI. It is a common Practice with Mountebanks and itinerant Oculiſts, to hold up their two Fingers extended, or elſe a Glaſs of Wine, before the Patient's Eye, as ſoon as the couching Needle is extracted; calling out to know

what the Object is, or of what Colour it appears; and if the Patient can diſtinguiſh, and anſwer rightly, they then conclude the Operation to have been well performed. But this is, by the more prudent Surgeons and Oculiſts, judged to be a pernicious Method; becauſe, by the Patient's ſtraining his Eye too ſoon to view the Objects, the Cataract is often rouſed and elevated again. It is therefore much better to defend the Eye immediately after Couching with a Compreſs dipt in ſome Collyrium, and ſecured by a Handkerchief, that the *Retina* may not be injured by a too ſtrong Action of the Light. It will be neceſſary to blind up both the Eyes, though you couched but one; becauſe if you leave the ſound Eye uncovered, it will be perhaps looking at Objects, and will conſequently draw or ſtrain the diſeaſed Eye in the ſame Direction; which may remove the Cataract, and cauſe it to aſcend again, or elſe induce an Inflammation, or other bad Accidents.

How to prevent the Cataract from riſing.

XXII. After your Dreſſing and Deligation, the Patient ſhould be laid on his Bed, upon his Back, with his Head elevated, and retained almoſt erect, by Pillows; and continue very quiet and compoſed for the Space of eight Days, without coughing, ſneezing, laughing, intenſe talking, or eating Food of a hard Digeſtion, in order to prevent the Cataract from riſing or being diſturbed. No Surgeon can aſſert that the Cataract ſhall continue ſuppreſſed after the firſt Time of couching; but the Patient has this Advantage, that if it aſcends it may be again ſuppreſſed, and his Sight recovered by the Operation. Accordingly MAITRE-JEAN writes, that a Patient, whom he couched in Autumn, had a Return of his Cataract in the Spring following; but it was happily removed again by repeating the Operation. We have even ſome Inſtances of the Cataracts having ſubſided again of themſelves, after they had riſen above the Pupil.

Further Treatment.

XXIII. A few Hours after the Operation, it will be convenient to bleed the Patient in proportion to his Strength and Fulneſs of Habit, to prevent an Inflammation in the wounded Eye; and to repeat the ſame, if neceſſary, with the Uſe of *Collyria* externally, and cooling Purges internally. 'Tis very remarkable, that the Patient is often troubled with a Vomiting an Hour or two after the Operation, as I have frequently obſerved, and imagine to ariſe from the Conſent of the Nerves, and their Irritation in the Operation, which ſoon goes off afterwards; which I find has been alſo obſerved by Mr. FREYTAGE. However, this Symptom of Vomiting is no good Preſage, becauſe the Patient's ſtraining in this Action, often cauſes the Cataract to aſcend. In the Evening, after the Operation, you ſhould order the Patient an anodyne Emulſion, to compoſe him to reſt; becauſe Watchings and Reſtleſſneſs very often occaſion the Cataract to aſcend again above the Pupil. The Diet and Regimen here muſt be ordered the ſame as we have directed in Wounds and inflammatory Diſorders. Laſtly, if the Patient does not go to Stool freely without ſtraining, it will be proper to help him with a Clyſter; nor ſhould he be permitted to diſturb his Head by riſing out of Bed for this Office; but, for the firſt few Days after the Operation, it will be more convenient to uſe a Bed-pan. All which Precautions are neceſſary, to prevent the lately depreſſed Cataract from being diſturbed or raiſed again above the Pupil.

XXIV. With regard to the Deligation and ſubſequent Dreſſings, it will be convenient to remove the Bandage very gently on the firſt Evening after the

Operation; and renewing the Compreſs dipt in ſome Collyrium, to apply the Bandage again as before. On the following Days this Proceſs muſt be repeated Morning and Evening at leaſt, and ſometimes four or five times in a Day; becauſe the Inflammation then becomes more intenſe, and the Compreſſes dry much ſooner: therefore the Operator ſhould, at this Time, be more ſollicitous to guard the Light from the Eye, eſpecially when the Inflammation runs high. If the Eye continues in a good Condition with but a ſlight Inflammation, you muſt continue this Method of Dreſſing till the eighth Day, when all the Danger will be over: you may then, by degrees, remove the Bandage, and admit the Light to the Eye, which ſhould be for ſome Time guarded at firſt with a Piece of green Silk hanging over the Forehead. On the tenth Day, if nothing forbids, the Patient may riſe and walk about his Chamber, provided his Window-curtains are drawn, and his Eyes defended with green Silk as before; which he may, by degrees, lay aſide, and return to his former Courſe of Life.

The Removal of Accidents.

XXV. That the young Surgeon may the better underſtand how to relieve the ſeveral Accidents which may attend this Operation, we ſhall conſider each of them ſeparately; and, 1. If a ſmall Portion of Blood ſhould be extravaſated, and eſcape into the aqueous Humour, ſo as to render it in ſome meaſure obſcure and turbid, you muſt diſpatch the Operation as faſt as poſſible, and dreſs up the Eye with a Compreſs dipt in the forementioned Collyrium; by which Means ſlight Extravaſations have been frequently obſerved to be diſperſed. But if a large Quantity of Blood is mixed with the aqueous Humour, it will then be almoſt impoſſible to avoid a Suppuration, termed *Hypopyum*, or other ill Conſequences, which endanger perpetual Blindneſs, or a total Deſtruction of the Eye. Yet even here you will find great Benefit from plentiful Bleeding, and from diſcutient Bags ſtuffed with Fennel, Sage, Hyſſop, and Roſemary, boiled in Wine, frequently to be applied warm to the Eye. 2. If the aqueous Humour itſelf eſcapes, or runs out of the Eye, in the Operation, ſo as to leave the *Cornea* flaccid, the Eye itſelf is not in any great Danger thereby; for the Humour will be reproduced ſo as to fill the *Cornea* again in a few Days. Laſtly, 3. If great Inflammation ſhould ariſe, you muſt omit nothing that will conduce to ſuppreſs it; as plentiful Bleeding, and drinking of Water, or other cooling and diluent Liquors, to bathe the Temples frequently with *Sp. Vini Camph.* to apply Bliſters behind the Ears, and clyſter the Patient as you ſhall ſee neceſſary.

Fabric of the couching Needle.

XXVI. From what has been ſaid, I think it is ſufficiently apparent how much the Moderns are improved above the Antients, as to their Knowledge of the true Nature or Diagnoſis, Prognoſis, and Method of curing this Diſorder: for, upon obſerving that a Cataract was rather conſtantly formed by an Opacity of the Cryſtalline, than from any Membrane, BRISSAC conſequently judged, that thoſe couching Needles would ſucceed beſt, which were made with a ſulcated and pretty broad Point, as in *Tab.* XVII. *Fig. 6. lit.* C. for by uſing thoſe ſlender-pointed Needles of the antient Surgeons, whether made of Gold, Silver, or Steel, it was almoſt impoſſible to avoid cutting or lacerating the Cataract in couching it: but the couching Needle of BRISSAC is made with an acuminated, as well as a broad and ſulcated Point, that it might the more readily perforate the Coats of the Eye. The Handle of the couching Needle

Needle A B is octangular; and the Side marked E E lying even with the Sulcus in its Point, is hatched, or otherwiſe particularly marked, that you may judge by the Poſition of the Handle how the Point of the Needle is directed, in reſpect of the Cataract in the Eye. Laſtly, the riſing or Protuberance of the Inſtrument, marked D, ſerves to indicate how deep it has entered into the Eye.

Other particular Needles.

XXVII. Thoſe Surgeons who have perſuaded themſelves, that a Cataract proceeds from a Membrane or Tunic, have alſo provided themſelves with an unciform Inſtrument, to extract the ſaid Membrane through the Puncture made in the Coats of the Eye by the Needle, and prevent the Diſorder from returning; as it might, if they were to leave the Cataract at the Bottom of the Eye. Some of their Inſtruments were made tubular, in order to ſuck out the Membrane from the Eye[a]; others were made like a Pair of ſmall Pliers in the Shape of a Needle; as in *Tab.* XVII. *Fig.* 10. and others again were like ſmall Hooks, which they introduced and extracted through a Cannula, together with the Tunic or Cataract, according to FREYTAGE. But their Methods and Inſtruments were as uſeleſs and miſchievous as their Notion of the Diſorder was falſe.

When the Cataract comes thro' the Pupil.

XXVIII. We have further to adviſe, that if the Cataract ſhould further extend itſelf, or ſlip through the Pupil, as it ſometimes may, it will then be proper to try if you can draw it back by paſſing the Needle through the Pupil; if not, to make a ſmall Inciſion in the lower Part of the *Cornea*, and thereby extract the Cataract by a ſmall Hook or Probe; an Inſtance of which Practice we have given in our profeſſed Treatiſe on this Diſorder.

TAYLOR's Method.

XXIX. The noted Oculiſt, TAYLOR, propoſes a new Method of his own, as he ſays, for couching Cataracts, in the ninth Chapter of his Treatiſe, which he deſcribes as follows: the Patient being ſeated as uſual, and his Eye held firm by the *Speculum Oculi*, he then makes a ſmall longitudinal Inciſion with a Lancet[b] of about half a Line in Length below the uſual Place; which Inciſion he continues through the external and internal Coats of the Eye into the vitreous Humour. He then takes a *Plano-convex* Needle of a very ſlender or thin make, and paſſes it through the Inciſion directly into the Eye, with its convex Part upwards and towards the Bottom of the cryſtalline Lens. He next gently elevates the Point of his Needle a little, till he finds a ſmall Reſiſtance on it from the cryſtalline Lens above it, which he alſo perceives to move, by looking through the Pupil. Being thus aſſured the Point of his Needle is under the Capſula of the Cryſtalline, he then guides his Needle downward towards the Bottom of the Eye, to divide the vitreous Humour, and make a Space for receiving the Cryſtalline, which he next depreſſes. In order to couch the Cryſtalline, after having divided the vitreous Humour, he draws his Needle about two Lines further out of the Eye, and then inſerts the Point of it into the lower Part of the Capſula of the Cryſtalline, which he thus incides or opens as he ſays, without injuring the *Ligamentum ciliare*; and in thus opening the Capſule, he alſo endeavours to enlarge the Space for receiving the Cryſtalline. Laſtly, in order to couch or depreſs the opake Cryſtalline, he again extracts his Needle almoſt three Lines

[a] Vid. SCHACCHI *Subſid. Medicin.* p. 54. & TH. FIENI, *Lib. Chirurg.* p. 30.

[b] He does not give us any Reaſon for uſing a Lancet, or for making his Inciſion longitudinal rather than oblique or tranſverſe, nor can I ſee any Reaſon for it; but it is a ſtanding Maxim in Surgery, never to uſe ſeveral Inſtruments for what may be done as well by one.

Lines more out of the Eye; then elevating its Point, and fixing the ſame into the upper Part of the Cryſtalline, he endeavours to depreſs and lodge it in the Space before made for its Reception in the vitreous Humour at the Fundus of the Eye, and then gently extracts his Needle. By this means, he aſſerts, that the *Uvea* and *Ligamentum ciliare* are not in the leaſt injured, but remain in their natural and ſound State; whereas in the common Method of couching they are uſually lacerated, to the great Detriment of the Eye, and its Office of Viſion. To conclude, the Subſtance of his Method of operating, which we have here briefly related, is ſo ſwelled and obſcured, by ſtuffing it with frivolous Cautions and Circumſtances, in his Treatiſe, from whence we have extracted it, that it there takes up more than three times the Compaſs in which we have here repreſented it, and yet we have omitted nothing but what was either inſignificant or unintelligible. There are even ſo many Circumſtances related, that it ſeems impoſſible the Author himſelf ſhould attend to all of them; and this may poſſibly be one Reaſon of his ill Succeſs in Practice, his Operation being followed with excruciating Pains, moſt violent Inflammation, and a Suppuration of the Eye, inſtead of recovering the Patient's Sight; as you may ſee related more at large in my Son's Treatiſe, on the unhappy couching of a Cataract in our Friend at *Amſterdam*, by TAYLOR, in 1735. However, the Practice deſerves to be conſidered and tried by the more prudent Oculiſts; and the Succeſs of it will, in Time, determine the Author's Merit.

His Treatment of the ſhaking Cataract.

XXX. When the Cataract moves, or when the opake cryſtalline Lens is ſlipt out of its Capſula, and fluctuates behind the Pupil, which TAYLOR then calls a *ſhaking Cataract*; the Diſorder, he ſays, will now require a different Method of Cure: to explain which he makes the Buſineſs of two diſtinct Chapters, which import no more than that he here paſſes his Needle as before, into the Eye, directing its Point to the upper and anterior Part of the Cataract, or opake Cryſtalline, to avoid injuring the ciliary Ligament; and then, with the plain Surface of his couching Needle, he depreſſes the ſame to the Bottom of the vitreous Humour.

His Treatment of the falſe Cataract and Glaucoma.

XXXI. In ſome Cataracts, which he terms *falſe*, he ſays, the Capſula of the Cryſtalline is vitiated, and become opake, as well as the Lens. The Method of couching both of which, and freeing them from the ciliary Ligament, is related by him in ſo prolix a Manner, that he again makes it the Buſineſs of two whole Chapters. Two other Chapters are again employed in explaining his Operation for the *Glaucoma*; by which Name he underſtands, contrary to all his Predeceſſors, an Opacity joined with an Expanſion of the cryſtalline Lens, which, with its vitiated Capſula, are extended or protruded forwards cloſe to the Margin of the Pupil; for the Cure of which he proceeds in the ſame Manner as before. But I know not what Right or Authority he has, more than his own Aſſurance, to impoſe this Name to a Diſorder, different from what it has been all along intended to ſignify by our Predeceſſors; for it will appear quite unwarrantable even to make, and much more to transfer Names, without an abſolute Neceſſity; ſince what he calls a *Glaucoma*, is, I think, a Species of the Cataract, and not a Diſorder of the vitreous Humour, ſeated much deeper in the Eye, as the Antients have all along underſtood by the Name.

His Extraction of Cataracts thro' the Cornea.

XXXII. We before obſerved, at *N.* XXVIII. that thoſe Cataracts which have eſcaped through the Pupil, may be extracted by an Inciſion made in the *Cornea*: but I have been aſſured from *England*, that this famous Oculiſt there boaſted, that

that he could, and does extract Cataracts in this Manner, which are even fixed behind the Pupil and *Uvea*. But I could never yet learn the Truth of his Aſſertion, or that he ever performed the Fact.

CHAP. LVI.

Of Dilating CONTRACTIONS *of the* PUPIL.

Deſcription of the Diſorder.

1. WE are now to treat of an Operation related to the foregoing, in which the Coats of the Eye are perforated by an Inſtrument, almoſt in the ſame Manner as in couching a Cataract, in order to open an imperforated or contracted Pupil. The Diſeaſe we are now ſpeaking of is therefore ſuch a total or cloſe Contraction of the Pupil, that it will not tranſmit Light enough to the Bottom of the Eye, to enable the Patient to ſee Objects diſtinctly. Sometimes this Diſorder has grown up from Infancy; and ſometimes it ariſes from an intenſe Inflammation of the Eye, or ſome ſudden and violent Conſtriction of the Pupil from other Cauſes, with a Palſy of the ſtraight Fibres in the *Uvea*; or when the internal Margin of that Membrane, which conſtitutes the Pupil, is concreted or joined to a Cataract, or to ſome Part of a Cataract, after the Operation. The Cure of the Diſorder is generally eſteemed extremely difficult, if not altogether impracticable. But the celebrated Mr. CHESELDEN has contrived a new Method of relieving this Diſorder, which he has not only tried ſeveral times with Succeſs, but alſo deſcribed his Proceſs in the *Philoſophical Tranſactions*, and in the *Appendix* to the fourth Edition of his Anatomy; which we ſhall therefore give a Place here in our Surgery, as follows:

This Operation.

II. The Eye-lids being held open by a *Speculum Oculi*, he then takes a narrow and ſingle-edged Scalpel, or Needle, *Tab.* XVII. *Fig.* 19. A A, almoſt like that for couching a Cataract; and paſſing it through the *Sclerotica* B, as in couching, he afterwards thruſts it forwards through the *Uvea* or *Iris*, and, in extracting it, cuts through the *Iris* in the Manner repreſented by *Fig.* 20. A. If the Diſorder is not accompanied with a Cataract, it will be beſt to perforate the *Iris* in the Middle, as you may perceive by *Fig.* 20. But when there is a Cataract, the Inciſion ſhould be made a little higher in the *Uvea*, that the Cataract may not obſtruct the Ingreſs of the Rays of Light. The Cataracts which ſometimes accompany this Diſorder, he ſays, are generally very ſmall; and ſometimes their Adheſion to the *Iris* is ſo firm, as to render it impracticable to couch or ſuppreſs them. In *Fig.* 21. the Inciſion or Aperture is repreſented lower than the Center of the *Cornea* and *Uvea*; becauſe in this Eye on which he performed the Operation there was an *Albugo*, or white Speck, upon the upper Part of the *Cornea*, which obliged him to inciſe lower than uſual. He does not indeed relate the Manner of treating the Patient afterwards, to ſuppreſs and guard againſt an Inflammation, and other Accidents; but 'tis reaſonable to ſuppoſe you muſt proceed in the ſame Method as after the Operation for a Cataract.

An EXPLANATION *of the* SEVENTEENTH PLATE.

Fig. 1. Demonſtrates the Poſition of the Patient, Surgeon, and Aſſiſtant, proper for couching a Cataract as explained in *Chap.* LV. *N.* XVIII.

Fig. 2, 3. Repreſents the ſilver couching Needles uſed by the Antients; the firſt having a ſlender and round Point like common Needles, and the laſt a triangular Point.

Fig. 4.

Plate 17

Vol. I. P. 440.

Fig. 2.

Fig. 3.

Fig. 4.








Fig.



16.

Plate 17


Fig. 2.
Fig. 3.
Fig. 4.
A
C
B
Fig. 5.
Fig. 6.
C
D
B
E
E
A
Fig. 7
Fig. 8.
Fig. 9
Fig. 20.
A
Fig. 15.


Fig. 8.
Fig. 9.
Fig. 10.
Fig. 11.
A
B
C
Fig. 12.
A
B
A
Fig. 14.
B A
C
B
Fig. 17.
Fig. 18.
A
A
Fig. 16.
B
B
C
Fig. 1.
A
B
C
Fig. 13.
C
D
E
B
D
Fig. 21.
A
Fig. 19.
A
A
B
C




Fig. 4. Repreſents a double-pointed couching Needle; that marked A being round and ſlender, and that at B a little broader or flatter: C denotes the Handle, which may be made of Silver, Braſs, Ivory, or Wood.

Fig. 5. Is another Needle with a ſtill broader Point, but ſharp-edged; with which a Cataract may be more commodiouſly held and couched than by a ſmaller Point.

Fig. 6. Denotes another couching Needle, almoſt like the former, only furniſhed with a *Sulcus* in its *Apex*, which is recommended by BRISSAC, and deſcribed more largely at N° XXVI. of *Chap.* LV.

Fig. 7, 8. Repreſent two Needles from SOLINGEN and NUCKE, which are ſaid to be invented by the *Dutch* Oculiſt SALMASIUS, and to be both uſed in one and the ſame Operation. That at *Fig.* 7. is ſulcated and ſharp pointed, almoſt like the preceding, and ſerves to perforate the Coats of the Eye; from whence BRISSAC ſeems to have taken his at *Fig.* 6. But that at *Fig.* 8. is obtuſe, and made ſo as to paſs through the *Sulcus* of the preceding Needle, while it continues in the Eye to depreſs the Cataract.

Fig. 9, and 10. Repreſent two Needles of pretty much the ſame Uſe with the two preceding, and are taken from BERN. ALBINUS's *Diſputatio de Cataracta, Francof. impreſſ.*

Fig. 11. Denotes the Needle propoſed by ALBINUS, in his ſaid Treatiſe, for extracting a membranous Cataract out of the Eye; being ſo contrived that the Point A opens like a Pair of Pliers, to extract the membranous Cataract, (if there be any in the Eye) by depreſſing the little Handle B; though I much doubt whether it was ever uſed with Succeſs.

Fig. 12, and 13. Repreſent the Parts of the preceding Needle ſeparate and aſunder. *Fig.* 12. is the ſulcated Point, in which is lodged the other Point *Fig.* 13. Theſe perforate the Eye the better, as they are more exactly fitted and poliſhed. They are connected by the Hinge B, C, D, *Fig.* 11, 12, and 13. E. *Fig.* 13. denotes a Spring to preſs the two Points cloſe together, till you open them by depreſſing it with your Thumb on the little Handle, B. *Fig.* 11. to apprehend and extract the Membrane.

Fig. 14. Repreſents the Method of holding open the Eye-lids with one Hand, and of paſſing the Needle with your other, for couching a Cataract, the Point B uſually appearing through the Pupil.

Fig. 15, and 16. Repreſent two *Specula Oculorum*, to hold the Eyes firm, and open their Lids in couching, and other Operations for the Eyes: the laſt is more correct than the firſt, as you may extend or contract the Circle A A, B B, by elevating or depreſſing the Button C. The Handle is denoted by D.

Fig. 17. Repreſents a Needle for couching a Cataract in the right Eye with the right Hand. A the Point of the Needle; B its Handle, in which is a particular Kind of Incurvation C to reſt upon the Noſe.

Fig. 18. Is a Cap or Sheath for including the Point of the ſame Needle.

Fig. 19. Is taken from the *Appendix* to the fourth Edition of Mr. CHESELDEN's *Anatomy*, to ſhew the Manner of directing his Cutting-Needle to open or inciſe the cloſed or contracted *Uvea*.

Fig. 20. Denotes the Manner of dividing the *Uvea* in its Middle by the ſame Inſtrument, to tranſmit the Rays of Light into the Eye.

Fig. 21. Repreſents the Manner in which Mr. CHESELDEN pierced the *Uvea* lower than uſual, on the Account of an *Albugo*, which infeſted the Middle of the *Cornea* in this Eye.

CHAP. LVII.

Of the PTERYGIUM, *or* Unguis Oculorum.

Deſcription. I. WHEN a preternatural Membrane is formed externally upon the Coats of the Eye, ſo as to extend itſelf over the *Cornea* and Pupil, and obſtruct the Sight, the Diſorder is then uſually denominated *Onyx* by the *Greeks*, and *Unguis*, or *Ungula*, by the *Latins*; it is alſo ſometimes named *Pterygium*, from its Reſemblance to the Wing of a Bat[a]. Sometimes the Pellicle or Film appears red, from the Number of ſmall Blood-veſſels, and then it is uſually denominated *Pannus*. It moſt frequently ariſes in the Angles of the Eyes from the Temples or Noſe, and ſometimes from above or below, extending itſelf, by degrees over the *Cornea*, (as in *Tab.* XVIII. *Fig.* 1 and 2. *a a.*) Sometimes it only adheres ſlightly to the *Cornea* by a few ſlender Fibres; and ſometimes again it is extended over the whole Eye, and continues moſt firmly and intimately attached to it, which uſually renders the Caſe much more difficult to cure.

Cure by Medicines. II. While the Pellicle is but recent and ſlightly attached, it may be removed by gentle Eſcharotics; ſuch as Powder *ex Sacchar. Canarienſ.* ʒj. *Vitrioli albi vel Aluminis uſti, vel etiam viridis Æris Gr.* iv. *vel* vj. which muſt be carefully ſprinkled at Intervals by a little at a Time upon the Membrane. Some uſe a Powder of the *Lapis ſciſſilis*, or of the *Os Sepiæ* mixed with Sugar. But as it will be difficult to uſe ſuch a Powder for Infants, it will be better to treat them with an Eye-water, as that of QUERCITAN, *cum Vitriol. alb. aut ſelle Muſtelæ piſcis, &c.* which may be alſo uſed to Advantage for Adults. If the Diſorder is accompanied with an Inflammation, it will be convenient for you to treat the Patient accordingly, by Bleeding, Bliſters, and cooling Medicines. M. ST. YVES ſets a great Value on the *Lapis medicamentoſus Crollii*, diſſolved in Water, and uſed to waſh the Eye; though, in my Opinion, a Solution of *Vitriol. alb.* ϑ ß. in *Aqua Chelidonii major.* ʒij. is little inferior, if at all.

Cure by the Scalpel. III. If the mild Eſcharotics, before propoſed, are inſufficient for deſtroying the Pellicle, you muſt then extirpate it. In order to which, the Patient muſt kneel down on his left Knee, if the right Eye be affected, and lean his Head back againſt the Light upon the Surgeon's Lap, or Knees, who then takes the ſmall Hook, *Tab.* XVIII. *Fig.* 3. or *Tab.* XV. *Fig.* 30. and, after the Eye-lids are held open by an Aſſiſtant, endeavours to paſs its Point under the thickeſt or looſeſt Part of the Pellicle, to elevate it a little. In the next place, he takes the Needle *a*, armed with a Thread, *Fig.* 1. *b b.* and paſſing it under the Pellicle, ties it with a double Knot; and then faſtening the two Ends in a Loop, *Fig.* 2 *b c.* he thereby attempts to make a gentle Elevation. This done, he now endeavours to ſeparate the upper and lower Margin of the Membrane with

[a] CELSUS, *Lib.* VII. *Cap.* 7. No. IV. and CASTEL, *Lex Med. per* BRUNO. *ſub tit.* ONYX.

with a Lancet, that he may afterwards cut off the reſt immediately in a ſtraight Line near the lacrymal Caruncle by a Pair of ſmall and ſtraight Sciſſars. He then draws back the Thread and Membrane towards the *Cornea*, and if it adheres any where to the Eye, frees it by degrees with a Scalpel or Sciſſars: in doing which, the Operator muſt have a principal Regard to two Things: 1. to avoid injuring the *Cornea*; and, 2. to obſerve that no Part of the Membrane be left adhering to the Eye; which laſt might occaſion a Return of the Diſorder. Yet it is better to leave ſome Part of the *Unguis* adhering to the *Cornea*, when its Separation is extremely difficult, than to wound the *Cornea*, and leave irremediable Scars in it; and this the rather, becauſe any ſmall Portion of the Membrane left behind may be taken off afterwards, by treating the Eye two or three Times in a Day with the gentle Eſcharotics before propoſed at N° II. Though there are ſome, who rather approve of the following *Collyrium* for removing the membranous Reliques:

℞ *Aq. Roſar. Damaſcenar. Plantag. ana.* ℥ j. *Matr. Perlar. ppt.* ℈ j. *Sacchari Saturni Gr.* vj. *Vitrioli albi Gr.* iij. *M. f. Collyr.*

M. ST. YVES approves of waſhing the Eye for three or four Days afterwards with *Sp. Vini* diluted with Water, and then to uſe a Solution of the *Lapis medicamentoſus* in Spring water. Laſtly, in extirpating the Pellicle, great Care muſt be taken not to cut off any Part of the lacrymal Caruncle, and much more not to remove the whole of it; for, if this Body be wanting in the greater *Canthus* of the Eye, where it ſtops and directs the Tears into the *Puncta lacrymalia*, the Patient will conſequently be troubled with a watery Eye, in which the lacrymal Humour will run down over his Cheek.

Other Methods of Cure.

IV. Some of theſe Pellicles which appear red, from the ſmall Blood-veſſels extended to them from the Corners of the Eyes, will wither or eaſily fall off with the Uſe of Medicines, upon ſcarifying and dividing thoſe Veſſels in the *Canthus* of the Eye which feed and nouriſh them. Sometimes the *Cornea* is incruſted over with a glutinous Matter, like Fat or a Membrane, which may be readily ſcowred off with the Gall of an Eel, Lamprey, or the Bile of ſome other Animal. This was probably the Caſe of TOBIAS, mentioned in the *Old Teſtament*. Sometimes indeed we meet with Membranes of this Nature, which are inſeparable from the *Cornea* by any means whatever. But this we cannot be aſſured of before Trial; and we ought rather to try the Operation in vain, than to relinquiſh the Diſorder, unjuſtly, as incurable. Laſtly, ſome Pellicles upon the Eye are extremely painful and ſtubborn, inclining to a cancerous Diſpoſition; and theſe it may be beſt for the Surgeon to relinquiſh as incurable.

Pterygium of the whole Eye.

V. When the *Pterygium* or *Unguis* is extended over the whole Eye, it will be convenient to divide it by a cruciform Inciſion into four Parts, according to M. St. YVES, and then to ſeparate each of them from the *Cornea* and Eye; (as we before directed for the *Unguis* in general) conducting the Remainder of your Dreſſing as we there preſcribed.

VI. Laſtly, when this Operation is to be performed upon the left Eye, the Patient ſhould riſe up from the Ground as ſoon as the Needle has been paſſed through the Membrane, and the Threads tied; for being placed in a Chair, the Operator

tor may have a better Command of the Eye than before; except he ſhould happen to be as active with his left Hand as with his right. If the Membrane appears to be thin and weak, Care ſhould be taken not to extend it too forcibly by the Thread, leaſt it ſhould break.

CHAP. LVIII.

Of the Albugo, Leucoma, Nebula, Nubecula, *and other* SPOTS *in the* Cornea *of the* EYE.

Deſcription. I. AS in ſeveral other Claſſes of Diſorders belonging to the Eye, ſo in this we meet with a great deal of Confuſion, by a Miſapplication and Reduplication of ſeveral Names, which are often uſed to import the ſame Diſeaſe; whence ariſe Difficulties and Miſtakes to the Learner, and Errors in the Method of Cure. However, we find that the moſt eminent Surgeons and Phyſicians intend, by theſe Names, Sorts of whitiſh Spots in the *Cornea*; though they appear not always alike, and of the ſame Kind, being ſometimes larger, or ſmaller, thicker or thinner, or more or leſs protuberant. According to their different State and Condition they more or leſs obſcure the Sight, and ſometimes wholly intercept it. Hence we have alſo a Reaſon why the Blemiſh was ſometimes called *Leucoma* by the *Greeks*, and *Albugo* by the *Latins*, or *Nebula* and *Nubecula*; according as it appeared darker or clearer.

Cauſes. II. The Cauſes of theſe Blemiſhes are various. For they may ariſe, 1. from an Obſtruction of the pellucid Veſſels in the *Tunica cornea*, and an Inſpiſſation of their contained Juices, proceeding from a violent Inflammation of the Eye: or. 2. from a Suppuration, and then an Induration, of theſe Juices, in the *Cornea*, after an Inflammation, ſo that it, by degrees, becomes more opake, as it hardens, and puts on a whitiſh Hue, being ſometimes miſtaken for an *Unguis* [a]. 3. Theſe Spots may ariſe from an external Eroſion or Ulcer in the *Cornea*; or, 4. from Puſtules, or *Veſiculæ* in various inflammatory Diſorders; particularly, 5. from thoſe which are occaſioned by the Small-Pox. 6. They may very often proceed from the Scars left after a Puncture in the *Cornea*, from a Sword, Knife, Fork, a Splinter, Glaſs, a Thorn, or the like; or, 7. from a Burn; or, 8. the corroding Acrimony of cauſtic Subſtances falling into the Eye. Laſtly, 9. they may ſometimes be formed of a peculiar Tunic growing to the Eye itſelf.

Prognoſis. III. Theſe Diſorders of the *Cornea* are ſome more and ſome leſs difficult to remove, according to their Duration, and the particular Cauſes from whence they proceed, with the Patient's Age, and other Circumſtances. Infants may be more eaſily freed from them than Adults, when they are not of any long ſtanding: but for thoſe which are Scars formed from Wounds, Burns, Punctures, or the like, there is little or no Hopes of removing them.

Methods of Cure. IV. If any one is deſirous to be ſucceſsful in removing theſe Spots, he muſt adapt his Method of Cure to the Cauſe of the Diſorder. For thoſe which ariſe from

[a] See *Chap.* LVII. and LX. Likewiſe MAUCHART's Treatiſe.

from inſpiſſated Humours betwixt the *Laminæ* of the *Cornea*, and are not of long ſtanding, may be beſt removed by a proper Regimen, attenuating Diet, and Medecines, eſpecially a plentiful Uſe of thoſe Decoctions and Infuſions which are ſudorific. But then at the ſame time muſt be uſed externally Phlebotomy, Scarification, Bliſters, and frequent waſhing of the Feet. Upon the Eye itſelf may be alſo applied diſcutient Bags *ex fol. Hyſſop. Roriſmarin. flor. Chamom. Sem. fænic. &c.* boiled in Wine or Water, and frequently laid on the Eye; or a Collyrium *ex Aq. fænic. cum. Sp. Vin. Camph.* Laſtly, it may be convenient for the Patient to hold his Eye ſometimes over the warm Vapours of Coffee, or a Decoction of the Woods. On the contrary, it will be here pernicious to uſe cold and aſtringing *Collyria*, eſpecially thoſe of white Vitriol, though they are much eſteemed; whereas warm Applications are found by Experience to be of the greateſt Service. When the Inflammation is diſperſed, the Patient may wet his Eye every Day with ſome of the *Aqua Ophthalmica Quercitani, cum Tutii pp.* made warm before uſing it. If any of the ſmall Veins proceeding to the Spot appear turgid on the White of the Eye, it will be proper to divide them by the double-edged and crooked Needle (*Tab.* I. *Fig.* 5. or *Tab.* XVI. *Fig.* 2.) a Lancet or Sciſſars. Laſtly, in ſome of them, which are of long ſtanding, you may rather expect any thing than their Cure.

VI. In thoſe whitiſh Spots which proceed from Abceſſes, or a Suppuration of Matter after an Inflammation betwixt the *Laminæ* of the *Cornea*, which they elevate like a Pea, or Pearl, (whence they are ſometimes called *Pearls*) you ſhould make an Inciſion into the *Cornea*, to diſcharge the included Matter; which might otherwiſe by degrees erode the *Cornea*, and deſtroy the Sight. Your Inciſion for this Purpoſe may be made either by the Lancet, or by a couching Needle, *Tab.* XVII. through the *Cornea* quite to the Abceſs; and repeated, if there be Occaſion; treating the Eye afterwards with ſome of the diſcutient Medicines propoſed at N° V. Others uſe Viper's Fat, to cleanſe or heal the Puncture or Inciſion, and by this Means, the Sight is ſometimes happily reſtored. But when the Matter is lodged deep, and not near the Outſide of the *Cornea*, it will be impoſſible to preſerve the Eye-ſight diſtinct and perfect, either, by this, or any other Means. Cure of Spots and Abceſſes.

VII. But when the *Cornea* is eroded externally, either from an Abceſs, Inflammation, or any other Cauſe, the following Method is taken by M. St. Yves. Firſt, he removes the Inflammation, and then orders the Patient to waſh his Eye frequently with the *Aqua viridis ophthalmica Hartmanni*, which is made weaker or ſtronger, according as the Patient can bear it; the admirable Virtues of which Water for removing Spots in the *Cornea*, are ſtrongly recommended by the ſame Author. Cure of an Albugo from an external Eroſion of the Cornea.

VIII. In ſome of thoſe ardent or inflammatory Puſtules of the *Cornea*, which appear afterwards whitiſh and protuberant, like a Pearl or Grain of Millet, the beſt and moſt expeditious Method of removing them is, by perforating with a Needle, ſo as to diſcharge their contained Matter. And in thoſe Puſtules ariſing from the Small-Pox, you ought to make an Apertion by a Needle or Lancet, immediately to diſcharge the eroding Matter, removing the Pellicle afterwards with ſome *Alumen uſtum cum Sacchar. cand. & Ovor. teſt. pp.* applied every Day to the *Cornea*. Others uſe Tinder, or burnt Lint dipt in Oil. By either of which Cure of thoſe from Burns, or the Small-Pox.

which the remaining Film will, by degrees, vaniſh, according to ST. YVES, (pag. 229.) The ſame Method of Cure muſt be taken for diſcharging the Matter in Puſtules formed in the *Cornea* from Burns; treating the Blemiſh afterwards with the Medicines we have directed in *Chap.* LVII. preceding.

Incurable Spots.

IX. Thoſe Spots of the *Cornea*, which ariſe from Wounds, Scars, or the Abuſe of the vitriolic *Collyria*, are ſeldom curable; as are thoſe alſo which render the *Cornea* quite opake, and are of very long ſtanding, or in which the natural Form of the Eye or *Cornea* are deſtroyed. In theſe Caſes it is therefore much better to leave the Patient to himſelf unmoleſted, than to torture his Eyes to no Purpoſe, by a tedious Courſe of Remedies and Operations.

CHAP. LIX.

Of the STAPHYLOMA.

A Staphyloma deſcribed.

I. UNDER the Term *Staphyloma*, (the *Grape*) are chiefly compriſed two Diſorders of the Eyes; one, in which the *Cornea* is more than uſually protuberant, as in *Tab.* XVIII. *Fig.* 4, 5, 6, 7, and 8; the other in which the Pupil or *Uvea* breaks forth and forms an unſightly Tumor on the *Cornea*, either from internal Cauſes, or from ſome wounding Inſtrument forced through the Coat; in which laſt Caſe the Sight of the Eye is uſually deſtroyed. See *Fig.* 8. *a a.*

Kinds.

II. There are various Species and Denominations of the *Staphyloma*, according to their Size and Shape: as the *Margarita*, *Myocephalus*, *Clavus*, *Mylon ſive Pomum*, and the *Staphyloma* or *Acinus* ſtrictly ſo called; of all which the biggeſt is the *Mylon*. But I have ſometimes obſerved not only the *Cornea*, but alſo the *Sclerotica* preternaturally diſtended, and enlarged to a great Degree after the ſame Manner as a *Hernia ventralis*, or, the lacrymal Sac, and then the Diſorder may be alſo denominated *Staphyloma*, becauſe thoſe two Coats, the *Cornea* and *Sclerotica*, are properly conſtituted but of one. However, it may be juſt to diſtinguiſh thoſe Tumors from each other, according to the different Parts affected, by denominating one of them *Staphyloma Scleroticæ*, and the other *Staphyloma Corneæ*.

Prognoſis.

III. A *Staphyloma* is a dangerous Diſorder, as well becauſe it greatly deforms the Eye, and deſtroys its Sight, as becauſe it often induces moſt violent Inflammations, Head-achs, Reſtleſſneſs, Abceſs, and ſometimes a Cancer in theſe Parts. The Cure of it is therefore generally undertaken, not ſo much to recover the Sight, as to preſerve or reſtore the Uniformity of the Eye, and prevent the malignant Symptoms before enumerated.

Cure of a recent Staphyloma.

IV. In the Cure of this Diſorder we muſt relieve the Tumor and Deformity of the *Sclerotica* and *Cornea*, by the Application of a Compreſs dipt in *Aqua Aluminis*, together with a Plate of Lead and Bandage, or ſome proper compreſſing Inſtrument. If the *Uvea* protrudes itſelf through a Wound in the *Cornea*, it ſhould be returned by a Probe: the Patient in the mean time muſt lie in a

ſupine Poſture, and the Wound be conſtantly dreſſed with the White of an Egg, or Mucilage of Quince-ſeeds, till it is healed; by which Means the Patient often recovers his Sight.

V. If the Diſorder is become inveterate, and inflexible to all Remedies, you muſt paſs a Needle with a double Thread through the Middle of the Tumor, as in *Fig.* 8. *Tab.* XVIII. then the two Ends of the Thread are to be tied together in a Knot, firſt on one Side, and then on the other: by which Means the Tumor will gradually wither, and, at length, fall off together with the Threads. Cure of an inveterate Staphyloma.

VI. But as this Ligature frequently occaſions violent Pain, Inflammation, and ſometimes a Suppuration of the Eye; it would ſeem to be a more ſafe and expeditious Method to extirpate the Tumor by the Sciſſars or Scalpel. In this Manner I myſelf once cut off a Protuberance of this Kind at the Root, from the Eye, of the Length of one's Finger, by a Pair of Sciſſars. Another Method.

VII. M. ST. YVES's Method of removing theſe Protuberances, (ſee his Treatiſe, p. 222) when they have not wholly covered and obſcured the *Cornea*, is to paſs a crooked Needle and Thread of Silk through the Middle of the *Staphyloma*; after removing the Needle, he twiſts together the Thread, and extends them with his left Hand; while, with a Scalpel or Lancet, he frees the Tumor under the Ligature, till he can, at length, totally extirpate it by the Sciſſars. Laſtly, he applies a Compreſs over the diſordered Eye, dipt in *Sp. Vini*, diluted with Water, as was obſerved in treating of the Cataract. And thus not only the *Staphyloma* is removed, but the *Cornea* itſelf becomes perfectly healed, or elſe leaves but a very ſmall Aperture in the Middle of the Wound; from whence indeed the aqueous Humour is continually diſcharged as faſt as it is ſecerned in the Eye, but without any Trouble or Uneaſineſs to the Patient; becauſe it flows gently with the Tears through the lacrymal Paſſages into the Noſe. The Method of M. ST. YVES.

VIII. When the whole *Cornea* is infeſted with a *Staphyloma*, as in *Fig.* 4, 5, 6, 7. the moſt expeditious Method of Cure is that of ST. YVES, by cutting out circularly not only the *Cornea*, but alſo the *Iris* or *Uvea*, all round within a Line of the Ring, by which it touches the *Albuginea*; after which, all the Humours of the Eye falling out, the remaining Coats contract themſelves into a ſmaller Compaſs, and the Wound itſelf will gradually heal up. You muſt then provide the Patient with an artificial Eye, adapted in Size, Shape, and Aſpect, to ſupply the Place of that which is wanting. In this Manner the artificial Eye may frequently be moved from one Side to the other by the remaining Muſcles of that Organ, ſo that many cannot diſcern it to be an artificial, but will take it for a true or natural Eye; and in this laſt Method I myſelf have cured the *Staphyloma*. A ſecond Method of ST. YVES.

CHAP.

CHAP. LX.

Of the HYPOPYON.

Deſcription. I. WE frequently meet with a Collection of purulent Matter immediately under the *Cornea*, in the Place of the aqueous Humour; which Diſorder is generally denominated *Hypopyon* or *Pyoſis* [a]. The *Hypopyon* ariſes from an Extravaſation of Blood or Matter in this Part, which may happen by a violent Inflammation, the Small-Pox, couching a Cataract, or from other external Injuries of the Eyes from Violence, as Contuſion, from a Blow or Fall, a Burn, *&c.* It is at the Beginning very often attended with excruciating Pains both of the Head and Eyes; and, according to the Degree of Injury, is ſoon after followed either with Blindneſs and a Deſtruction of the Eye, or Death itſelf.

Cure. II. There are chiefly three Methods of treating this Diſorder. The firſt and mildeſt is by diſperſing the Matter with diſcutient Remedies; ſuch as the Application of Compreſſes dipt in a Decoction of Sage, Eye-bright, Hyſſop, and Fennel-ſeeds in Wine, or of little Bags ſtuffed with the ſame Ingredients and boiled in Wine, which are to be frequently renewed; by which Means, when the Blood or Matter is in no great Quantity, the Eye recovers its former Integrity and Action, as I have frequently experienced. Therefore you ſhould continue the Patient in this Method ſo long as you find any Benefit from it, even till the corrupt Matter or Blood is all diſſipated or diſperſed. But, if the Pain and other Symptoms are rendered more intenſe by theſe Applications, you muſt proceed immediately to the Operation. Otherwiſe, there will be great Danger of the contained Matter's eroding the *Cornea*, and deſtroying the internal Parts of the Eye, which will induce Blindneſs after the moſt intenſe Pains.

Cure by agitating the Head. III. But before we treat of the Operation, it may be proper to deſcribe the Method of Cure, which, we read, was formerly uſed with Succeſs by JUSTUS, an eminent Oculiſt in the Time of GALEN, who himſelf was an Eye-witneſs of his Practice, as he writes in the End of his XIVth Book *De Methodo medendi*. In the firſt place, he ſeated the Patient on a Sort of Chair over-againſt himſelf; then taking Hold of his Head with both Hands, he ſhook it about very aſſiduouſly, till all the purulent Matter diſappeared; in which Operation it is very remarkable, that GALEN himſelf teſtifies, the Spectators could perceive the corrupt Matter gradually ſubſiding to the Bottom of the Eye. Moſt People will be apt to reject this Method, as uſeleſs and ridiculous; but my Opinion is, that it may be often very effectual in removing the *Hypopyon*. In this I am confirmed, not only by the Authority of GALEN, but alſo from my own Experience in

[a] Indeed M. ST. YVES names this Diſorder of the Eyes *Onyx*; the *Hypopyon*, according to him, being a Suppuration in the *Tunica Cornea* itſelf: ſo that an *Onyx*, or *Unguis*, may ariſe from an *Hypopyon*, when the Matter of the laſt erodes into the *Cornea*, by deſtroying its internal *Lamella*. See his Treatiſe, *De Morb. Oculor.* Part II. Cap. 9. pag. 221, *& ſeq.* Hence we may ſee how much even ſome of our modern Surgeons and Oculiſts are at Variance in their aſcertaining the Diſorders of the Eyes and their Names.

in a Patient who, being juſt entered under my Care for an Hypopyon, was obliged to take a Journey in a Chariot; by the repeated ſhaking and jolting of which, upon his Return the next Day, I found all the purulent Matter diſperſed; and, without doubt, it was ſubſided or thrown down behind the Uvea. It may therefore not be improper to try this Practice before the chirurgical Operation by the Hand and Inſtruments. But before you ſhake the Head, it will be proper to diſpoſe it, or the Patient's whole Body, in a ſupine Poſture, and to preſs the Eye, firſt, with the Fingers, in order to looſen and remove the Matter. But, when the Diſorder is great and obſtinate, the purulent Matter being too copious, or too firmly fixed to be diſperſed in this Manner, Recourſe muſt then be had to the Operation long ago deſcribed and recommended by GALEN, ÆTIUS, and others of the Antients; which has met with ſo much Neglect among our Modern Surgeons and Oculiſts, that it would ſcarce have been known or heard of at preſent, if it had not been reſtored in the laſt Century by RIVERIUS, MEEKREN, NUCKE, and BIDLOW.

IV. Preparatory to the Operation, your Patient muſt be placed and ſeated againſt the Light, with his Head and Hands firmly ſecured each by an Aſſiſtant, as in couching a Cataract. Then the Surgeon himſelf depreſſes the lower Eyelid, while an Aſſiſtant elevates the upper. The Operator now takes a Lancet, and therewith cautiouſly incides through the *Cornea*, below the Pupil, and about the Space of a Line from the *Albuginea*, making his Apertion big enough to diſcharge the Matter with the aqueous Humour; but with Caution at the ſame time to avoid wounding the *Uvea* behind the Matter. If the Matter does not diſcharge freely of itſelf, you muſt aſſiſt it by a gentle Preſſure and Agitation with your Fingers; and in about three or four Hours after the Operation, you muſt dreſs the Eye with a Compreſs dipt in a Collyrium *ex Aq. Plantaginis vel Roſar. & Albo Ovor.* or a Mucilage *ex Sem. Cydonior.* prepared, either of them, with or without Camphor. By this means you will find the Wound in the *Cornea* quickly healed, and the aqueous Humour ſoon after reſtored, with the Patient's Sight, if none of the internal Parts are injured. And though there may remain a ſmall *Cicatrix* in the *Cornea*, yet that being made lower than the Pupil, will cauſe very little, if any, Impediment to the Sight. In the mean time, to perform this Operation with the Lancet ſafely, you ought to involve that Inſtrument in Lint, or a Piece of Plaſter, ſo as to leave not above a Straw's Breadth of its Point uncovered, that it may not run too far into the Eye. MEEKREN has on this Account invented an Inſtrument purpoſely for the Operation, publiſhed in the tenth Chapter of his chirurgical Operations, and delineated in our *Tab.* XVIII. *Fig.* 10.

Method of operating.

V. Sometimes the purulent Matter is found too much inſpiſſated to be eaſily diſcharged through the Inciſion made by the Lancet in the *Cornea*; and in that Caſe it will be more convenient to uſe the Needle, *Tab.* XVIII. *Fig.* 12. which we have elſewhere propoſed for making Setons. For the recurve Point of this Needle is not only leſs apt to wound the *Uvea*, but by its triangular Figure it alſo makes a larger Aperture, which will more readily diſcharge the inſpiſſated Matter; but then we uſually involve this Needle almoſt up to its Point in a Slip of ſome Plaſter, as I before adviſed you to do the Lancet. PLATNERUS has given

Another Method of Cure.

us the Figure of a particular Inſtrument for this Purpoſe[a], having a Sort of triangular *Apex*, the Invention of which he aſcribes to Mr. WOOLHOUSE. See our *Tab.* XVIII. *Fig.* 13. When the Matter included under the *Cornea* is too thick to flow out of itſelf, or by Preſſure, M. ST. YVES[b] propoſes to waſh it out by injecting with a ſmall Syringe, repeating the Operation every Day, till it be all removed; and then you may proceed to heal the Wound in the *Cornea*. If any Inflammation appears, the Patient ſhould be bled, bliſtered, ſcarified, and the affected Parts treated with a diſcutient Fomentation, and other proper Medicines.

CHAP. LXI.

Of Inciding the CORNEA, *to diſcharge Extravaſated* BLOOD.

When the Operation is neceſſary.

I. BLOOD extravaſated in but a ſmall Quantity from external Violence, or Injuries offered to the Eye, may be generally diſperſed and carried off by the diſcutient Remedies before propoſed at N° II. of the preceding Chapter. But when the Quantity is larger than can be thus removed, you ought immediately to open the *Cornea* by Inciſion, as we directed in the preceding Chapter, to prevent the ſtagnant Blood from ſuppurating and deſtroying the Eye.

An Inſtance of this Practice.

II. But leſt any body ſhould think I propoſe of my own Head a raſh and unheard-of Practice, I ſhall give the Reader an Inſtance of it (from the *Hiſt. Acad. Pariſ. An.* 1709. pag. 16. *Edit. Amſtel.*) in which it ſucceeded very well. Therefore, whenever any Perſon has, by ſome external Violence, had ſo much Blood extravaſated in his Eye, as to deſtroy his Sight, and be incapable of Diſperſion, it is the Advice of the Phyſician GANDOLPHUS, to have recourſe to this Practice. He therefore inſtantly made a tranſverſe Inciſion through the *Cornea*, and, by that means, happily diſcharged the extravaſated Blood, in ſuch a Manner, that the Patient was cured with hardly any Pain, and without any deforming *Cicatrix*, ſo that he recovered his former Sight without any Defect; and yet he was obliged to perforate the *Cornea* three times, by reaſon of the Quantity and ſtrong Adheſion of the Blood. To promote the healing of the Inciſion, he, for the Space of eight Days, applied Compreſſes dipt in a Mixture of *Aq. Plantag.* ℥iv. *& Aq. Vulneraria* ℥ij. In little more than a Week's Time, the Cure was ſo well performed, that one could perceive no Difference betwixt the Eye that had undergone the Operation, and the other which had not; excepting only that it's Pupil was a little larger than the other, which ſeems to have been rather the Effect of the Blow than of the Operation.

[a] *Diſſert. de Fiſtulâ lacrymali.* [b] *De Morb. Ocul.* p. 227.

CHAP.

CHAP. LXII.

Of the DISTENTION *and* PROLAPSUS OCULI, *alſo the* FUNGUS *and* CANCER.

I. SOMETIMES the Eye is ſo violently inflamed and ſwelled, that it cannot be contained in its Orbit or Socket by the Lids, but protrudes itſelf out of its natural Seat. This is a Diſorder attended not only with great Deformity, but alſo with intenſe Pains, and frequently Blindneſs or an obſtinate Cancer. How ghaſtly the Diſorder appears, may be perceived, I think, from the Figures we have given of it in *Tab.* XVIII. *Fig.* 14, 15[a]. PAREY mentions a Caſe he ſaw, in which the Eye was ſo vehemently diſtended by pernicious Humours, that it at laſt burſt out of its proper Coats; and the like may be alſo ſeen in MUYS, *Dec.* II. *Obſ.* I. This is termed by the *Greeks* a *Proptoſis*, and by the *Latins* a *Prolapſus Oculi*; but by ſome it is denominated an *Hydrophthalmia*, and by others *Eſcophthalmia*[b], when the Eye is very much diſtended with a watery Humour; but the more modern Authors have, from its Similitude, named the Diſorder, *Oculus Bovinus aut Elephantinus:* though I muſt confeſs that many of theſe Names are rather intended to ſignify different Diſeaſes than one and the ſame; whence Error and Confuſion. The Cauſes of this Diſorder are various, being ſometimes from a violent Inflammation, or a Redundancy of Humours in the Eye, from an Obſtruction of the reductory Veſſels; ſometimes from a *Scirrhus*, Cancer, or ſome external Violence. The Inſtances given us by HILDANUS, *Cent.* I. *Obſ.* I. MUYS, *Dec.* XII. *Obſ.* 1. and by me, in *Tab.* XVIII. *Fig.* 15. ſeem to have been from a Cancer; and more Inſtances of the ſame Kind may be ſeen in STALPART, VANDER WIEL, *Part* II. *Obſ.* 9. and in the other Writers of Obſervations. Laſtly, there are ſome Surgeons and Phyſicians who denominate this Diſorder *Ficus* or *Fungus*, which are in reality different Diſeaſes.

Nature of the Diſorder.

II. When the Diſorder is recent, and the Figure of the Eye is not yet deformed, thoſe Humours, producing the *Hydrophthalmia*, may be generally diſperſed by Bleeding, Purging, and Veſicatories, with internal Attenuants and Diluents, and external diſcutient Fomentations. But, if the Caſe is too obſtinate to yield to Remedies, you muſt have Recourſe to the chirurgical Operation of *Paracenteſis*, as in other dropſical Caſes. This *Paracenteſis* muſt be made either with a Lancet, or a ſmall *Trocar*, to diſcharge the offending Humours, repeating the Diſcharge every Day, or every other Day, or as often as ſhall be found neceſſary. At every Dreſſing, a concave Plate of Lead, with a Compreſs dipped in ſome diſcutient Liquor, (ſee *Chap.* LX. *N.* II.) ſhould be firmly ſecured upon the Eye, to recover its natural Figure. By carefully obſerving

Cure by Diſcutients or Puncture.

[a] *Vid.* BARTISCHIUS in *Chirurg. Ocular.* p. 218. and HILDAN. *Obſ.* 1.
[b] *Vid.* NUCK *De Duct. Aquoſ.* p. 119. & 120. STALPART. VANDER WIEL, *&c.*

obſerving this Method, NUCKE[a] cured a Patient of an *Hydrophthalmia*, though he made his *Paracenteſis* in the *Cornea* itſelf: but as that may leave an ugly *Cicatrix* in the *Cornea*, I rather make my Perforation with a Lancet in the *Sclerotica* than in the *Cornea*. After diſcharging the Humours, I dreſs the Eye with Lint dipt in *Aq. Roſar. & Album. Ovor. permiſt.* defend it with a leaden Plate, and then apply my Compreſs dipt in *Sp. Vini*; and laſtly, my Bandage, not neglecting Internals at the ſame Time, till the Eye is cured, and recovers its State.

Cure by the Scalpel.

III. When the natural Figure of the Eye and its Office of Viſion are deſtroyed, and the Pains become more and more intenſe, there then remains but one, and a lamentable Method of relieving the Patient, by making a tranſverſe Inciſion through the Coats of the Eye, and diſcharging the contained Humours; which done, and the Eye deterged as in other Ulcers, you muſt cover the Eye-lids with Compreſs and Bandage. But if, after the Humours are diſcharged, the Eye remains larger than can be eaſily covered with the Eye-lids, it will be neceſſary to cut off ſo much as is redundant with the Scalpel or Sciſſars; by which Means the Deformity may be afterwards the better concealed by an artificial Eye. Sometimes the Surgeon may cut out the *Cornea* by a circular Inciſion, in this Diſorder, as we propoſed in the *Staphyloma*, Chap. LVIII. preceding.

Another Method of Cure.

IV. BARTISCHIUS, HILDANUS, and MUYS, have contrived a crooked Scalpel, like a Spoon, for extirpating the Eye when it is thus diſordered: but, upon mature Conſideration, I believe the Surgeon will not ſtand in need of any ſuch Inſtrument. For, to ſay nothing of the Difficulty you will meet with in ſharpening and uſing ſuch an Inſtrument, it will be found, in moſt Caſes, ſufficient to extirpate only the redundant or tumified Part of the Eye, which prevents the Eye-lids from cloſing: to which you may add, the Danger there will be of wounding and uncovering the thin Bones which compoſe the Orbit, by this crooked Scalpel. But if ever the Surgeon ſhall find it neceſſary to extirpate the whole Eye for a *Scirrhus*, or cancerous Diſorder of it, he may perform the ſame with equal Advantage by the ſtraight Scalpel, *Tab.* XII. *Fig.* 14. which is the ſame I uſed in extirpating thoſe ghaſtly Tumors of this Kind, repreſented in *Tab.* XVIII. *Fig.* 14, and 15. Though there are ſome Surgeons who think it the mildeſt Practice to free the Eye ſo far from its Orbit by a Scalpel, till you can make a Ligature about the protuberant Part in order to remove it, by that Means, like other Excreſcences. But the more prudent in the Profeſſion generally prefer any Method to this, becauſe of the intenſe Pain, Inflammation, and Convulſions, which, by this Means, torture and often kill the Patient. Therefore whenever you meet with the Eye infeſted, even to its Root, with a *Scirrhus*, or Cancer, there is no ſafer Method of relieving the Patient from his painful Diſorder, than by extirpating it clean out from the Orbit, in the Manner performed by HILDANUS and MUYS; deterging and healing the Wound afterwards in the uſual Method.

5

V. It

[a] Lib. *De Duct. Oculor. aquoſ.* pag. 120. and VALENTINE in *Miſc. Nat. Cur.* Ann. VI. Obſ. 70.

V. It ſometimes happens in this Diſorder, that after having performed the Operation, a new fleſhy Excreſcence ſprouts up over the Eye, and forms a freſh Tumor; to prevent which, you muſt dreſs with Lint dipt in *Aq. Phagedænica*, and make a pretty tight Deligation over the leaden Plate with which you are to cover the Eye. It may be here alſo obſerved, that Cancers of the Eye, like the ſame Diſorder in other Parts, will very often return, after they have been ſeemingly cured by the Operation and Treatment here propoſed, and may be again removed by the ſame Practice; as appears from the Obſervation of MUYS, before cited. Laſtly, when the Diſorder ariſes from a *Caries*, or *Spina ventoſa* of the Bones themſelves compoſing the Orbit, if it will not give way to Mercury (as it often does) the Phyſician muſt then be content to palliate the Diſorder, relieve the Pains, and prevent its bad Conſequences; for a total Removal thereof is frequently altogether impracticable.

CHAP. LXIII.

Of ARTIFICIAL EYES.

I. THE Loſs of an Eye is frequently occaſioned by a Wound, an Abceſs in the Small-Pox, or an Operation in Surgery; and then the unhappy Patient is deſirous of concealing his Misfortune by an artificial Eye, which is contrived to hide the Deformity ariſing from this Accident. The modern artificial Eyes are made of concave Plates of Silver, Gold, or Glaſs ſtained or enamelled, ſo as to reſemble the natural Eye. See *Tab.* VII. *Fig.* 1. The nearer it approaches the ſound Eye in Size and Appearance, the more firmly it will ſtay under the Eye-lids, and the more eaſily deceive the Spectator. But it will be frequently neceſſary for the Patient to wipe his artificial Eye clean, leſt if any Gum or Sordes ſhould gather upon it, the Fallacy might be thereby diſcovered; to prevent which, it may be alſo proper for him to be provided with ſeveral of theſe artificial Eyes, that if one ſhould happen to be loſt, broke, or diſfigured, its Place may be immediately ſupplied with another. Upon going to Bed, it is proper to diſmount the artificial Eye, and to replace it again under the Eye-lids, after he wakes in the Morning. But then, that the artificial Eye may be taken out and put in with Neatneſs and Conveniency, the Surgeon muſt take care to remove ſo much of the diſordered Eye, as will make Room for receiving the artificial. Their Compoſition.

II. It is here to be obſerved, that the more cloſely the artificial Eye is compreſſed by the Eye-lids, and by the diſeaſed Eye, the more perfectly it will perform the Motions of the natural Eye, which it will receive from the remaining Muſcles which agitate the diſeaſed Globe. It is therefore not without Reaſon that we before adviſed the Surgeon to remove no more of the Eye than what was preternaturally projected beyond its anterior Part; except when a *Scirrhus* Motion of the artificial Eye.

or Cancer ſhould require an Extirpation of the whole; and then indeed it cannot be expected that the artificial Eye ſhould have any other Motion than what it receives from the Lids.

It is ſometimes better to be without them.

III. I have ſeveral times obſerved ſome of theſe artificial Eyes produce Pain, Inflammation, Tears, and other Inconveniencies, by irritating the Parts which are not of a proper Conformation, or when the artificial is not right ſhaped; ſo that they will often inflame, weaken, and deſtroy the Sight of the ſound Eye. In ſuch Caſes, it will be beſt for the Patient either to provide himſelf with an artificial Eye which is better adapted, or elſe totally to relinquiſh the Uſe of them, rather than loſe the Uſe of both Eyes.

CHAP. LXIV.

Of the STRABISMUS, *or* SQUINTING.

Cauſes.

I. WE frequently meet with Perſons whoſe Eyes, when they look upon any thing, are diſtorted, or turned towards the outer or inner Corners of their Eye-lids, inſtead of being directed towards the Object; which is the Diſorder commonly termed *Strabiſmus*, or *Squinting*. Sometimes only one Eye, but more frequently both are thus affected. The Diſorder is frequently cauſed in Infants, from letting them conſtantly ſuck at one and the ſame Breaſt, or placing them in the Cradle, ſo that they always look the ſame Way towards the Light or Window. By this repeated Action, the Muſcles on that Side become too ſtrong and powerful to be balanced by the reſt which counter-act them on the other Side of the Eye; whence it is contorted, or looks obliquely. But this Diſorder is more frequently cauſed in Infants from convulſive and epileptic Motions; to which the Muſcles of their Eyes, as well as of their other Limbs, are extremely ſubject. Laſtly, it may proceed as well in Adults as Infants, from a Spaſm and Rigor, or from a Palſy in one or two of the Muſcles of the Eye, as alſo from a Defect or Inſenſibility in ſome Part of the *Retina*. For when that Part of the *Retina* which is oppoſite to the Pupil, and receives the Impreſſion of the Object, is from any Cauſe rendered inſenſible, the Patient is then obliged to turn his Eye obliquely, till the Pupil directs the Rays from the Object upon ſome other ſound Part of the *Retina*, in order to ſee the ſame.

When and how it may be cured.

II. Squinting is a Diſorder which is hardly ever cured without Difficulty, more eſpecially when in Adults, and cauſed by ſome Defect in the Muſcles or *Retina* of the Eye (eſpecially if the Diſorder ariſes from a bad Habit only.) But in young Infants you will probably ſucceed, according to the Advice of M. ST. YVES, by frequently placing them before a Looking-glaſs, that their Eyes may be directed towards the Image of their own Face. Thoſe more advanced in Years may be aſſiſted by reading very ſmall Writing, or inſpecting very minute Objects, provided you direct them to turn their Eyes even, and to bathe them at Times with *Aq. Hungar.* or anoint them with the *Balſamum Floventi*. There are others who propoſe to cure this Diſorder with a Sort of



Plate 18.




Fig. 14.
A
C
D
B
E
Fig. 1.
Fig. 2.
Fig. 3.
Fig. 4.
Fig. 5.
Fig. 6.
Fig. 7.
Fig. 13.
B
A
Fig. 11.


Fig. 15.
Fig. 13.
C
A
A
B
Fig. 10.
Fig. 12.
Fig. 11.
A
Fig. 16.
B
B
A
A
Fig. 9.
C


A
b
Fig. 12.
Fig. 16.
Fig. 1.
B
A
Fig. 9.
b
Fig. 2.
C

Maſk or Eye-ſwath, as in *Tab.* XVIII. *Fig.* 16. taken from SOLINGEN, and deſcribed more particularly in the Explanation of the following Table. This Method is alſo recommended by BARTISCHIUS in his *Ophthalmodulcia*, pag. 15, 16 and 17. But, leſt Infants ſhould look ſtraight through the Aperture with only one Eye, and ſquint in the mean time with the other, it will be beſt to bind up one Eye till the other is rectified, and then to correct the other in the ſame Manner; which is ſeldom practicable, through the Unrulineſs of Infants, and other Impediments.

An EXPLANATION *of the* EIGHTEENTH PLATE.

Fig. 1. Denotes an *Unguis a* on the Eye, with the Method of paſſing a Needle and Thread under it *b b*, for its Removal.

Fig. 2. Repreſents another *Unguis*, or *Pterygium a a*, with a Thread tied round it *b b*, and at their Extremities tied in the Knot *c*, to form a Loop for extending and elevating the ſame: but that the Thread may not ſlide upon the Film, it is firſt tied with the double Knot *a*.

Fig. 3. Repreſents a Hook uſed in ſeparating Films, and other Tubercles, from the Eye.

Fig. 4. Denotes a front View of a *Staphyloma*, or Protuberance of the *Cornea*, which I cured.

Fig. 5. Gives a lateral View of the ſame *Staphyloma*.

Fig. 6. Repreſents a front View of another larger and more depending *Staphyloma*, which I cured.

Fig. 7. Gives a lateral View of the ſame.

Fig. 8. Is a leſſer *Staphyloma*, marked *a a*, with a double Thread paſſed under it, from SOLINGEN.

Fig. 9. A *Scalprum*, to ſcrape or exfoliate carious Bones in the *Fiſtula lacrymalis*, from PLATNERUS's Diſſertation *De Fiſt. lacr.*

Fig. 10. Repreſents MEEKREN's Inſtrument for perforating the *Cornea* in an *Hypopyon*. A A the Handle; B the Scalpel, or rather the Point of a double-edged Scalpel, having a Button or Protuberance at its Baſis, to prevent the Point from entering too deep into the Eye: C the Screw by which the *Capſula* or Caſe, *Fig.* 11. is faſtened on.

Fig. 12. Denotes a large Needle which may ſerve to make Setons; but is here deſigned to perforate the *Cornea* if you ſecure it from entering too deep, by involving it in a Slip of Plaſter up to A.

Fig. 13. Repreſents an Inſtrument deſigned to perforate the *Cornea* in an *Hypopyon*. A denotes the Handle, B the triangular Point a little crooked, almoſt like the preceding Needle; which ſhould, like that, be involved in a Slip of Plaſter up to the Point, to prevent its entering too far beyond the *Cornea*.

Fig. 14. The Letters A B denote a ſcirrhous Eye, enlarged to the Size of an Hen's Egg, upon which is a blackiſh Tubercle, like a Grape, marked C; and D denotes the vitiated Pupil and *Cornea*. E the lower Eye-lid depreſſed by the Tumor.

Fig. 15.

Fig. 15. Denotes a larger Fungus of the left Eye, weighing half a Pound, which, with the preceding, I extirpated and cured in 1721, the particular Nature and Treatment of which I ſhall deſcribe in my *Chirurgical Obſervations*, which I intend ſhortly to publiſh.

Fig. 16. Repreſents a Bandage to cure the Diſorder of Squinting in Children. L. AA. two concave Plates of Silver, Ivory, or Ebony, perforated in the Middle: L. CC the Bandage by which they are fixed to the Eyes. Children are, by this Method, practiſed to direct their Eyes ſtraight forward through the Apertures, and thereby acquire a regular Habit.

The END *of the* FIRST VOLUME.

